Published Online: 2024 June 22

Research Article



A Prospective Study of Right Ventricular Function After Coronary Artery Bypass Grafting: Are Usual Echocardiographic Parameters Useful?

Firoozeh Abtahi (1)¹, Zohreh Bazyar¹, Khalil Zarrabi², Mani Hassanzadeh (1)¹, Zahra Mehdipour Namdar (1)³, Amir Aslani (1)¹, Seyed Ali Hosseini (1)³, Armin Attar (1)^{1,*}

¹ Department of Cardiovascular Medicine, Shiraz University of Medical Sciences, Shiraz, Iran

² Department of Cardiac Surgery, Shiraz University of Medical Sciences, Shiraz, Iran

³ Student Research Committee, Shiraz University of Medical Sciences, Shiraz, Iran

Corresponding Author: Department of Cardiovascular Medicine, Shiraz University of Medical Sciences, Shiraz, Iran. Email: attar_armin@yahoo.com

Received: 6 April, 2024; Revised: 25 May, 2024; Accepted: 5 June, 2024

Abstract

Background: The best method for accurately assessing the right ventricle (RV) after coronary artery bypass grafting (CABG) remains controversial. In this study, we aimed to investigate several echocardiographic RV function indices to determine which are most effective in evaluating RV function post-CABG.

Objectives: Here, we aimed to assess RV functional parameters after CABG using echocardiography.

Methods: This prospective before-after study was conducted on 60 patients scheduled for CABG. The patients were assessed before and six weeks after CABG using standard echocardiography and Speckle-Tracking Echocardiography (STE). The significance of changes in study parameters was evaluated using the paired *t*-test or the non-parametric Wilcoxon test. Two-sided P-values \leq 0.05 were considered statistically significant. Statistical analysis was performed using SPSS software, version 16.0.

Results: Six weeks after CABG, we found significant changes in two parameters: Tricuspid annular plane systolic excursion (TAPSE) and S' tissue Doppler imaging (20.82 ± 2.53 vs. 14.60 ± 1.84 , P = 0.01; 11.79 ± 1.99 vs. 8.80 ± 0.88 , P = 0.01, respectively). However, no significant changes occurred in the right ventricular global 2D longitudinal strain or free wall longitudinal strain (-18.83 ± 2.88 vs. -18.79 ± 2.66, P = 0.88; -21.29 ± 3.36 vs. -20.86 ± 3.17, P = 0.12, respectively).

Conclusions: Tricuspid annular plane systolic excursion and S' tissue Doppler imaging indices change following CABG and cannot be considered standard echocardiographic parameters for assessing RV function after the procedure. However, the two major parameters for right ventricular functional assessment, global 2D longitudinal strain and free wall longitudinal strain, may be less affected by CABG and thus could be more reliable as standard parameters.

Keywords: Speckle Tracking Echocardiography, CABG, Right Ventricle, Doppler Imaging

1. Background

Coronary artery disease (CAD) is a leading cause of death. Recent studies have demonstrated an upward trend in its prevalence due to sedentary lifestyles, poor diets, and a consequent lack of control over related risk factors (1). Applying different therapeutic approaches, including lifestyle modification, various medications, and revascularization in severe cases, can improve disease-related outcomes, long-term patient survival, and various components of quality of life (2, 3). Regarding revascularization, coronary artery bypass grafting (CABG) surgery and percutaneous coronary intervention (PCI) have significantly reduced mortality and morbidity in patients by improving blood flow to the myocardial tissue, balancing tissue supply and demand, and ultimately enhancing ventricular systolic and diastolic function (4, 5).

Copyright © 2024, Abtahi et al. This open-access article is available under the Creative Commons Attribution 4.0 (CC BY 4.0) International License (https://creativecommons.org/licenses/by/4.0/), which allows for unrestricted use, distribution, and reproduction in any medium, provided that the original work is properly cited.

How to Cite: Abtahi F, Bazyar Z, Zarrabi K, Hassanzadeh M, Mehdipour Namdar Z, et al. A Prospective Study of Right Ventricular Function After Coronary Artery Bypass Grafting: Are Usual Echocardiographic Parameters Useful?. Int Cardiovasc Res J. 2024; 18 (1): e146326. https://doi.org/10.5812/icrj-146326.

Traditionally, dynamic indicators such as left ventricular ejection fraction (LVEF) or the diameters of the heart muscles, cavities, and vessels have been used to assess ventricular function. However, the emphasis has recently shifted toward more accurate parameters, such as longitudinal myocardial strain, in evaluating myocardial changes in function following revascularization. Speckle tracking echocardiography (STE), a new diagnostic modality, provides an opportunity to assess deformation in the myocardium's circumferential, longitudinal, and radial planes, correlating well with changes in mechanical function (6). Speckle motion analysis can assess the velocity and strain of myocardial tissue independently of cardiac translation (7). The quantitative evaluation of ventricular functional state by STE offers a significant advantage.

Several studies on the echocardiographic assessment of RV function after cardiac surgery have evaluated these parameters. Tricuspid annular plane systolic excursion (TAPSE) and s' appear to be reduced in subjects undergoing surgery for both congenital and acquired heart diseases (8). However, these changes may not necessarily indicate RV dysfunction (9). Here, we aimed to investigate various echocardiographic RV function indices to determine which are most useful for evaluating RV function after CABG.

2. Objectives

The present study assessed right ventricular functional changes using STE following CABG.

3. Methods

3.1. Study Methodology

This prospective before-and-after study was conducted on patients with a definite diagnosis of CAD (confirmed by coronary angiography) who were candidates for on-pump CABG at Faghihi and Namazi hospitals in Shiraz, Iran, in 2019 and 2020.

3.2. Target Population

The study population included all CABG candidates aged 30 to 70 years without a previous history of angioplasty or other coronary interventions and without a history of acute coronary syndrome or myocardial pacing. All patients had severe coronary artery stenosis (> 70% stenosis) on angiography. We excluded subjects with any evidence of congenital heart diseases, chronic obstructive lung disease, valvular heart diseases (in initial echocardiography assessment), or reduced LVEF (less than 35%). Additionally, patients with a poor echocardiographic window were excluded. A total of 60 patients completed the study.

3.3. Sampling and Methods

All cases underwent a detailed medical history assessment. Before scheduling CABG, all patients were evaluated using speckle-tracking echocardiography (STE) (General Electric Vivid E9, Norway) and subsequently underwent CABG using the same technique. All patients received standard post-operative care and medications in line with current European guidelines (reference). Six weeks after CABG, echocardiography was performed again.

All echocardiographic evaluations were conducted using General Electric's (GE) E9 machine with an M5ScD probe and 1.5 - 4.6 MHz bandwidth. Each patient underwent two rounds of echocardiography performed by two expert operators who were blinded to the treatment to reduce bias and inter-observer variability. The mean of these examinations is reported. Echocardiography was performed in the left lateral decubitus position, and the following parameters were assessed by STE before and after surgery: Fractional area change (FAC), left ventricular end-diastolic diameter (LVEDD), right ventricular end-diastolic diameter (RVEDD), tricuspid annular plane systolic excursion (TAPSE), S' tissue Doppler imaging (S'TDI), global 2D longitudinal right ventricular strain, and right ventricle free wall 2D longitudinal strain. The primary endpoint was to assess the changes in these parameters following CABG.

3.4. Statistical Analysis

Results are presented as mean \pm standard deviation (SD) for quantitative variables and as frequency (percentage) for categorical variables. The significance of changes in study parameters was assessed using the paired *t*-test or the non-parametric Wilcoxon test. Two-sided P-values \leq 0.05 were considered statistically significant. Statistical analyses were performed using SPSS version 16.0 for Windows (IBM, Armonk, New York).

4. Results

This study involved 60 CABG candidates with a mean age of 66 \pm 9 years. Most patients (71.7%) had a prior history of myocardial infarction. Nearly two-thirds (63.3%) of patients had three-vessel coronary artery disease. The most prominent risk factor was hypertension (88.3%). The mean number of grafts was 3.0 \pm 0.8, with a bypass time of 123 \pm 39 minutes. Table 1

Table 1. Baseline and Operative Characteristics of the Study Population ^a	
Variables	Values
Age, y	66±9
Male sex	48 (80)
Prior myocardial infarction	43 (71.7)
Hypertension	53 (88.3)
Diabetes mellitus	25 (41.6)
Atrial fibrillation	7 (11.7)
Current smoker	14 (23.3)
Body mass index, kg/m ²	30.2±5.8
COPD	20 (33.3)
Cerebrovascular accident	17 (28.3)
Peripheral artery disease	27 (46.7)
Prior cardiac surgery	3 (5.0)
Laboratory values	
GFR, mL/min/1.73 m ²	85±36
Hemoglobin, g/dL	13.4 ± 1.8
Medications	
Beta blocker	55 (91.7)
ACEI/ARB	43 (71.7)
Spironolactone	4 (6.7)
Digoxin	5 (8.3)
Diuretics	26 (43.3)
Operative parameters	
Number of grafts	3.0 ± 0.8
Bypass time, min	123 ± 39
Ischemic time, min	80 ± 27
Number of diseased coronary vessels	
1 vessel	3 (5.0)
2 vessels	19 (31.7)
3 vessels	38 (63.3)
Left main disease	20 (33.3)

Abbreviations: COPD, chronic obstructive pulmonary disease; GFR, glomerular filtration rate. $^{\rm a}$ Values are expressed as No. (%) or Mean ± SD.

presents the baseline and operative characteristics of the patients.

Echocardiographic parameters before and after CABG are summarized in Table 2. Among all myocardial parameters assessed by STE before and six weeks after CABG, significant changes were found in the two parameters of TAPSE and S' tissue Doppler imaging. However, other parameters such as left ventricular endsystolic diameter, right atrial area, right ventricular enddiastolic area, right ventricular end-systolic area, right ventricular fractional area change, right ventricular end-diastolic diameter, right ventricular global 2D longitudinal strain (Figures 1 and 2), and right ventricular free wall longitudinal strain did not change significantly. Nonetheless, significant improvements in left ventricular end-diastolic diameter (from 47.11 \pm 4.60 mm to 47.80 \pm 4.26 mm, P = 0.01) and ejection fraction (from 55.00 \pm 3.83% to 56.14 \pm 4.03%, P = 0.03) were recorded. None of the patients had severe valve diseases requiring intervention.

5. Discussion

The present study assessed right ventricular functional changes using STE following CABG. The 2010 ASE guidelines recommend standard echocardiography to evaluate the right heart in adults by assessing parameters such as TAPSE, S' tissue Doppler imaging index, right ventricular FAC, and the myocardial performance index. Measuring multiple indicators provides a more reliable estimate of right ventricular

Table 2. Echocardiographic Parameters Before and six Weeks After Coronary Artery Bypass Grafting (CABG)			
Parameter	Before	After	P-Value
Left ventricular end-diastolic diameter	47.11±4.60	47.80 ± 4.26	0.01
Left ventricular end-systolic diameter	30.42 ± 4.57	30.91 ± 3.97	0.13
Left ventricular ejection fraction	56.14 ± 4.03	55.00 ± 3.83	0.03
Right atrial area	12.31 ± 2.31	12.56 ± 2.24	0.16
Right ventricular end-diastolic area	13.88 ± 3.67	13.96 ± 3.92	0.77
Right ventricular end-systolic area	6.53±2.30	6.51 ± 2.46	0.84
Right ventricular fractional area change	53.08 ± 6.80	53.47 ± 6.32	0.42
Right ventricular end-diastolic diameter	26.34 ± 3.10	26.62 ± 2.93	0.77
Right ventricular global 2D longitudinal strain	-18.83 ± 2.88	$\textbf{-18.79} \pm 2.66$	0.88
Right ventricular free wall longitudinal strain	-21.29 ± 3.36	-20.86 ± 3.17	0.12
Tricuspid annular plane systolic excursion	20.82 ± 2.53	14.60 ± 1.84	0.01
S' tissue Doppler imaging	11.79 ± 1.99	8.80 ± 0.88	0.01



Figure 1. Right ventricular global 2D longitudinal strain before coronary artery bypass grafting. FR, frame rate; HR, heart rate; GS, global strain.

function (10). In this study, three of these indicators were examined, showing significant changes in TAPSE and S' tissue Doppler imaging indices. However, the change in FAC did not reach statistical significance. This index represents the percentage of surface change within the right ventricle between diastole and systole, providing an estimate of the overall right ventricular systolic function. The FAC marker helps independently indicate heart failure and can predict sudden death from myocardial infarction, stroke, and mortality following pulmonary embolism (11). However, our study

indicates that it remains unchanged following CABG, at least in the first weeks after treatment.

Myocardial deformity assessed according to strain is more valuable than the study of wall motion (velocity and displacement) in diagnosing regional cardiac abnormalities (12). Several recent studies have shown that two-dimensional signaling (2D-STI) imaging is a practical method for assessing right ventricular function. Longitudinal strain is independent of the overall movement of the heart, thus allowing the analysis of regional deformation of the heart in different parts of the right ventricle (13-15). However, our



Figure 2. Right ventricular global 2D longitudinal strain after coronary artery bypass grafting. FR, frame rate; HR, heart rate; GS, global strain.

study showed no significant changes in the two related parameters, namely global 2D longitudinal right ventricular strain and right ventricular free longitudinal strain, after CABG.

In a study by Abdelmoneum et al. in 2019 (16), a significant decrease in peak right ventricle systolic velocity and TAPSE and an increase in right ventricle fractional area change were reported. In their study, and contrary to our observation, a significant decrease in right ventricle global longitudinal strain and right ventricle free wall longitudinal strain was also detected. However, similar to our study and Korshin et al. (17), although TAPSE and RV longitudinal displacement were significantly reduced after CABG, RV speckle tracking strain did not change significantly. Khani et al. (18) recorded a statistically significant decrease in RVGLS from (-19 to -11) one week after CABG. Factors such as operator experience in assessing myocardial dynamic parameters, patient evaluation time, and cardiovascular comorbidities may explain this discrepancy in the findings of various studies. Hence, it remains unclear whether surgery can affect the right ventricular myocardial infarction indices, highlighting the need for further studies.

As a secondary outcome, we found the change in LVEDD following treatment to be 0.684 mm, which was statistically significant although not clinically significant. While the left ventricular diameter at the end of systole increased after treatment, this did not reach statistical significance. Left ventricular diameters are among the main indices for assessing ventricular function and can be easily measured bv echocardiography during LVEF measurement. Previous studies have highlighted the importance of LV size in relation to cardiac mortality. A study of patients with LV disorders showed that increased LV diastolic diameter was associated with increased cardiovascular mortality (19). Similarly, another study showed that a reduced risk of death or heart failure was directly related to decreasing left ventricular diastolic diameter (20). Some studies in specific populations suggest that LV diameter can be useful in assessing the risk of unpredictable sudden cardiac death (21). Hence, while our study primarily focused on RV parameters, the value of indices related to LV function is undeniable.

The limitations of this study included its singlecenter nature and lack of long-term follow-up. Furthermore, we did not investigate pathophysiological mechanisms that could alter standard echocardiographic parameters after CABG. Additionally, we omitted patients with a limited echocardiographic window as strain imaging and speckle tracking depend on an optimal acoustic window, which is not always available. In contrast, TAPSE and S' are less dependent on the acoustic window.

5.1. Conclusions

This study assessed right ventricular functional changes using STE over six weeks following CABG. Our findings indicate that CABG induces changes in TAPSE and S' tissue Doppler imaging indices but does not affect right ventricular global 2D longitudinal strain or free wall longitudinal strain.

Footnotes

Authors' Contribution: Study concept and design: F.A, Z. B; acquisition of data: K. Z, M. H, Z. M.N, S.A.H.; analysis and interpretation of data: Z. M.N, A. Aslani, S.A.H.; drafting of the manuscript: F.A, Z.B, M. H, Z. M.N; critical revision of the manuscript for important intellectual content: K. Z, A. Aslani, A. Attar, S.A.H.; statistical analysis: F.A, A. Aslani, A. Attar; administrative, technical, and material support: A. Attar; study supervision: F.A, A. Aslani, A. Attar.

Conflict of Interests Statement: The authors have no financial interests related to the material in the manuscript.

Data Availability: The data supporting this study are available from Armin Attar upon reasonable request.

Ethical Approval: Shiraz University of Medical Sciences approved the study protocol with an ethical code of IR.SUMS.MED.REC.1398.061.

Funding/Support: This study was performed by a grant from the Vice Chancellery of Research of Shiraz University of Medical Sciences, and the authors wish to thank the institute. The funding body has no role in the design or reporting of this study. (Grant number: 18614)

Informed Consent: All participants provided written informed consent before enrolment.

References

- Lozano R, Naghavi M, Foreman K, Lim S, Shibuya K, Aboyans V, et al. Global and regional mortality from 235 causes of death for 20 age groups in 1990 and 2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet*. 2012;**380**(9859):2095-128. [PubMed ID: 23245604]. [PubMed Central ID: PMC10790329]. https://doi.org/10.1016/S0140-6736(12)61728-0.
- Kaambwa B, Gesesew HA, Horsfall M, Chew D. Quality of Life Changes in Acute Coronary Syndromes Patients: A Systematic Review and Meta-Analysis. Int J Environ Res Public Health. 2020;17(18). [PubMed ID: 32967168]. [PubMed Central ID: PMC7558854]. https://doi.org/10.3390/ijerph17186889.
- 3. Schmidt-RioValle J, Abu Ejheisheh M, Membrive-Jimenez MJ, Suleiman-Martos N, Albendin-Garcia L, Correa-Rodriguez M, et al.

Quality of Life After Coronary Artery Bypass Surgery: A Systematic Review and Meta-Analysis. *Int J Environ Res Public Health*. 2020;**17**(22). [PubMed ID: 33202650]. [PubMed Central ID: PMC7697861]. https://doi.org/10.3390/ijerph17228439.

- 4. Holm NR, Makikallio T, Lindsay MM, Spence MS, Erglis A, Menown IBA, et al. Percutaneous coronary angioplasty versus coronary artery bypass grafting in the treatment of unprotected left main stenosis: updated 5-year outcomes from the randomised, non-inferiority NOBLE trial. *Lancet.* 2020;**395**(10219):191-9. [PubMed ID: 31879028]. https://doi.org/10.1016/S0140-6736(19)32972-1.
- Bianco V, Kilic A, Mulukutla S, Gleason TG, Kliner D, Allen CC, et al. Percutaneous coronary intervention versus coronary artery bypass grafting in patients with reduced ejection fraction. *J Thorac Cardiovasc Surg.* 2021;**161**(3):1022-1031 e5. [PubMed ID: 33059935]. https://doi.org/10.1016/j.jtcvs.2020.06.159.
- Ebrahimi F, Gharedaghi MH, Zubair M, Kohanchi D, Aghajani K, Candido K. Speckle- Tracking Echocardiography for the Staging of Diastolic Dysfunction: The Correlation Between Strain-Based Indices and the Severity of Left Ventricular Diastolic Dysfunction. J Cardiothorac Vasc Anesth. 2021;35(1):216-21. [PubMed ID: 32753329]. https://doi.org/10.1053/j.jvca.2020.06.081.
- Blessberger H, Bergler-Klein J, Graf S, Syeda B, Wagner H, Kammler J, et al. Speckle Tracking-Derived Longitudinal Strain: Validation and Influence of Scanner Settings. *Ultrasound Med Biol.* 2021;47(1):154-62. [PubMed ID: 33077338]. https://doi.org/10.1016/j.ultrasmedbio.2020.09.005.
- Mandoli GE, Cameli M, Novo G, Agricola E, Righini FM, Santoro C, et al. Right ventricular function after cardiac surgery: the diagnostic and prognostic role of echocardiography. *Heart Fail Rev.* 2019;24(5):625-35. [PubMed ID: 30982175]. https://doi.org/10.1007/s10741-019-09785-2.
- Singh A, Huang X, Dai L, Wyler D, Alfirevic A, Blackstone EH, et al. Right ventricular function is reduced during cardiac surgery independent of procedural characteristics, reoperative status, or pericardiotomy. J Thorac Cardiovasc Surg. 2020;159(4):1430-1438 e4. [PubMed ID: 31133354]. https://doi.org/10.1016/j.jtcvs.2019.04.035.
- 10. Daskalov I, Ivanchev M, Miletieva M. Guidelines for the Echocardiography Assessment of the Right Heart in adults: a report from the American Society of Echocardiography endorsed by the European Association of Echocardiography, a registered branch of the European Society of Cardiology, and the Canadian Society of Echocardiography. S'rdechno-s' dovi Zabolyavaniya/Med Univ Sofia -Central Medical Library. 2012;**43**(2):48-52.
- Lang RM, Badano LP, Mor-Avi V, Afilalo J, Armstrong A, Ernande L, et al. Recommendations for cardiac chamber quantification by echocardiography in adults: an update from the American Society of Echocardiography and the European Association of Cardiovascular Imaging. Eur Heart J Cardiovasc Imaging. 2015;16(3):233-70. [PubMed ID: 25712077]. https://doi.org/10.1093/ehjci/jev014.
- Grignola JC, Gines F, Guzzo D. Comparison of the Tei index with invasive measurements of right ventricular function. *Int J Cardiol.* 2006;**113**(1):25-33. [PubMed ID: 16325940]. https://doi.org/10.1016/j.ijcard.2005.10.012.
- Li Y, Wang Y, Ye X, Kong L, Zhu W, Lu X. Clinical study of right ventricular longitudinal strain for assessing right ventricular dysfunction and hemodynamics in pulmonary hypertension. *Medicine (Baltimore)*. 2016;95(50). e5668. [PubMed ID: 27977616]. [PubMed Central ID: PMC5268062]. https://doi.org/10.1097/MD.000000000005668.
- Pirat B, McCulloch ML, Zoghbi WA. Evaluation of global and regional right ventricular systolic function in patients with pulmonary hypertension using a novel speckle tracking method. *Am J Cardiol.* 2006;**98**(5):699-704. [PubMed ID: 16923465]. https://doi.org/10.1016/j.amjcard.2006.03.056.

- Dragulescu A, Mertens LL. Developments in echocardiographic techniques for the evaluation of ventricular function in children. *Arch Cardiovasc Dis.* 2010;**103**(11-12):603-14. [PubMed ID: 21147445]. https://doi.org/10.1016/j.acvd.2010.09.004.
- Abdelmoneum MS, Elmeligy NA, Eldarky EA, Mohamad MM. Evaluation of the effect of coronary artery bypass grafting on the right ventricular function using speckle tracking echocardiography. *J Cardiol Cardiovascular Med.* 2019;4(3):236-41. https://doi.org/10.29328/journal.jccm.1001075.
- Korshin A, Gronlykke L, Holmgaard F, Kjoller SM, Gustafsson F, Nilsson JC, et al. Right ventricular transverse displacement increases following cardiac surgery: possibly compensating loss in tricuspid annular plane systolic excursion (TAPSE). J Clin Monit Comput. 2020;34(6):1139-48. [PubMed ID: 31983013]. https://doi.org/10.1007/s10877-020-00466-2.
- Khani M, Hamidzad M, Bayat F, Esfahani MA, Saffarian F, Bakhshande H, et al. 2D Speckel Tracking of RV Function after CABG and CPB Time. *Res Cardiovascular Med.* 2020;9(4):77-82. https://doi.org/10.4103/rcm.rcm_26_20.

- Quinones MA, Greenberg BH, Kopelen HA, Koilpillai C, Limacher MC, Shindler DM, et al. Echocardiographic predictors of clinical outcome in patients with left ventricular dysfunction enrolled in the SOLVD registry and trials: significance of left ventricular hypertrophy. Studies of Left Ventricular Dysfunction. J Am Coll Cardiol. 2000;35(5):1237-44. [PubMed ID: 10758966]. https://doi.org/10.1016/s0735-1097(00)00511-8.
- Solomon SD, Foster E, Bourgoun M, Shah A, Viloria E, Brown MW, et al. Effect of cardiac resynchronization therapy on reverse remodeling and relation to outcome: multicenter automatic defibrillator implantation trial: cardiac resynchronization therapy. *Circulation*. 2010;**122**(10):985-92. [PubMed ID: 20733097]. https://doi.org/10.1161/CIRCULATIONAHA.110.955039.
- 21. Watanabe J, Shiba N, Shinozaki T, Koseki Y, Karibe A, Komaru T, et al. Prognostic value of plasma brain natriuretic peptide combined with left ventricular dimensions in predicting sudden death of patients with chronic heart failure. *J Card Fail*. 2005;**11**(1):50-5. [PubMed ID: 15704064]. https://doi.org/10.1016/j.cardfail.2004.06.434.