



# Prediction of Quality of Life Based on Anger, Depression, Anxiety, spousal Intimacy, and Sexual Satisfaction in Iranian Patients Undergoing Coronary Artery Bypass Surgery

Zeynab Mahdipour <sup>1</sup>, Amir Sam Kianimoghadam <sup>2,\*</sup>, Mohammad Hasani <sup>3</sup>, Abbas Masjedi Arani <sup>2</sup>

<sup>1</sup> Department of Psychology, Shahid Beheshti University of Medical Sciences, Tehran, Iran

<sup>2</sup> Department of Clinical Psychology, Faculty of Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran

<sup>3</sup> Vascular Surgery Department of Taleghani Hospital, Shahid Beheshti University of Medical Sciences, Tehran, Iran

\*Corresponding Author: Department of Clinical Psychology, Faculty of Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran. Email: [kianimoghadam@sbmu.ac.ir](mailto:kianimoghadam@sbmu.ac.ir)

Received: 4 March, 2025; Revised: 19 July, 2025; Accepted: 20 July, 2025

## Abstract

**Background:** Psychological consequences following coronary artery bypass graft (CABG) surgery, such as anger, depression, anxiety, spousal intimacy, and sexual satisfaction, significantly impact patients' quality of life (QOL). Given the increasing prevalence of heart disease, understanding these factors is essential for post-surgical care and recovery.

**Objectives:** This study aimed to investigate the predictive role of psychological and relational factors in determining the QOL of patients undergoing CABG.

**Methods:** A cross-sectional study was conducted with 130 patients who had recently undergone CABG at Ayatollah Taleghani Hospital in Tehran. Standardized self-report questionnaires assessed psychological and relational factors, including the Quality of Life Scale (SF-36), Aggression Questionnaire (AQ), Depression Anxiety Stress Scale-21 (DASS-21), Walker-Thompson Intimacy Scale (MIS), and Larson's Sexual Satisfaction Questionnaire (LSSQ). Pearson's correlation and multiple regression analyses examined the relationships among variables, controlling for demographic factors such as age and education level.

**Results:** Depression ( $R = -0.637$ ), anxiety ( $R = -0.669$ ), and sexual satisfaction ( $R = 0.59$ ) were significant predictors of QOL, while anger ( $R = -0.371$ ) and spousal intimacy ( $R = 0.459$ ) did not demonstrate a significant impact. The regression model accounted for 59.5% of the variance in QOL ( $P < 0.01$ ). Findings suggest that psychological distress plays a stronger role in predicting QOL than relational factors in post-CABG patients.

**Conclusions:** Findings indicate that depression, anxiety, and sexual satisfaction significantly play key roles in determining the QOL of CABG patients. Implementing psychological interventions aimed at reducing depression and anxiety while enhancing sexual satisfaction may significantly improve postoperative well-being. Integrating these strategies into patient care could enhance recovery and QOL after CABG.

**Keywords:** Quality of Life, Coronary Artery Disease, Coronary Artery Bypass Graft, Depression, Anxiety, Anger, Sexual Satisfaction, Spousal Intimate Relationship

## 1. Background

Cardiovascular disease (CVD), which affects the heart and blood vessels, is becoming more prevalent each year and is a major public health concern, serving as one of the primary causes of death, disability, and reduced quality of life (QOL) globally (1). In 2019, the global

prevalence of CVD reached nearly 500 million people (1). According to conducted studies, the prevalence of CVDs in Iran has increased by 20 to 45 percent in recent years (2). About 17.9 million deaths are attributed to CVDs, which are the leading cause of death worldwide (3). These diseases also have a substantial financial impact, causing 7.28 trillion dollars in economic losses (3, 4).

Copyright © 2025, Mahdipour et al. This open-access article is available under the Creative Commons Attribution 4.0 (CC BY 4.0) International License (<https://creativecommons.org/licenses/by/4.0/>), which allows for unrestricted use, distribution, and reproduction in any medium, provided that the original work is properly cited.

**How to Cite:** Mahdipour Z, Kianimoghadam A S, Hasani M, Masjedi Arani A. Prediction of Quality of Life Based on Anger, Depression, Anxiety, spousal Intimacy, and Sexual Satisfaction in Iranian Patients Undergoing Coronary Artery Bypass Surgery. Int Cardiovasc Res J. 2025; 19 (1): e160976. <https://doi.org/10.5812/icrj-160976>.

Coronary artery bypass graft (CABG) is a widely used treatment for coronary disease, and recent advancements have significantly improved postoperative and long-term outcomes, including enhanced cardiac activity and QOL for heart disease patients (5, 6). One important outcome of treatments like CABG is QOL (7). The QOL signifies the overall well-being of individuals, which includes aspects such as health, emotional well-being, and social connections (8). In Iran, approximately 30 - 40 thousand people have undergone cardiac surgery, with CABG accounting for 50 - 60% of the total (9). The CABG is a stressful procedure that can improve life expectancy but may lead to mood disorders, physical dysfunction, and hinder participation in rehabilitation, negatively impacting QOL (10). Depression and heart disease are highly comorbid, affecting 14% to 47% of patients, many of whom experience depressive symptoms before and after CABG surgery (11). Depression is a prevalent psychiatric disorder characterized by persistent sadness, disinterest in activities, and a variety of emotional, behavioral, and cognitive symptoms (12). Patients undergoing CABG often experience anxiety and panic due to various stresses like environmental changes, pain, waiting, and fear of death (13). Anxiety is an emotional response to perceived threats, involving worry, tension, and physiological changes like increased heart rate and blood pressure (14). This surgery may also lead to anger and suppressed rage (15). Anger is an emotional response to obstacles or provocation, often shown in social interactions (16). The CABG surgery may also impact spousal intimacy (17). Improved intimacy and spousal support aid in treatment, recovery, and reduce the risk of complications, mortality, and readmission (18). The CABG can also affect sexual satisfaction, which in turn impacts the overall QOL (19). Prior studies have explored the role of psychological factors, demographic variables, and clinical characteristics in predicting recovery following CABG, but several critical gaps remain. Most studies have been conducted in various cultural or social contexts, and their results cannot be generalized to Iranian populations. Additionally, a comprehensive and interactive investigation of these factors has not been conducted in Iran, highlighting an important research gap. Previous studies have reported contradictory results regarding QOL after CABG. Some studies report significant improvements (6, 20), while others emphasize patients' difficulties in daily activities post-surgery, indicating the need for further interventions (9, 21). These discrepancies emphasize the necessity of additional research to achieve consistent and reliable

findings. The relationships between these variables and QOL remain underexplored in a systematic manner.

## 2. Objectives

This study aims to explore the predictive role of depression, anxiety, marital intimacy, and sexual satisfaction on the QOL of CABG patients, aiming to resolve contradictions in previous findings and provide clearer insights into their psychological and functional recovery.

## 3. Methods

### 3.1. Study Methods and Participants

The study employed a cross-sectional correlation method to assess the effects of anger, depression, anxiety, spousal intimacy, and sexual satisfaction on QOL in post-CABG patients. A collaboration letter for participant recruitment from Ayatollah Taleghani Hospital was obtained from the university's research vice presidency after project approval and ethics code issuance by Shahid Beheshti University of Medical Sciences. The inclusion criteria were patients aged 20 to 65 years, diagnosed with CAD, who underwent CABG within the last 12 months and are currently married. Exclusion criteria included significant comorbidities (e.g., diabetes, cancer) and incomplete or patterned questionnaire responses. Participants with chronic physical illnesses were excluded to prevent independent effects of chronic pain, fatigue, treatment side effects, and physical limitations, which could directly impact psychological states and the quality of spousal relationships. This approach enhanced sample homogeneity and improved the internal validity of the study.

The minimum sample size for regression analysis in this investigation was 91, estimated with a median effect size ( $f^2$ ) = 0.15,  $P = 0.05$ , power  $(1 - \beta) = 0.80$  (calculation made using G\*Power version 3.1), and five significant predictors that were suggested in previous research. The final analysis included 130 responses. The questionnaire, consisting of 113 questions, assessed sociodemographic variables such as gender, age, education, alcohol consumption, marital status, and smoking behavior. Data were collected from October 01, 2023, to November 01, 2024, at Hospital Center Taleghani, with approval from the Ethics Committee of Shahid Beheshti University. Participants were recruited based on inclusion criteria and completed a face-to-face structured questionnaire. They were informed about the study's purpose, data confidentiality, and their

voluntary participation was confirmed through written consent. Participants were briefed on study procedures, potential benefits, and risks. The researcher ensured participants understood their rights, including the option to withdraw at any time without consequences.

### 3.2. Measures

#### 3.2.1. Quality of Life Scale

The Quality of Life Scale (SF-36) is a 36-item questionnaire that measures QOL, with scores ranging from 26 to 130, where higher scores indicate better health (22). It has shown high reliability, with Cronbach's alpha values  $\geq 0.70$  for all eight subscales (23). The Persian version has been rigorously validated in both clinical and non-clinical populations in Iran, with Cronbach's alpha coefficients ranging from 0.70 to 0.85 for each domain. Asghari Moghaddam and Faqihi confirmed its reliability and validity for assessing QOL in Iranian populations (24).

#### 3.2.2. Aggression Questionnaire

Bass and Perry developed the Aggression Questionnaire (AGQ) in 1992, a 30-item self-report tool. It includes 14 items on anger, 8 on aggression, and 8 on hostility. In this study, only the 14 anger-related items were used. Participants select from four options: "Never", "rarely", "sometimes", and "always", scored 0 to 3. The anger score ranges from 0 to 42, with scores above the average indicating high anger (25). Soltani and Hosseini reported a Cronbach's alpha of 0.83, showing good reliability for this scale (26).

#### 3.2.3. Depression, Anxiety, and Stress Scale

The Depression Anxiety Stress Scale-21 (DASS-21) Questionnaire, developed by Lovibond and Lovibond, measures depression and anxiety through three subscales. Each subscale has a score range of 0 to 21, with higher scores indicating greater severity. Scores are reported separately, and the overall score is not calculated (27). Validity in Iran has been confirmed through factor analysis, criterion validity, and forward-backward translation. Reliability coefficients are 0.73 for anxiety and 0.81 for depression, making it a valuable tool for both clinical and research purposes (28).

#### 3.2.4. Walker-Thompson Intimacy Scale

The 17-item Walker-Thompson Intimacy Questionnaire, developed by Walker and Thompson in 1983, measures intimacy and affection among married

couples. Its Cronbach's alpha ranges from 0.91 to 0.97 (29). Validity was assessed through face and content validity. Researchers concluded it effectively measures marital intimacy. In comparison to Bagarozzi's Intimacy Scale, Babaei et al. reported a concurrent validity of 0.61 and an overall reliability of 0.9 in Iran (30).

#### 3.2.5. Larson's Sexual Satisfaction Questionnaire

A reliable method for assessing participants' sexual experiences was Larson's Sexual Satisfaction Questionnaire (LSSQ). It includes 25 items on a 5-point Likert scale from 1 = never to 5 = always, with scores ranging from 1 to 125. High sexual satisfaction has been associated with a high score (31). This questionnaire's Cronbach's  $\alpha$  was 0.93, and its validity and reliability were assessed to be 0.90 and 0.98 in Iran, respectively (32).

### 3.3. Data Analysis

The data were analyzed using IBM SPSS Statistics version 24.0. Descriptive statistics were used to summarize the demographic characteristics, including age, gender, education level, alcohol consumption, smoking habits, and disease-related factors. To check the normality of the data, skewness and kurtosis were used. We assessed the relationships between independent variables and QOL using independent *t*-tests, ANOVA, and Pearson or Spearman correlation tests, depending on the distribution of the data. A multiple linear regression analysis was performed to explore how independent variables, such as anger, depression, spousal intimacy, functional anxiety, and marital sexual satisfaction, affected QOL while adjusting for potential confounders like age and gender. The results were presented with  $R^2$  values, regression coefficients, and *P*-values, with a significance level set at  $P < 0.05$ .

## 4. Results

A total of 130 post-CABG patients (60% male, 40% female) with a mean age of  $51.9 \pm 7.9$  years participated in the study. All participants were married. Regarding education, 19.2% had a middle school degree, 18.5% held a high school diploma, and 62.3% had a college or university degree. Among them, 38.5% were smokers, 31.5% reported alcohol use, and 74.6% had a family history of heart disease. Sociodemographic and clinical characteristics are detailed in Table 1.

Independence of residuals was confirmed with the Durbin-Watson value of 1.83, indicating independent observations. Multicollinearity was not an issue, with VIF values below 10 and Tolerance above 0.1. Residual

**Table 1.** Sociodemographic and Clinical Characteristics of the Participants (N = 130) <sup>a</sup>

| Variables              | Values    |
|------------------------|-----------|
| <b>Age</b>             |           |
| 20 - 30                | 0         |
| 30 - 40                | 10 (7.7)  |
| 40 - 50                | 48 (36.9) |
| ≥ 50                   | 70 (53.8) |
| <b>Gender</b>          |           |
| Men                    | 78 (60)   |
| Women                  | 52 (40)   |
| <b>Education level</b> |           |
| ≤ Middle school        | 25 (19.2) |
| High school            | 24 (18.5) |
| ≥ College              | 81 (62.3) |
| <b>Marital status</b>  |           |
| Single                 | 0         |
| Married                | 130 (100) |
| <b>Smoking</b>         |           |
| Current smoker         | 50 (38.5) |
| Non-smoker             | 80 (61.5) |
| <b>Drinking</b>        |           |
| Current drinker        | 41 (31.5) |
| Non-drinker            | 89 (68.5) |
| <b>Family history</b>  |           |
| Yes                    | 97 (74.6) |
| No                     | 33 (25.4) |

<sup>a</sup> Values are expressed as No (%).

analysis and skewness/kurtosis values were within acceptable ranges, supporting the normality of the data for further analysis.

The participants' mean QOL scores was  $98.61 \pm 19.93$  points. The mean depression and anxiety scores were  $10.44 \pm 5.58$  and  $9.36 \pm 5.2$  points, respectively. The mean anger score was  $24.93 \pm 8.77$  points. The mean spousal intimacy and sexual satisfaction scores were  $80.86 \pm 24.07$  and  $88.60 \pm 14.72$  points, respectively (Table 2).

Table 3 presents the descriptive statistics of the study variables and the correlations between them. As demonstrated, all variables were statistically correlated with each other. The QOL score was negatively correlated with depression, anxiety, and anger, and positively associated with spousal intimacy and sexual satisfaction.

Multiple regression analysis was conducted to identify the factors influencing the participants' QOL scores. Before testing the regression model, the underlying assumptions for carrying out multiple regression analyses were examined and verified. This

ensured the validity and reliability of the regression results.

The results, as shown in Table 4, indicated that anger ( $B = -0.012$ ,  $P = 0.938$ ) did not have a significant impact on QOL. In contrast, depression ( $B = -1.082$ ,  $P = 0.0001$ ) and anxiety ( $B = -1.435$ ,  $P = 0.0001$ ) had significant negative effects on QOL. Spousal intimacy ( $B = -0.039$ ,  $P = 0.566$ ) did not exhibit a significant effect, while sexual satisfaction ( $B = 0.391$ ,  $P = 0.0001$ ) showed a significant positive effect.

The standardized coefficients ( $\beta$ ) reflect the relative impact of the variables. Here, anxiety ( $\beta = -0.375$ ) had the most significant negative effect, while sexual satisfaction ( $\beta = 0.289$ ) had the most significant positive effect on QOL. The confidence intervals (CI) for anger (CI:  $-0.329$  to  $0.304$ ) and spousal intimacy (CI:  $-0.171$  to  $0.094$ ) include zero, indicating that these variables do not have a significant effect on QOL. Conversely, depression (CI:  $-1.662$  to  $-0.501$ ), anxiety (CI:  $-1.985$  to  $-0.885$ ), and sexual satisfaction (CI:  $0.182$  to  $0.600$ ) exhibit significant effects as their confidence intervals do not include zero.

**Table 2.** Variable Means, Standard Deviations, and Normality

| Variables           | Mean $\pm$ SD     | Skewness | Kurtosis |
|---------------------|-------------------|----------|----------|
| QOL                 | 98.61 $\pm$ 19.93 | -0.27    | -0.27    |
| Depression          | 10.44 $\pm$ 5.58  | -0.17    | -0.83    |
| Anxiety             | 9.36 $\pm$ 5.2    | 0.4      | -0.53    |
| Anger               | 24.93 $\pm$ 8.77  | -0.34    | -0.1     |
| Spousal intimacy    | 80.86 $\pm$ 24.07 | -0.54    | -0.33    |
| Sexual satisfaction | 88.60 $\pm$ 14.72 | -0.56    | 1.03     |

Abbreviation: QOL, quality of life.

**Table 3.** Pearson Correlation Coefficient of the Research Variables (Significant at  $P < 0.01$ )

| Variables           | 1 | 2      | 3      | 4      | 5      | 6      |
|---------------------|---|--------|--------|--------|--------|--------|
| QOL                 | 1 | -0.637 | -0.669 | -0.371 | 0.459  | 0.59   |
| Depression          |   | 1      | 0.571  | 0.544  | -0.466 | -0.493 |
| Anxiety             |   |        | 1      | 0.347  | -0.415 | -0.479 |
| Anger               |   |        |        | 1      | -0.466 | -0.318 |
| Spousal intimacy    |   |        |        |        | 1      | 0.631  |
| Sexual satisfaction |   |        |        |        |        | 1      |

Abbreviation: QOL, quality of life.

**Table 4.** Multiple Regression of Quality of Life

| Variables           | B (Unstandardized Coefficients) | Std. Error | $\beta$ (Standardized Coefficients) | t      | P-Value | 95% CI for B (Unstandardized Coefficients) |
|---------------------|---------------------------------|------------|-------------------------------------|--------|---------|--|
| Depression          | -1.082                          | 0.293      | -0.303                              | -3.69  | 0       | (-1.662, -0.501)                           |
| Anxiety             | -1.435                          | 0.278      | -0.375                              | -5.166 | 0       | (-1.985, -0.885)                           |
| Anger               | -0.012                          | 0.16       | -0.005                              | -0.077 | 0.938   | (-0.329, 0.304)                            |
| Spousal intimacy    | -0.039                          | 0.067      | -0.047                              | -0.576 | 0.566   | (-0.171, 0.094)                            |
| Sexual satisfaction | 0.391                           | 0.106      | 0.289                               | 3.699  | 0       | (0.182, 0.600)                             |

As shown in Table 5, the model's R was 0.771, indicating a strong correlation between the predictors and the dependent variable.  $R^2$  is 0.595, showing that 59.5% of the variance in the dependent variable is explained by the model. Adjusted  $R^2$  is 0.578, indicating that 57.8% of the variability is explained after accounting for the predictors.  $\Delta R^2$  is 0.08. The F statistic is 36.368, and the P-value is 0.000, showing the model's overall statistical significance ( $P < 0.05$ ).

## 5. Discussion

This study examines the factors affecting the QOL in Iranian patients after CABG. Although the surgery may impact QOL, other factors also play a role in its outcome (33). Among the research variables, anxiety, depression, and sexual satisfaction had the most explanatory power

in predicting the QOL of patients undergoing CABG. Higher depression levels were associated with lower QOL ( $P < 0.001$ ), consistent with previous research (34, 35). Even with successful surgery, depression negatively impacts QOL (36). Limited social support, reduced activity, and job loss due to prolonged recovery can increase a sense of loss of control, raising depression risk (37-39). The post-surgical period disrupts daily routines, making future planning difficult and further increasing depression likelihood (40). Additionally, depression activates the hypothalamus-pituitary-adrenal axis, boosting pro-inflammatory cytokines (41). In this study, a significant relationship was found between QOL and anxiety ( $P < 0.001$ ).

Although anxiety may decrease post-surgery, it often does not return to subclinical levels. Factors such as a pessimistic outlook, catastrophizing, and a lack of



**Table 5.** Model Summary for Predicting Quality of Life

| R     | R <sup>2</sup> | Adj R <sup>2</sup> | $\Delta R^2$ | F      | P |
|-------|----------------|--------------------|--------------|--------|---|
| 0.771 | 0.595          | 0.578              | 0.08         | 36.368 | 0 |

control exacerbate anxiety (42). Physiological changes like hypothalamus-pituitary-adrenal (HPA) axis activation and increased cortisol (43), along with financial problems, treatment costs, and the inability to return to work, further intensify anxiety (44). Insufficient information from healthcare providers, fear of health status without medical supervision, difficulty adapting to life after discharge, and changes in social roles also contribute to heightened anxiety (42, 45).

In this study, sexual satisfaction showed a positive and significant relationship ( $P < 0.001$ ) with QOL. Some previous studies are consistent with the results of this research (46, 47), while others obtained results contrary to our study (48). The CVD contributes to impaired sexual satisfaction, and CABG can significantly address these issues (49). The CABG, by reducing angina, improves social, physical, and sexual activities, ultimately leading to sexual satisfaction (46).

In this study, contrary to the findings of Rapelli et al., no significant relationship ( $P = 0.566$ ) was found between marital intimacy and QOL in CABG patients (17). This discrepancy may arise from cultural differences in expressing intimacy; in collectivist Iranian culture, spousal support is often implicit, involving family duties or nonverbal emotional support, while explicit verbal support is more common in Western cultures (50). Instruments like the Walker-Thompson Scale, developed in Western contexts, may not fully capture these implicit forms of intimacy in Iran. Additionally, CABG patients, due to fatigue and physical limitations after surgery, may prioritize physical recovery or social support over emotional intimacy (51). Therefore, the nonsignificant relationship could be due to tool limitations or differing patient priorities. Future studies should use culturally adapted instruments or mixed-method approaches.

Additionally, in this study, anger had no significant impact ( $P = 0.938$ ). As a transient emotion, it may not have yielded significant results due to the study's cross-sectional design. Overall, this study reaffirms the negative effects of depression and anxiety while highlighting the positive role of sexual satisfaction in improving QOL in cardiac patients. The discrepancies in the influence of anger and spousal intimacy call for further research to understand the complex

psychological and social factors affecting the QOL in this population.

The current findings are particularly significant given their practical implications. Depression and anxiety are critical targets for postoperative care, and psychological interventions should be integrated into rehabilitation programs to address these issues. Evidence-based therapies such as cognitive-behavioral therapy (CBT) and mindfulness-based stress reduction (MBSR) have shown efficacy in reducing psychological distress and improving QOL in cardiac patients (52, 53). Moreover, couple-based sexual counseling, encompassing educational and interactive sessions with both the patient and their spouse, can enhance sexual satisfaction by improving communication skills (46).

This study's strengths include an adequate sample size, enhancing statistical power, and a diverse sample of male and female patients from various age and education levels, allowing for generalizability. Using valid, standardized tools and multiple regression analysis ensures accurate, precise results by examining the simultaneous effects of multiple variables. However, the study has some limitations.

### 5.1. Cross-Sectional Design Limitations

The cross-sectional design of this study does not allow for causal inference or determination of temporal directionality between variables. This approach was chosen due to time and resource limitations; however, the findings are restricted to correlational relationships. Future longitudinal studies are recommended to more accurately examine causal and temporal associations.

### 5.2. Exclusion of Patients with Chronic Comorbidities

Excluding patients with chronic comorbidities limits the generalizability of the findings and narrows the sample to a subgroup of CABG patients without such conditions. Consequently, the results may not fully represent the broader CABG population. Future studies could enhance generalizability by including these patients or statistically controlling for relevant variables, providing a more comprehensive understanding of the psychological and relational factors involved.

### 5.3. Limited Scope of the Anger Subscale

The use of the 14-item Anger subscale from the AGQ may not capture other dimensions of aggression, such as hostility or physical aggression. This selection was based on the primary focus on the emotional experience of anger, but it may not fully reflect the role of aggression in the QOL of CABG patients. Therefore, employing the full AGQ or additional subscales is recommended for a more comprehensive assessment of various aggression dimensions.

### 5.4. Focus on Sexual Satisfaction

This study focused on sexual satisfaction as a cognitive construct and therefore used the LSSQ, which does not assess sexual functioning. Consequently, the results may have overlooked the role of sexual functioning. Future studies should employ more comprehensive tools like the International Index of Erectile Function (IIEF) and the Female Sexual Function Index (FSFI) to better examine sexual functioning as a mediator in the relationship with QOL.

### 5.5. Multicollinearity and Factor Analysis

The VIF and Tolerance indices confirmed that multicollinearity was not an issue. However, the lack of factor analysis, despite the validated structure of DASS-21 in Iran, and the choice to analyze depression and anxiety separately may have limited insights into their overlap. Future studies with larger samples could examine their structure via factor analysis and consider using a composite 'psychological distress' Index to reflect shared variance while maintaining clinical distinctions.

### 5.6. Reliance on Self-Reported Data

Reliance on self-reported data may introduce response bias. Future studies should consider adding physiological measures, such as cortisol, to improve assessment accuracy. This approach would provide a more objective measure of psychological states and enhance the robustness of the findings by correlating self-reported data with physiological indicators.

## Footnotes

**Authors' Contribution:** Study concept, design, analysis, interpretation of data, drafting of the manuscript, critical revision of the manuscript for important intellectual content, statistical analysis: Z. M.;

Corresponding author and supervision: A. S. K.; Resources: M. H.

**Conflict of Interests Statement:** The authors declare no conflict of interest.

**Data Availability:** The dataset presented in the study is available on request from the corresponding author during submission or after publication.

**Ethical Approval:** This study was approved under the ethical approval code [IR.SBMU.MSP.REC.1402.513](#).

**Funding/Support:** The present study received no funding/support.

**Informed Consent:** Informed consent was obtained from all participant.

## References

1. Rethemiotaki I. Global prevalence of cardiovascular diseases by gender and age during 2010-2019. *Arch Med Sci Atheroscler Dis.* 2023;**8**:e196-205. [PubMed ID: [38283927](#)]. [PubMed Central ID: [PMC10811545](#)]. <https://doi.org/10.5114/amsad/176654>.
2. Pourghane P, Ghalandari M, Shahroudi P, Rahimi F, Godarzwand L. A Comparative Study on Demographic, Clinical and Laboratory Characteristics of Patients Undergoing Coronary Artery Bypass Graft in a Hospital in Iran: 2018-2019 and 2020-2021. *J Guilan Univ Med Sci.* 2024;**33**(3):280-91. <https://doi.org/10.32598/jgums.33.3.469.4>.
3. Hossain MS, Dulal DD, Faysal DF. Prevalence of Cardiovascular Disease and Associated Risk Factors among Adults. *Dinkum Journal of Medical Innovations.* 2024;**3**(5):379-90.
4. Xu L, Zhai X, Shi D, Zhang Y. Depression and coronary heart disease: mechanisms, interventions, and treatments. *Front Psychiatry.* 2024;**15**:1328048. [PubMed ID: [38404466](#)]. [PubMed Central ID: [PMC10884284](#)]. <https://doi.org/10.3389/fpsy.2024.1328048>.
5. Harik L, Perezgrovas-Olaria R, Soletti GJ, Dimagli A, Alzghari T, An KR, et al. Graft thrombosis after coronary artery bypass surgery and current practice for prevention. *Front Cardiovasc Med.* 2023;**10**:1125126. [PubMed ID: [36970352](#)]. [PubMed Central ID: [PMC10031065](#)]. <https://doi.org/10.3389/fcvm.2023.1125126>.
6. Schmidt-RioValle J, Abu Ejheisheh M, Membrive-Jimenez MJ, Suleiman-Martos N, Albendin-Garcia L, Correa-Rodriguez M, et al. Quality of Life After Coronary Artery Bypass Surgery: A Systematic Review and Meta-Analysis. *Int J Environ Res Public Health.* 2020;**17**(22). [PubMed ID: [33202650](#)]. [PubMed Central ID: [PMC7697861](#)]. <https://doi.org/10.3390/ijerph17228439>.
7. Kim H, Bae SH, Lim SH, Park JH. Predictors of health-related quality of life after coronary artery bypass graft surgery. *Sci Rep.* 2022;**12**(1):16119. [PubMed ID: [36167788](#)]. [PubMed Central ID: [PMC9515078](#)]. <https://doi.org/10.1038/s41598-022-20414-1>.
8. van Leeuwen KM, van Loon MS, van Nes FA, Bosmans JE, de Vet HCW, Ket JCF, et al. What does quality of life mean to older adults? A thematic synthesis. *PLoS One.* 2019;**14**(3). e0213263. [PubMed ID: [30849098](#)]. [PubMed Central ID: [PMC6407786](#)]. <https://doi.org/10.1371/journal.pone.0213263>.
9. Jannati M, Aslani A. Management of anxiety and depression in post coronary artery bypass graft surgery. *J Mind Med Sci.* 2023;**10**(1):34-41. <https://doi.org/10.22543/2392-7674.1360>.

10. Correa-Rodriguez M, Abu Ejheisheh M, Suleiman-Martos N, Membrive-Jimenez MJ, Velando-Soriano A, Schmidt-RioValle J, et al. Prevalence of Depression in Coronary Artery Bypass Surgery: A Systematic Review and Meta-Analysis. *J Clin Med*. 2020;**9**(4). [PubMed ID: 32225052]. [PubMed Central ID: PMC7230184]. <https://doi.org/10.3390/jcm9040909>.
11. Raven S, Bader C, Azar A, Rudolph JL. Depressive symptoms after CABG surgery: a meta-analysis. *Harv Rev Psychiatry*. 2013;**21**(2):59-69. [PubMed ID: 23656830]. [PubMed Central ID: PMC3652273]. <https://doi.org/10.1097/HRP.0b013e31828a3612>.
12. Chen X. The diversity of clinical features of depressive disorders and their diagnostic challenges. *Transactions on Materials, Biotechnology and Life Sciences*. 2024;**2**:90-5. <https://doi.org/10.62051/tfy16r18>.
13. Wu L, Yao Y. Exploring the effect of music therapy as intervention to reduce anxiety pre- and post-operatively in CABG surgery: A quantitative systematic review. *Nurs Open*. 2023;**10**(12):7544-65. [PubMed ID: 37823363]. [PubMed Central ID: PMC10643824]. <https://doi.org/10.1002/nop2.2024>.
14. Kogan CS, Wong C-W, Emmelkamp PMG. Anxiety and fear-related disorders. *American Psychol Association*. 2024:137-154. <https://doi.org/10.1037/0000392-008>.
15. Dubyts DC. *The experience of waiting for coronary artery bypass graft surgery*. Kelowna, British Columbia: University of British Columbia; 1988.
16. Silva L. Is Anger a Hostile Emotion? *Rev Philosophy Psychol*. 2021;**15**(2):383-402. <https://doi.org/10.1007/s13164-021-00557-2>.
17. Rapelli G, Donato S, Giusti EM, Pietrabissa G, Parise M, Pagani AF, et al. Recognizing and Appreciating the Partner's Support Protects Relationship Satisfaction during Cardiac Illness. *J Clin Med*. 2024;**13**(4). [PubMed ID: 38398491]. [PubMed Central ID: PMC10890017]. <https://doi.org/10.3390/jcm13041180>.
18. Barbod B, Rezapour MY, Choobfroushzadeh A, Mirhosseini J, Koohestani F. [Effect of spousal support and companionship intervention on quality of life, marital intimacy and stress in patients after coronary artery bypass graft surgery (CABG): A multiple baseline single-case study]. *J Cardiovascular Nurs J*. 2020;**9**(1):56-67. FA.
19. Inayat S, Aziz F, Younas A, Durante A. Determinants of Sexual Health and Sexual Quality of Life after Cardiovascular Surgeries: An Integrative Review. *Heart Lung Circ*. 2024;**33**(10):1414-26. [PubMed ID: 38969608]. <https://doi.org/10.1016/j.hlc.2024.05.010>.
20. Pacaric S, Turk T, Eric I, Orkic Z, Petek Eric A, Milostic-Srb A, et al. Assessment of the Quality of Life in Patients before and after Coronary Artery Bypass Grafting (CABG): A Prospective Study. *Int J Environ Res Public Health*. 2020;**17**(4). [PubMed ID: 32098322]. [PubMed Central ID: PMC7068373]. <https://doi.org/10.3390/ijerph17041417>.
21. Emamzadehashemi KR, Khanghah AG, Azizi A, Paryad E, Noveiri MJS. Quality of life and activities of daily living one year after Coronary Artery Bypass Graft (CABG) surgery: a cross-sectional study. *J Cardiothorac Surg*. 2024;**19**(1):367. [PubMed ID: 38915074]. [PubMed Central ID: PMC11637172]. <https://doi.org/10.1186/s13019-024-02848-y>.
22. Hunt JO, Hendrata MV, Myles PS. Quality of life 12 months after coronary artery bypass graft surgery. *Heart Lung*. 2000;**29**(6):401-11. [PubMed ID: 11080320]. <https://doi.org/10.1067/mhl.2000.110578>.
23. Yarlas A, Bayliss M, Cappelleri JC, Maher S, Bushmakini AG, Chen LA, et al. Psychometric validation of the SF-36(R) Health Survey in ulcerative colitis: results from a systematic literature review. *Qual Life Res*. 2018;**27**(2):273-90. [PubMed ID: 28849311]. <https://doi.org/10.1007/s11136-017-1690-6>.
24. Asghari Moghaddam M, Faghehi S. [Validity of the sf-36 health survey questionnaire in two Iranian samples]. *Clin Psychol Personality*. 2003;**1**(1):1-10. FA.
25. Buss AH, Perry M. The aggression questionnaire. *J Pers Soc Psychol*. 1992;**63**(3):452-9. [PubMed ID: 1403624]. <https://doi.org/10.1037/0022-3514.63.3.452>.
26. Soltani F, Hosseini F. [The relationship of social support with aggression and hopelessness in nurses]. *Iran J Nurs Res*. 2018;**13**(2):74-80. FA.
27. Antoni MH, Wimberly SR, Lechner SC, Kazi A, Sifre T, Urcuyo KR, et al. Reduction of cancer-specific thought intrusions and anxiety symptoms with a stress management intervention among women undergoing treatment for breast cancer. *Am J Psychiatry*. 2006;**163**(10):1791-7. [PubMed ID: 17012691]. [PubMed Central ID: PMC5756627]. <https://doi.org/10.1176/ajp.2006.163.10.1791>.
28. Sahebi A, Asghari MJ, Salari RS. [Validation of depression anxiety and stress scale (DASS-21) for an Iranian population]. *J Iran Psychol*. 2005;**1**(4). FA.
29. Walker AJ, Thompson L. Intimacy and Intergenerational Aid and Contact among Mothers and Daughters. *J Marriage Family*. 1983;**45**(4). <https://doi.org/10.2307/351796>.
30. Babaei R, Yazdi SM, Ghasemzadeh S. [Prediction of Marital Satisfaction based on Perceived Support, Alexithymia and Empresment]. *J Nurse Physician Within War*. 2019;**6**(18):43-53. FA.
31. Larson JH, Anderson SM, Holman TB, Niemann BK. A longitudinal study of the effects of premarital communication, relationship stability, and self-esteem on sexual satisfaction in the first year of marriage. *J Sex Marital Ther*. 1998;**24**(3):193-206. [PubMed ID: 9670124]. <https://doi.org/10.1080/00926239808404933>.
32. Shams Mofarrah Z, Shahsiah M, Mohebi S, Tabaraee Y. [The effect of marital counseling on sexual satisfaction of couples in Shiraz city]. *J Health System Res*. 2011;**6**(3):0. FA.
33. Alzahrani AA, AlAssiri AK, Al-Ebrahim KE, Ganbou ZT, Alsudaiss MM, Khafagy AM. Impact of Clinical and Sociodemographic Factors on Quality of Life Following Coronary Artery Bypass Grafting: A Mixed-Methods Study. *Cureus*. 2024. <https://doi.org/10.7759/cureus.56781>.
34. AbuRuz ME, Al-Dweik G. Depressive symptoms, perceived control and quality of life among patients undergoing coronary artery bypass graft: a prospective cohort study. *BMC Nurs*. 2022;**21**(1):87. [PubMed ID: 35410251]. [PubMed Central ID: PMC9004116]. <https://doi.org/10.1186/s12912-022-00857-7>.
35. Cyganrowicz B, Milaniak I. The effect of a preoperative education program on the quality of life and the reduction of symptoms of anxiety and depression in patients undergoing a coronary artery bypass graft surgery – a quasi-experimental study. *Problemy Pielęgniarstwa*. 2024;**32**(2):77-84. <https://doi.org/10.5114/ppiel.2024.140688>.
36. Stafford L, Berk M, Reddy P, Jackson HJ. Comorbid depression and health-related quality of life in patients with coronary artery disease. *J Psychosom Res*. 2007;**62**(4):401-10. [PubMed ID: 17383491]. <https://doi.org/10.1016/j.jpsychores.2006.12.009>.
37. Alsubaie MM, Stain HJ, Webster LAD, Wadman R. The role of sources of social support on depression and quality of life for university students. *Int J Adolescence Youth*. 2019;**24**(4):484-96. <https://doi.org/10.1080/02673843.2019.1568887>.
38. Mortensen M, Sandvik R, Svendsen OS, Haaverstad R, Moi AL. Return to work after coronary artery bypass grafting and aortic valve replacement surgery: A scoping review. *Scand J Caring Sci*. 2022;**36**(4):893-909. [PubMed ID: 34057755]. <https://doi.org/10.1111/scs.13006>.
39. Mehrdad R, Ghadiri Asli N, Pouryaghoub G, Saraei M, Salimi F, Nejatian M. Predictors of early return to work after a coronary artery bypass graft surgery (CABG). *Int J Occup Med Environ Health*. 2016;**29**(6):947-57. [PubMed ID: 27869245]. <https://doi.org/10.13075/ijom.1896.00798>.



40. LaPier TK, Wintz G, Holmes W, Cartmell E, Hartl S, Kostoff N, et al. Analysis of Activities of Daily Living Performance in Patients Recovering from Coronary Artery Bypass Surgery. *Physical & Occupational Therapy In Geriatrics*. 2009;27(1):16-35. <https://doi.org/10.1080/02703180802206215>.
41. AbuRuz ME, Momani A, Shajrawi A. The Association Between Depressive Symptoms and Length of Hospital Stay Following Coronary Artery Bypass Graft is Moderated by Perceived Control. *Risk Manag Healthc Policy*. 2021;14:1499-507. [PubMed ID: 33883956]. [PubMed Central ID: PMC8053611]. <https://doi.org/10.2147/RMHP.S306162>.
42. Robley L, Ballard N, Holtzman D, Cooper W. The experience of stress for open heart surgery patients and their caregivers. *West J Nurs Res*. 2010;32(6):794-813. [PubMed ID: 20696847]. <https://doi.org/10.1177/0193945910361469>.
43. Hinds JA, Sanchez ER. The Role of the Hypothalamus-Pituitary-Adrenal (HPA) Axis in Test-Induced Anxiety: Assessments, Physiological Responses, and Molecular Details. *Stresses*. 2022;2(1):146-55. <https://doi.org/10.3390/stresses2010011>.
44. Caumo W, Schmidt AP, Schneider CN, Bergmann J, Iwamoto CW, Adamatti LC, et al. Risk factors for postoperative anxiety in adults. *Anaesthesia*. 2001;56(8):720-8. [PubMed ID: 11493233]. <https://doi.org/10.1046/j.1365-2044.2001.01842.x>.
45. Bijttebier P, Vanoost S, Delva D, Ferdinande P, Frans E. Needs of relatives of critical care patients: perceptions of relatives, physicians and nurses. *Intensive Care Med*. 2001;27(1):160-5. [PubMed ID: 11280629]. <https://doi.org/10.1007/s001340000750>.
46. Lai YH, Hsieh SR, Ho WC, Chiou AF. Factors associated with sexual quality of life in patients before and after coronary artery bypass grafting surgery. *J Cardiovasc Nurs*. 2011;26(6):487-96. [PubMed ID: 21263341]. <https://doi.org/10.1097/JCN.0b013e3182050269>.
47. Caglar M, Yesiltepe Oskay U, Arican E, Akyol E. Relationship between self-efficacy and sexual function after open heart surgery. *Eur J Cardiovasc Nurs*. 2021;20(4):376-82. [PubMed ID: 33620463]. <https://doi.org/10.1093/eurjcn/zvaa012>.
48. Reese JB, Shelby RA, Taylor KL. Sexual quality of life in patients undergoing coronary artery bypass graft surgery. *Psychol Health*. 2012;27(6):721-36. [PubMed ID: 22149897]. <https://doi.org/10.1080/08870446.2011.623781>.
49. Yuksel MB, Gecit I, Karakose A, Pirincci N, Gunes M, Aydin C, et al. Coronary Artery Bypass Surgery and Sexual Function. *Medical Science and Discovery*. 2015;2(4). <https://doi.org/10.17546/msd.36693>.
50. Keshtiar N, Kuhlmann M. The effects of culture and gender on the recognition of emotional speech: Evidence from Persian speakers living in a collectivist society. *Int J Society, Culture & Language*. 2016;4(2):71-86.
51. Dolapoglu N, Dolapoglu A, Altunoz S. Caregiver burden, family adaptation, partnership, growth, affection, and resolve, anxiety levels, and perceived social support in relatives of patients with open heart surgery. *Rev Assoc Med Bras (1992)*. 2024;70(2). e20230494. [PubMed ID: 38198299]. [PubMed Central ID: PMC10768673]. <https://doi.org/10.1590/1806-9282.20230494>.
52. Holdgaard A, Eckhardt-Hansen C, Lassen CF, Kjesbu IE, Dall CH, Michaelsen KL, et al. Cognitive-behavioural therapy reduces psychological distress in younger patients with cardiac disease: a randomized trial. *Eur Heart J*. 2023;44(11):986-96. [PubMed ID: 36649937]. <https://doi.org/10.1093/eurheartj/ehac792>.
53. Wu K, Wan M, Zhou H, Li C, Zhou X, Li E, et al. Mindfulness-based stress reduction combined with early cardiac rehabilitation improves negative mood states and cardiac function in patients with acute myocardial infarction assisted with an intra-aortic balloon pump: a randomized controlled trial. *Front Cardiovasc Med*. 2023;10:1166157. [PubMed ID: 37324635]. [PubMed Central ID: PMC10265675]. <https://doi.org/10.3389/fcvm.2023.1166157>.