Published online: 2024 March 26.

Research Article



Reduction of Surgical Bleeding During Lumbar Laminectomy with Pedicular Screw Using Oral Clonidine as a Premedication: A Randomized Comparative Study

Bahman Naghipour¹, Seyedeh Sorour Zahedi^{2,*}, Eisa Bilehjani¹, Farhad Mirzaei³, Sepideh Sadat Zahedi², Sahar Zahedi⁴

¹ Department of Anesthesiology, Madani Heart Center, Faculty of Medicine, Tabriz University of Medical Sciences, Tabriz, Iran

² Department of Anesthesiology, Faculty of Medicine, Tabriz University of Medical Sciences, Tabriz, Iran

³ Department of Neurosurgery, Imam Reza Hospital, Faculty of Medicine, Tabriz University of Medical Sciences, Tabriz, Iran

⁴ Department of ENT, Faculty of Medicine, Tabriz Azad University of Medical Sciences, Tabriz, Iran

corresponding author: Department of Anesthesiology, Faculty of Medicine, Tabriz University of Medical Sciences, Tabriz, Iran. Email: sururzahedi@yahoo.com

Received 2024 February 28; Accepted 2024 March 26.

Abstract

Background: Spinal fusion surgeries are often associated with major blood loss. The aim of this study was to investigate the effect of oral clonidine as premedication on the amount of blood loss and the need for blood transfusion during lumbar laminectomy spine surgery with pedicle screw insertion, achieved by controlled hypotension.

Methods: In this double-blind randomized clinical trial, 100 patients scheduled for lumbar spine laminectomy with pedicle screw insertion (as a fusion procedure) were included. Patients were classified into two groups: The intervention group received a 0.2 mg clonidine tablet one hour before entering the operating room, and the control group received a placebo. The patients were compared with respect to hemodynamic variables, quality of the surgical field, duration of the surgery, estimated amount of blood loss, and any blood transfusion.

Results: Oral clonidine was found to be significantly better in terms of maintaining stable dynamics and having a lower amount of blood loss, thereby requiring fewer blood transfusions compared to the control group.

Conclusions: Premedication with 0.2 mg oral clonidine in patients undergoing lumbar spine laminectomy with pedicle screw insertion results in improved perioperative hemodynamic stability and a reduction in the amount of intraoperative blood loss, thereby reducing the need for blood transfusion.

Keywords: Clonidine, Blood Loss, Control of Blood Pressure, Alpha-2 Adrenergic, Fusion Surgery

1. Background

hemorrhage Intraoperative is а potential complication in surgical procedures, posing significant challenges for surgeons and anesthesiologists (1). Attention to intraoperative and postoperative hemorrhage is crucial, and its source should be promptly controlled (2). Heavy bleeding typically occurs after more extensive operations (3). Spinal surgery, often characterized bv a significant intraoperative hemorrhage, may be attributed partly to factors such as large surgical incisions, prolonged operation duration, and increased involvement of cancellous bone (4). Intraoperative and postoperative hemorrhage can lead to heightened coagulation, postoperative hematoma, anemia, and the need for allogeneic blood transfusion, which may result in complications like hemolytic and non-hemolytic reactions, acute lung injury, transmission of viral and bacterial infections, hypothermia, and coagulation disorders (5), incurring costs for both families and society. The economic burden of blood loss includes direct costs, such as blood products and intraoperative hemorrhage control technology, as well as indirect costs like long-term patient hospitalization and complication management (6, 7).

Numerous studies have aimed to reduce hemorrhage during spine fusion surgeries using various

Copyright © 2024, Naghipour et al. This open-access article is available under the Creative Commons Attribution 4.0 (CC BY 4.0) International License (https://creativecommons.org/licenses/by/4.0/), which allows for unrestricted use, distribution, and reproduction in any medium, provided that the original work is properly cited.

medications, including aprotinin, aminocaproic acid, tranexamic acid, and modified factor VII (8, 9). Standard anesthesia techniques and arterial blood pressure control play significant roles in preventing excessive intraoperative hemorrhage. The correlation between intraoperative hypotension and reduced intraoperative hemorrhage has been extensively documented, highlighting the importance of lowering blood pressure to minimize blood loss (10-12). Alpha-2 adrenergic agonists such as clonidine and dexmedetomidine, with distinct mechanisms of action compared to other anesthetics, have been studied for their effects on hemodynamic stability. Clonidine, an imidazole derivative, exerts agonistic effects on alpha-2 adrenergic receptors, is well absorbed orally, reaches peak effect in 60 - 90 minutes, and has an elimination half-life of 9 - 12 hours. Low-dose clonidine infusion intravenously reduces mean arterial pressure by activating central and peripheral alpha-2 receptors, while high doses through intravenous infusion or rapid intravenous bolus may elevate systemic blood pressure by activating peripheral alpha-1 adrenergic receptors post-binding (9).

Singh et al. (13) assessed the effects of clonidine versus atenolol on improving surgical visibility, reducing intraoperative blood loss, and maintaining hemodynamic stability in rhinoplasty under general anesthesia. Their study involved 60 ASA grade I and II patients undergoing rhinoplasty, randomly assigned to two groups of 30 people. Two hours before surgery, group A received 50 mg of oral atenolol, while group B received 100 µg of oral clonidine. Results showed that heart rate (HR) and blood pressure remained within normal ranges in both groups throughout the surgery. Patients who received clonidine experienced less blood loss, and surgeons reported better visibility during the operation. Bajaj et al. (14) investigated clonidine's efficacy and safety in reducing blood loss during pituitary adenoma surgery. The study included 50 patients with pituitary adenoma randomly divided into two groups. Group A (25 patients) received oral clonidine (200 µg), while group B (25 patients) received a placebo. Clonidine was found to be a safe and effective agent for reducing hemorrhage in microscopic transsphenoidal pituitary adenoma surgeries.

Ebneshahidi and Mohseni (9) examined the impact of oral clonidine on reducing intraoperative hemorrhage and maintaining hemodynamic stability during cesarean section under general anesthesia. They observed significantly lower blood loss and reduced narcotic requirements in the clonidine-treated group compared to the placebo group. Deepa and Shekhar (15) investigated the effects of pre-medication with intravenous clonidine on modulating hemodynamic responses during laparoscopic surgeries in 60 patients. Patients were divided into two groups: One group received 2 µg/kg of intravenous clonidine, while the second group received a placebo. Their findings indicated that patients pre-medicated with intravenous clonidine exhibited stable hemodynamics and improved analgesia and sedation compared to those who did not receive clonidine.

Overall, pre-medication with clonidine has been shown to significantly enhance patient stability during and after surgery across various procedures. Given its efficacy in reducing hemorrhage in multiple surgeries and its positive effects in thoracic spine trauma surgery, we aimed to evaluate its impact on bleeding volume in patients undergoing lumbar laminectomy with pedicle screw fixation for lumbar disc herniation. By mitigating side effects and reducing hospitalization duration and associated costs for families and society, widespread adoption of this approach can be justified based on the findings of this field study.

1.1. Fusion Surgery

Spinal fusion is performed to correct deformities or relieve back pain and is recommended for diseases such as degenerative scoliosis, spondylolisthesis, and vertebral fractures causing spinal instability. Additionally, fusion surgery is used in conditions such as tumors, spinal canal stenosis, disc herniation, and spinal weakness or instability caused by diseases like arthritis (spondylosis).

1.2. Clonidine

Clonidine is an imidazole derivative drug that has an agonistic effect on alpha-2 adrenergic receptors. Clonidine is well absorbed orally, with an onset of effect around 30 - 60 minutes, a peak effect at 60 - 90 minutes, and a half-life of 9 - 12 hours. In the central nervous system and in the presynaptic location of nerves, clonidine inhibits the secretion and release of the neurotransmitter norepinephrine, thereby having an inhibitory effect on the sympathetic system.

1.3. Dexmedetomidine

It is an alpha-2 adrenergic agonist with sedative, anxiolytic, hypnotic, analgesic, and sympatholytic effects. It is metabolized in the liver and excreted through urine and feces.

2. Methods

2.1. Study Population, Clinical Design, and Disposition

This study was conducted at the educationaltherapeutic center of Imam Reza, Tabriz, between January 2020 and July 2020, involving patients referred to the hospital's operating room for laminectomy surgery with pedicle screw insertion (as a fusion procedure) due to lumbar intervertebral disc herniation. Written and informed consent was obtained from all patients, and their information remained confidential. The study received approval from the local ethics committee of Tabriz University of Medical Sciences (IR.TBZMED.REC.1398.936) and was registered at Iranian Registry of Clinical the Trials (IRCT20140107016117N4).

The sample size was estimated to be 100 patients aged 20 - 60 years, selected through a convenience consecutive method based on the order of patients visiting the operating room. Inclusion criteria were ASA class I - II patients aged 20 - 60 years, candidates for laminectomy with pedicle screw fixation. Exclusion criteria included patients with systemic diseases (cardiovascular, cerebral, renal, liver, and kidney), blood loss exceeding 10% in hypertensive patients, diabetic patients, and non-use of medications that interfere with clonidine, such as ibuprofen, losartan, beta-blockers, diazepam, and vitamin B12 (16, 17).

Patients were randomly assigned to two groups (intervention and control). The intervention group received a clonidine tablet (0.2 mg) one hour before entering the operating room with half a glass of water in the ward and then were transferred to the operating room. No pill was administered to the control group. Both patients and anesthesiologists were blinded to the drug (clonidine). All procedures were managed by a resident anesthesiologist. In the operating room, all patients received intravenous midazolam (2 mg) and fentanyl (100 μ g), followed by induction of anesthesia using propofol (2 mg/kg), cis-atracurium (0.15 mg/kg), and lidocaine (1 mg/kg). Three minutes after anesthesia induction y monitoring HR, blood pressure, O₂ sat, and electrocardiogram (ECG), patients were intubated with a cuffed spiral tube (inner diameter 8 mm in men and 7.5 mm in women). Anesthesia was maintained with nitrous oxide 3 L/min and O₂ 3 L/min, isoflurane 1%, and remifentanil infusion (0.05 µg/kg/min). Blood pressure and HR were recorded before anesthesia, after anesthesia induction, 1 minute and 3 minutes after intubation, 1 minute after surgical incision, and then every 15 minutes until recovery. If blood pressure exceeded 150/100 despite anesthesia, intravenous

nitroglycerin (TNG) infusion was used to reduce blood pressure, and the amount used was recorded. The hemodynamic status of the patient and the amount of TNG used were compared between the two groups.

2.2. Statistical Analysis

The data were expressed using descriptive statistics (mean \pm standard deviation and frequency percentage) and analyzed using paired-samples *t*-test, repeated-measures ANOVA, and independent *t*-test. Normal distribution of data was assessed using the Kolmogorov–Smirnov test. SPSS 22 software was used for data analysis, and a P-value < 0.05 was considered statistically significant.

3. Results

Patient demographics are displayed in Table 1, indicating no statistically significant difference between the two groups. Additionally, regarding sex distribution, sixty cases (60%) were male, and 40 cases (40%) were female, with males showing the highest frequency in both groups. Concerning physical condition and ASA class, 88% of the patients were classified as class I, and 12% as class II, with no significant difference between the two groups (P = 0.067).

Table 1. Demographic Characteristics During the Study in two Groups				
Variables	No.	Mean ± SD	P-Value	
Height (cm)				
Control	50	164.7 ± 4.1	0.560	
Clonidine	50	163.8 ± 4.1	0.560	
Weight (kg)				
Control	50	79.2 ± 5.82	0.245	
Clonidine	50	81.4 ± 4.80	0.245	
BMI (kg/m ²)				
Control	50	27.3 ± 2.05	0.468	
Clonidine	50	27.5 ± 1.92	0.468	
Age (y)				
Control	50	43.6 ± 6.73	0.237	
Clonidine	50	41.6 ± 3.39	0.237	

Surveys indicated that the surgery duration in the clonidine group lasted 110 minutes, while the average for the control group was 170 minutes, and this difference was statistically significant. The total dose of remifentanil infusion in the clonidine and control groups was 50.7 ± 69 and 623 ± 13 , respectively, and this difference was statistically significant (P = 0.01). TNG was not used in either of the two groups.

Figure 1A shows that mean systolic blood pressure (SBP) in the clonidine group decreased after induction



of anesthesia, after intubation, and during surgery compared to the control group, and this decrease was significant. Figure 1B show that similar to SBP, in the clonidine group, the mean diastolic blood pressure (DBP) significantly reduced after induction of anesthesia, after intubation and during surgery. According to Figure 1C, in the clonidine group, the average blood pressure (ABP) values were significantly reduced compared to the control group. Figure 1D shows that the average HR in the clonidine group decreased significantly after intubation, during surgery and recovery compared to the control group. Also, Table 2 shows the average standard deviations for SBP, DBP, ABP and HR values.



Figure 2. A, the amount of blood transfusion; B, surgeon satisfaction; C, total hemorrhage; and D, amount of blood in the drains on the first and second days after surgery

Table 2. Mean Standard Deviation for Studied Values					
Result	Average Deviation Control Group	Average Deviation Clonidine Group	P- Value		
SBP	±14.2	±15.1	0.04		
DBP	±12.2	±13.4	0.03		
ABP	±14.5	±15.5	0.04		
HR	±12.3	±13.8	0.03		

The average amount of blood transfusion in the clonidine and control groups was 7 and 16 units, respectively, and a significant difference was observed between the two groups (P = 0.02) (Figure 2A). Regarding surgeon satisfaction, 84% of surgeons in the clonidine group and 32% in the control group were satisfied, and there was a significant difference between the two groups (Figure 2B) (P = 0.04). Additionally, the results showed that hemorrhage in the clonidine group decreased by 34.5% (Figure 2C) (P = 0.05). According to

Figure 2D, there was no significant difference between the groups in the amount of blood in the drains on the first and second days after the surgery (P = 0.80 and P = 0.73).

4. Discussion

Using alpha-2 agonist drugs, such as clonidine and dexmedetomidine, is one of the methods to prevent hemodynamic instability and reduce intraoperative hemorrhage. Alpha-2 agonist drugs are sympatholytic and affect central receptors to reduce peripheral norepinephrine release (5). Only a few studies have been conducted on the effect of alpha-2 agonist drugs, especially clonidine, on hemodynamic response and intraoperative hemorrhage.

In this study, we investigated the effect of oral clonidine on hemorrhage in patients undergoing fusion surgery using the data of 100 patients undergoing

fusion surgery. The effectiveness of clonidine was evidenced by a significant reduction in operation time and the amount of hemorrhage. This reduction in hemorrhage can be explained by the pharmacological effect of the drug. Clonidine is a selective alpha-2 adrenergic receptor agonist and when consumed orally, it is rapidly absorbed and reaches its maximum level in 60 to 90 minutes, and its half-life is between 9 and 12 hours. This drug causes vasoconstriction by acting on alpha-2 receptors through its sympatholytic effect. It also has receptors in the brain stem, which causes a decrease in blood pressure and bradycardia (14).

The results showed that in the clonidine group, the mean SBP, DBP, mean arterial pressure (MAP), and HR significantly decreased after induction of anesthesia, after intubation, at seven different time points from 15 minutes after intubation to 90 min after the operation, and also in the recovery. Also, total blood loss was reduced in the clonidine group. The amount of blood and fluid collected from the drains was less in the patients who received clonidine, and as a result, the amount of blood transfusion in the patients in the clonidine group was lower than in the control group.

Regarding previous studies, clonidine reduced intraoperative hemodynamic stability more than dexmedetomidine (18). Pandazi et al. (19) studied low-dose clonidine premedication and hypotension after carotid artery surgery and concluded that intravenous premedication with low-dose clonidine (1 μ g/kg) is effective in preventing hypertensive attacks during carotid endarterectomy under general anesthesia. However, clonidine seemed to increase the incidence of postoperative hypotension possibility. Similar results were also reported by other researchers (20). They declared that clonidine as a selective alpha-2 agonist, is a central sympatholytic, which can significantly cause hemodynamic stability and dose-dependently reduce HR and blood pressure.

The effectiveness of clonidine in reducing hemorrhage and blood transfusion reduced doubts and concerns about blood transfusion complications. No complications or adverse outcomes were observed in our study. In this study, SBP, DBP, ABP, and HR were lower in the clonidine group, so the amount of remifentanil given to control BP and HR was significantly lower in the clonidine group. This finding is consistent with that of Okuyama et al. (21), who observed that clonidine has a favorable effect on hemodynamic stability and analgesia. Also, the duration of surgery was significantly shorter in the clonidine group, which may be due to the favorable hemorrhage control leading to the improved field of vision. In addition to estimating the amount of blood loss, the satisfaction of the surgical team was also assessed (22, 23). The surgeon was satisfied with the bloodless operative field in 42 cases (84%) in the clonidine group compared to 16 cases (32%) in the control group, and this difference was statistically significant (P = 0.05).

4.1. Conclusions

In this study, the effect of using clonidine as a preanesthetic medicine on patients who underwent lumbar laminectomy with pedicular screws was studied, and the most remarkable findings are listed below:

- The preventive use of oral clonidine before surgery was effective in reducing blood loss during and after spine surgery.

- Clonidine reduced the amount of blood transfusion to patients and its related complications.

- Surgeon satisfaction was significantly higher in the intervention group compared to the control group.

- SBP, DBP, ABP, and mean HR were significantly reduced, indicating higher hemodynamic stability during surgery and recovery.

Footnotes

Authors' Contribution: Seyedeh Sorour Zahedi: Data curation, writing original draft, visualization, investigation; Bahman Naghipour: Conceptualization, methodology and supervision; Eisa Bilehjani and Farhad Mirzaei: Conceptualization and methodology; Sepideh Sadat Zahedi and Sahar Zahedi: Writing and review.

Clinical Trial Registration: IRCT20140107016117N4.

Conflict of Interests: The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Data Reproducibility: The dataset presented in the study is available on request from the corresponding author during submission or after publication.

Ethical Approval: This study was approved by the local ethics committee of Tabriz University of Medical Sciences (IR.TBZMED.REC.1398.936).

Funding/Support: The authors state that they had not received any financial support.

Informed Consent: Written and informed consent was obtained from all patients and their information remained confidential.

References

- Ferraris VA, Ferraris SP, Saha SP, Hessel EA, Haan CK; Society of Thoracic Surgeons Blood Conservation Guideline Task Force, et al. Perioperative blood transfusion and blood conservation in cardiac surgery: the Society of Thoracic Surgeons and The Society of Cardiovascular Anesthesiologists clinical practice guideline. Ann Thorac Surg. 2007;83(5 Suppl):S27-86. [PubMed ID: 17462454]. https://doi.org/10.1016/j.athoracsur.2007.02.099.
- Koh MB, Hunt BJ. The management of perioperative bleeding. *Blood Rev.* 2003;17(3):179-85. [PubMed ID: 12818228]. https://doi.org/10.1016/s0268-960x(02)00062-0.
- Chand NK, Subramanya HB, Rao GV. Management of patients who refuse blood transfusion. *Indian J Anaesth.* 2014;**58**(5):658-64. [PubMed ID: 25535432]. [PubMed Central ID: PMC4260316]. https://doi.org/10.4103/0019-5049.144680.
- Elmekaty M, Kotani Y, Mehy EE, Robinson Y, Tantawy AE, Sekiguchi I, et al. Clinical and Radiological Comparison between Three Different Minimally Invasive Surgical Fusion Techniques for Single-Level Lumbar Isthmic and Degenerative Spondylolisthesis: Minimally Invasive Surgical Posterolateral Fusion versus Minimally Invasive Surgical Transforaminal Lumbar Interbody Fusion versus Midline Lumbar Fusion. Asian Spine J. 2018;12(5):870-9. [PubMed ID: 30213170]. [PubMed Central ID: PMC6147873]. https://doi.org/10.31616/asj.2018.12.5.870.
- Winter SF, Santaguida C, Wong J, Fehlings MG. ystemic and topical use of tranexamic acid in spinal surgery: a systematic review. *Global spine J.* 2016;6(3):284-95. [PubMed ID: 27099820]. [PubMed Central ID: PMC4836933]. https://doi.org/10.1055/s-0035-1563609.
- Tate DE, Friedman RJ. Blood conservation in spinal surgery. Review of current techniques. Spine. 1992;17(12):1450-6. [PubMed ID: 1471002]. https://doi.org/10.1097/00007632-199212000-00002.
- Hofmann A, Ozawa S, Farrugia A, Farmer SL, Shander A. Economic considerations on transfusion medicine and patient blood management. *Best Pract Res Clin Anaesthesiol.* 2013;27(1):59-68. [PubMed ID: 23590916]. https://doi.org/10.1016/j.bpa.2013.02.001.
- Willner D, Spennati V, Stohl S, Tosti G, Aloisio S, Bilotta F. Spine Surgery and Blood Loss: Systematic Review of Clinical Evidence. *Anesth Analg.* 2016;**123**(5):1307-15. [PubMed ID: 27749350]. https://doi.org/10.1213/ANE.000000000001485.
- Ebneshahidi A, Mohseni M. Premedication with oral clonidine decreases intraoperative bleeding and provides hemodynamic stability in cesarean section. *Anesth Pain Med.* 2011;1(1):30-3. [PubMed ID: 25729653]. [PubMed Central ID: PMC4335754]. https://doi.org/10.5812/kowsar.22287523.1337.
- Wong J, El Beheiry H, Rampersaud YR, Lewis S, Ahn H, De Silva Y, et al. Tranexamic Acid reduces perioperative blood loss in adult patients having spinal fusion surgery. *Anesth Analg.* 2008;**107**(5):1479-86. [PubMed ID: 18931202]. https://doi.org/10.1213/ane.0b013e3181831e44.
- Camarasa MA, Olle G, Serra-Prat M, Martin A, Sanchez M, Ricos P, et al. Efficacy of aminocaproic, tranexamic acids in the control of bleeding during total knee replacement: a randomized clinical trial. Br J

Anaesth. 2006;**96**(5):576-82. [PubMed ID: 16531440]. https://doi.org/10.1093/bja/ael057.

- Thompson GH, Florentino-Pineda I, Poe-Kochert C. The role of amicar in decreasing perioperative blood loss in idiopathic scoliosis. *Spine*. 2005;**30**(17 Suppl):S94-9. [PubMed ID: 16138073]. https://doi.org/10.1097/01.brs.0000175188.05542.a9.
- Singh A, Ruchi G, Tripat B, Aujla KS, Anuradha A, Vanita S. A Study to Evaluate the Effectiveness of Clonidine vs Atenolol in Providing Optimal Surgical Field in Nasal Surgeries under General Anaesthesia. *J Anesth Clin Res.* 2014;5(3):1-4. https://doi.org/10.4172/2155-6148.1000396.
- Bajaj J, Mittal RS, Sharma A. Preoperative clonidine use in transsphenoidal pituitary adenoma surgeries - a randomized controlled trial. Br J Neurosurg. 2017;31(1):2-4. [PubMed ID: 27535352]. https://doi.org/10.1080/02688697.2016.1220501.
- 15. Deepa C, Shekhar R. Effect of Premedication with Intravenous Clonidine in Modulating the Haemodyanamic Responses during Laparoscopic Surgeries. *Indian J Anaesth Analg.* 2018:545.
- de Padua AI, de Castro M, Schmidt A, Coutinho Neto J, Terra Filho J, Martinez JA. Clonidine as a pre-anesthetic agent for flexible bronchoscopy. *Respir Med.* 2004;**98**(8):746-51. [PubMed ID: 15303639]. https://doi.org/10.1016/j.rmed.2004.01.015.
- Taghipour Anvari Z, Afshar-Fereydouniyan N, Imani F, Sakhaei M, Alijani B, Mohseni M. Effect of clonidine premedication on blood loss in spine surgery. *Anesth Pain Med.* 2012;1(4):252-6. [PubMed ID: 24904810]. [PubMed Central ID: PMC4018704]. https://doi.org/10.5812/aapm.2197.
- Bharti D, Saran J, Kumar C, Nanda HS. Comparison of clonidine and dexmedetomidine on cardiovascular stability in laparoscopic cholecystectomy. *Int J Sci Study*. 2016;4(1):43-50.
- Pandazi A, Karamanis P, Sidiropoulou T, Matsota P, Papasideris C, Niokou D, et al. Low-dose (1 microg/kg) clonidine premedication and hypotension after carotid artery surgery. *Vasc Endovascular Surg.* 2011;**45**(7):614-8. [PubMed ID: 21984028]. https://doi.org/10.1177/1538574411414300.
- Lee J, Lovell AT, Parry MG, Glaisyer HR, Bromley LM. I.v. clonidine: does it work as a hypotensive agent with inhalation anaesthesia? *Br J Anaesth.* 1999;**82**(4):639-40. [PubMed ID: 10472237]. https://doi.org/10.1093/bja/82.4.639.
- 21. Okuyama K, Inomata S, Toyooka H. The effects of prostaglandin E1 or oral clonidine premedication on blood loss during paranasal sinus surgery. *Can J Anaesth*. 2005;**52**(5):546-7. [PubMed ID: 15872138]. https://doi.org/10.1007/BF03016539.
- 22. Moghadam MJ, Ommi D, Mirkheshti A, Shadnoush M, Dabbagh A. The effect of pretreatment with clonidine on propofol consumption in opium abuser and non-abuser patients undergoing elective leg surgery. J Res Med Sci: Offic J Isfahan Univ Med Sci. 2012;17(8):728.
- 23. Jabbary Moghaddam M, Ommi D, Mirkheshti A, Dabbagh A, Memary E, Sadeghi A, et al. Effects of clonidine premedication upon postoperative shivering and recovery time in patients with and without opium addiction after elective leg fracture surgeries. *Anesth Pain Med.* 2013;**2**(3):107-10. [PubMed ID: 24244918]. [PubMed Central ID: PMC3821124]. https://doi.org/10.5812/aapm.7143.