












Maternal Education is a Major Factor in Growth Regulation in Twins and Singletons

Novina Novina ^{1,2,*}, Binu Dorjee ³, Michael Hermanussen ⁴, Christiane Scheffler ⁵, Barry Bogin ⁶, Aman Bhakti Pulungan ⁷, Madarina Julia ⁸, Yoyos Dias Ismiarto ⁹, Budi Setiabudiawan ²

¹ Doctoral Programme, Universitas Padjadjaran, Bandung, West Java, Indonesia

² Department of Child Health, Faculty of Medicine, Universitas Padjadjaran/Dr. Hasan Sadikin General Hospital, Bandung, West Java, Indonesia

³ Department of Anthropology, University of North Bengal, Raja Rammohunpur, Darjeeling, India

⁴ Christian-Albrecht University of Kiel, Aschauhof, Eckernförde-Altenhof, Germany

⁵ University of Potsdam, Human Biology, Germany

⁶ School of Sport, Exercise and Health Sciences, Loughborough University, United Kingdom

⁷ Department of Child Health, Faculty of Medicine, Universitas Indonesia/ Dr. Ciptomangunkusumo General Hospital Jakarta, Indonesia

⁸ Department of Child Health, Faculty of Medicine, Public Health and Nursing, Universitas Gadjah Mada, Yogyakarta, Indonesia

⁹ Department of Orthopaedy and Traumatology Faculty of Medicine, Universitas Padjadjaran/Dr. Hasan Sadikin General Hospital, Bandung, West Java, Indonesia

* Corresponding author: Doctoral Programme, Universitas Padjadjaran, Bandung, West Java, Indonesia. Email: novina@unpad.ac.id

Received 2024 January 20; Revised 2024 March 21; Accepted 2024 March 26.

Abstract

Background: The social, economic, political, and emotional (SEPE) environment plays a crucial role in shaping human development both pre- and post-birth, with direct and independent effects on growth, as research shows.

Objectives: This study examines the growth of twins and singletons and their association with maternal education as one of the social-economic (SE) variables.

Methods: This cross-sectional study utilized Nutritional Status Monitoring for Children under 5 Years Old in Bandung District in September 2019. A total of 158 healthy children aged < 5 years, involving 35 twins (70 children) and 88 singletons, were included. Data on weight and length at birth and weight and length/height taken at age 8.6 to 60 months were plotted according to WHOCS 2006. Statistical analyses and visualizations, including SE variables analysis, were conducted using density plots, correlation plots, St. Nicolas House Analysis (SNHA), and ANOVA.

Results: At birth, z-scores of the body length, weight, and body mass index (BMI) of twins were lower than those of singletons ($P < 0.05$). After birth, z-scores of twins' length/height, weight, and BMI adjusted to those of singletons, with maternal education being the strongest among SE variables of early childhood growth adjustment.

Conclusions: Twins were shorter, lighter, and had a lower average BMI at birth than singletons. There was no apparent association between birth measurements and SE variables. However, after birth, maternal education starts to gain a central position in regulating child growth.

Keywords: Child, Growth, Educational Status, Maternal, St. Nicolas House Analysis, Twins

1. Background

At birth, twins are smaller and lighter than singletons. However, the differences in length and weight often disappear as early as one year of life (1). Some studies have observed differences in head circumference, weight, and length up to 2.5 years of age (2). Another study showed growth differences up to four

years of age, except for twins born small-for-gestational-age (3). Twins attain normal adult height but are slightly shorter than their singleton siblings (2, 4). Growth differences at birth are attributed to the average shorter length of twin pregnancies and the intrauterine environment. After birth, twins catch up in body size (2).

The social-economic-political-emotional (SEPE) environment provides important material and moral

conditioning for human growth before and after birth. This concept is broader than the traditional view, as it embraces political and emotional environments (5). Social-economic-political-emotional factors directly and independently influence human growth (6, 7). Education, earnings, occupation, and the social status of parents and grandparents are well-known SEPE determinants of children's growth and health (8, 9). Mainly, maternal education as one of the SEPE factors is important for children's growth and health (10-12). However, studies on the association between maternal education and the growth of twins are rare.

Conducting research on the growth of twins in Indonesia while considering social-economic, political, and emotional factors is paramount for understanding the complex interplay of influences shaping their development. By examining how variables such as parental education, family income, access to healthcare, number of siblings, and others impact the growth trajectories of twins compared to singletons, this research can uncover disparities and inform targeted interventions to mitigate them. Such research not only enriches our understanding of twin development but also underscores the importance of holistic support systems to ensure the optimal growth and flourishing of twin children in Indonesia. We hypothesized that:

(1) After birth, twins' growth catches up, and singletons' growth remains constant.

(2) Mothers' educational level is associated with anthropometric indices at birth of twins (birth weight and birth length).

(3) Mothers' educational level is associated with the early postnatal growth of twins.

2. Objectives

This study examines the growth of twins and singletons and their association with maternal education as one of the social-economic (SE) variables.

3. Methods

The study was conducted among children from 31 Sub-Districts of Bandung District, Indonesia, including urban and rural areas, who are required to participate in the Nutritional Status Monitoring for Children under 5 Years Old program at the Bandung District Health Office. This monitoring aims to collect information and monitor follow-up actions for the growth and development of healthy children in Bandung District. Screening/monitoring of infant growth and development is recommended to be conducted monthly. For children aged 12 to 24 months, it is

recommended every 3 months, and for children aged 24 months to 72 months, it is recommended every 6 months. Demographic data of parents and place of residence are also collected. Bandung District has an area of 1 762.40 km² and comprises 31 sub-districts with a population of 3 575 982. The sex ratio of the total population is 102.4 males to 100 females. Most women were employed as laborers/employees (42.5%) or family workers (18.9%) (13).

The inclusion criteria for this study stipulated the utilization of all secondary data sources, provided that they were complete. Only datasets with comprehensive information were considered eligible for analysis. The exclusion criteria for this study involved excluding incomplete datasets and outlier data points. Any secondary data that lacked comprehensive information or contained outliers were omitted from the analysis.

Initially, 178 twins and siblings of twins were included. In the final analysis, the number was reduced to 158 children after excluding outliers and incomplete data and consisted of 35 twin pairs and 88 singletons (Figure 1). Their ages ranged from 8.6 months to 60 months. From the singletons group, 33 were female, and 55 were male. From the twins' group, 33 were females, and 37 were males.




Figure 1. Flow Chart of Study Subjects

Social-economic (SE) variables were selected based on data availability, including anthropometric measurements such as length/height and weight during the newborn period and later ages up to 60 months.

These variables are contained within the Nutritional Status Monitoring for Children under 5 Years Old at the Bandung District Health Office. Measurements were performed by healthcare providers using standardized tools during routine visits to primary healthcare in Bandung District. The measurement technique and nutritional status evaluation were performed according to Child Anthropometric Standards in Indonesia (14).

Weight measurements were conducted with the Indonesian Dacin scale, a widely used tool in primary maternal and child health care for children, providing accuracy to the nearest 100 grams. For children aged two years or younger, length was measured using an infantometer in a supine position. Height measurements were taken with a microtoise or stadiometer in children older than two years, ensuring precision to the nearest 1.0 millimeter. Sanes Sumber Makmur manufactured the Dacin scale, and GEA Medical produced both the infantometer and microtoise.

The z-scores of length at birth (LAZbirth), weight at birth (WAZbirth), height-for-age (HAZ), weight-for-age (WAZ), and BMI-for-age (BAZ) at ages between 8 and 60 months were calculated using the WHO Anthro plus software (15).

Statistical analyses were performed using R Gui (<https://cran.r-project.org>). Pearson's correlation was employed to examine associations between variables, St. Nicolas' House Analysis (SNHA) with a threshold value of $r > 0.3$, and one-way analysis of variance (ANOVA). The SNHA technique is a novel non-parametric statistical method that helps translate correlation matrices into network graphs by tracing "association chains." A series of coefficients of determinants characterized by the symmetry of ranks of R^2 both in the forward and backward direction is named "association chains." Thus, the association chains formed are ranked according to the magnitude of the correlation coefficients (R^2), for example, $c[A*B]$, $c[B*C]$, $c[C*D]$, with the property $c[A*B] > c[A*C] > c[A*D]$, and $c[D*C] > c[D*B] > c[D*A]$. Performance metrics, including balanced classification rate and F1-score, indicated that SNHA outperformed approaches employing intricate correlation value thresholds and those relying on partial correlations when analyzing bands and hubs (16). This method is well-suited for managing the correlations commonly encountered in anthropometric, SE, and sociodemographic variables (16).

This innovative method converts the correlation matrix into a network graph, serving as a valuable visual tool and aiding in data exploration. Researchers currently utilize SNHA to document associations among

diverse growth, SE, and sociodemographic variables, representing them as a chain (17, 18). Principally, St. Nicolas's house forms a network comprising "nodes" and "edges" (16). Corplot is usually presented along with SNHA graphs since they complement each other for better visualization. In addition, density plots and ANOVA were used to assess and visualize the associations and distribution of growth parameters.

The investigated variables are as follows: (1) Mother's age at birth (MaB) refers to the age of the mother at the time when she gave birth to the subject of this research; (2) father's occupation (FaOccu) refers to the profession of the father at the time when the subject of this research was born; (3) father's education (FaEdu) refers to the level of education attained by the father at the time when the subject of this research was born; (4) mother's occupation (MOccu) refers to the profession of the mother at the time when the subject of this research was born; (5) mother's education (MEdu) refers to the level of education attained by the mother at the time when the subject of this research was born; (6) family income refers to the total earnings or financial resources acquired by a household within a given month.

4. Results

Social-economic variables investigated in this research are included in Table 1. The education levels of fathers and mothers were significantly different among SE variables in twins and singletons ($P < 0.05$).

Table 1. Classifications of Socio-economic (SE) Variables of Twins and Singletons^a

Variables and Category	No. (%)		P-Value ^a
	Singletons	Twin	
Father age, mean (SD)	36.13 (6.5)	35.43 (7.5)	0.482
Father's education (FaEdu)			0.006
Elementary (1 - 6 y)	21 (23.9)	34 (37)	
Junior high (7 - 9 y)	24 (27.3)	8 (8.7)	
Senior high (10 - 12 y)	30 (34.1)	20 (21.7)	
College (13 - 16 y)	13 (14.8)	8 (8.7)	
Father's occupation (FaOccu)			0.87
Entrepreneurs	34 (38.6)	27 (38.6)	
Labors	54 (61.4)	43 (61.4)	
Mother age (MoAge), mean (SD)	32.4 (6.0)	33.06 (7.5)	0.497
Mother's education (MoEdu)			0.006
Elementary (1 - 6 y)	18 (20.5)	31 (44.3)	
Junior high (7 - 9 y)	29 (33.0)	16 (22.9)	
Senior high (10-12 yrs.)	27 (30.7)	19 (20.7)	
College (13 - 16 y)	14 (15.2)	4 (5.7)	
Mother's occupation (MoOccu)			0.31
Entrepreneurs	12 (13.6)	9 (12.9)	
Labors	7 (8.0)	11 (15.7)	

Not working	69 (78.4)	50 (71.4)	
Family's monthly income (Income)^b			0.151
Low	23 (26.1)	22 (31.4)	
Middle	53 (60.2)	32 (45.7)	
High	12 (13.6)	16 (22.9)	
Type of health insurance (HIn)^c			0.32
Premium	24 (27.3)	11 (15.7)	
Non-premium	56 (63.6)	50 (73.4)	
None	8 (9.1)	9 (12.9)	
Number of siblings (Sibs)			0.359
1	9 (9.9)	12 (17.1)	
2	74 (81.3)	50 (71.4)	
3	8 (8.8)	8 (11.4)	

^a Chi-square test, the value of < 0.05 is set as the level of significance.

^b Low if below Rp. 1.000.000, intermediate if it is between Rp.1.000.000-Rp. 3.000.000, High if more than Rp. 3.000.000. (classification is slightly adjusted based on the Indonesian Central Bureau of Statistics) (19).

^c Premium if BPJS (Indonesia Social Security Agency of Health), non-premium if JKN-KIS, and SKTM (complimentary Indonesian National Health Insurance provided explicitly for individuals with low income) (20).

At birth, twins were shorter, lighter, and had a lower BMI than singletons (Table 2). After birth, twins' height, weight, and BMI z-scores adjusted to those of singletons, with maternal education being the strongest predictor of early childhood growth adjustment. In this process, the height and weight z-scores of singletons can be observed to decline, while twins' Z-scores remain relatively stable. The z-scores of length at birth, WAZbirth, HAZ, WAZ, and BAZ at ages between 8 and 60 months are presented in Figure 2. These differences became smaller at later ages. This was also supported by ANOVA results (Table 2).

Table 2. Age and Gender of Twins and Singletons, Age of Mother at Birth, and Anthropometric Measurements^a

Variables	Singleton; N = 88	Twin 1; N = 35	Twin 2; N = 35	P-Value ^b
Children age (mo)	37.22 ± 14.48	31.94 ± 13.64	31.94 ± 13.64	0.07
Gender				
Boy	55 (62.5)	14 (40)	23 (65.7)	0.22
Girl	33 (37.5)	21 (60)	12 (34.3)	
MaB (y)	29.27 ± 6.05	29.91 ± 7.04	29.91 ± 7.04	0.83
WAZbirth	-0.85 ± 1.06	-2.21 ± 1.30	-2.52 ± 1.23	< 0.001
BAZbirth	-0.87 ± 1.44	-1.83 ± 1.74	-2.43 ± 1.68	< 0.001
LAZbirth	-0.6 ± 1.16	-2.12 ± 1.75	-2.08 ± 1.49	< 0.001
WAZ	-1.59 ± 1.10	-1.67 ± 1.0	-1.68 ± 1.12	0.9
BAZ	-0.45 ± 1.15	-0.44 ± 1.21	-0.47 ± 1.47	0.99
HAZ	-1.94 ± 1.35	-2.17 ± 1.23	-2.12 ± 1.27	0.62

Abbreviations: MaB, mother age at birth; LAZbirth, length-for-age z-scores at birth; BAZbirth, BMI-for-age z-scores at birth; WAZbirth, weight-for-age z-scores at birth; HAZ, Growth after birth is represented by height-for-age z-scores; WAZ, weight-for-age z-scores; BAZ, BMI-for-age z-scores.

birth; HAZ, Growth after birth is represented by height-for-age z-scores; WAZ, weight-for-age z-scores; BAZ, BMI-for-age z-scores.

^a Values are expressed as mean ± SD or No. (%).

^b ANOVA test, the value of 0.05 is set as level of significance.




Figure 2. Density distributions. A, Weight-for-age z-scores of twins and singletons at birth (WAZbirth) and age 8.6 - 60 months (WAZ); B, BMI-for-age z-scores of twins and singletons at birth (BAZbirth) and age 8.6 - 60 months (BAZ); C, Length-for-age z-scores of twins and singletons at birth (LAZbirth) and age 8.6 - 60 months (HAZ).

Spearman correlations are presented as complots, and associations of variables are further elucidated using SNHA, a network graph. Figure 3A presents associations of SE variables such as the mother's age at birth (MaB), father's occupation (FaOccu), father's education (FaEdu), mother's education (MEdu), mother's occupation (MOccu), family's monthly income (Income), type of health insurance (Ins), number of siblings (Sibs), and twin birth (STT) with birth measurements LAZbirth and WAZbirth using complot and SNHA. There were no reliable associations ($r > 0.3$) except for twin birth (STT). The association chain, SNHA, indicates that twin birth is the only variable linked with birth measurements. The other nodes either formed separate networks or remained disconnected. Among the factors investigated, only twin birth was related to birth measurements. Statistically significant associations between STT and LAZbirth ($F = 22.898$, df :

2.0, $P < 0.05$) and BAZbirth ($F = 13.858$, $df: 2.0$, $P < 0.05$) were confirmed by ANOVA.




Figure 3. Correlation plot on the left and the St. Nicolas House Analysis (SNHA) graph on the right show the association of socio-economic (SE) variables with body measurements at birth (A) and age 8.6 - 60 months (B).

Association analysis of SE variables with growth at age 8.6 to 60 months showed that maternal education (Medu) was the only variable prominently correlated with the linear growth of the children (Figure 3B). Another reliable correlation was observed between BAZ and maternal age at birth (MaB). Compared to MaB, Medu was reliably associated with many other SE variables. The associations are depicted in SNHA. Association chains are formed between several variables, with maternal education being centrally located and directly connected to children's linear growth (HAZ). The plot highlights the pivotal position of the mother's education. A statistically significant association between the mother's education and HAZ ($F = 3.15$, $df = 3.0$, $P < 0.05$) was confirmed by ANOVA.

5. Discussion

The present study revealed that twins were shorter, lighter, and had a lower BMI at birth than singletons. These differences disappear later in infancy and childhood. Many studies have reported such observations from different countries (1-3) and were supported by this present study from Bandung District,

Indonesia. Twins usually reach an average final height and BMI as adults (4). In contrast to the general perception that twins tend to catch up in height, weight, and BMI during early infancy, Bandung District's data showed that this catch-up was only relative. The findings of this study reject the first hypothesis. Similar to what has been observed in many low- and middle-income countries, Z-scores for height and weight tend to decline during the first two years of life (21). In this study, singleton babies showed a decline in height and weight Z-scores. However, twins do not participate in this decline and maintain their birth Z-scores in height, weight, and BMI. This timing of growth faltering, which is high in South Asian countries and often referred to as a critical window for nutritional intervention, may not be due to nutrition, as twin growth remains relatively stable (22).

Our findings also reject the 2nd hypothesis and accept the 3rd hypothesis about the association between maternal education and birth measurements and growth after that. In this study, twin pregnancy had a direct effect on fetal growth. After birth, maternal education was directly associated with children's growth. Studies have reported that maternal education is an important predictor of children's linear growth (16, 23-25). Postnatal growth rates up to 2 years vary with maternal educational status (26). Offspring of educated mothers exhibited increased birth length and experienced accelerated growth from birth to three months and up to 12 - 34 months (27).

Moreover, an earlier study reported similar observations regarding the physical growth of children in Indonesia (16). Mothers' education in Indonesia is linked to their children's health through the role of social capital. Mothers who have completed primary education or lower have less social capital than those with a higher level of education. The mechanisms are explained in terms of social networking and transfer of knowledge. Mothers' social capital is positively and significantly associated with children's health via improvement in mothers' knowledge, which affects their parenting behavior (28).

In India, a woman's decision-making authority within the family is considered a significant predictor of early childhood growth (29). Maternal autonomy related to visiting the market and financial access is associated with the linear growth of children (30). Other studies emphasize that literate mothers have better access to healthcare infrastructure than illiterate mothers. The decision-making power in the family, maternal autonomy, and access to health infrastructure could be reasons for the better growth of children of

educated mothers. Maternal education is also known to impact prenatal care and ultimately reduce the incidence of low birth weight (31). Other social advantages conferred by better education are less counter to prejudices, discrimination, and violence during the early formative years of offspring. Studies indicate that emotional adversities faced by mothers during pregnancy and the neonatal period can impact the health and growth of children and may last until adulthood (32).

Regarding social status, findings indicate that within a traditional horticultural community, mothers with education and proficiency in two additional languages, including English alongside their local language, tend to have children exhibiting improved linear growth (33). Educated women are more likely to find partners with high status based on earnings, occupation, or any other traditional marker of social status, such as caste or ethnic group. Highly educated women are more likely to characterize a high-status society or a group within a country based on traditional social status markers. Thus, social status in different forms is self-perpetuating. This reflects the idea of strategic growth adjustment, which occurs when social dominance stimulates growth among peers. In humans, this phenomenon was studied among German boarding school boys educated in the Grand Ducal Carlsschule in Stuttgart. The schoolboys were aware of their social position, which resulted in their final height variation (34, 35).

5.1. Limitations

The present study indicates that maternal education plays a central role in the regulation of child growth among various SE variables, such as the mother's age at birth, father's education, father's occupation, mother's education, mother's occupation, family monthly income, type of health insurance, and the number of siblings. However, these were not a complete set of SEPE factors. This study used secondary data from Nutritional Status Monitoring for Children under 5 Years Old in Bandung District. There was no information on the political and emotional environments of the population under study, nor information regarding the IVF status of the twins, parental anthropometrics, or mothers' weight gain during pregnancy.

5.2. Conclusions

Twins displayed lesser height, weight, and a lower mean BMI at birth compared to singletons. Initially, there seemed to be no clear link between

anthropometric measures and SE factors in twins, including parental education and occupation, family income, health insurance type, and the number of siblings. However, this changed postnatally, as maternal education emerged as a significant SE factor influencing children's growth.

These findings suggest potential implications for healthcare providers and policymakers, highlighting the importance of considering maternal education in interventions aimed at improving early childhood growth among twins. Furthermore, future research could delve deeper into understanding the mechanisms through which maternal education influences growth outcomes in twins and explore additional social economic factors that may impact their development. Additionally, investigating the long-term effects of early growth disparities between twins and singletons on health and well-being outcomes could provide valuable insights for public health initiatives and clinical practice.

Footnotes

Authors' Contribution: Study concept and design: NN, ABP, YDI, BS, and BD; acquisition of data: NN, ABP, YDI, BS; analysis and interpretation of data: NN, BD; drafting of the manuscript: NN, BD, MH, CS, BB, ABP, MJ, YDI, BS; critical revision of the manuscript for important intellectual content: NN, BD, MH, CS, BB, ABP, MJ, YDI, BS; statistical analysis: NN, BD, MH, CS, ABP, YDI, BS; administrative, technical, and material support: NN, YDI, BS; Study supervision: ABP, YDI, BS; all authors read and approved the final manuscript.

Conflict of Interests: The authors affirm the absence of any conflicting interests. The funding sources did not play a role in the study's design, data collection, analysis, interpretation, manuscript writing, or the decision to publish the findings.

Data Availability: The data supporting the conclusions in the study are included in the article, and further inquiries can be directed to the corresponding author.

Ethical Approval: This study was approved by the Ethics Committee of the Faculty of Medicine, Universitas Padjadjaran (Ethical Approval number 1170/UN6.KEP/EC/2019) for studies involving humans and conformed to the ethical guidelines of the Declaration of Helsinki.

Funding/Support: The publication APC (Article Processing Charge) was covered by Universitas Padjadjaran's Research Grant: Riset Disertasi Dosen

Unpad (RDDU), <https://drpm.unpad.ac.id/research-support/bantuan-article-processing-charge-apc/>.

Informed Consent: Written consent for participating in the Nutritional Status Monitoring was obtained from the parents according to the Bandung District Health Office policy.

References

- Mazkereth R, Miron E, Leibovitch L, Kuint J, Strauss T, Maayan-Metzger A. Growth parameters of discordant preterm twins during the first year of life. *J Matern Fetal Neonatal Med*. 2014;27(17):1795-9. [PubMed ID: 24397374]. <https://doi.org/10.3109/14767058.2014.880688>.
- Van Dommelen P, De Gunst M, Van Der Vaart A, Van Buuren S, Boomsma D. Growth references for height, weight and body mass index of twins aged 0-2.5 years. *Acta Paediatr*. 2008;97(8):1099-104. [PubMed ID: 18460042]. <https://doi.org/10.1111/j.1651-2227.2008.00853.x>.
- Chaudhari S, Bhalerao MR, Vaidya U, Pandit A, Nene U. Growth and development of twins compared with singletons at ages one and four years. *Indian pediatrics*. 1997;34:1081-6.
- Eriksen W, Sundet JM, Tambs K. Adult body height of twins compared with that of singletons: a register-based birth cohort study of Norwegian males. *Am J Epidemiol*. 2013;177(9):1015-9. [PubMed ID: 23543161]. <https://doi.org/10.1093/aje/kws341>.
- Bogin B. Social-Economic-Political-Emotional (SEPE) factors regulate human growth. *Human Biology and Public Health*. 2021;1.
- Goudet S, Griffiths P, Bogin B, Madise N. Interventions to tackle malnutrition and its risk factors in children living in slums: a scoping review. *Ann Hum Biol*. 2017;44(1):1-10. [PubMed ID: 27356853]. <https://doi.org/10.1080/03014460.2016.1205660>.
- Hermanussen M, Bilogub M, Lindl AC, Harper D, Mansukoski L, Scheffler C. Weight and height growth of malnourished school-age children during re-feeding. Three historic studies published shortly after World War I. *Eur J Clin Nutr*. 2018;72(12):1603-19. [PubMed ID: 30166640]. <https://doi.org/10.1038/s41430-018-0274-z>.
- Koziel S, Zareba M, Bielicki T, Scheffler C, Hermanussen M. Social mobility of the father influences child growth: A three-generation study. *Am J Hum Biol*. 2019;31(4). e23270. [PubMed ID: 31190434]. <https://doi.org/10.1002/ajhb.23270>.
- Li Z, Kim R, Vollmer S, Subramanian SV. Factors Associated With Child Stunting, Wasting, and Underweight in 35 Low- and Middle-Income Countries. *JAMA Netw Open*. 2020;3(4). e203386. [PubMed ID: 32320037]. [PubMed Central ID: PMC7177203]. <https://doi.org/10.1001/jamanetworkopen.2020.3386>.
- Elverud IS, Stordal K, Chiduo M, Klingenberg C. Factors Influencing Growth of Children Aged 12-24 Months in the Tanga Region, Tanzania. *J Trop Pediatr*. 2020;66(2):210-7. [PubMed ID: 31504993]. <https://doi.org/10.1093/tropej/fmz056>.
- Jaros E, Gugushvili A. Parental education, health literacy and children's adult body height. *J Biosoc Sci*. 2020;52(5):696-718. [PubMed ID: 31722763]. <https://doi.org/10.1017/S0021932019000737>.
- Oyekale AS, Maselwa TC. Maternal Education, Fertility, and Child Survival in Comoros. *Int J Environ Res Public Health*. 2018;15(12). [PubMed ID: 30544762]. [PubMed Central ID: PMC6313670]. <https://doi.org/10.3390/ijerph15122814>.
- Kabupaten Bandung. *Badan Pusat Statistik Kabupaten Bandung*. Indonesia: Sosial dan Populasi; 2022. Available from: <https://bandungkab.bps.go.id/>.
- Menteri Kesehatan Republik Indonesia. *Menteri Kesehatan Republik Indonesia Nomor 2 Tahun 2020 tentang Standar Antropometri Anak*. Peraturan Menteri Kesehatan Republik Indonesia; 2020. Available from: http://hukor.kemkes.go.id/uploads/produk_hukum/PMK_No__2_Th_2020_ttg_Standar_Antropometri_Anak.pdf.
- World Health Organization. *Anthro Survey Analyser and other tools*. WHO; 2023. Available from: <https://www.who.int/tools/child-growth-standards/software>.
- Groth D, Scheffler C, Hermanussen M. Body height in stunted Indonesian children depends directly on parental education and not via a nutrition mediated pathway - Evidence from tracing association chains by St. Nicolas House Analysis. *Anthropol Anz*. 2019;76(5):445-51. [PubMed ID: 30990515]. <https://doi.org/10.1127/anthranz/2019/1027>.
- Dorjee B, Bogin B, Scheffler C, Groth D, Sen J, Hermanussen M. Association of anthropometric indices of nutritional status with growth in height among Limboo children of Sikkim, India. *Anthropol Anz*. 2020;77(5):389-98. [PubMed ID: 32405638]. <https://doi.org/10.1127/anthranz/2020/1174>.
- Martin L, Dorjee B, Groth D, Scheffler C. Positive influence of parental education on growth of children - statistical analysis of correlation between social and nutritional factors on children's height using the St. Nicolas House Analysis. *Anthropol Anz*. 2020;77(5):375-87. [PubMed ID: 32405637]. <https://doi.org/10.1127/anthranz/2020/1177>.
- Badan Pusat Statistik Kabupaten Bandung. *Upah minimum Kabupaten Bandung tahun 2018-2020*. Sosial dan Populasi; 2020, [cited 2023]. Available from: <https://bandungkab.bps.go.id/indicator/19/345/1/upah-minimum-kabupaten-bandung-tahun.html>.
- Badan Penyelenggara Jaminan Sosial (BPJS) Kesehatan. *Panduan layanan bagi peserta jaminan kesehatan nasional kartu Indonesia sehat (JKN-KIS)*. Sosial (BPJS) Kesehatan; 2023. Available from: <https://www.bpjs-kesehatan.go.id/bpjs/arsip/view/1477>.
- Prentice AM, Ward KA, Goldberg GR, Jarjou LM, Moore SE, Fulford AJ, et al. Critical windows for nutritional interventions against stunting. *Am J Clin Nutr*. 2013;97(5):911-8. [PubMed ID: 23553163]. [PubMed Central ID: PMC3628381]. <https://doi.org/10.3945/ajcn.112.052332>.
- Victora CG, de Onis M, Hallal PC, Blossner M, Shrimpton R. Worldwide timing of growth faltering: revisiting implications for interventions. *Pediatrics*. 2010;125(3):e473-80. [PubMed ID: 20156903]. <https://doi.org/10.1542/peds.2009-1519>.
- Scheffler C, Krutzfeldt LM, Dasgupta P, Hermanussen M. No association between fat tissue and height in 5019 children and adolescents, measured between 1982 and in 2011 in Kolkata/India. *Anthropol Anz*. 2018;74(5):403-11. [PubMed ID: 29543314]. <https://doi.org/10.1127/anthranz/2018/0827>.
- Devakumar D, Chaube SS, Wells JC, Saville NM, Ayres JG, Manandhar DS, et al. Effect of antenatal multiple micronutrient supplementation on anthropometry and blood pressure in mid-childhood in Nepal: follow-up of a double-blind randomised controlled trial. *Lancet Glob Health*. 2014;2(11):e654-63. [PubMed ID: 25442690]. [PubMed Central ID: PMC4224012]. [https://doi.org/10.1016/S2214-109X\(14\)70314-6](https://doi.org/10.1016/S2214-109X(14)70314-6).
- Abuya BA, Ciera J, Kimani-Murage E. Effect of mother's education on child's nutritional status in the slums of Nairobi. *BMC Pediatr*. 2012;12:80. [PubMed ID: 22721431]. [PubMed Central ID: PMC3444953]. <https://doi.org/10.1186/1471-2431-12-80>.
- Matijasevich A, Howe LD, Tilling K, Santos IS, Barros AJ, Lawlor DA. Maternal education inequalities in height growth rates in early childhood: 2004 Pelotas birth cohort study. *Paediatr Perinat Epidemiol*. 2012;26(3):236-49. [PubMed ID: 22471683]. [PubMed Central ID: PMC3491696]. <https://doi.org/10.1111/j.1365-3016.2011.01251.x>.

27. Patel R, Lawlor DA, Kramer MS, Smith GD, Bogdanovich N, Matush L, et al. Socio-economic position and adiposity among children and their parents in the Republic of Belarus. *Eur J Public Health*. 2011;**21**(2):158-65. [PubMed ID: 20418336]. [PubMed Central ID: PMC3451194]. <https://doi.org/10.1093/eurpub/ckq041>.
28. Sujarwoto S, Tampubolon G. Mother's social capital and child health in Indonesia. *Soc Sci Med*. 2013;**91**:1-9. [PubMed ID: 23849232]. <https://doi.org/10.1016/j.socscimed.2013.04.032>.
29. Das S, Chanani S, Shah More N, Osrin D, Pantvaidya S, Jayaraman A. Determinants of stunting among children under 2 years in urban informal settlements in Mumbai, India: evidence from a household census. *J Health Popul Nutr*. 2020;**39**(1):10. [PubMed ID: 33246506]. [PubMed Central ID: PMC7693500]. <https://doi.org/10.1186/s41043-020-00222-x>.
30. Shroff M, Griffiths P, Adair L, Suchindran C, Bentley M. Maternal autonomy is inversely related to child stunting in Andhra Pradesh, India. *Matern Child Nutr*. 2009;**5**(1):64-74. [PubMed ID: 19161545]. [PubMed Central ID: PMC3811039]. <https://doi.org/10.1111/j.1740-8709.2008.00161.x>.
31. Laksono AD, Wulandari RD, Ibad M, Kusriani I. The effects of mother's education on achieving exclusive breastfeeding in Indonesia. *BMC Public Health*. 2021;**21**(1):14. [PubMed ID: 33402139]. [PubMed Central ID: PMC7786474]. <https://doi.org/10.1186/s12889-020-10018-7>.
32. Bogin B, Varea C. COVID-19, crisis, and emotional stress: A biocultural perspective of their impact on growth and development for the next generation. *Am J Hum Biol*. 2020;**32**(5). e23474. [PubMed ID: 32672890]. [PubMed Central ID: PMC7404495]. <https://doi.org/10.1002/ajhb.23474>.
33. King SE, Nicholas Mascie-Taylor CG. Nutritional status of children from Papua New Guinea: associations with socioeconomic factors. *Am J Hum Biol*. 2002;**14**(5):659-68. [PubMed ID: 12203820]. <https://doi.org/10.1002/ajhb.10080>.
34. Hermanussen M, Bogin B, Scheffler C. The impact of social identity and social dominance on the regulation of human growth: A viewpoint. *Acta Paediatr*. 2019;**108**(12):2132-4. [PubMed ID: 31420898]. <https://doi.org/10.1111/apa.14970>.
35. Hermanussen M, Scheffler C. Stature signals status: The association of stature, status and perceived dominance - a thought experiment. *Anthropol Anz*. 2016;**73**(4):265-74. [PubMed ID: 27643683]. <https://doi.org/10.1127/anthranz/2016/0698>.