



Evaluation of *Scrophularia striata* Hydroalcoholic Extract's Hematomodulatory Effects: An Approach to Traditional Ethnomedicine

Masoud Shamohammadi¹, Mehrdad Pooyanmehr^{2,*} and Ali Maleki³

¹Faculty of Veterinary Medicine, Razi University, Kermanshah, Iran

²Department of Basic Sciences, Faculty of Veterinary Medicine, Razi University, Kermanshah, Iran

³Medical Biology Research Center, Health Technology Institute, Kermanshah University of Medical Sciences, Kermanshah, Iran

*Corresponding author: Department of Basic Sciences, Faculty of Veterinary Medicine, Razi University, Kermanshah, Iran. Email: m.pooyanmehr@razi.ac.ir

Received 2021 June 13; Revised 2022 February 05; Accepted 2022 May 14.

Abstract

Background: Silver nanoparticles (AgNPs) have broad biomedical applications and may result in a high risk of human and environmental toxicity. *Scrophularia striata* possesses a strong antioxidant capacity. **Objectives:**

This study seeks to examine the modulatory influence of *Scrophularia striata* hydroalcoholic extract on the toxicity of silver nanoparticles on blood granulocytes, agranulocytes, and general parameters in male Wistar rats.

Methods: Thirty male rats weighing an average of 200 ± 20 were divided into five experimental groups of six. In 30 days, rats were given 2 mL of distilled water in group 1 (negative control), 200 ppm AgNPs (hepatotoxicity dosage) in group 2, as well as 20, 60, and 180 mg/kg *Scrophularia striata* extract and 200 ppm AgNPs in groups 3, 4, and 5. Rats' blood samples were taken 24 hours following their last partial anesthesia treatment. One-way ANOVA was used to evaluate the data, followed by Tukey's post hoc test.

Results: When compared to the control groups, oral treatment of AgNPs resulted in substantial changes in WBC, NE, LY, MO, EO, BA, RBC, Hb, HCT, MCV, MCH, MCHC, PLT, and MPV levels. In comparison to the positive and negative groups, the extract's modulatory effects on hematological parameters demonstrated substantial variations, especially in groups 4 and 5 (60 and 180 mg/kg).

Conclusions: Due to the presence of some unique components, such as flavonoids, the extract of the *Scrophularia striata* plant was discovered to be an efficient hematomodulatory, compensating to some extent for the cytotoxic side effects of AgNPs within the body.

Keywords: Silver Nanoparticle, Toxicity, Hematomodulatory, Traditiona Ethnomedicine, *Scrophularia striata*

1. Background

Nanotechnology research and development is an expanding discipline, and the global output of new nanoparticles is rising. The NPs have application in various fields, including biomedicine, diseases diagnosis, drug, agriculture, food industry, antimicrobial therapy, cosmetics, toothpastes, shampoos, paints, food supplements, water treatment, etc (1). Despite the benefits, NPs have led to potential toxicities in both humans and animals. Among the nanomaterials, Silver nanoparticles (AgNPs) has gained interest because of their distinctive properties, such as chemical stability, antibacterial activity, antifungal, antiviral, and anti-inflammatory characteristics (2). In recent years, there has been a surge in the production and use of silver nanoparticles (AgNPs), which will account for more than half of all global nanomaterial products by 2024

(3). Passing through cell membranes and interacting with biomolecules, silver nanoparticles induce DNA and protein damage and cell modification (4). Toxicity induced by silver nanoparticles (AgNPs) and the role that oxidative stress plays in this process were demonstrated in human cells (5). Medicinal plants and natural products, on the other hand, have been utilized as traditional therapies for a variety of ailments for centuries (6). The phytochemical elements of medicinal plants are responsible for the majority of their pharmacological effects (7). In the western states of Iran, a plant of Snapdragon with the scientific name of *Scrophularia striata* Boiss has traditional medical usage. The different extracts of this plant with the local name of *Teshneh Dary* are used in treatment of many infections. *Scrophularia striata*'s wound-healing properties (8), antibacterial effects (9), anti-inflammatory activities and anti-pain (10), anti-cancer properties (11), antioxi-

tion (12), ranging reduced edema, infiltration, T-cells proliferation and their nitric oxide production anti-activity have been studied (13).

2. Objectives

Based on a review of the unique characteristics of *Scrophularia striata* and the fact that no research has been done on the plant's protective properties against AgNP toxicity, the goal of this study is to assess the hemathological modulatory and protective effects of a hydroalcoholic extract of *Scrophularia striata* in the rat model due to AgNP hepatotoxicity as a traditional ethnomedical method.

3. Methods

3.1. Plant Sample Collection

In the spring, *Scrophularia striata*'s fresh leaves were harvested in the province of Kermanshah and its environs. A botanist validated the plant's specifications with plant code No 42801, Kuh.

3.2. Preparation of the Extract

The leaves of *S. striata* were washed, chopped, and air-dried at room temperature in the shade before being processed into powder with a laboratory grinder. To create hydroalcoholic extract, the powder was extracted three times for 48 h using a 70/30 (V/V) combination of ethanol and water. We filtered the extract with Kartuosh filter paper. At a temperature of 40°C, we used a rotary evaporator to evaporate and concentrate the solvent. The amount of dry extract obtained was 0.22 mg. The dried extract was kept at 4°C until further usage. In this study, the correct doses of the extract were made by mixing the dried extract with normal saline (14, 15).

3.3. Preparation of Nanoparticles

The Pishgaman Iranian Nanomaterial Company provided a silver nanoparticle solution with a concentration of 4000 p. The AgNPs have a diameter of 5 to 8 nm. UV/Vis spectrophotometry (Biotek Epoch, USA) and inductively coupled plasma optical emission spectrometry (ICP-OES, Cambridge, UK) were used to filter the AgNPs, and scanning electron microscopy (SEM) and transmission electron microscopy (TEM) techniques were used to confirm their original physical and chemical properties. Pishgaman Iran Nanomaterials Company was in charge of standards. Silver nanoparticles were prepared and characterized by swirling 100 mM silver nitrate into a 1 percent (w/v) tannic acid solution (pH adjusted to 8 by adding 150 mM potassium carbonate) of polyvinyl pyrrolidone (PVP). The

solution turned a pale-yellow color (AgNPs). Following the manufacturer's directions, we diluted the original solution to 20 units to reach the toxicity concentration (200 ppm). After that, the animals were gavaged every day for 30 days (16, 17).

3.4. Experimental Animals

Based on the inclusion and exclusion criteria, thirty adult Wistar rats were bought from the Animal Care Unit of the Faculty of Veterinary Medicine at Razi University in Kermanshah, Iran. Prior to beginning the investigation, it was crucial to identify these variables. The following are the requirements for inclusion: Male Wistar rats aged 8 to 10 weeks, weighing 200 to 210 g, with typical behavior and activity. The following are the conditions for being excluded: female Wistar rats, male Wistar rats who were sick or frail, and rats that had been used in any previous inquiry were excluded from the study. Six rats per cage were housed in stainless steel cages under regulated environmental conditions of temperature ($22 \pm 2^\circ\text{C}$), relative humidity ($55 \pm 5\%$), and illumination (12-h light/12-h dark cycle). All of the rats were fed regular lab pellets and had access to clean water at all times.

3.5. Experimental Design

This experimental research study was carried out between June 2019 and August 2019 at the biomedical laboratory of the Department of Basic Science sector of immunohematology at the Faculty of Veterinary Medicine, Razi University. After that, the rats were divided into five groups, each with six animals. The normal control group consisted of rats that were given normal saline orally and regularly. Animals in group 2 were only given AgNPs (200 ppm) orally and on a regular basis. The hydroalcoholic extract of *S. striata* was given to rats in groups 3, 4, and 5 by oral gavage at dosages of 20, 60, and 180 mg/kg body weight, respectively, with AgNPs. The dosages of AgNPs and *S. striata* extracts used in this investigation were calculated using previous publications and our own research (18, 19). The investigation lasted 30 days. In accordance with the National Institutes of Health Guidelines for the care and use of laboratory animals in biomedical research, the Animal Ethics Committee at Razi University gave its approval for the study to be conducted (Animal Ethical Approval Number: 396-2-038).

3.6. Animals Euthanize

At the end of the treatment period of thirty days, the animals were euthanized by giving them an intraperitoneal injection of 1 mg/kg Xylazine HCl (Xylazine 2%; Alfasan) and 0.5 mg/kg Ketamine HCl. This combination

caused the animals to die quickly (Ketamine 5 percent; Trittau, Germany) (20).

3.7. Sample Collection

The Whole blood was extracted from the heart and to a standard laboratory tubes with anticoagulation (EDTA) and one minute were shaken to prevent possible blood clotting. The hematology factors including, NE% (percent of neutrophils), WBC (white blood cell count), MO% (percent of monocytes), LY% (percent of lymphocytes), EO% (percent of eosinophils), RBC (red blood cell count), BA% (percent of basophils), HCT (hematocrit), Hb (hemoglobin concentration), MCH (mean corpuscular hemoglobin), MCV (mean corpuscular volume), MCHC (mean corpuscular hemoglobin concentration), PLT (platelet count), and MPV (mean platelet volume) were analyzed using a blood cell counter auto analyzer (Sysmex KX-21N Technology, Japan) (21, 22).

3.8. Statistical Analysis

The acquired data were analyzed with SPSS software for Windows (Version 21, Chicago, IL, USA), using one-way analysis of variance (ANOVA) and Tukey's HSD post-hoc test. The significance level between the groups was fixed at $P < 0.05$ and the findings were presented as mean \pm SEM.

4. Results

4.1. Granulocyte & a-granulocyte Parameters

The results showed, in all studied factors (granulocyte & agranulocyte parameters) administration of Ag nanoparticles and induction hepatotoxicity with a dose of 200 ppm, changed in the positive control group compared to the negative control group. It was also demonstrated that administration of *Scrophularia striata* hydroalcoholic extract in treatment groups (20, 60, and 180 mg/kg) together with AgNPs 200 ppm (Dose of hepatotoxicity) in a dose-dependent, especially group 4 (60 mg/kg) and group 5 (180 mg/kg) had hematomodulations of granulocyte and a-granulocyte parameters in comparison with the positive control groups and could modulate the cytotoxicity effects of silver nanoparticles in rats. Consequently, extract dosage at concentrations 20 mg/kg, 60 mg/kg, and 180 mg/kg (groups 3, 4, and 5) caused the levels of WBC, Neutrophils, Lymphocyte, monocyte, eosinophils, and basophils to get close to the negative control ($P < 0.05$), (Figures 1-6). In order, the highest effect of the extract on indices WBC (8.3 ± 0.4 , 8.3 ± 0.37) was related to the concentrations of 60&180 mg/kg (Figure 1). The highest dosage of the extract of *Scrophularia striata* at concentrations of

180 mg/kg (group 5) showed hematomodulation of Neutrophils (49 ± 0.77) and the highest dosage of the extract at concentrations of 60&180 mg/kg (groups 4 & 5) showed hematomodulation of Lymphocyte (33 ± 0.94 & 33 ± 1.23) levels compared with the positive and negative control ($P < 0.05$), (Figures 2,3). The highest effect of the extract on indices monocyte (12 ± 0.65 & 12 ± 0.68), was related to the concentration of 60 & 180 mg/kg (groups 4 & 5) (Figure 4). The results showed that hematomodulated levels of eosinophils (0.83 ± 0.01) in 5group & basophils (0.08 ± 0.05 , 0.08 ± 0.01 & 0.07 ± 0.61) in 3, 4 & 5 groups in comparison with the positive and negative control (Figures 5,6), ($P < 0.05$).

4.2. General Parameters

The evaluation of the data revealed that the treatment of 200 ppm AgNPs altered the range of general blood parameters in rats (RBC, Hb, HCT, MCV, MCH, MCHC, PLT, and MPV) in both the positive and negative control groups. In addition to this, it was demonstrated that the administration of *Scrophularia striata* hydroalcoholic extract in treatment groups (20, 60, and 180 mg/kg) along with AgNPs 200 ppm, especially group 5 (180 mg/kg) and group 4 (60 mg/kg) had hematomodulations on blood general parameters in comparison with the positive control groups and could modulate the cytotoxicity effects of silver nanoparticles ($P < 0.05$), (Table 1). In order, the highest effect of the extract on indices RBC (8.4 ± 0.69), was related to the concentrations of 180 mg/kg. The maximum dose of *Scrophularia striata* extract at concentrations of 180 mg/kg (group 5) exhibited hematomodulation of Hb (11.15 ± 1.39) and HCT (38.8 ± 0.43) levels relative to the positive and negative control ($P < 0.05$). The effect of the extract on indices MCV (46.3 ± 1.01 & 46.93 ± 0.63) was related to the concentration of 60 & 180 mg/kg. The highest effect of the extract on indices MCH (13.2 ± 0.57 & 13.2 ± 0.99) & MCHC (30.4 ± 0.6), was related to the concentration of 60 & 180 mg/kg. The findings demonstrated hematomodulated levels of PLT (1075.3 ± 14.1 & 1075.73 ± 9.98) in groups 4 and 5 and MPV (10.88 ± 0.86) in group 5 relative to the positive and negative control ($P < 0.05$). According to collected results, treatment with *S. striata* extract substantially dose-dependently regulated hematomodulation levels of the aforementioned parameters compared to AgNPs-treated rats ($P < 0.05$) (Table 1).

5. Discussion

Fresh and green components of *Scrophularia striata* have been shown to be employed in traditional medicine studies. *S. striata* contains anti-inflammatory




Figure 1. WBC parameter after 30 Days administration the AgNP & *Scrophularia striata* extract with significantly different the three treatment groups in comparison with the control groups ($P < 0.05$).




Figure 2. Neutrophil parameter after 30 Days administration the AgNP & *Scrophularia striata* extract with significantly different the three treatment groups in comparison with the control groups ($P < 0.05$).




Figure 3. Lymphocyte parameter after 30 Days administration the AgNP & *Scrophularia striata* extract with significantly different the three treatment groups in comparison with the control groups ($P < 0.05$).




Figure 4. Monocyte parameter after 30 Days administration the AgNP & *Scrophularia striata* extract with significantly different the three treatment groups in comparison with the control groups ($P < 0.05$).




Figure 5. Eosinophil parameter after 30 Days administration the AgNP & *Scrophularia striata* extract with significantly different the three treatment groups in comparison with the control groups ($P < 0.05$).




Figure 6. Basophil parameter after 30 Days administration the AgNP & *Scrophularia striata* extract with significantly different the three treatment groups in comparison with the control groups ($P < 0.05$). Notice that the values of the different small letters a, b, and c indicate a significant difference between the examined groups and the PC* and NC# groups, respectively ($P < 0.05$). In comparison to the control NC, each value is presented as the mean \pm SD of the three replications: PC stands for positive control, and PC stands for negative control. 200 PPM AgNP hepatotoxicity dosage: A: 20 mg/kg, B: 60 mg/kg, C: 180 mg/kg.

Table 1. The Hematological Parameters Value After 30 Days Administration the AgNP & *Scrophularia striata* Extract^a

Parameter	NC#	PC*	A	B	C
RBC (10^3 mm^3)	9.3 \pm 0.54	7.5 \pm 0.43	7.7 \pm 0.56 ^c	8.03 \pm 0.48 ^b	8.4 \pm 0.69 ^a
HB(g/dL)	13.35 \pm 1.73	10.5 \pm 0.79	10.75 \pm 1.2 ^c	10.95 \pm 0.99 ^b	11.15 \pm 1.39 ^a
HCT (%)	44.6 \pm 1.57	35.2 \pm 0.87	37.4 \pm 0.84 ^c	38.5 \pm 0.76 ^b	38.8 \pm 0.43 ^a
M CV (fl)	50.8 \pm 1.64	44.95 \pm 1.92	45.95 \pm 0.97 ^c	46.3 \pm 1.01 ^a	46.93 \pm 0.63 ^b
M CH (pg)	15.1 \pm 1.06	12.5 \pm 0.99	12.8 \pm 1.6 ^b	13.2 \pm 0.57 ^a	13.2 \pm 0.99 ^a
M CH C(g/dL)	31.58 \pm 0.89	29.1 \pm 1.17	30.1 \pm 0.83 ^b	30.4 \pm 0.6 ^a	30.13 \pm 0.91 ^b
PLT (10^3 mm^3)	1298.61 \pm 61.15	1050 \pm 96.25	915.5 \pm 37.65 ^b	1075.3 \pm 14.1 ^a	1075.73 \pm 9.98 ^a
MPV (fl)	10.64 \pm 0.56	11.64 \pm 0.19	11.26 \pm 0.21 ^b	11.21 \pm 0.44 ^b	10.88 \pm 0.86 ^a

^aNotice that the values of the different small letters a, b, and c indicate a significant difference between the examined groups and the PC* and NC# groups, respectively ($P < 0.05$). In comparison to the control NC, each value is presented as the mean \pm SD of the three replications: PC stands for positive control, and PC stands for negative control. 200 PPM AgNP hepatotoxicity dosage: A: 20 mg/kg, B: 60 mg/kg, C: 180 mg/kg.

flavonoid components (iridoids, phenolic acids, quercetin, isorhamnetin-3-O-rutinoside, cinnamic acid, polyphenolic acid, nepitrin, chlorogenic acid, phenylpropanoid glycoside, and other glycoterpenoids) according to phytochemical analysis (10, 23). Therefore, in this study, the possible hematomodulation of *Scrophularia striata* extract against AgNPs-induced hepatotoxicity was investigated. The CBC test, which examines the morphological characteristics and number of different types of blood cells (Red blood cells, Leukocytes, and Platelets) per cubic millimeter of blood, is one of the most common tests and provides valuable infor-

mation about different types of blood cell disorders, such as anemia (24).

The WBC disorders are mostly seen as proliferative, leukopenia, and cell dysfunction (25). Experimental studies have shown that long-term oral administration of AgNP's toxic dose in mice (28 - 30 days), increases spleen size and weight, total number of T, B, Mon and Nut cells (26), serum levels of IgM, IgG, proinflammatory cytokines and inflammation such as IL-1 β , TNF- α , IL-6, IL-4, IL-10, IL-12, and TGF- β (27), granulocytes and a-granulocytes blood (28). An increase in the numbers of white blood cells, may indicate

an antigen response or dysfunction of the bone marrow, liver, and spleen (29). Furthermore, earlier studies have shown that the hazardous potential of Ag-NPs affects some of the primary processes in inflammation (oxygen reactive species (ROS) formation and complement system, nitric oxide (NO) and COX-2 pathways) (30).

In the present study, the levels of granulocyte and a-granulocyte parameters (WBC, Neutrophil, lymphocyte, monocyte, eosinophils, and basophils) were increased in the positive control group due to the hepatotoxicity. The above results were consistent with the studies of Lee (31), Dosoky (32) and Pinzaru (33). The several studies have demonstrated the antioxidant and immunomodulatory properties of *S. striata* extract by using its anti-inflammatory mechanisms and with the help of Th2s that suppressed IL-10, IL-13, TGF- β , IL-1 β , TNF- α , NO and PGE2, reduced inflammation and thus the amount of white blood cells modulated (18, 34, 35). The hemathomo modulatory effects of *Scrophularia striata* extract with AgNPs 200 ppm were observed in treatment groups A, B, and C at the levels of granulocytes and a-granulocytes blood parameters, which is consistent with the findings of Abedi (36), Rostami (37), and Mahdavi (38) studies on hemathomo modulatory activity and growth performance. Therefore, based on the chemical analysis of *S. striata*, the hemathomodulation properties of the plant extract are related to the presence of several important anti-inflammatory compounds (phenolic acids, flavonoids, iridoids and phenylpropanoid glycosides).

Determination of RBC and the whole blood HGB level is a suitable parameter to assess the ability of blood to deliver oxygen to tissues and organs and to transport CO₂ to the lungs. The HCT indicates the percentage of red blood cells in total blood volume and decreases as the number of red blood cells decreases. The MCV shows red blood cell volume and has diagnostic value in anemia. The MCH index shows the average amount of hemoglobin in red blood cells and has diagnostic value in anemia. The MCHC indicates the average concentration of hemoglobin in red blood cells. Also, obtained results on CBC in this study showed an alteration in quantity of red blood cells parameters. Several mechanisms have been described for changes in blood parameters after nanoparticle consumption. For example, by their toxicological properties, Nanoparticles disrupt hemoglobin synthesis by increasing ROS, free radicals, oxidative stress (39, 40), thus there'll be liver damage, low metabolic activity, negative effects on bio kinetics (41), lipid peroxidation increase (42), and bone marrow suppression during erythrocyte maturation, in addition to disrupting RBC membrane integrity and erythrocyte hemolysis (43, 44). Several authors have suggested that silver nanoparticles link to sulfur and phosphorus

contained in biomolecules like DNA or other biological compounds, thus causing cell toxicity (45). Increased RBC degradation and decreased RBC count due to iron, cobalamin, or folic acid deficiency have been reported with oral administration and intraperitoneal injection of AgNP (46). The decrease in the red blood cell and HGB may have been resulted from the suppression of circulating erythropoietin hormone (a glycoprotein which stimulates the process of erythropoiesis) (47, 48). Inflammation of the gut and impaired absorption of iron, resulting in the production of inflammatory cytokines such as IL-6, which increase the production of hepcidin. With its inhibitory effect on ferroprotein-1, Hepsidine reduces the release of iron in the intestinal epithelial cells into the plasma, which in turn reduces the body's ferritin level and prevents the transformation of iron to pronormoblasts (49).

The significant decreases in the levels of RBC, HGB, HCT, MCH, MCHC and MCV with the dosage of 200PPM Ag-NP for 30 days reflects the hematotoxic effects of Ag-NP which could be attributed to the impaired absorption of iron, increase in the production of hepcidin & free radicals that compromise the integrity of the membrane. In this study, RBC and HGB levels increased and afterwards increase of HCT, MCV, MCH and MCHC levels were observed in three treatments groups (A, B & C) in comparison with the PC groups. The results of the present study correlate with several reports on the hematotoxic of AgNPs (50, 51), and this results were consistent with the Rostami, and Mahdavi on hemathomodulatory effects of *Scrophularia striata* studies (37, 38).

The PLT shows the quantity of platelet cells in the blood, and too much or too little leads to blood clotting disease and MPV shows the volume of platelets and increased or decreased platelet volume provides information about bone marrow platelet production (52). Also, the significant decreases in the levels of PLT counts and MPV (In treatment groups A, B & C) under the effect of 200PPM Ag-NP for 30 days reflects the hematotoxic effects of Ag-NP which could be attributed to the production of free radicals that compromise the integrity of the membrane. The results show the modulatory effects of *Scrophularia striata* extract on PLT counts and MPV, in three treatments groups (A, B & C) in comparison with the PC groups. Therefore, in the analysis of the data, the hematomodulation effects of *Scrophularia striata* are probably due to the presence of antioxidant compounds such as phenolic acids, flavonols and flavonoids (14, 53), which reduces the free radical damage of silver nanoparticles and thus reduces hemolysis and the suppressant effects of Bone marrow which leads to a blood-modulating effect on red blood cells (23).

5.1. Conclusions

The reported changes in white and red blood cells following the gavage of nanoparticles are possibly due to an immunogenic response (increased inflammation & ROS) and disturbances in signaling pathways and maturation of cells (dysfunction of the bone marrow, liver, and mitochondrial damage). These factors can have an effect on blood cells as well as the division and development of other cells. Also, silver nanoparticles increase inflammation and produce hepcidin which causes impaired absorption of iron and blood precursors and also they decrease the production of thrombopoietin which causes decrease in platelet count. It seems by reducing liver damage and inflammatory factors (free radicals) and thus reducing the production of hepcidin and increasing the production of thrombopoietin, the extract of *S. striata* modulates the levels of blood parameters (blood cells and platelets) due to the flavonoid compounds. Nevertheless, more research is needed to figure out the cellular and molecular processes that make *S. striata* extract protective.

Acknowledgments

The results described in this research is related to the graduate student thesis of veterinary medicine (D.V.M) with No: 7087460, Date: 19.11. 2018, and was supported with grant of the Faculty of Veterinary Medicine, Razi University, Iran. The researchers are extremely grateful to the laboratory experts involved in the experiments.

Footnotes

Authors' Contribution: Study concept and design: M.P developed the original idea and the protocol, abstracted and analyzed data, wrote the manuscript, and is a guarantor. A. M. and M. Sh. contributed to the development of the protocol, abstracted data, and prepared the manuscript.

Conflict of Interests: The authors declare that they have no conflict of interest. The authors alone are responsible for the content of the paper.

Data Reproducibility: It was not declared by the authors.

Ethical Approval: In accordance with the National Institutes of Health Guidelines for the care and use of laboratory animals in biomedical research, the Animal Ethics Committee at Razi University gave its approval for the study to be conducted (Animal Ethical Approval Number: 396-2-038).

Funding/Support: A part of the data in this research is related to the graduate student thesis of veterinary medicine

with No: 7087460, Date: 19.11. 2018, in the Faculty of Veterinary Medicine, Razi University, Iran. The researchers are extremely grateful to the laboratory experts involved in the experiments.

References

1. Stark WJ, Stoessel PR, Wohlleben W, Hafner A. Industrial applications of nanoparticles. *Chem Soc Rev*. 2015;**44**(16):5793-805. doi: [10.1039/c4cs00362d](https://doi.org/10.1039/c4cs00362d). [PubMed: [25669838](https://pubmed.ncbi.nlm.nih.gov/25669838/)].
2. Ivask A, Juganson K, Bondarenko O, Mortimer M, Aruoja V, Kasemets K, et al. Mechanisms of toxic action of Ag, ZnO and CuO nanoparticles to selected ecotoxicological test organisms and mammalian cells in vitro: a comparative review. *Nanotoxicology*. 2014;**8** Suppl 1:57-71. doi: [10.3109/17435390.2013.855831](https://doi.org/10.3109/17435390.2013.855831). [PubMed: [24256211](https://pubmed.ncbi.nlm.nih.gov/24256211/)].
3. Franci G, Falanga A, Galdiero S, Palomba L, Rai M, Morelli G, et al. Silver nanoparticles as potential antibacterial agents. *Molecules*. 2015;**20**(5):8856-74. doi: [10.3390/molecules20058856](https://doi.org/10.3390/molecules20058856). [PubMed: [25993417](https://pubmed.ncbi.nlm.nih.gov/25993417/)]. [PubMed Central: [PMC6272636](https://pubmed.ncbi.nlm.nih.gov/PMC6272636/)].
4. Maneewattanapinyo P, Banlunara W, Thammacharoen C, Ekgasit S, Kaewamatawong T. An evaluation of acute toxicity of colloidal silver nanoparticles. *J Vet Med Sci*. 2011;**73**(11):1417-23. doi: [10.1292/jvms.11-0038](https://doi.org/10.1292/jvms.11-0038).
5. Adeyemi OS, Shittu EO, Akpor OB, Rotimi D, Batiha GE. Silver nanoparticles restrict microbial growth by promoting oxidative stress and DNA damage. *EXCLI J*. 2020;**19**:492-500. doi: [10.17179/excli2020-1244](https://doi.org/10.17179/excli2020-1244). [PubMed Central: [PMC7214780](https://pubmed.ncbi.nlm.nih.gov/PMC7214780/)].
6. Cock IE, Winnett V, Sirdaarta J, Matthews B. The potential of selected Australian medicinal plants with anti-Proteus activity for the treatment and prevention of rheumatoid arthritis. *Pharmacogn Mag*. 2015;**11**(Suppl 1):S190-208. doi: [10.4103/0973-1296.157734](https://doi.org/10.4103/0973-1296.157734). [PubMed: [26109767](https://pubmed.ncbi.nlm.nih.gov/26109767/)]. [PubMed Central: [PMC4461961](https://pubmed.ncbi.nlm.nih.gov/PMC4461961/)].
7. Baharvand-Ahmadi B, Asadi-Samani M. A mini-review on the most important effective medicinal plants to treat hypertension in ethnobotanical evidence of Iran. *J Nephro pharmacology*. 2017;**6**(1):3-8. [PubMed Central: [PMC5295656](https://pubmed.ncbi.nlm.nih.gov/PMC5295656/)].
8. Haddadi R, Tamri P, Javani Jooni F. In vitro wound healing activity of *Scrophularia striata* hydroalcoholic extract. *S Afr J Bot*. 2019;**121**:505-9. doi: [10.1016/j.sajb.2019.01.002](https://doi.org/10.1016/j.sajb.2019.01.002).
9. Mohammadi M, Aelaei M, Saidi M. Antibacterial properties of *Scrophularia striata* Boiss. (Tashenehdari) extract on vase life improvement in "Stanza" and "Pink Elegance" gerbera cut flowers. *Biocatal Agric Biotechnol*. 2020;**28**. doi: [10.1016/j.bcab.2020.101738](https://doi.org/10.1016/j.bcab.2020.101738).
10. Tamri P. A mini-review on phytochemistry and pharmacological activities of *Scrophularia striata*. *J HerbMed Pharmacol*. 2019;**8**(2):85-9. doi: [10.15171/jhp.2019.14](https://doi.org/10.15171/jhp.2019.14).
11. Azadmehr A, Hajiaghaee R, Mazandarani M. Induction of apoptosis and G2/M cell cycle arrest by *Scrophularia striata* in a human leukaemia cell line. *Cell Prolif*. 2013;**46**(6):637-43. doi: [10.1111/cpr.12074](https://doi.org/10.1111/cpr.12074). [PubMed: [24460717](https://pubmed.ncbi.nlm.nih.gov/24460717/)]. [PubMed Central: [PMC6496267](https://pubmed.ncbi.nlm.nih.gov/PMC6496267/)].
12. Zargoosh Z, Ghavam M, Bacchetta G, Tavili A. Effects of ecological factors on the antioxidant potential and total phenol content of *Scrophularia striata* Boiss. *Sci Rep*. 2019;**9**(1):1-15. doi: [10.1038/s41598-019-52605-8](https://doi.org/10.1038/s41598-019-52605-8). [PubMed: [31690810](https://pubmed.ncbi.nlm.nih.gov/31690810/)]. [PubMed Central: [PMC6831573](https://pubmed.ncbi.nlm.nih.gov/PMC6831573/)].
13. Giessrigl B, Yazici G, Teichmann M, Kopf S, Ghassemi S, Atanasov AG, et al. Effects of *Scrophularia* extracts on tumor cell proliferation, death and intravasation through lymphoendothelial cell barriers. *Int J Oncol*. 2012;**40**(6):2063-74. doi: [10.3892/ijo.2012.1388](https://doi.org/10.3892/ijo.2012.1388).
14. Mahboubi M, Kazempour N, Boland Nazari AR. Total phenolic, total flavonoids, antioxidant and antimicrobial activities of *scrophularia striata* boiss extracts. *Jundishapur J Nat Pharm Prod*. 2013;**8**(1):15-9. doi: [10.17795/jjnpp-7621](https://doi.org/10.17795/jjnpp-7621). [PubMed: [24624181](https://pubmed.ncbi.nlm.nih.gov/24624181/)]. [PubMed Central: [PMC3941883](https://pubmed.ncbi.nlm.nih.gov/PMC3941883/)].

15. Zangeneh MM, Najafi F, Tahvilian R, Zangeneh A, Moradi R. Assessment of In Vitro Antibacterial Properties of the Hydroalcoholic Extract of *Scrophularia striata* Against *Staphylococcus aureus* (ATCC No. 25923). *Int J Pharmacogn Phytochem Res*. 2017;**9**(1):40-4. doi: [10.25258/ij-paprv9i1.8037](https://doi.org/10.25258/ij-paprv9i1.8037).
16. Xie J, Dong W, Liu R, Wang Y, Li Y. Research on the hepatotoxicity mechanism of citrate-modified silver nanoparticles based on metabolomics and proteomics. *Nanotoxicology*. 2018;**12**(1):18-31. doi: [10.1080/17435390.2017.1415389](https://doi.org/10.1080/17435390.2017.1415389). [PubMed: 29251223].
17. Naguib M, Mahmoud UM, Mekawy IA, Sayed AEH. Hepatotoxic effects of silver nanoparticles on *Clarias gariepinus*; Biochemical, histopathological, and histochemical studies. *Toxicol Rep*. 2020;**7**:133-41. doi: [10.1016/j.toxrep.2020.01.002](https://doi.org/10.1016/j.toxrep.2020.01.002). [PubMed: 31956514]. [PubMed Central: PMC6962648].
18. Shamohammadi M, Pooyanmehr M, Maleki A, Alimohammadi S. Evaluation of Protective and Immunomodulatory Effects of Hydroalcoholic Extract of *Scrophularia striata* on Silver Nanoparticle-Induced Toxicity in Male Rats. *Arch Adv Biosci*. 2021;**12**(1):7-17. doi: [10.22037/aab.v12i1.32832](https://doi.org/10.22037/aab.v12i1.32832).
19. Shariatzadeh S, Kazempour N, Nazifi S. Investigation of Protective Effect of Hydroalcoholic Extract of Ginger on Cytotoxicity of Silver Nanoparticles on Hepatic Enzymes, Hemathologic Factors, Blood Oxidative Stress Markers and Hepatic Apoptosis In Balb-C Mice. *J Arak Uni Med Sci*. 2016;**19**(9):78-82. doi: [10.22059/ijym.2021.308066.1005119](https://doi.org/10.22059/ijym.2021.308066.1005119).
20. Wellington D, Mikaelian I, Singer L. Comparison of ketamine-xylozine and ketamine-dexmedetomidine anesthesia and intraperitoneal tolerance in rats. *J Am Assoc Lab Anim Sci*. 2013;**52**(4):481-7. [PubMed Central: PMC3725934].
21. Ilies M, Iuga CA, Loghin F, Dhople VM, Thiele T, Volker U, et al. Impact of blood sample collection methods on blood protein profiling studies. *Clin Chim Acta*. 2017;**471**:128-34. doi: [10.1016/j.cca.2017.05.030](https://doi.org/10.1016/j.cca.2017.05.030). [PubMed: 28577959].
22. Kumar M, Dandapat S, Prasad Sinha M, Kumar A, Raipat BS. Different blood collection methods from rats: A review. *Balneo Res J*. 2017;**8**(2):46-50. doi: [10.12680/balneo.2017.141](https://doi.org/10.12680/balneo.2017.141).
23. Kerdar T, Moradkhani S, Dastan D. Phytochemical and Biological Studies of *Scrophularia striata* from Ilam. *Jundishapur J Nat Pharm Prod*. 2018;**13**(3). doi: [10.5812/jjnpp.62705](https://doi.org/10.5812/jjnpp.62705).
24. Abdelmahmud E, Yassin MA, Ahmed M. Iron Deficiency Anemia-Induced Neutropenia in Adult Female. *Cureus*. 2020. doi: [10.7759/cureus.8899](https://doi.org/10.7759/cureus.8899).
25. Yang Y, Hu T, Li J, Xin M, Zhao X. Structural characterization and effect on leukopenia of fucoidan from *Durvillaea antarctica*. *Carbohydr Polym*. 2021;**256**:117529. doi: [10.1016/j.carbpol.2020.117529](https://doi.org/10.1016/j.carbpol.2020.117529). [PubMed: 33483047].
26. De Jong WH, Van Der Ven LT, Sleijffers A, Park MV, Jansen EH, Van Loveren H, et al. Systemic and immunotoxicity of silver nanoparticles in an intravenous 28 days repeated dose toxicity study in rats. *Biomaterials*. 2013;**34**(33):8333-43. doi: [10.1016/j.biomaterials.2013.06.048](https://doi.org/10.1016/j.biomaterials.2013.06.048). [PubMed: 23886731].
27. Devanabanda M, Lathief SA, Madduri R. Immunotoxic effects of gold and silver nanoparticles: Inhibition of mitogen-induced proliferative responses and viability of human and murine lymphocytes in vitro. *J Immunotoxicol*. 2016;**13**(6):897-902. doi: [10.1080/1547691X.2016.1234522](https://doi.org/10.1080/1547691X.2016.1234522). [PubMed: 27754724].
28. Munger MA, Radwanski P, Hadlock GC, Stoddard G, Shaaban A, Falconer J, et al. In vivo human time-exposure study of orally dosed commercial silver nanoparticles. *Nanomedicine*. 2014;**10**(1):1-9. doi: [10.1016/j.nano.2013.06.010](https://doi.org/10.1016/j.nano.2013.06.010). [PubMed: 23811290]. [PubMed Central: PMC3877176].
29. Heydrnejad MS, Samani RJ, Aghaeivanda S. Toxic Effects of Silver Nanoparticles on Liver and Some Hematological Parameters in Male and Female Mice (*Mus musculus*). *Biol Trace Elem Res*. 2015;**165**(2):153-8. doi: [10.1007/s12011-015-0247-1](https://doi.org/10.1007/s12011-015-0247-1). [PubMed: 25637567].
30. Barcinska E, Wierzbicka J, Zauszkiewicz-Pawlak A, Jacewicz D, Dabrowska A, Inkielwicz-Stepniak I. Role of Oxidative and Nitro-Oxidative Damage in Silver Nanoparticles Cytotoxic Effect against Human Pancreatic Ductal Adenocarcinoma Cells. *Oxid Med Cell Longev*. 2018;**2018**:8251961. doi: [10.1155/2018/8251961](https://doi.org/10.1155/2018/8251961). [PubMed: 30186549]. [PubMed Central: PMC6116403].
31. Lee JH, Gulumian M, Faustman EM, Workman T, Jeon K, Yu JJ. Blood Biochemical and Hematological Study after Subacute Intravenous Injection of Gold and Silver Nanoparticles and Co-administered Gold and Silver Nanoparticles of Similar Sizes. *Biomed Res Int*. 2018;**2018**:8460910. doi: [10.1155/2018/8460910](https://doi.org/10.1155/2018/8460910). [PubMed: 30140702]. [PubMed Central: PMC6081526].
32. Dosoky WM, Fouda MMG, Alwan AB, Abdelsalam NR, Taha AE, Ghareeb RY, et al. Dietary supplementation of silver-silica nanoparticles promotes histological, immunological, ultrastructural, and performance parameters of broiler chickens. *Sci Rep*. 2021;**11**(1):1-15. doi: [10.1038/s41598-021-83753-5](https://doi.org/10.1038/s41598-021-83753-5). [PubMed: 33603060]. [PubMed Central: PMC7892842].
33. Pinzaru I, Coricovac D, Dehelean C, Moaca EA, Mioc M, Baderca F, et al. Stable PEG-coated silver nanoparticles - A comprehensive toxicological profile. *Food Chem Toxicol*. 2018;**111**:546-56. doi: [10.1016/j.fct.2017.11.051](https://doi.org/10.1016/j.fct.2017.11.051). [PubMed: 29191727].
34. Rostami F, Taherpour K, Ghasemi HA, Akbari Gharaei M, Shirzadi H. Effects of *Scrophularia striata* hydroalcoholic extract in comparison to salinomycin on growth performance, intestinal health and immunity in broiler chickens following a mixed-species *Eimeria* challenge. *Vet Parasitol*. 2021;**293**:109417. doi: [10.1016/j.vetpar.2021.109417](https://doi.org/10.1016/j.vetpar.2021.109417). [PubMed: 33819905].
35. Goudarzvand M, Saadat P, Ebrahimi H, Hajiaghaee R, Sadat Miri N, Fallahnezhad S, et al. Immunomodulatory and anti-inflammatory effects of *Scrophularia megalantha* ethanolic extract on an experimental model of multiple sclerosis. *Re J Pharmacogn*. 2019;**6**(1):43-50. doi: [10.22127/RJP.2018.80370](https://doi.org/10.22127/RJP.2018.80370).
36. Abedi A, Bamdad K, Yaghoubi Shahir H, Dehghany R, Moradi A, Saadati H. Comparison of the protective effects of *Scrophularia striata* extract with vitamin E on cognitive function, anxiety and pain sensitivity in diazinon-exposed male rats. *J Physiol Pharmacol*. 2020;**25**(1):47-56. doi: [10.32598/ppj.25.1.30](https://doi.org/10.32598/ppj.25.1.30).
37. Rostami F, Taherpour K, Akbari GM, Shirzadi Hassan, Ghasemi HA. Effects of hydroalcoholic extract of *Scrophularia striata* in comparison with antibiotic, probiotic and a multivitamin and mineral supplement on growth performance and blood parameters of broilers. *Anim Sci Res*. 2019;**28**(4):197-201.
38. Mahdavi A, Bahrami AM, Yousefzadeh S. The effect of hydroalcoholic extract of *Scrophularia striata* on the changes of testis tissue in hypercholesterolemic rats. *Vet Clin Pathol Q Sci J*. 2021;**14**(56):347-64.
39. Mirzaei S, Hadadi Z, Attar F, Mousavi SE, Zargar SS, Tajik A, et al. ROS-mediated heme degradation and cytotoxicity induced by iron nanoparticles: hemoglobin and lymphocyte cells as targets. *J Biomol Struct Dyn*. 2018;**36**(16):4235-45. doi: [10.1080/07391102.2017.1411832](https://doi.org/10.1080/07391102.2017.1411832). [PubMed: 29195491].
40. Sarkar PK, Halder A, Adhikari A, Polley N, Darbar S, Lemmens P, et al. DNA-based fiber optic sensor for direct in-vivo measurement of oxidative stress. *Sens Actuators B Chem*. 2018;**255**:2194-202. doi: [10.1016/j.snb.2017.09.029](https://doi.org/10.1016/j.snb.2017.09.029).
41. Jo MR, Bae SH, Go MR, Kim HJ, Hwang YG, Choi SJ. Toxicity and Biokinetics of Colloidal Gold Nanoparticles. *Nanomaterials (Basel)*. 2015;**5**(2):835-50. doi: [10.3390/nano5020835](https://doi.org/10.3390/nano5020835). [PubMed: 28347039]. [PubMed Central: PMC5312906].
42. Paciorek P, Zuberek M, Grzelak A. Products of Lipid Peroxidation as a Factor in the Toxic Effect of Silver Nanoparticles. *Materials (Basel)*. 2020;**13**(11). doi: [10.3390/ma13112460](https://doi.org/10.3390/ma13112460). [PubMed: 32481688]. [PubMed Central: PMC7321096].
43. Park EJ, Bae E, Yi J, Kim Y, Choi K, Lee SH, et al. Repeated-dose toxicity and inflammatory responses in mice by oral administration of silver nanoparticles. *Environ Toxicol Pharmacol*. 2010;**30**(2):162-8. doi: [10.1016/j.etap.2010.05.004](https://doi.org/10.1016/j.etap.2010.05.004). [PubMed: 21787647].

44. Kim MJ, Shin S. Toxic effects of silver nanoparticles and nanowires on erythrocyte rheology. *Food Chem Toxicol.* 2014;**67**:80–6. doi: [10.1016/j.fct.2014.02.006](https://doi.org/10.1016/j.fct.2014.02.006). [PubMed: [24534065](https://pubmed.ncbi.nlm.nih.gov/24534065/)].
45. Wang E, Huang Y, Du Q, Sun Y. Silver nanoparticle induced toxicity to human sperm by increasing ROS(reactive oxygen species) production and DNA damage. *Environ Toxicol Pharmacol.* 2017;**52**:193–9. doi: [10.1016/j.etap.2017.04.010](https://doi.org/10.1016/j.etap.2017.04.010). [PubMed: [28433807](https://pubmed.ncbi.nlm.nih.gov/28433807/)].
46. Fang W, Chi Z, Li W, Zhang X, Zhang Q. Comparative study on the toxic mechanisms of medical nanosilver and silver ions on the antioxidant system of erythrocytes: from the aspects of antioxidant enzyme activities and molecular interaction mechanisms. *J Nanobiotechnology.* 2019;**17**(1):1–13. doi: [10.1186/s12951-019-0502-2](https://doi.org/10.1186/s12951-019-0502-2). [PubMed: [31101056](https://pubmed.ncbi.nlm.nih.gov/31101056/)]. [PubMed Central: [PMC6524268](https://pubmed.ncbi.nlm.nih.gov/PMC6524268/)].
47. Goldwasser E, Kung CK, Eliason J. On the Mechanism of Erythropoietin-induced Differentiation: XIII. The role of sialic acid in erythropoietin action. *J Biol Chem.* 1974;**249**(13):4202–6. doi: [10.1016/s0021-9258\(19\)42503-9](https://doi.org/10.1016/s0021-9258(19)42503-9).
48. Qian Y, Zhang J, Hu Q, Xu M, Chen Y, Hu G, et al. Silver nanoparticle-induced hemoglobin decrease involves alteration of histone 3 methylation status. *Biomaterials.* 2015;**70**:12–22. doi: [10.1016/j.biomaterials.2015.08.015](https://doi.org/10.1016/j.biomaterials.2015.08.015). [PubMed: [26295435](https://pubmed.ncbi.nlm.nih.gov/26295435/)].
49. Cenci L, Andreetto E, Vestri A, Bovi M, Barozzi M, Iacob E, et al. Surface plasmon resonance based on molecularly imprinted nanoparticles for the picomolar detection of the iron regulating hormone Hepcidin-25. *J Nanobiotechnology.* 2015;**13**:1–15. doi: [10.1186/s12951-015-0115-3](https://doi.org/10.1186/s12951-015-0115-3). [PubMed: [26311037](https://pubmed.ncbi.nlm.nih.gov/26311037/)]. [PubMed Central: [PMC4549936](https://pubmed.ncbi.nlm.nih.gov/PMC4549936/)].
50. Al-Baker A, AlKshab AA, Ismail H. Effect of silver nanoparticles on some blood parameters in rats. *Iraqi J Vet Sci.* 2020;**34**(2):389–95. doi: [10.33899/ijvs.2020.165812](https://doi.org/10.33899/ijvs.2020.165812).
51. Khan MS, Qureshi NA, Jabeen F, Shakeel M, Asghar MS. Assessment of Waterborne Amine-Coated Silver Nanoparticle (Ag-NP)-Induced Toxicity in *Labeo rohita* by Histological and Hematological Profiles. *Biol Trace Elem Res.* 2018;**182**(1):130–9. doi: [10.1007/s12011-017-1080-5](https://doi.org/10.1007/s12011-017-1080-5). [PubMed: [28634814](https://pubmed.ncbi.nlm.nih.gov/28634814/)].
52. Gasparyan AY, Ayvazyan L, Mikhailidis DP, Kitas GD. Mean platelet volume: a link between thrombosis and inflammation? *Curr Pharm Des.* 2011;**17**(1):47–58. doi: [10.2174/138161211795049804](https://doi.org/10.2174/138161211795049804). [PubMed: [21247392](https://pubmed.ncbi.nlm.nih.gov/21247392/)].
53. Monsef-Esfahani HR, Hajiaghah R, Shahverdi AR, Khorramizadeh MR, Amini M. Flavonoids, cinnamic acid and phenyl propanoid from aerial parts of *Scrophularia striata*. *Pharm Biol.* 2010;**48**(3):333–6. doi: [10.3109/13880200903133829](https://doi.org/10.3109/13880200903133829). [PubMed: [20645822](https://pubmed.ncbi.nlm.nih.gov/20645822/)].