



Interleukin-4, Interleukin-1beta, and Creatine Kinase Changes to the DeLorme and Oxford Resistance Training Techniques

Asad Mardani ¹, Hossein Abednatanzi ^{1,*}, Mandana Gholami¹, Farshad Ghazalian ¹ and Kamal Azizbeigi ²

¹Department of Physical Education and Sport Science, Sciences and Research Branch, Islamic Azad University, Tehran, Iran

²Department of Physical Education, Sanandaj Branch, Islamic Azad University, Sanandaj, Iran

*Corresponding author: Department of Physical Education and Sport Science, Sciences and Research Branch, Islamic Azad University, Tehran, Iran. Email: abednazari@gmail.com

Received 2021 September 21; Revised 2022 August 01; Accepted 2022 August 10.

Abstract

Background: Exercise intensity sequence of resistance training (RT) has a key role in the physiological response and adaptations.

Objectives: The aim of the present study was to compare the effect of DeLorme and Oxford resistance training techniques on the concentration of interleukin-4 (IL-4), interleukin-1 beta (IL-1beta) and creatine kinase (CK) enzyme in overweight men.

Methods: Thirty overweight young men ($25 \leq \text{BMI} < 30$) voluntarily participated in the present study and were randomly assigned to the DeLorme (Del; $n = 10$), Oxford (OXF; $n = 10$) resistance training techniques and control (Con; $n = 10$). Del performed 4-5 sets at intensity of 50% of one-maximum repetition (1RM) for warm up, the second set with 85% of 1RM, the third set at 90% of 1RM, and the fourth set at 95 of 1RM, and finally the fifth set at 100 1RM of exercise, while the OXF performed mentioned protocol exactly reverse order for eight weeks. RT was done three day/week in nonconsecutive for eight week. Blood sampling was gathered before RT, and repeated 72 hour after the last session of RT, and the levels of CK activity as well as the concentration of IL-4 and IL-1beta concentration were measured in the plasma.

Results: The results showed that there was no significant difference between the Del and OXF in biochemical variables ($P > 0.05$). Also, in the IL-4, IL-1beta and CK, there were no improvement was found in the Del and OXF compared to the control ($P > 0.05$).

Conclusions: Finally, it can be said that none of the DeLorme and Oxford resistance training technique are preferred in improving systemic inflammatory factors, and the sequence of resistance training intensity is more or less and vice versa has no effect on improving systemic inflammatory factors.

Keywords: Strength Training, Exercise Intensity, Systemic Inflammation, Cytokine

1. Background

The positive effect of resistance training (RT) on many diseases and physiological systems is well established (1). RT can increase body mass and improve body composition, improving blood pressure and insulin resistance, as well as reduce oxidative stress and improve systemic inflammation (1). Systemic inflammation has been reported to be associated with many diseases, including cardiovascular disease (CVD) (2). Plasma indices of inflammation include an increase in some pre-inflammatory cytokines (3). Interleukin-6 (IL-6), interleukin-4 (IL-4), and interleukin-1beta (IL-1beta) are cytokines that show more inflammation than normal values, and are associated with cytokines and pre-inflammatory factors (4).

However, the effect of RT on inflammation has been more or less studied, and studies have reported somewhat

a reduction in inflammation systemic by RT (5, 6). Some studies suggest that RT may well improve systemic inflammatory factors (7), while several studies have suggested that RT has not been shown to improve pre-inflammatory factors (5).

Anyway, RT has many factors and variables compared to other types of exercise training mode such as aerobic training, and it seems that the change in each variable of RT has a special response and adaptation to training (8). Accordingly, this issue has been considered by studies and in this regard, some variables of RT such as intensity on changes in pre-inflammatory cytokines have been investigated (9).

However, the intensity of exercise is an important factor in influencing many biochemical changes and adaptation (8). But how to apply the intensity of the exercises, especially in RT, can also be considered. In this regard, it has

been reported that the intensity of training from high to low and or from low to high during resistance training improves strength, and there is no difference in the amount of cell damage between the two types of training (10).

However, considering the aspect of the issue and the lack of information on the effect of different intensity sequences on resistance training on inflammation, it is necessary to examine the sequence effect of resistance training on systemic changes in the likelihood of systemic inflammation.

2. Objectives

The aim of the present study is to investigate the effect of resistance training intensity from high to low intensity training sequence and from low to high intensity training sequence in the form of DeLorme and Oxford resistance training on some systemic inflammatory factors and cell damage, and we attempted to answer the question of whether there is a significant difference between DeLorme and Oxford resistance training technique after eight weeks of training program on systemic inflammatory factors or not?

3. Methods

3.1. Subjects

30 overweight men ($25 \leq$ Body Mass Index (BMI) \leq 29.5) voluntarily participated in the current study, and randomly assigned in the Delorme resistance training (Del; n = 10), Oxford resistance training (OXF; n = 10) and control (Con; n = 10).

All participants were required to be eligible for the following research prior to enrollment. (1) None experience in regular exercise training programs. (2) No history of common chronic health problems. (3) No smoking. (4) Have not taken any antioxidant supplements for the past 6 months. After explaining the full description of the topic, goals, study methods, and completing and obtaining the consent form health questionnaire and sports history of all volunteers after fulfilling the mentioned conditions. The present study was carried out with the approval of the ethics committee of Sport Sciences Research Institute of Iran with the ID number (IR.SSRI.REC.1399.797).

3.2. Preliminary Assessments

First, anthropometric indices were measured in the first session. Initially, the subject's weight and height were assessed by scale and audiometer (Seca, Mod 285, Germany) according to the instruction. Then, based on the

weight and height, body mass index (BMI: kg/m²), all subjects were calculated. After that, subjects were trained in a separate session on how to properly perform the resistance exercise program and safety tips. Firstly, each exercise was performed by the coach, and sufficient explanations were presented for subjects. The subjects then performed the exercise without load. During the performing exercises, the subjects were given the necessary explanation and feedback on exercise technique.

Also, the general and specific resistance training warm up were explained to the subjects. Then, in the last session of training, the measurement of maximum repetition (1RM) in Leg press (LP) and Bench press (BP) exercise in all three groups was estimated through the Brzycki equation (11). In this way, the subjects selected a weight by initial estimation of their maximum strength, and performed the exercise to the exhaustion (the number of exercise need to be less than 10 repetitions). Then, by including the amount of weight and the number of repetitions in the formula for the estimation equation, a 1RM was calculated by the Brzycki formula. $(0.278/0 \times \text{Number of repetitions to the point of fatigue}) - 0.278/1$ || amount of weight transferred = one more repetition.

3.3. Resistance Training Program

The resistance training was performed three times a week in nonconsecutive days for eight weeks. All training sessions were done simultaneously in the Del and the OXF at 5 PM under the supervision of the expert coaches. The exercises for the upper limbs were chest press (CP), lat pull down (LP), bicep (BC) and triceps curls (TC) barbell. Also Leg press (LP), knee flexion (KF) and extension (EF) with machine were included in the lower limb exercises. Sit-ups exercise are also used to improve the abdominal and trunk muscles. Also, 5-10 minute warm-up training was included at the beginning and end of the each session. (10)

3.3.1. Delorme Resistance Training Program

Del resistance training program performed resistance training by a gradual increase in intensity, and a reduction in the number of repetitions until the exercise intensity reached to 1 RM. Thus, the subjects trained the first 2-3 sets at intensity of 50% of 1 RM for warm up, the second set was started at 85% of 1RM (6 repetitions), then the third set was performed at 90% of 1 RM (3-4 repetitions), and the fourth set was done at 95% of 1RM (2-3 repetitions), and finally the fifth set was performed and reached 100% 1RM (10).

3.3.2. Oxford Training Program

The OXF also performed the exercise training program using the method of gradual reduction of intensity and increasing the number of repetitions (10). The control group

did not participate in any physical activities or exercise training, and continued their daily activities without any change (10).

3.4. Blood Sampling and Biochemical Analysis

In order to evaluate the biochemical variables, 5 mL blood sampling was gathered in two phase, before any intervention, and also after 2 months of resistance training in fasting status with 48 hour interval with the last exercise training session. Subjects were first asked to avoid any physical activity 2 days before the test. Then 5 mL of blood sample was gathered from the antecubital vein in setting position after 15 rests at 8 - 10 am. After that, the serum was separated from the cell by centrifuging (Hitek, Germany) in the 2500 rpm for 7-10 min. IL-4 concentration serum IL-4 and IL-1beta were determined by an enzyme-linked immunosorbent assay (ELISA) with special kits according to the specifications of the manufacturer (abcam, Co, UK). Kits [(ab215089, UK); (Sensitivity: 1.08 pg/mL; Intra-assay: 6 %; and Inter-assay: 7%) and [(ab46052, UK) (Sensitivity: 6.5 pg/mL; Intra-assay: 4.5%; and Inter-assay: 8.7%)] were used for IL-4 and IL-1beta assay; respectively. Also, the CK activity of enzyme was measured by the kit (Pishtaz Teb, Iran). The activity of CK in the (U/L) and the concentration values of interleukin-4 and interleukin-1beta were expressed as (pg/mL).

3.5. Statistical Methods

All data were expressed as (Mean \pm SD). Kolmogorov-Smirnov test was first used to test the normality of data distribution. For the comparison of before and after the intervention, ANOVA with repeated measure was used. All statistical analyzing were performed at a significance level of $P \leq 0.05$ using SPSS-16 software.

4. Results

The anthropometric characteristics of the subjects are presented in [Table 1](#).

We found that BP and LP significantly were increased in the Del and OXF ($P = 0.001$), while there was no significant difference between Del and OXF in BP and LP ($P > 0.05$). Also, BMI had not significant differences between the two groups ($P > 0.05$).

The results showed that there was no significant difference in the concentration of IL-4, IL-1 beta and also CK before the start of the RT in the three groups ($P > 0.05$). Also we found that IL-4 and IL-1 beta were not affected by Del and OXF, and also the changes of these cytokines were not significant considering the effect of time. Also, the results showed that the CK was not affected by Del and OXF and

also the changes in this cellular damage index were not significant considering the effect of time ([Table 2](#)).

5. Discussion

The aim of the present study was to investigate the effect of DeLorme and Oxford resistance training techniques on changes in IL-4, IL-1 beta and CK in overweight men.

Accordingly, 20 overweight male underwent DeLorme and Oxford resistance training techniques for eight weeks to investigate the effectiveness of the mentioned resistance training techniques on IL-4, IL-1 beta and CK and physiological and functional variables.

We found that the strength increased significantly after eight week of RT in the Del and OXF. We found that 1RM was significantly increased in CP and SQ exercises in the Del as well as in OXF; however, there were no significant difference between groups. This showed that both RT protocol were similar efficient in strength increment.

The strength changes indicate that resistance training protocol in the current study could challenge the physiological systems, and therefore it has been effective on stress and stimulate systems.

Results showed that there was no significant difference in BMI between the Del and OXF before and after eight weeks resistance training.

The results showed that both Del and OXF did not significantly change IL-4 and IL-1beta concentration after 48 hours of the last session of resistance training. Also, we found that intra-group changes (time effect) were not statistically significant; this means that, regardless of the resistance training technique, both RT protocols had not a significant effect on IL-4 and IL-1beta. Therefore, it seems that the oxidative stress and damage caused by both protocols is similar, because the amount of these cytokines as inflammatory systemic indices had similarly behavior after both resistance exercises. It has been reported that exercise leads to a robust inflammatory response mainly characterized by the mobilization of leukocytes and an increase in circulating inflammatory mediators produced by immune cells and directly from the active muscle tissue (12). However, the relationship between CK and inflammatory responses to exercise and physical activity suggests that resistance training will disrupt myofibrils (13), especially during the eccentric phase. The rupture of the fibers leads to inflammatory responses that are modulated by cytokines (14). Cytokines have been shown to play an important role in muscle repair and repair processes (15). Consistent with the results of the present study it has been reported that resistance training for four months did not affect on IL-6 and TNF-alpha levels in healthy middle-aged

Table 1. Characteristics of the Subjects Before Resistance Training Intervention^a

Groups	Age (y)	Height (cm)	Weight (kg)	BMI (kg/m ²)
Del	20.5 ± 2.1	174.8 ± 1.5	97.4 ± 7.4	28.2 ± 1.6
OXF	22.8 ± 1.11	175.2 ± 4.678	96.3 ± 7.2	27.9 ± 1.4
Control	21.5 ± 2.3	173 ± 1.22	97.7 ± 6.4	28.3 ± 1.5

^a Values are presented as mean ± standard deviation.

Table 2. Statistical Analysis and Variables Changes Before and After the Resistance Training^a

Variables	Time		Time	Group × Time
	Pre Test	Post Test		
IL-1 beta (pg./m L)			0.849	0.407
Del	0.69 ± 0.77	0.45 ± 0.76		
OXF	0.41 ± 0.84	0.46 ± 0.90		
Control	0.51 ± 0.21	0.61 ± 0.19		
IL-4 (pg./m L)			0.654	0.155
Del	2.08 ± 0.25	1.98 ± 0.82		
OXF	1.49 ± 0.54	2 ± 0.73		
Control	2.3 ± 0.71	131 ± 11.65		
CK (U/L)			0.12	0.250
Del	134.6 ± 1.16	1.47 ± 0.19		
OXF	145 ± 13.73	135.9 ± 11.13		
Control	131 ± 14.63	128.8 ± 11.07		

Abbreviations: Del, Delorme; OXF, Oxford; IL-1beta, Interleukin-1beta; IL-4, Interleukin-4; CK, creatine kinase; BP, bench press; LP, leg press.

^a Values are presented as mean ± standard deviation.

men (16). However, it has recently been reported that moderate to high resistance training improve inflammatory factors in inactive elderly women (17). With considering the intensity of resistance training, the present resistance training protocol is similar with the resistance training protocols mentioned in the study, however, it seems that some other factors such as subject's sex, age and resistance training duration may affect the discrepancy of these results (18, 19), because gender hormones such as estrogen may affect inflammatory factors (20). It doesn't seem reasonable to introduce that the duration of the current resistance training (8 week) is the main cause for the unchanging IL-6 and IL-1beta (21). However, in refuting this issue, it should be noted that studies that have used the same duration (10-12 weeks) have reported a significant reduction in IL-6 and TNF-alpha (22). Therefore, the different cytokines behaviors do not appear to be due to the different of the training protocols used in the studies, and physiological factors such as body fat percentage should be considered.

In the present study, both Del and OXF resistance training technique did not effect on body fat percentage. It seems that the only studies that have reported a reduc-

tion in body fat, they have reported reduction of concentrations of pre-inflammatory cytokines after resistance training (20).

However, in comparison between the effects of exercise training on changes in inflammatory indices, some inconsistencies are related to the type of sample from which the cytokine was measured (tissue sample vs. plasma samples). On the other hand, biochemical methods of measuring cytokines (ELISA vs. Flow Cytometry) and human factors are also involved. However, because IL-4 and IL-1beta, as well as the activity of CK, were not affected by resistance training, it was difficult to compare intergroup and the susceptibility of these cytokines to the two types of resistance training technique with varying sequences intensity, it will be inexplicable.

5.1. Conclusions

Finally it can be concluded that Delorme and Oxford resistance training technique did not have favorable effects on systemic inflammatory factors, and it can be said that the sequence of resistance training intensity did not

influence on the inflammation and injury and there is a need for further research in this context.

Footnotes

Authors' Contribution: Asad Mardani designed the research and collected the data. Hossein Abednatanzi designed the research and supervised the work. Mandana Gholami analyzed data. Farshad Ghazalian contributed to the design and implementation of the research. Kamal Azizbeigi wrote the article.

Conflict of Interests: The authors declare no competing interests.

Ethical Approval: The present study was carried out with the approval of the ethics committee of Sport Sciences Research Institute of Iran with the ID number IR.SSRI.REC.1399.797 (Link: ethics.ssrc.ac.ir/article_3212.html).

Funding/Support: The authors received no funding.

Informed Consent: Informed consent has been obtained from subjects.

References

- Westcott WL. Resistance training is medicine: effects of strength training on health. *Curr Sports Med Rep*. 2012;**11**(4):209-16. doi: [10.1249/JSR.0b013e31825dabb8](https://doi.org/10.1249/JSR.0b013e31825dabb8). [PubMed: [22777332](https://pubmed.ncbi.nlm.nih.gov/22777332/)].
- Mason JC, Libby P. Cardiovascular disease in patients with chronic inflammation: mechanisms underlying premature cardiovascular events in rheumatologic conditions. *Eur Heart J*. 2015;**36**(8):482-9c. doi: [10.1093/eurheartj/ehu403](https://doi.org/10.1093/eurheartj/ehu403). [PubMed: [25433021](https://pubmed.ncbi.nlm.nih.gov/25433021/)]. [PubMed Central: [PMC4340364](https://pubmed.ncbi.nlm.nih.gov/PMC4340364/)].
- Jaffer U, Wade RG, Gourlay T. Cytokines in the systemic inflammatory response syndrome: a review. *HSR Proc Intensive Care Cardiovasc Anesth*. 2010;**2**(3):161-75. [PubMed: [23441054](https://pubmed.ncbi.nlm.nih.gov/23441054/)]. [PubMed Central: [PMC3484588](https://pubmed.ncbi.nlm.nih.gov/PMC3484588/)].
- Wojdasiewicz P, Poniatowski LA, Szukiewicz D. The role of inflammatory and anti-inflammatory cytokines in the pathogenesis of osteoarthritis. *Mediators Inflamm*. 2014;**2014**:561459. doi: [10.1155/2014/561459](https://doi.org/10.1155/2014/561459). [PubMed: [24876674](https://pubmed.ncbi.nlm.nih.gov/24876674/)]. [PubMed Central: [PMC4021678](https://pubmed.ncbi.nlm.nih.gov/PMC4021678/)].
- Azizbeigi K, Atashak S, Khorshidi D. Effect of progressive resistance training on interleukin-6 and C-reactive protein in untrained males. *Int Med J*. 2016;**23**(2):195-7.
- Abd El-Kader SM, Al-Shreef FM. Inflammatory cytokines and immune system modulation by aerobic versus resisted exercise training for elderly. *Afr Health Sci*. 2018;**18**(1):120-31. doi: [10.4314/ahs.v18i1.16](https://doi.org/10.4314/ahs.v18i1.16). [PubMed: [29977265](https://pubmed.ncbi.nlm.nih.gov/29977265/)]. [PubMed Central: [PMC6016983](https://pubmed.ncbi.nlm.nih.gov/PMC6016983/)].
- Zobairy M, Matinhomaei H, Hatamian H, Azizbeigi K, Azarbayjani MA. Effect of Elastic Resistance Training and Vitamin D on Systemic Inflammation Indices in Untrained Men: A Clinical Trial. *Casp J Neurol Sci*. 2017;**3**(11):196-205. doi: [10.29252/nirp.cjns.3.11.196](https://doi.org/10.29252/nirp.cjns.3.11.196).
- Bird SP, Tarpenning KM, Marino FE. Designing resistance training programmes to enhance muscular fitness: a review of the acute programme variables. *Sports Med*. 2005;**35**(10):841-51. doi: [10.2165/00007256-200535100-00002](https://doi.org/10.2165/00007256-200535100-00002). [PubMed: [16180944](https://pubmed.ncbi.nlm.nih.gov/16180944/)].
- Azizbeigi K, Azarbayjani MA, Atashak S, Stannard SR. Effect of moderate and high resistance training intensity on indices of inflammatory and oxidative stress. *Res Sports Med*. 2015;**23**(1):73-87. doi: [10.1080/15438627.2014.975807](https://doi.org/10.1080/15438627.2014.975807). [PubMed: [25630248](https://pubmed.ncbi.nlm.nih.gov/25630248/)].
- da Silva DP, Curty VM, Areas JM, Souza SC, Hackney AC, Machado M. Comparison of DeLorme with Oxford resistance training techniques: effects of training on muscle damage markers. *Biol Sport*. 2010;**27**(2):77-81. doi: [10.5604/20831862.913066](https://doi.org/10.5604/20831862.913066).
- Brzycki M. Strength Testing—Predicting a One-Rep Max from Reps-to-Fatigue. *J Phys Educ Recreat Dance*. 1993;**64**(1):88-90. doi: [10.1080/07303084.1993.10606684](https://doi.org/10.1080/07303084.1993.10606684).
- Cerqueira E, Marinho DA, Neiva HP, Lourenco O. Inflammatory Effects of High and Moderate Intensity Exercise—A Systematic Review. *Front Physiol*. 2019;**10**:1550. doi: [10.3389/fphys.2019.01550](https://doi.org/10.3389/fphys.2019.01550). [PubMed: [31992987](https://pubmed.ncbi.nlm.nih.gov/31992987/)]. [PubMed Central: [PMC6962351](https://pubmed.ncbi.nlm.nih.gov/PMC6962351/)].
- Ahmadi A, Agha AH, Gharakhanlou R, Zarifi A. Study of relationship between serum interleukin 6 (IL-6) and creatine kinase (CK) changes response in active women after sub maximal eccentric and concentric exercise. *Olympic J*. 2009;**17**:63-72.
- Proske U, Morgan DL. Muscle damage from eccentric exercise: mechanism, mechanical signs, adaptation and clinical applications. *J Physiol*. 2001;**537**(Pt 2):333-45. doi: [10.1111/j.1469-7793.2001.00333.x](https://doi.org/10.1111/j.1469-7793.2001.00333.x). [PubMed: [11731568](https://pubmed.ncbi.nlm.nih.gov/11731568/)]. [PubMed Central: [PMC2278966](https://pubmed.ncbi.nlm.nih.gov/PMC2278966/)].
- Philippou A, Maridaki M, Theos A, Koutsilieris M. Cytokines in muscle damage. *Adv Clin Chem*. 2012;**58**:49-87. doi: [10.1016/b978-0-12-394383-5.00010-2](https://doi.org/10.1016/b978-0-12-394383-5.00010-2). [PubMed: [22950343](https://pubmed.ncbi.nlm.nih.gov/22950343/)].
- Libardi CA, De Souza GV, Cavaglieri CR, Madruga VA, Chacon-Mikahil MP. Effect of resistance, endurance, and concurrent training on TNF-alpha, IL-6, and CRP. *Med Sci Sports Exerc*. 2012;**44**(1):50-6. doi: [10.1249/MSS.0b013e318229d2e9](https://doi.org/10.1249/MSS.0b013e318229d2e9). [PubMed: [21697747](https://pubmed.ncbi.nlm.nih.gov/21697747/)].
- Phillips MD, Flynn MG, McFarlin BK, Stewart LK, Timmerman KL. Resistance training at eight-repetition maximum reduces the inflammatory milieu in elderly women. *Med Sci Sports Exerc*. 2010;**42**(2):314-25. doi: [10.1249/MSS.0b013e3181b11ab7](https://doi.org/10.1249/MSS.0b013e3181b11ab7). [PubMed: [19927028](https://pubmed.ncbi.nlm.nih.gov/19927028/)].
- Levinger I, Goodman C, Peake J, Garnham A, Hare DL, Jerums G, et al. Inflammation, hepatic enzymes and resistance training in individuals with metabolic risk factors. *Diabet Med*. 2009;**26**(3):220-7. doi: [10.1111/j.1464-5491.2009.02679.x](https://doi.org/10.1111/j.1464-5491.2009.02679.x). [PubMed: [19317815](https://pubmed.ncbi.nlm.nih.gov/19317815/)].
- Monteiro R, Teixeira D, Calhau C. Estrogen signaling in metabolic inflammation. *Mediators Inflamm*. 2014;**2014**:615917. doi: [10.1155/2014/615917](https://doi.org/10.1155/2014/615917). [PubMed: [25400333](https://pubmed.ncbi.nlm.nih.gov/25400333/)]. [PubMed Central: [PMC4226184](https://pubmed.ncbi.nlm.nih.gov/PMC4226184/)].
- Selvin E, Paynter NP, Erlinger TP. The effect of weight loss on C-reactive protein: a systematic review. *Arch Intern Med*. 2007;**167**(1):31-9. doi: [10.1001/archinte.167.1.31](https://doi.org/10.1001/archinte.167.1.31). [PubMed: [17210875](https://pubmed.ncbi.nlm.nih.gov/17210875/)].
- Penninx BW, Kritchevsky SB, Newman AB, Nicklas BJ, Simonsick EM, Rubin S, et al. Inflammatory markers and incident mobility limitation in the elderly. *J Am Geriatr Soc*. 2004;**52**(7):1105-13. doi: [10.1111/j.1532-5415.2004.52308.x](https://doi.org/10.1111/j.1532-5415.2004.52308.x). [PubMed: [15209648](https://pubmed.ncbi.nlm.nih.gov/15209648/)].
- Donges CE, Duffield R, Drinkwater EJ. Effects of resistance or aerobic exercise training on interleukin-6, C-reactive protein, and body composition. *Med Sci Sports Exerc*. 2010;**42**(2):304-13. doi: [10.1249/MSS.0b013e3181b117ca](https://doi.org/10.1249/MSS.0b013e3181b117ca). [PubMed: [20083961](https://pubmed.ncbi.nlm.nih.gov/20083961/)].