Published online 2023 October 16.

Effect of Core Stability Trainings on Functional Movement Screening Scores and Trunk Muscle Endurance in Female Kyokushin Karate Athletes

Mohammad Rahimi ¹, Hadi Samadi ¹, ^{*}, Zohre Nikzade Abbasi ¹ and Afsane Rahnama ¹

¹Faculty of Sport Sciences, Shahid Rajaee Teacher Training University, Tehran, Iran

Corresponding author: Faculty of Sport Sciences, Shahid Rajaee Teacher Training University, Tehran, Iran. Email: hadi.samadi@gmail.com

Received 2023 May 29; Revised 2023 August 13; Accepted 2023 August 27.

Abstract

Background: The objective of this study was to examine the impact of six weeks of core instability training on functional movement screening (FMS) scores and core trunk endurance in female Kyokushin karate athletes.

Methods: Thirty female Kyokushin Karate athletes aged 14 - 18 years were recruited and randomly assigned to either an experimental group or a control group (n = 15 per group). Functional movement patterns, including the FMS, Sorenson, and McGill tests, were assessed before and after the intervention. The experimental group underwent six weeks of Kyokushin Karate training, while the control group performed core stability exercises. The statistical analysis involved paired *t*-tests and analysis of covariance (ANCOVA), with a significance level set at P ≤ 0.05 .

Results: The paired t-test results indicated a significant difference in pre- and post-test scores in both the control and experimental groups (P < 0.05). However, the ANCOVA showed no significant differences between the groups (P > 0.05).

Conclusions: Both Kyokushin training and core stability exercises have been found to increase core stability and FMS scores. Therefore, it can be suggested that Kyokushin Karate athletes may not need to perform separate core stability exercises as part of their training routine.

Keywords: Kyokushin Karate, Core Stability, FMS, Core Muscle Endurance, Movement Pattern

1. Background

Kyokushin karate is widely regarded as the first official full-contact karate style, founded by Mas Oyama (1, 2). The combination of aerobic, balance, and coordination exercises in Kyokushin may improve cognitive functions and motor coordination (3). Since the emphasis is on close-range fighting, it may affect body composition, the musculoskeletal system, and physical fitness (1, 4). One of the important elements contributing to better combat is having strong trunk muscles. The core can be described as a muscular power box, with abdominal muscles, paraspinal and gluteal muscles in the back, the diaphragm as the roof, and pelvic floor muscles and pelvic girdle as the floor (4). The core muscles can be considered a muscle belt to stabilize the spine (5). Kibler et al. defined core stability as the ability to control the position and motion of the trunk over the pelvis to allow for optimum

production, transfer, and control of force and motion to the terminal segments in integrated athletic activities (6). Core stability, strength, and endurance ensure spine stability for force production. Numerous authorities assert that incorporating core stability exercises into one's workout routine can improve motor control and functional movement patterns, ultimately leading to enhanced athletic performance (7).

The core exercise increases strength, endurance, and neuromuscular control of the trunk; it also reduces ligament tension and stabilizes the vertebral column in a normal anatomical position. Moreover, it can improve spine control, intra-abdominal pressure, and trunk movements (8). Muscle dysfunction, including loss of strength and endurance in the deep stabilizer muscle's, may lead to incorrect movement patterns and make athletes prone to injury (9). Optimal core function involves both trunk mobility and stability. When the

Copyright © 2023, Journal of Clinical Research in Paramedical Sciences. This is an open-access article distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (CC BY-NC 4.0) (https://creativecommons.org/licenses/by-nc/4.0/) which permits copy and redistribute the material just in noncommercial usages, provided the original work is properly cited. core is functioning efficiently, normal length-tension relationships are maintained, allowing the athlete to produce powerful movements in the extremities (10). Core stability exercises may also affect trunk neuromuscular stability and greatly influence upper and lower extremity performance (6). Core endurance is defined as the ability to maintain a position or perform multiple repetitions. If the core muscular system does not have enough stability to maintain the vertebral column in a normal position, then trunk motion will be distorted, and the risk of instability and postural misalignment may be increased (11). Therefore, optimal core stability can prevent postural disorders and increase physical performance (12).

Functional movement screening (FMS) is a tool used to identify movement deficiencies, instability, or limitations that may increase the risk of injury for athletes. Functional movement screening evaluates the coordination, flexibility, and endurance of the trunk (13), and has been used to predict the risk of injury in various sports (14, 15). There are several types of training methods aimed at improving sports performance (16), and several researchers have reported the importance of core stability training in lower extremity movements (17, 18). To improve the performance of the trunk muscles, stabilization training is generally employed along with muscle strength and endurance trainings (17, 18). However, Akuthota and Nadler found no significant relationship between core stability and FMS scores (19). Despite this, literature reviews have shown that a number of proprioceptive training programs, along with balance, strength, and specific exercise training for a particular sport, can increase the FMS score. For example, Ciemiński reported that a ten-day proprioceptive training program increased the FMS score in young female volleyball players (20).

Due to the high prevalence of sport-related injuries among athletes, it is necessary to conduct reliable assessments of physical fitness and functional condition. Previous research studies have shown that FMS can be a reliable tool for predicting sports-related injuries. A lower FMS score can predict injuries in athletes. However, no research has been found on the effect of core stability exercise on FMS in Kyokushin karate athletes.

2. Objectives

The main purpose of this study was to investigate the effect of core stability exercise on the FMS score and trunk muscle endurance in female Kyokushin karate athletes.

3. Methods

This semi-experimental research involved 30 female participants aged 14 - 18, were recruited and randomly assigned to either an experimental group or a control group. The statistical population of the study was female Kyokushin Karate in Karaj city. Participants in the control group (n = 15) were included in the Kyokushin Karate special exercises, while those in the experimental group (n = 15) were included in the core stability training exercises in addition to the s Kyokushin Karate special exercises. The inclusion criteria include regular participation in karate for least two years, be healthy women with no recent surgery on extremities or neuromuscular diseases within the last 6 months, and not have had any serious injuries within the past six months. Exclusion criteria included missing three or two consecutive training sessions or having an obvious musculoskeletal deformity or injury during the study (21).

In this study, the sample size was tested with g power analysis at the beginning of the study. G power was determined with α at 0.05 and 1- β error probability is 0.80 for the 30 participants (22). We performed an a priori estimation of power and sample size through the G-Power software (version 3.1.9.4) programme written by Kiel University, made in Germany.

3.1. Functional Movement Screen

A functional assessment was conducted using the FMS test. In adolescents aged 11 to 15 years, the intra-rater reliability and inter-rater reliability of the FMS were reported to be 0.11 to 0.83 and 0.23 to 0.88, respectively, by Wright et al. (23). Parenteau et al. reported an FMS score of 0.96 (confidence interval 0.92 - 0.98) in 28 youth hockey players aged 13 - 16 (24). The FMS test is a quick and easy-to-administer test that can be completed in 5 -10 minutes by coaches. The test comprises seven physical tasks, each component test was scored on an ordinal scale (0 to 3 points), based on the quality of movement, with 3 being the maximum score. A score of 2 indicated that the participant required some type of compensation or was unable to complete the entire movement. A score of 1 was given if the individual was unable to remain in the movement position throughout the movement, lost balance during the test, or did not meet the minimum criteria to score a 2. Pain during any of the FMS component tests or during any of the clearing tests indicated a score of 0(14). The seven tasks in the FMS test are deep squat, hurdle step, in-line lunge, shoulder mobility, active straight leg raise, trunk stability push-up, and rotary stability. The maximum total score for the test is 21, and a score below 14 indicates susceptibility to injury (25).

3.2. Evaluation of the Core Muscles Endurance

The endurance of the core muscles was evaluated using the Biering-Sorenson and McGill tests. Each test was performed three times with a 5-minute rest period in between, and the assessment was conducted in all anatomical planes, including posterior, anterior, and lateral (26). To evaluate the endurance of the anterior trunk flexor muscles, the trunk flexor test was performed, and the time was recorded. Similarly, the trunk extensor test was conducted to measure the endurance of the posterior trunk muscles. To evaluate the endurance of the lateral trunk flexor muscles, the side plank test was performed, and the time was recorded (Figure 1). The reliability of the trunk flexor test, the trunk extensor test and the side plank test have been reported 0.93, 0.98, and 0.95, respectively (27).

3.3. Exercise Training Program

The training sessions took place at the Azadi gym in Karaj city. The training sessions were conducted under the supervision of an instructor with a first-class national coaching degree in this field led the training sessions and was present to supervise all exercises in three sessions per week. The Kyokushin Karate special exercises in this study were designed using the help of Mitt Pa, boxing bags, and other necessary equipment in this field. The Kyokushin Karate special exercises included kicking, punching, striking, and blocking moves in both stationary and variable positions. The researcher, with a first-class national coaching degree and 14 years of coaching experience, compiled the Kyokushin karate exercises in the study. Each session began with a 10 min warm-up program, followed by regular Kyokushin Karate special exercises for approximately 40 - 50 min 3 times per week, for 6 weeks. The participants in the experimental group was performed core stability training for approximately 20 - 30 in addition to the Kyokushin Karate special exercises. The core stability exercises included lower abdominal exercises, half sit-ups, side bridge, sit-ups with rotation, and prone bridge (Table 1). The exercise intensity was gradually increased. Proper breathing patterns and correct posture were emphasized and taught before the start of each session. Each subject underwent a secondary measurement (post-test) up to one week after finishing the exercises. The conditions of each secondary measurement session were exactly the same as

the primary measurement session and all them conducted by the same researcher.

Normality of data distribution was evaluated using the Shapiro-Wilk test. To assess the effect of the procedure, differences between all variables in the experimental and control groups were analyzed using paired sample t-tests. The analysis of covariance (ANCOVA) was used to further analyze the results, with a significance level set at P \leq 0.05.

4. Results

Table 2 presents the characteristics of both groups, and no significant differences were found between the two groups (P > 0.05). The paired *t*-test was used for within-group comparisons, and the ANCOVA test was used for between-group comparisons of the study variables in the experimental and control groups (Tables 3 and 4).

5. Discussion

The objective of this study was to investigate the effect of core stability exercises on the FMS score and trunk muscle endurance in Kyokushin karate players. The findings revealed significant improvements in the core stability and FMS scores in both the experimental and control groups (P \geq 0.05). There were no statistically significant differences between the groups in the post-test outcomes (P > 0.05). Therefore, it can be concluded that Kyokushin karate training and core stability exercises have similar effects on core muscle endurance and FMS score in Kyokushin karate players.

Kyokushin karate is renowned as a full-contact style and is considered one of the toughest forms of karate (28). In this style, competitors engage in fights without any weapons using maximum speed and strength (29). Due to this close-range fighting style, it affects the body composition and musculoskeletal system (1, 4). Karate training demands high-level functional abilities, including speed, muscle strength, and coordination (6, 7). Rzepko found that Kyokushin karate training can improve balance and posture in teenage karate athletes (30). Additionally, research has demonstrated that karate training can enhance neuromuscular control strategy and movement patterns (31).

Core stability exercises are effective in enhancing physical fitness by promoting a neutral spine position during daily activities and increasing spinal muscle endurance (32). Core stability involves the control



Figure 1. Left to right - Biering-Sorenson test, trunk flexor test, and side plank test

Table 1. Core Stability Training Protocol

Stages	Exercise Trainings
Week 1	Lower abdominal exercises, half sit-ups, side bridge, sit-up with rotation (2 sets, 10 reps)
Week 2	Lower abdominal exercises, half sit-ups, side bridge, sit-up with rotation (3 sets, 10 reps)
Week 3	Lower abdominal exercises (2 sets, 10 reps), half sit-ups, side bridge, sit-up with rotation (2 sets, 15 reps)
Week 4	Lower abdominal exercises, half sit-ups, side bridge, sit-up with rotation (2 sets, 15 reps), prone bridge (2 sets in 10 s)
Week 5	Lower abdominal exercises (2 sets, 10 reps), half sit-ups, side bridge, sit-up with rotation (2 sets, 20 reps), prone bridge (3 sets in 10 s)
Week 6	Lower abdominal exercises (3 sets, 10 reps), half sit-ups, side bridge, sit-up with rotation (2 sets, 20 reps), prone bridge (2 sets in 15 s)

Table 2. Descriptive Characteristics of Subjects in the Research Groups at Baseline $\left(\text{Pre-test}\right)^a$

Variables	Mean \pm Standard Deviation			
Control				
Height (cm)	158.27 ± 5.32			
Weight (kg)	55.53 ± 6.18			
Age (y)	16.53 ± 1.45			
BMI (kg.m ⁻²)	22.11 ± 1.42			
Experimental				
Height (cm)	157.40 ± 5.48			
Weight (kg)	56.60 ± 5.56			
Age (y)	16.60 ± 1.24			
BMI (kg.m ⁻²)	22.78 ± 0.84			

 a No significant difference was found between the 2 groups in terms of outcome measures (P $> \,$ 0.05).

of the hip and pelvic girdle in three different levels, including local vertebral control, lumbar-pelvic control, and postural control, all of which affect the kinetic chain. Impairment of vertebral and lumbar-pelvic control can negatively impact postural control and body balance. Poor postural control can lead to falls, increasing the risk of injury. Peate et al. found that Swiss ball exercises have a positive effect on improving core strength (33). The role of trunk stabilizers is to provide stability, support, and control to the trunk or core region of the body and improve overall functional performance, enhance athletic performance, reduce the risk of injuries, and support proper movement mechanics.

Poor endurance of core muscles can lead to fatigue during intensive training, reducing coordination and muscle strength. If the core muscle strength-to-endurance ratio becomes greater than four, it is likely to result in lower back pain. There is no linear relationship between strength and endurance, and with an increase in strength, the endurance increases to some extent, after which it remains constant. As a result, the reduction of trunk muscle endurance may follow a reduction in muscle strength, or vice versa, reduced strength may also affect the strength of the thigh muscles, which is one of the effective and integral factors of core stability (6). Mitchell et al. found a positive relationship between FMS scores and core muscle endurance in children (34). Core muscle weakness can decrease core muscle force production in limbs, which affects the kinetic chain (35). FMS is a popular screening tool in sports medicine that evaluates muscle strength, balance, core stability, coordination, motor control, flexibility, range of motion, and kinetic chain (13). The FMS test focuses on motor limitations and asymmetry. A high FMS score indicates high core stability and balance. Additionally, the FMS score can demonstrate altered movement patterns and bilateral asymmetries in movement, and athletes with lower FMS scores are more prone to injury. Kiesel et al. showed that athletes with FMS scores of less than 14 have a 1.78 times greater risk

Table 3. Results of Paired t-Tests for Within-Group Differences								
Groups		Mean \pm Standard Deviation		P.Value				
		Post-test Pre-test		1 Value				
Control								
	Right plank	40.48 ± 14.98	33.14 ± 15.57	0.001				
	Left plank	41.80 ± 13.37	34.05 ± 14.28	0.001				
	Flexion	42.57±15.63	34.36 ± 13.92	0.001				
	Sorenson	49.61 ± 16.47	42.91 ± 16.02	0.001				
	FMS	17.13 ± 1.40	16.13 ± 1.24	0.001				
Expe	rimental							
	Right plank	33.88 ± 8.96	24.71± 7.33	0.001				
	Left plank	33.46 ± 7.24	24.03 ± 7.62	0.001				
	Flexion	36.32 ± 10.01	27.60 ± 10.09	0.001				
	Sorenson	43.86 ± 15.13	35.21±13.36	0.001				
	FMS	17.20 ± 1.21	15.93 ± 1.33	0.001				

Table 4. The Results of ANCOVA for Between-Group Differences ^a										
Variables	Sum of Squares	Degree of Freedom	Mean Squares	F	Р	Effect Size	Test Power			
Right plank	15.82	1	15.81	1.05	0.31	0.04	0.17			
Left plank	2.01	1	2.01	0.16	0.69	0.006	0.07			
Flexion	4.16	1	4.16	0.31	0.58	0.01	0.08			
Sorenson	37.43	1	37.43	3.07	0.09	0.10	0.39			
FMS	0.39	1	0.39	0.61	0.44	0.02	0.12			

^a There were no significant differences in any of the variables between the experimental and control groups (P > 0.05).

of injury (13). Similarly, Lehr et al. found that the high-risk-injury group had lower FMS scores compared to the low-risk-injury group, with a 3.4 times higher risk of injury (36).

The current research has some limitations. The Kyokushin karate special exercises was not monitored, which does not allow for how the control and experimental groups trained besides for the core intervention. Moreover, in this study focuses only on female Kyokushin karate athletes aged 14 - 18. Therefore, the results may not be applicable to male athletes or individuals participating in different sports and age categories. Furthermore, a small sample size or a specific sample demographic may limit the generalizability of the study's findings to a broader population of Kyokushin karate athletes or other sports. Despite the limitations, this study presents some evidence that Kyokushin karate special exercises can enhance improvements in core endurance and FMS scores in female Kyokushin karate athletes.

5.1. Conclusions

Both Kyokushin training and core stability exercises have been found to increase trunk stability and FMS scores. Therefore, it can be suggested that Kyokushin karate athletes may not need to perform separate core stability exercises as part of their training routine.

Footnotes

Authors' Contribution: M Rahimi, study concept and design, critical revision; H Samadi, statistical analysis; Z Nikzade Abbasi, collected the data; A Rahnama, drafting of the manuscript.

Conflict of Interests: The authors have no conflict of interest.

Data Reproducibility: The dataset presented in the study is available on request from the corresponding author during submission or after publication.

Ethical Approval: The ethics are approved by the Committee of the Research Institute of Physical

Education and Sports Sciencesthe ethics code number of SSRI.REC-2112-1421(R1) was received.

Funding/Support: The costs of this research were covered by the authors.

Informed Consent: Informed consent was obtained.

References

- 1. Drzał-Grabiec J, Truszczyńska A. Evaluation of selected postural parameters in children who practice kyokushin karate. *Biomed Hum Kinet*. 2014;**6**(1). https://doi.org/10.2478/bhk-2014-0013.
- Ali N, Rouhi G, Robertson G. Gender, Vertical Height and Horizontal Distance Effects on Single-Leg Landing Kinematics: Implications for Risk of non-contact ACL Injury. J Hum Kinet. 2013;37:27-38. [PubMed ID: 24146702]. [PubMed Central ID: PMC3796838]. https:// doi.org/10.2478/hukin-2013-0022.
- 3. Zabjek M. Reconsidering the Relationship between Japanese Martial arts and Religion: Case Study of Mt. Mitsumine and Kyokushin Karate. "Religious Cultures in Asia: Mutual Transformations through Multiple Modernities" International Research Forum. Tokyo, Japan. Institute for Japanese Culture and Classics; 2020. 39 p.
- 4. Kuznetsova Z, Kalinina I, Dyakova Y, Polovnikova M, Kudryashova Y, Polovnikova M, et al. Morphotypological characteristics of primary school children engaged in karate Kyokushin. *First International Scientific-Practical Conference "Actual Issues of Physical Education and Innovation in Sports" (PES 2020)*. Naberezhyne Chelny, Russia. BIO Web of Conferences; 2020. 75 p.
- 5. Tse MA. Exploring the impact of core stability on performance [dissertation]. Hong Kong, Japan: University of Hong Kong; 2009.
- Kibler WB, Press J, Sciascia A. The role of core stability in athletic function. *Sports Med*. 2006;**36**(3):189–98. [PubMed ID: 16526831]. https: //doi.org/10.2165/00007256-200636030-00001.
- Hill J, Leiszler M. Review and role of plyometrics and core rehabilitation in competitive sport. *Curr Sports Med Rep.* 2011;10(6):345-51. [PubMed ID: 22071395]. https://doi.org/10.1249/ [SR.0b013e31823b3b94.
- McCaskey A. The effects of core stability training on star excursion balance test and global core muscular endurance [master's degree]. Toledo, Ohio: University of Toledo; 2011.
- Sato K, Mokha M. Does core strength training influence running kinetics, lower-extremity stability, and 5000-M performance in runners? J Strength Cond Res. 2009;23(1):133-40. [PubMed ID: 19077735]. https://doi.org/10.1519/JSC.0b013e31818eb0c5.
- Lederman E. The myth of core stability. J Bodyw Mov Ther. 2010;14(1):84–98. [PubMed ID: 20006294]. https://doi.org/10.1016/ j.jbmt.2009.08.001.
- Akuthota V, Ferreiro A, Moore T, Fredericson M. Core stability exercise principles. *Curr Sports Med Rep.* 2008;7(1):39–44. [PubMed ID:18296944]. https://doi.org/10.1097/01.CSMR. 0000308663.13278.69.
- Allen BA, Hannon JC, Burns RD, Williams SM. Effect of a core conditioning intervention on tests of trunk muscular endurance in school-aged children. J Strength Cond Res. 2014;28(7):2063-70. [PubMed ID: 24378666]. https://doi.org/10.1519/ JSC.0000000000000352.
- Kiesel K, Plisky PJ, Voight ML. Can Serious Injury in Professional Football be Predicted by a Preseason Functional Movement Screen? *N Am J Sports Phys Ther.* 2007;2(3):147–58. [PubMed ID: 21522210]. [PubMed Central ID: PMC2953296].

- Cook G, Burton L, Hoogenboom B. Pre-participation screening: the use of fundamental movements as an assessment of function part 1. *N Am J Sports Phys Ther*. 2006;1(2):62–72. [PubMed ID: 21522216]. [PubMed Central ID: PMC2953313].
- Cook G, Burton L, Hoogenboom B. Pre-participation screening: the use of fundamental movements as an assessment of function - part
 N Am J Sports Phys Ther. 2006;1(3):132–9. [PubMed ID: 21522225]. [PubMed Central ID: PMC2953359].
- Girold S, Maurin D, Dugue B, Chatard JC, Millet G. Effects of dry-land vs. resisted- and assisted-sprint exercises on swimming sprint performances. *J Strength Cond Res.* 2007;21(2):599–605. [PubMed ID: 17530963]. https://doi.org/10.1519/R-19695.1.
- Leetun DT, Ireland ML, Willson JD, Ballantyne BT, Davis IM. Core stability measures as risk factors for lower extremity injury in athletes. *Med Sci Sports Exerc.* 2004;36(6):926-34. [PubMed ID: 15179160]. https://doi.org/10.1249/01.mss.0000128145. 75199.c3.
- Willson JD, Ireland ML, Davis I. Core strength and lower extremity alignment during single leg squats. *Med Sci Sports Exerc.* 2006;38(5):945-52. [PubMed ID: 16672849]. https://doi.org/10.1249/01. mss.0000218140.05074.fa.
- Akuthota V, Nadler SF. Core strengthening. Arch Phys Med Rehabil. 2004;85(3 Suppl 1):S86–92. [PubMed ID: 15034861]. https://doi.org/10. 1053/j.apmr.2003.12.005.
- Ciemiński K. The influence of 10-day proprioceptive training on the FMS test results in young female volleyball players-a pilot study. Age (years). 2018;15:0.95.
- Boling M, Padua D, Blackburn J, Petschauer M, Hirth C. Hip Adduction Does not Affect VMO EMG Amplitude or VMO:VL Ratios during a Dynamic Squat Exercise. J Sport Rehabil. 2006;15(3):195–205. https:// doi.org/10.1123/jsr.15.3.195.
- Kabadayi M, Karadeniz S, Yilmaz AK, Karaduman E, Bostanci O, Akyildiz Z, et al. Effects of Core Training in Physical Fitness of Youth Karate Athletes: A Controlled Study Design. *Int J Environ Res Public Health*. 2022;**19**(10). [PubMed ID: 35627351]. [PubMed Central ID: PMC9141055]. https://doi.org/10.3390/ijerph19105816.
- Wright MD, Portas MD, Evans VJ, Weston M. The effectiveness of 4 weeks of fundamental movement training on functional movement screen and physiological performance in physically active children. *J Strength Cond Res.* 2015;**29**(1):254–61. [PubMed ID: 25072666]. https: //doi.org/10.1519/JSC.000000000000602.
- Parenteau GE, Gaudreault N, Chambers S, Boisvert C, Grenier A, Gagne G, et al. Functional movement screen test: a reliable screening test for young elite ice hockey players. *Phys Ther Sport*. 2014;15(3):169–75. [PubMed ID: 24291023]. https://doi.org/10.1016/j.ptsp.2013.10.001.
- Chorba RS, Chorba DJ, Bouillon LE, Overmyer CA, Landis JA. Use of a functional movement screening tool to determine injury risk in female collegiate athletes. *N Am J Sports Phys Ther.* 2010;5(2):47-54. [PubMed ID: 21589661]. [PubMed Central ID: PMC2953387].
- Nesser TW, Huxel KC, Tincher JL, Okada T. The relationship between core stability and performance in division I football players. *J Strength Cond Res*. 2008;22(6):1750–4. [PubMed ID: 18978631]. https://doi.org/10. 1519/JSC.0b013e3181874564.
- McGill SM, Childs A, Liebenson C. Endurance times for low back stabilization exercises: clinical targets for testing and training from a normal database. Arch Phys Med Rehabil. 1999;80(8):941-4. [PubMed ID: 10453772]. https://doi.org/10.1016/s0003-9993(99)90087-4.
- Woodward TW. A review of the effects of martial arts practice on health. WMJ. 2009;108(1):40–3. [PubMed ID: 19326635].
- 29. Saienko V. Improvement and control of the development level of special endurance in athletes of high qualification in kyokushin kaikan karate. *Int J Pharm Technol*. 2016;**8**(3):18026–42.

- 30. Rzepko M. The influence of training on static and functional balance in adolescent karateka. *Sci Martial Arts.* 2019;**15**:275.
- Blaszczyszyn M, Szczesna A, Pawlyta M, Marszalek M, Karczmit D. Kinematic Analysis of Mae-Geri Kicks in Beginner and Advanced Kyokushin Karate Athletes. *Int J Environ Res Public Health*. 2019;**16**(17). [PubMed ID: 31470588]. [PubMed Central ID: PMC6747486]. https:// doi.org/10.3390/ijerph16173155.
- Reimann F, Maziarz M, Flock G, Habib AM, Drucker DJ, Gribble FM. Characterization and functional role of voltage gated cation conductances in the glucagon-like peptide-1 secreting GLUTag cell line. J Physiol. 2005;563(Pt 1):161–75. [PubMed ID:15611035]. [PubMed Central ID: PMC1665554]. https://doi.org/10.1113/jphysiol.2004.076414.
- Peate WF, Bates G, Lunda K, Francis S, Bellamy K. Core strength: a new model for injury prediction and prevention. J Occup Med Toxicol. 2007;2:3. [PubMed ID: 17428333]. [PubMed Central ID: PMC1865378].

https://doi.org/10.1186/1745-6673-2-3.

- 34. Mitchell UH, Johnson AW, Adamson B. Relationship between functional movement screen scores, core strength, posture, and body mass index in school children in Moldova. J Strength Cond Res. 2015;29(5):1172-9. [PubMed ID: 25719919]. https://doi.org/10.1519/JSC.000000000000722.
- Tse MA, McManus AM, Masters RS. Development and validation of a core endurance intervention program: implications for performance in college-age rowers. J Strength Cond Res. 2005;19(3):547-52. [PubMed ID: 16095402]. https://doi.org/10.1519/15424.1.
- Lehr ME, Plisky PJ, Butler RJ, Fink ML, Kiesel KB, Underwood FB. Field-expedient screening and injury risk algorithm categories as predictors of noncontact lower extremity injury. *Scand J Med Sci Sports.* 2013;23(4):e225–32. [PubMed ID: 23517071]. https://doi.org/10. 1111/sms.12062.