



The Effect of Circuit Resistance Training on Plasma Concentration of Endothelin-1, Nitric Oxide and Vascular Diameter in Elderly Men

Hassan Sanian¹, Hassan Matinhomaei^{2,*} and Maghsoud Peeri²

¹Faculty of Physical Education and Sport Sciences, Central Tehran Branch, Islamic Azad University, Tehran, Iran

²Department of Exercise Physiology, Faculty of Physical Education and Sport Sciences, Central Tehran Branch, Islamic Azad University, Tehran, Iran

*Corresponding author: Associate Professor, Department of Exercise Physiology, Faculty of Physical Education and Sport Sciences, Central Tehran Branch, Islamic Azad University, Tehran, Iran. Email: hasan.matinhomaei@gmail.com

Received 2018 June 17; Revised 2019 January 01; Accepted 2019 January 05.

Abstract

Background: aging is associated with an increase in vascular dysfunction but it can be prevented by exercise training. However, the effect of circuit resistance training as a useful training method in improving cardio-respiratory and muscular strength is unclear.

Objectives: The aim of this study was to investigate the effect of CRT on plasma endothelin-1, nitric oxide (NO) and vascular size in aged men.

Methods: Twenty healthy aged men (55 ± 5 year) were randomly divided into control and CRT groups. CRT group performed circuit resistance training with 40 - 60 one repetition maximum for twelve weeks (3 times per week). Vascular size, plasma endothelin-1 and NO, blood pressure and heart rate were measured forty eight hours before and after the training period.

Results: CRT increased the vascular size and decreased endothelin-1 and systolic blood pressure compared to the pre-test; however, it had no effect on NO concentration and heart rate. In comparison with the control group, CRT significantly increased the vascular size and decreased endothelin-1 and blood pressure but there was no significant difference in NO concentration and heart rate between the two groups.

Conclusions: Circuit resistance training as a proper training method could induce the increase in endothelial function in aged individual and prevents vascular dysfunction related to aging.

Keywords: Circuit Resistance Training, Endothelin-1, Nitric Oxide, Vascular Diameter

1. Background

In industrialized countries, vascular diseases such as myocardial infarction, heart attack, and obesity become more prevalent with age (1). The prevalence of these diseases is associated with increased activity of the endothelial system during aging (1). At endothelial level, aging is associated with increased endothelial cell disorders, decreased endothelium-dependent vasodilation, increased endothelium-dependent vasodilatation, and decreased nitric oxide activity (2). Endothelin-1, an endothelium-derived vasoconstrictor, affects the function of various tissues such as the heart, pancreas and the nervous system (3). Endothelin causes increased vascular contraction, increased cell proliferation, blood coagulation, and inflammation (4) and is associated with the development of atherosclerosis and increased blood pressure with aging (5, 6) which can play an important role in disease and aging processes. However, increase in other substrates can inhibit the harmful effects of endothelin-1. Nitric oxide is

one of these inhibitors which acts as a vasodilator and anti-inflammatory and anticoagulant substrate (7). The imbalance between nitric oxide and endothelin-1 plays a role in the pathology of renal and pulmonary and previous diseases (5).

Exercise is recommended as a treatment for improvement of vascular endothelial dysfunction due to age (7, 8). Regular exercise usually inhibits endothelin-1 expression by increasing nitric oxide bioavailability (7). It has been shown that aerobic exercise reduces arterial stiffness in patients with hypertension (9, 10). On the other hand, high-intensity resistance exercises lead to increased arterial stiffness (11) but low-intensity exercises reduce arterial stiffness (12). Although resistance training is recommended by the American College of Sports Medicine as a necessary exercise for individuals (13), its high intensity can have a detrimental effect on vascular function (3). In this regard, it has been shown that moderate-intensity resistance training increases nitric oxide in elderly people but has no effect on the plasma concentration of endothelin-1 (14). It

has also been reported that moderate-intensity resistance training has no effect on the stiffness of arteries in the elderly, but aerobic exercise reduces it (15). Therefore, it appears that the training model can be one of the factors affecting endothelial function. Resistance exercises have been shown to increase arterial stiffness, but in combination with aerobic training, arterial stiffness decreases (16). Among various training models, circuit resistance training as a training model has the effects of both aerobic training and resistance training (17). However, the effect of this type of exercise on factors affecting vascular function in the elderly is still unknown.

2. Objectives

The purpose of this study was to investigate the effect of circuit resistance training on endothelin-1 and nitric oxide plasma concentrations in the elderly.

3. Methods

The population of this study was elderly volunteers. They were selected through call in public and administrative centers. Twenty healthy elderly men with an average age of 55 years were randomly divided into two control and circuit resistance training groups. Then, after measuring 1 maximum repetition, the resistance training group performed 10 movements (bench press, pec deck flies, barbell curls, low pulley curls, pushdowns, back lat pulldowns, seated rows, leg extensions, lying leg curls and, angled leg press) with the intensity of 40% - 65% 1maximum repetition 3 sessions per week for 3 months. The control group continued their routine daily activities during the three-month study period. The vascular diameter was measured 48 hours before the exercise program and 48 hours after the last training session using the SUN Doppler method. To perform sono-doppler, every participant was laid flat to the back on the bench with hand located beside the body and a qualified cardiologist performed the test using a 5 - 13 MHz (linear) multi-frequency probe. Blood samples were collected from the brachial vein before and 48 hours after the last training session, and after plasma isolation, the samples were stored at -20°C for measuring levels of Endothelin-1 and nitric oxide.

3.1. Statistical Method

Data analysis was done using SPSS 21 software. After verifying data normality, using Shapiro-Wilk test, a *t*-test was used to compare intra-group changes. Independent *t*-test was used to compare pre-test and post-test changes between the two groups. The significance level was 0.05.

4. Results

An intra-group analysis showed no significant difference between the pre-test and post-test in the control group ($P < 0.05$) in terms of vascular diameter, endothelin-1 plasma concentration, nitric oxide, systolic blood pressure and heart rate. However, the systolic blood pressure decreased significantly ($P = 0.44$) (Table 1 and Figure 1).

The results of intra-group analysis in circuit resistance training group showed that, compared to the pre-test, exercise significantly increased vascular diameter ($P = 0.021$), and decreased endothelin-1 ($P < 0.001$), and systolic blood pressure ($P < 0.001$); however, it has no significant effect on plasma concentration of nitric oxide, diastolic blood pressure and heart rate ($P < 0.05$) (Table 1 and Figure 1).


The results of the inter-group analysis showed that resistance training significantly increased the vascular diameter in the resistance training group compared to the control group ($P = 0.11$). Circuit resistance training also significantly decreased endothelin-1 ($P < 0.001$) and systolic blood pressure ($P < 0.001$) of the resistance training group compared with control group. However, no significant difference was found between the two groups in plasma concentration of nitric oxide, diastolic blood pressure and heart rate ($P < 0.05$) (Table 1 and Figure 1).

5. Discussion

Elderly is not a disease, but is a biological process associated with changes and problems in the body, including cardiovascular disease, hypertension and coronary artery problems (18). In general, inappropriate performance of endothelial cells has been observed to be associated not only with diseases such as high blood pressure, high cholesterol, and atherosclerosis but it is also associated with age. Vascular endothelial cells play an important role in regulating vascular activity by producing vascular activating agents, such as endothelin-1 and nitric oxide (19). Endothelin-1 is the strongest vascular vasoconstrictor, that its contractile effect is ten times higher than angiotensin-2, vasopressin and neuropeptide Y. It acts through two receptors A and B that located in the membrane of the cell (19). In relation to the effect of physical activity on endothelin-1, it has been determined that exercise increases endothelin B receptors in smooth muscle and ultimately bronchial contraction. Also, endothelin B receptor in endothelial cells leads to vasodilatation by increasing nitric oxide. Type A receptors are found mostly in cardiac muscle and vascular smooth muscle, and the main mediator of vascular contraction is by endothelin, while type B is found in the kidney, uterus, central nervous system and vascular endothelial cells, and Its stimulation produces nitric oxide (20).

Table 1. Blood Pressure and Heart Rate Changes in the Control and Circuit Resistance Training Groups^a

	Control Group		Training Group	
	Pre	Post	Pre	Post
Systolic blood pressure	125 ± 0.31	123 ± 0.44	142.4 ± 0.57	127 ± 0.53
Diastolic blood pressure	88 ± 0.41 ^b	83.9 ± 0.36	100 ± 0.71 ^{b,c}	95.5 ± 0.43
Heart rate	83.6 ± 2.01	85 ± 1.67	82.6 ± 3.03	81.9 ± 3.12

^aValues are expressed as mean ± SD.^bSignificant difference with post-test.^cSignificant difference between the two groups ($P < 0.05$).**Figure 1.** Changes in vascular diameter and endothelin-1 and nitric oxide plasma concentrations in circuit resistance training and control groups (mean ± SD). * Significant difference with pre-test, \$ significant difference with the control group ($P < 0.05$).

Various factors contribute to the production of this index various factors in the production of this index include rhe-

ological and neurohumorrhagic factors which these factors are also affected by exercise. Endothelin-1 produces ni-

tric oxide and eventually artery vasodilatation with effect on type B receptors (19).

Aging is associated with elevated endothelin-1 plasma concentration levels (21) which increases the contraction tone in the peripheral muscle in the elderly (22). Endothelin causes increased vascular contraction, increased cell proliferation, blood coagulation, and inflammation (4) and is associated with the development of atherosclerosis and increased blood pressure with aging (5, 6). Therefore, it can play an important role in increasing blood pressure. But exercise lowers endothelin-1 plasma and intramuscular levels in the elderly (23), which is consistent with the findings of the present study. The results of this study indicated that circuit resistance training can decrease endothelin-1 vasoconstrictive plasma concentration and systolic blood pressure in elderly people. In addition, it increased the vascular diameter but had no effect on nitric oxide plasma concentration. Contrary to this study, Maeda et al. reported that moderate-intensity resistance exercises increases nitric oxide levels in the elderly, but has no effect on endothelin-1 plasma concentrations (14). However, in agreement with the current study, one study showed a decrease in the concentration of endothelin-1 in healthy young people (24). Also, decreased endothelin-1 concentration has been reported as a result of aerobic exercise (25).

In addition, it has been reported that exercise lowers vascular tone induced by endothelin-1 (22). In this study, the plasma concentration of nitric oxide increased but not significantly. One of the reasons for lack of change in nitric oxide can be due to a significant decrease in endothelin-1 since these two substrates are opposed to each other. Given that decreased endothelin-1 lowered its contractile effect on the arteries, there was no need for a compensatory mechanism to increase nitric oxide for vasodilatation. The decrease in endothelin-1 was also associated with a reduction in systolic blood pressure. Hence, with a reduction in endothelin-1, vascular smooth muscle might have returned to their resting condition, resulting in a reduction in systolic blood pressure. Research has also shown that circuit resistance training, especially when light weights used (40% - 60% 1MR), are useful for controlling body weight (17). In addition, during circuit resistance exercises, heart rate, metabolic cost, and energy consumption are higher than those in traditional resistance exercises (26) and can be a desirable exercise strategy to increase cardiovascular fitness and strength (17). Results of this study showed that circuit resistance training can increase vascular diameter and thereby contribute to increased cardiac respiratory endurance.

The major mechanism of the main changes of the effect of vascular training is not clear, but it has been

shown that 12-week circuit resistance training reduces arterial stiffness without altering blood pressure in the elderly (27). Previous studies have reported that changes in the endothelium of arteries are due to chronic changes in endothelium- and nitric oxide-induced blood flow (28, 29). As the current study showed, the reduction of arterial stiffness due to exercise was because of an increase in vascular diameter and a decrease in endothelin-1 concentration, which reduces its contractile effect. In addition, it has been shown that muscle fibers affect the sensitivity of endothelin-1 so that aging is associated with a decrease in endothelin sensitivity in fast twitch fibers (20). In the current study, the resistance type of exercise was used that could have a positive effect on increasing fast contractions. Therefore, in this study, the increased sensitivity of endothelin-1 is one of the reasons for the decrease of endothelin concentration, but this finding requires more research because due to limitation, the percentage of the fibers were not examined in this study.

The exact mechanism of nitric oxide and endothelin-1 changes is not clear from the exercise, but it has been shown that increased blood flow during the exercise can increase the shear pressure to the vascular wall, which can change the expression of genes in the endothelial (30, 31). An increase in shear pressure leads to an increase in nitric oxide and a decrease in endothelin-1 (32, 33). One of the possible reasons for lack of change in the plasma concentration of nitric oxide can be due to its base level. People with endothelial dysfunction have been reported to be more exposed to nitric oxide performance but it has no effect on people with normal endothelial function (34). Therefore, in the present study, the participants might have a normal endothelial function with no need for an increase in nitric oxide concentration, and that only reduction of endothelin-1 has been able to increase endothelial function and subsequently decrease systolic blood pressure. It has been shown that low-intensity resistance training inhibits hypertension (35), which is in agreement with the findings of the present study.

In this regard, Thijssen et al. In relation to the role of endothelin-1, on the contraction of the elderly's leg arteries, showed that increasing the cross-sectional area in leg vascular contraction with age, partly done by endothelin. Also 8 weeks of bicycle training in elderly people without mobility reduced leg vascular contraction and somewhat decreased endothelin-1 (22). In examining the decrease in the diameter of the blood vessels with endothelin-1 and its increase with age in healthy people and reducing it by exercise, Van Guilder et al. found that endothelin-1 increased blood pressure by decreasing vascular diameter; This index increases with age and decreases due to the regular aerobic exercise activity (36).

In one study on the effects of short-term leg resistance training on arterial performance in older men, artery stiffness did not change with resistance training, and plasma nitric oxide concentration increased after these exercises and no changes in endothelin-1 plasma concentrations were observed (14). The exact mechanism for reducing endothelin-1 plasma after exercise resistance is uncertain. The regulation body tropical hormones by physical activity or changes in body weight and total fat mass, as well as increased strength and power of skeletal muscle around blood vessels, is likely to reduce the body's need to the vascular endothelial cells to function, as a result, the secreted material of these cells in the plasma also decreases (15, 37).

Overall, the results of the present study for the first time showed that circuit resistance training as an appropriate training method can be used in the elderly's fitness program to reduce endothelial dysfunction due to aging. This improvement occurs as a result of a decrease in endothelin-1 plasma concentrations without altering the plasma concentration of nitric oxide. Also, the results confirmed that the circuit resistance training with decreased endothelin-1 concentration can reduce blood pressure.

Footnotes

Conflict of Interests: The authors declare that there is no conflict of interest in the current research.

Funding/Support: This study was not supported by any funding.

Patient Consent: There is no patient consent.

References

- Barton M. Aging and endothelin: Determinants of disease. *Life Sci*. 2014;**118**(2):97-109. doi: [10.1016/j.lfs.2014.09.009](https://doi.org/10.1016/j.lfs.2014.09.009). [PubMed: [25239727](https://pubmed.ncbi.nlm.nih.gov/25239727/)].
- El Assar M, Angulo J, Vallejo S, Peiro C, Sanchez-Ferrer CF, Rodriguez-Manas L. Mechanisms involved in the aging-induced vascular dysfunction. *Front Physiol*. 2012;**3**:132. doi: [10.3389/fphys.2012.00132](https://doi.org/10.3389/fphys.2012.00132). [PubMed: [22783194](https://pubmed.ncbi.nlm.nih.gov/22783194/)]. [PubMed Central: [PMC3361078](https://pubmed.ncbi.nlm.nih.gov/PMC3361078/)].
- Li Y, Hanssen H, Cordes M, Rossmeißl A, Endes S, Schmidt-Trucksass A. Aerobic, resistance and combined exercise training on arterial stiffness in normotensive and hypertensive adults: A review. *Eur J Sport Sci*. 2015;**15**(5):443-57. doi: [10.1080/17461391.2014.955129](https://doi.org/10.1080/17461391.2014.955129). [PubMed: [25251989](https://pubmed.ncbi.nlm.nih.gov/25251989/)].
- Barton M, Yanagisawa M. Endothelin: 20 years from discovery to therapy. *Can J Physiol Pharmacol*. 2008;**86**(8):485-98. doi: [10.1139/Y08-059](https://doi.org/10.1139/Y08-059). [PubMed: [18758495](https://pubmed.ncbi.nlm.nih.gov/18758495/)].
- Bourque SL, Davidge ST, Adams MA. The interaction between endothelin-1 and nitric oxide in the vasculature: New perspectives. *Am J Physiol Regul Integr Comp Physiol*. 2011;**300**(6):R1288-95. doi: [10.1152/ajpregu.00397.2010](https://doi.org/10.1152/ajpregu.00397.2010). [PubMed: [21368267](https://pubmed.ncbi.nlm.nih.gov/21368267/)].
- Trinity JD, Barrett-O'Keefe Z, Ives SJ, Morgan G, Rossman MJ, Donato AJ, et al. Endogenous endothelin-1 and femoral artery shear rate: Impact of age and implications for atherosclerosis. *J Hypertens*. 2016;**34**(2):266-73. doi: [10.1097/HJH.0000000000000777](https://doi.org/10.1097/HJH.0000000000000777). [PubMed: [26599223](https://pubmed.ncbi.nlm.nih.gov/26599223/)]. [PubMed Central: [PMC5380231](https://pubmed.ncbi.nlm.nih.gov/PMC5380231/)].
- Barton M. Cholesterol and atherosclerosis: Modulation by oestrogen. *Curr Opin Lipidol*. 2013;**24**(3):214-20. doi: [10.1097/MOL.0b013e3283613a94](https://doi.org/10.1097/MOL.0b013e3283613a94). [PubMed: [23594711](https://pubmed.ncbi.nlm.nih.gov/23594711/)].
- Traupe T, Ortmann J, Munter K, Barton M. Endothelial therapy of atherosclerosis and its risk factors. *Curr Vasc Pharmacol*. 2003;**1**(2):111-21. doi: [10.2174/1570161033476763](https://doi.org/10.2174/1570161033476763). [PubMed: [15320838](https://pubmed.ncbi.nlm.nih.gov/15320838/)].
- Beck DT, Martin JS, Casey DP, Braith RW. Exercise training reduces peripheral arterial stiffness and myocardial oxygen demand in young prehypertensive subjects. *Am J Hypertens*. 2013;**26**(9):1093-102. doi: [10.1093/ajh/hpt080](https://doi.org/10.1093/ajh/hpt080). [PubMed: [23736111](https://pubmed.ncbi.nlm.nih.gov/23736111/)]. [PubMed Central: [PMC3741227](https://pubmed.ncbi.nlm.nih.gov/PMC3741227/)].
- Guimaraes GV, Ciolac EG, Carvalho VO, D'Avila VM, Bortolotto LA, Bocchi EA. Effects of continuous vs. interval exercise training on blood pressure and arterial stiffness in treated hypertension. *Hypertens Res*. 2010;**33**(6):627-32. doi: [10.1038/hr.2010.42](https://doi.org/10.1038/hr.2010.42). [PubMed: [20379194](https://pubmed.ncbi.nlm.nih.gov/20379194/)].
- Miyachi M, Kawano H, Sugawara J, Takahashi K, Hayashi K, Yamazaki K, et al. Unfavorable effects of resistance training on central arterial compliance: A randomized intervention study. *Circulation*. 2004;**110**(18):2858-63. doi: [10.1161/01.CIR.0000146380.08401.99](https://doi.org/10.1161/01.CIR.0000146380.08401.99). [PubMed: [15492301](https://pubmed.ncbi.nlm.nih.gov/15492301/)].
- Okamoto T, Masuhara M, Ikuta K. Effects of low-intensity resistance training with slow lifting and lowering on vascular function. *J Hum Hypertens*. 2008;**22**(7):509-11. doi: [10.1038/jhh.2008.12](https://doi.org/10.1038/jhh.2008.12). [PubMed: [18337757](https://pubmed.ncbi.nlm.nih.gov/18337757/)].
- Garber CE, Blissmer B, Deschenes MR, Franklin BA, Lamonte MJ, Lee IM, et al. American College of Sports Medicine position stand. Quantity and quality of exercise for developing and maintaining cardiorespiratory, musculoskeletal, and neuromotor fitness in apparently healthy adults: Guidance for prescribing exercise. *Med Sci Sports Exerc*. 2011;**43**(7):1334-59. doi: [10.1249/MSS.0b013e318213febf](https://doi.org/10.1249/MSS.0b013e318213febf). [PubMed: [21694556](https://pubmed.ncbi.nlm.nih.gov/21694556/)].
- Maeda S, Otsuki T, Iemitsu M, Kamioka M, Sugawara J, Kuno S, et al. Effects of leg resistance training on arterial function in older men. *Br J Sports Med*. 2006;**40**(10):867-9. doi: [10.1136/bjsm.2006.029538](https://doi.org/10.1136/bjsm.2006.029538). [PubMed: [16920770](https://pubmed.ncbi.nlm.nih.gov/16920770/)]. [PubMed Central: [PMC2465061](https://pubmed.ncbi.nlm.nih.gov/PMC2465061/)].
- Maeda S, Miyauchi T, Iemitsu M, Sugawara J, Nagata Y, Goto K. Resistance exercise training reduces plasma endothelin-1 concentration in healthy young humans. *J Cardiovasc Pharmacol*. 2004;**44** Suppl 1:S443-6. doi: [10.1097/01.fjc.0000166319.91623.bo](https://doi.org/10.1097/01.fjc.0000166319.91623.bo). [PubMed: [15838344](https://pubmed.ncbi.nlm.nih.gov/15838344/)].
- Ciccolo JT, Carr LJ, Krupel KL, Longval JL. The role of resistance training in the prevention and treatment of chronic disease. *Am J Lifestyle Med*. 2009;**4**(4):293-308. doi: [10.1177/1559827609354034](https://doi.org/10.1177/1559827609354034).
- Alcaraz PE, Sanchez-Lorente J, Blazevich AJ. Physical performance and cardiovascular responses to an acute bout of heavy resistance circuit training versus traditional strength training. *J Strength Cond Res*. 2008;**22**(3):667-71. doi: [10.1519/JSC.0b013e31816a588f](https://doi.org/10.1519/JSC.0b013e31816a588f). [PubMed: [18438256](https://pubmed.ncbi.nlm.nih.gov/18438256/)].
- Ahmadiasl N, NIKNAZAR S, FARAJNIA S, Alipour MR. [Effect of three months exercise on expression of endothelin-1 mRNA in the lung tissue]. *Pharmaceut Sci*. 2008;**14**(3):56-62. Persian.
- Qassemian A, Salesi M. [The effect of an 8-week concurrent training on plasma endothelin_1 level and blood pressure of old women]. *J Keraman Univ Med Sci*. 2014;**21**(1). Persian.
- Donato AJ, Lesniewski LA, Delp MD. The effects of aging and exercise training on endothelin-1 vasoconstrictor responses in rat skeletal muscle arterioles. *Cardiovasc Res*. 2005;**66**(2):393-401. doi: [10.1016/j.cardiores.2004.10.023](https://doi.org/10.1016/j.cardiores.2004.10.023). [PubMed: [15820208](https://pubmed.ncbi.nlm.nih.gov/15820208/)].
- Donato AJ, Gano LB, Eskurza I, Silver AE, Gates PE, Jablonski K, et al. Vascular endothelial dysfunction with aging: Endothelin-1 and endothelial nitric oxide synthase. *Am J Physiol Heart Circ Physiol*. 2009;**297**(1):H425-32. doi: [10.1152/ajpheart.00689.2008](https://doi.org/10.1152/ajpheart.00689.2008). [PubMed: [19465546](https://pubmed.ncbi.nlm.nih.gov/19465546/)]. [PubMed Central: [PMC2717331](https://pubmed.ncbi.nlm.nih.gov/PMC2717331/)].

22. Thijssen DH, Rongen GA, van Dijk A, Smits P, Hopman MT. Enhanced endothelin-1-mediated leg vascular tone in healthy older subjects. *J Appl Physiol (1985)*. 2007;**103**(3):852-7. doi: [10.1152/jappphysiol.00357.2007](https://doi.org/10.1152/jappphysiol.00357.2007). [PubMed: [17556493](https://pubmed.ncbi.nlm.nih.gov/17556493/)].
23. Nyberg M, Mortensen SP, Hellsten Y. Physical activity opposes the age-related increase in skeletal muscle and plasma endothelin-1 levels and normalizes plasma endothelin-1 levels in individuals with essential hypertension. *Acta Physiol (Oxf)*. 2013;**207**(3):524-35. doi: [10.1111/apha.12048](https://doi.org/10.1111/apha.12048). [PubMed: [23227981](https://pubmed.ncbi.nlm.nih.gov/23227981/)].
24. Maeda S, Miyauchi T, Kakiyama T, Sugawara J, Iemitsu M, Irukayama-Tomobe Y, et al. Effects of exercise training of 8 weeks and detraining on plasma levels of endothelium-derived factors, endothelin-1 and nitric oxide, in healthy young humans. *Life Sci*. 2001;**69**(9):1005-16. doi: [10.1016/S0024-3205\(01\)01192-4](https://doi.org/10.1016/S0024-3205(01)01192-4). [PubMed: [11508642](https://pubmed.ncbi.nlm.nih.gov/11508642/)].
25. Maeda S, Tanabe T, Miyauchi T, Otsuki T, Sugawara J, Iemitsu M, et al. Aerobic exercise training reduces plasma endothelin-1 concentration in older women. *J Appl Physiol (1985)*. 2003;**95**(1):336-41. doi: [10.1152/jappphysiol.01016.2002](https://doi.org/10.1152/jappphysiol.01016.2002). [PubMed: [12611765](https://pubmed.ncbi.nlm.nih.gov/12611765/)].
26. Pichon CE, Hunter GR, Morris M, Bond RL, Metz J. Blood pressure and heart rate response and metabolic cost of circuit versus traditional weight training. *J Strength Cond Res*. 1996;**10**(3):153-6. doi: [10.1519/00124278-199608000-00004](https://doi.org/10.1519/00124278-199608000-00004).
27. Miura H, Nakagawa E, Takahashi Y. Influence of group training frequency on arterial stiffness in elderly women. *Eur J Appl Physiol*. 2008;**104**(6):1039-44. doi: [10.1007/s00421-008-0860-1](https://doi.org/10.1007/s00421-008-0860-1). [PubMed: [18751997](https://pubmed.ncbi.nlm.nih.gov/18751997/)].
28. Rudic RD, Shesely EG, Maeda N, Smithies O, Segal SS, Sessa WC. Direct evidence for the importance of endothelium-derived nitric oxide in vascular remodeling. *J Clin Invest*. 1998;**101**(4):731-6. doi: [10.1172/JCI1699](https://doi.org/10.1172/JCI1699). [PubMed: [9466966](https://pubmed.ncbi.nlm.nih.gov/9466966/)]. [PubMed Central: [PMC508619](https://pubmed.ncbi.nlm.nih.gov/PMC508619/)].
29. Tronc F, Wassef M, Esposito B, Henrion D, Glagov S, Tedgui A. Role of NO in flow-induced remodeling of the rabbit common carotid artery. *Arterioscler Thromb Vasc Biol*. 1996;**16**(10):1256-62. doi: [10.1161/01.ATV.16.10.1256](https://doi.org/10.1161/01.ATV.16.10.1256). [PubMed: [8857922](https://pubmed.ncbi.nlm.nih.gov/8857922/)].
30. Laughlin MH, Newcomer SC, Bender SB. Importance of hemodynamic forces as signals for exercise-induced changes in endothelial cell phenotype. *J Appl Physiol (1985)*. 2008;**104**(3):588-600. doi: [10.1152/jappphysiol.01096.2007](https://doi.org/10.1152/jappphysiol.01096.2007). [PubMed: [18063803](https://pubmed.ncbi.nlm.nih.gov/18063803/)]. [PubMed Central: [PMC3289055](https://pubmed.ncbi.nlm.nih.gov/PMC3289055/)].
31. Tuttle JL, Nachreiner RD, Bhuller AS, Condict KW, Connors BA, Herring BP, et al. Shear level influences resistance artery remodeling: Wall dimensions, cell density, and eNOS expression. *Am J Physiol Heart Circ Physiol*. 2001;**281**(3):H1380-9. doi: [10.1152/ajpheart.2001.281.3.H1380](https://doi.org/10.1152/ajpheart.2001.281.3.H1380). [PubMed: [11514310](https://pubmed.ncbi.nlm.nih.gov/11514310/)].
32. Miller VM, Burnett JC Jr. Modulation of NO and endothelin by chronic increases in blood flow in canine femoral arteries. *Am J Physiol*. 1992;**263**(1 Pt 2):H103-8. doi: [10.1152/ajpheart.1992.263.1.H103](https://doi.org/10.1152/ajpheart.1992.263.1.H103). [PubMed: [1636749](https://pubmed.ncbi.nlm.nih.gov/1636749/)].
33. Nadaud S, Philippe M, Arnal JF, Michel JB, Soubrier F. Sustained increase in aortic endothelial nitric oxide synthase expression in vivo in a model of chronic high blood flow. *Circ Res*. 1996;**79**(4):857-63. doi: [10.1161/01.RES.79.4.857](https://doi.org/10.1161/01.RES.79.4.857). [PubMed: [8831511](https://pubmed.ncbi.nlm.nih.gov/8831511/)].
34. Green DJ, Maiorana A, O'Driscoll G, Taylor R. Effect of exercise training on endothelium-derived nitric oxide function in humans. *J Physiol*. 2004;**561**(Pt 1):1-25. doi: [10.1113/jphysiol.2004.068197](https://doi.org/10.1113/jphysiol.2004.068197). [PubMed: [15375191](https://pubmed.ncbi.nlm.nih.gov/15375191/)]. [PubMed Central: [PMC1665322](https://pubmed.ncbi.nlm.nih.gov/PMC1665322/)].
35. Tanimoto M, Ishii N. Effects of low-intensity resistance exercise with slow movement and tonic force generation on muscular function in young men. *J Appl Physiol (1985)*. 2006;**100**(4):1150-7. doi: [10.1152/jappphysiol.00741.2005](https://doi.org/10.1152/jappphysiol.00741.2005). [PubMed: [16339347](https://pubmed.ncbi.nlm.nih.gov/16339347/)].
36. Van Guilder GP, Westby CM, Greiner JJ, Stauffer BL, DeSouza CA. Endothelin-1 vasoconstrictor tone increases with age in healthy men but can be reduced by regular aerobic exercise. *Hypertension*. 2007;**50**(2):403-9. doi: [10.1161/HYPERTENSIONAHA.107.088294](https://doi.org/10.1161/HYPERTENSIONAHA.107.088294). [PubMed: [17576858](https://pubmed.ncbi.nlm.nih.gov/17576858/)].
37. Maeda S, Miyauchi T, Iemitsu M, Tanabe T, Yokota T, Goto K, et al. Effects of exercise training on expression of endothelin-1 mRNA in the aorta of aged rats. *Clin Sci (Lond)*. 2002;**103** Suppl 48:118S-23S. doi: [10.1042/CS103S118S](https://doi.org/10.1042/CS103S118S). [PubMed: [12193068](https://pubmed.ncbi.nlm.nih.gov/12193068/)].