

Comparison of Pregnancy Outcomes in Diabetic and Healthy Women

Mostafa Jafari^{1,*}, Hadis Fattahi², Leila Sheikhy¹, Sodabeh Eskandari¹ and Mehdi Mirzaei-Alavijeh³

¹Health Faculty, Kermanshah University of Medical Sciences, Kermanshah, Iran

²Kermanshah University of Medical Sciences, Kermanshah, Iran

³Social Development & Health Promotion Research Center, Health Institute, Kermanshah University of Medical Sciences, Kermanshah, Iran

*Corresponding author: Health Faculty, Kermanshah University of Medical Sciences, Kermanshah, Iran. Email: mostafajafari66@yahoo.com

Received 2021 September 04; Revised 2021 December 28; Accepted 2022 January 23.

Abstract

Background: Gestational diabetes is the most prevalent metabolic disorder during pregnancy. Therefore, it is essential to recognize its outcomes and complications.

Objectives: The present study aimed to compare pregnancy outcomes in diabetic and healthy women.

Methods: This case-control study was conducted on 380 diabetic and healthy women, who were divided into two groups of 190, at Moatazedi Hospital in Kermanshah, Iran in 2019. The participants were selected via convenience sampling. Data were collected using a researcher-made checklist designed based on the family records of the participants. Data analysis was performed in SPSS version 22 using chi-square, Fisher's exact test, and the Mann-Whitney U test.

Results: A significant correlation was observed between gestational age, maternal body mass index, birth weight, infant's height, mode of delivery ($P < 0.01$), preeclampsia ($P = 0.01$), and the family history of diabetes ($P = 0.03$) in the diabetic and healthy pregnant women. However, no significant correlation was denoted between the five- and 10-minute Apgar scores, maternal disorders, and maternal care during pregnancy.

Conclusions: According to the results, gestational diabetes is associated with an increase rate of complications, which must be diminished through prevention and disease control to promote maternal and neonatal health.

Keywords: Diabetes, Complication, Pregnancy, Metabolic Disorders

1. Background

Gestational diabetes is a type of glucose intolerance, which occurs during pregnancy and often resolves spontaneously after pregnancy (1, 2). It is the most common metabolic disorder during pregnancy (3), with the prevalence rate reported to be 1 - 14% in different countries (4). The lowest prevalence of gestational diabetes has been estimated at less than 1% in Singapore and Tanzania, while the highest rate is reported to be 14% in India (5). According to a study in Iran, the prevalence of gestational diabetes is 1.3 - 8.9% (6). Notably, the difference in the prevalence of gestational diabetes in different countries is due to ethnic/racial differences, study populations, and the method used for screening and diagnosis (7).

Gestational diabetes is a public health concern, currently affecting a large population of pregnant women and causing short-term and long-term maternal and neonatal consequences (8). The short-term complications of gestational diabetes include cesarean section, preeclampsia, congenital anomalies, fetal macrosomia, stunted growth, underweight, stillbirth, intrauterine injuries, and neonatal metabolic disorders. In addition, the late complications in the mother and infant include the increased risk of type II diabetes mellitus, impaired glucose tolerance, and obe-

sity (2, 7, 9-11). Maternal risk factors for gestational diabetes include obesity, family history of diabetes, infertility treatment, current urinary tract infections, fetal macrosomia, sudden unexpected infant death, premature birth, preeclampsia, gestational diabetes, and advanced maternal age (12-14).

Diagnosis of gestational diabetes is important in terms of preventing diseases and perinatal disorders, while it also has a positive effect on the long-term health consequences of the mother and the child (15). The disease is asymptomatic and associated with multiple complications, which are not identified by the patients due to the absence of physical complications and lack of referral. Therefore, it is essential to diagnose gestational diabetes using effective screening methods (4).

Iran is a developing country with a young population and has a larger proportion of women of childbearing age who are exposed to the disease (16-18). The present study aimed to compare the pregnancy outcomes of pregnant women with gestational diabetes and healthy women. Conducting the study on the target population of the selected hospital is an innovation.

2. Objectives

The present study aimed to compare pregnancy outcomes in diabetic and healthy women.

3. Methods

This case-control study was conducted on the pregnant women referring to Moatazedi Hospital in Kermanshah, Iran in 2019. The sample population consisted of 380 pregnant women who were selected via census sampling from the available hospital files, including 190 diabetic and 190 non-diabetic pregnant women. Data collection was carried out based on the family records of the subjects.

Data were extracted from the medical records available in the hospital without any rejections. The control group was matched in terms of age and place of residence. The sample size of each group was determined to be 190 cases (total: 380). Data were collected using a checklist including maternal age, body mass index (BMI), gestational age, education level, employment status, and place of residence. The inclusion criterion was diabetic pregnant women.

(1) Neonatal variables: height, weight, and five- and 10-minute Apgar scores

(2) Maternal variables: preeclampsia, mode of delivery, delivery outcome, prenatal care, family history of diabetes, history of smoking, prior pregnancies, and congenital anomalies

All ethical issues such as obtaining a consent form were considered by the researchers, the necessary coordination was made with the hospital officials, and the issue of confidentiality was also observed in the agenda. After completing and encoding the data, data analysis was performed in SPSS version 22. Chi-square was used to compare the frequency distribution of the qualitative variables by a frequency table, and the correlations between the variables in the case and control groups were assessed using Fisher's exact test. In addition, the Mann-Whitney U test was applied to compare the means of the quantitative variables in the case and control groups. In all the statistical analyses, the significance level was set at 0.05.

4. Results

In total, 190 women with gestational diabetes and 190 healthy pregnant women were enrolled in the study (Table 1).

The comparison of the mean maternal age showed a significant difference between the case and control groups ($P < 0.01$). (Table 2)

Table 1. Comparison of Demographic and Fertility Characteristics in Case and Control Groups^a

Variables	Case	Control	P-Value
Mothers' age	27.5 ± 62	27.49 ± 4.17	0.80
Maternal BMI	26.05 ± 2.97	24.79 ± 2.96	< 0.01
Gestational age (w)	38.08 ± 1.72	37.09 ± 1.57	< 0.01

^a Values are expressed as mean ± SD.

Table 2. Comparison of Frequency Distribution and Percentage of Demographic Variables in Case and Control Groups^a

Variables	Case	Control	P-Value
Place of residence			< 0.01
City	114 (60)	121 (63.7)	
Village	76 (40)	69 (36.3)	
Level of education			< 0.01
Diploma and less	159 (83.7)	155 (81.6)	
Academic education	31 (16.3)	35 (18.4)	
Occupational status			< 0.01
Housewife	184 (96.8)	183 (96.3)	
Employee	6 (3.2)	7 (3.7)	

^a Values are expressed as No. (%).

The comparison of the case and control groups indicated significant differences in terms of the place of residence ($P < 0.01$), education level ($P < 0.01$), and occupation status ($P < 0.01$). (Table 3)

Table 3. Comparison of Mean Gestational Age, BMI, Apgar Scores, and Neonatal Height and Weight in Case and Control Groups

Variables	Mothers		P-value
	Diabetes	Healthy	
	Average Rating	Average Rating	
Gestational age	222.39	158.61	< 0.01
Maternal BMI	211.32	169.68	< 0.01
Neonatal Apgar score 5 minutes	195.15	185.85	0.23
Neonatal Apgar score 10 minutes	191.86	189.14	0.61
Neonatal weight	256.39	124.61	< 0.01
Neonatal height	210.89	168.11	< 0.01

The comparison of the mean gestational age ($P < 0.01$), maternal BMI ($P < 0.01$), neonatal weight ($P < 0.01$), and neonatal height ($P < 0.01$) also indicated significant differences between the case and control groups (Table 4).

Our findings also indicated a significant correlation between preeclampsia ($P < 0.01$), mode of delivery ($P < 0.01$),

Table 4. Comparison of Women with Gestational Diabetes with Healthy Mothers^a

Variables	Mothers		P-value
	Diabetes	Healthy	
Pre-eclampsia			
Yes	16 (8.4)	5 (2.6)	0.01
No	174 (91.6)	185 (97.4)	
Type of delivery			
Normal	28 (14.7)	60 (31.6)	< 0.01
Cesarean section	162 (85.3)	130 (68.4)	
Result of childbirth			
Alive	185 (97.4)	187 (94.4)	0.72
Dead	5 (2.6)	3 (1.6)	
Pregnancy care			
Yes	183 (96.3)	182 (95.8)	0.79
No	7 (3.7)	8 (4.2)	
Family history of diabetes			
Yes	11 (5.8)	3 (1.6)	0.03
No	179 (94.2)	187 (98.4)	
Congenital anomalies			
Yes	183 (96.3)	187 (98.4)	0.34
No	7 (3.7)	3 (1.6)	
History of pregnancy			
Yes	62 (32.6)	74 (38.9)	< 0.01
No	128 (67.4)	116 (61.6)	
History of smoking			
Yes	9 (4.7)	8 (4.2)	< 0.01
No	181 (95.3)	182 (95.8)	

^a Values are expressed as No. (%).

family history of diabetes ($P = 0.03$), prior pregnancies ($P < 0.01$), and history of smoking ($P < 0.01$).

5. Discussion

According to the results of the present, the gestational age of the healthy mothers was significantly different from the gestational age of the diabetic mothers whose gestational age was lower than the healthy mothers. This is consistent with the previous findings in this regard (17). Furthermore, the BM of the pregnant women had a significant difference between the case and control groups, which is also consistent with the previous studies in this regard (19-21).

According to the current research, the outcomes of birth weight and neonatal height were significantly different between the mothers with gestational diabetes and the healthy women, and the neonates of the mothers with gestational diabetes were overweight and taller than the infants of the healthy mothers. In addition, the ratio of cesarean sections of mothers in this category confirms this reason. The findings of the current research in this regard are consistent with previous studies (22).

The results of the present study indicated a significant difference between the diabetic and healthy women in terms of preeclampsia, which is consistent with another research in this regard (20). Furthermore, the diabetic and healthy women significantly differed in terms of the mode of delivery as the women with gestational diabetes showed a higher rate of cesarean delivery compared to the healthy women; this is also in line with the previous findings in this regard (22, 23). Since most of the women with gestational diabetes give birth to overweight neonates and due to the anatomical conditions of these women, the rate of cesarean section is often higher among these women than normal subjects.

The results of the present study indicated that the family history of type II diabetes mellitus had a significant difference between the women with gestational diabetes and the healthy women. This finding is consistent with the previous studies in this regard. According to the current research, the family history of diabetes was identified as a risk factor for gestational diabetes, which is in line with other studies. This finding also emphasizes the role of genetics in susceptibility to gestational diabetes. On the other hand, some studies have indicated genetics to be an independent risk factor, especially in individuals aged more than 30 years (24, 25).

The results of the present study showed no significant differences in the five- and 10-minute Apgar scores of infants between the healthy women and the women with gestational diabetes, which is consistent with the results of a similar study (1). Moreover, the two groups in our research were compared in terms of congenital malformations, and no significant difference was observed in this regard. This finding is also in line with some of the previous studies in this respect (23, 26).

The findings of the current research indicated a significant difference between the case and control groups in terms of smoking, which is inconsistent with other studies (27). On the other hand, a significant difference was observed between the two groups in terms of prior pregnancies as the proportion of primiparous healthy women was higher than those with gestational diabetes who experienced their second or multiple pregnancies. This is in line with the findings of similar studies (20).

5.1. Limitations of the Study

The main limitation was that the present study was conducted in only one hospital. Therefore, it is suggested that similar studies target a larger number of hospitals.

5.2. Suggestions

- Taking educational measures to enhance general maternal and neonatal health
- Investment of authorities in maternal and neonatal health
- Interventions to reduce and control gestational diabetes
- Studies on larger sample sizes to investigate gestational diabetes

5.3. Conclusions

According to the results, gestational diabetes was associated with an increased rate of complications, which must be diminished through disease prevention and control to promote maternal and neonatal health.

Acknowledgments

Hereby, we extend our gratitude to Moatazedi Hospital of Kermanshah for assisting us in this research project.

Footnotes

Authors' Contribution: M.J.: study concept and design, M.R.: Data analysis and interpretation, M.J., H.F., and L.S.: Drafting of the manuscript, M.J.: Critical revision of the manuscript for important intellectual content. All the authors provided comments and approved the final manuscript.

Conflict of Interests: None declared.

Funding/Support: This study did not have the financial support of any organization.

References

1. Gasim T. Gestational diabetes mellitus: maternal and perinatal outcomes in 220 Saudi women. *Oman Med J*. 2012;**27**(2):140-4. doi: [10.5001/omj.2012.29](https://doi.org/10.5001/omj.2012.29). [PubMed: [22496940](https://pubmed.ncbi.nlm.nih.gov/22496940/)]. [PubMed Central: [PMC3321340](https://pubmed.ncbi.nlm.nih.gov/PMC3321340/)].
2. Petersmann A, Nauck M, Muller-Wieland D, Kerner W, Muller UA, Landgraf R, et al. Definition, Classification and Diagnosis of Diabetes Mellitus. *Exp Clin Endocrinol Diabetes*. 2018;**126**(7):406-10. doi: [10.1055/a-0584-6223](https://doi.org/10.1055/a-0584-6223). [PubMed: [29975979](https://pubmed.ncbi.nlm.nih.gov/29975979/)].
3. Yang H, Wei Y, Gao X, Xu X, Fan L, He J, et al. Risk factors for gestational diabetes mellitus in Chinese women: a prospective study of 16,286 pregnant women in China. *Diabet Med*. 2009;**26**(11):1099-104. doi: [10.1111/j.1464-5491.2009.02845.x](https://doi.org/10.1111/j.1464-5491.2009.02845.x). [PubMed: [19929987](https://pubmed.ncbi.nlm.nih.gov/19929987/)].
4. Rashidi H, Kalantari K, Shahbazian H, Noughaj S. The relationship between fasting plasma glucose in the first trimester of pregnancy and the incidence of gestational diabetes in Iran. *Diabetes Metab Syndr*. 2021;**15**(4):102193. doi: [10.1016/j.dsx.2021.102193](https://doi.org/10.1016/j.dsx.2021.102193). [PubMed: [34229225](https://pubmed.ncbi.nlm.nih.gov/34229225/)].
5. Reece EA, Leguizamón G, Wiznitzer A. Gestational diabetes: the need for a common ground. *Lancet*. 2009;**373**(9677):1789-97. doi: [10.1016/S0140-6736\(09\)60515-8](https://doi.org/10.1016/S0140-6736(09)60515-8). [PubMed: [19465234](https://pubmed.ncbi.nlm.nih.gov/19465234/)].
6. Larijani B. A review on the prevalence of gestational diabetes mellitus (GDM) in different regions of Iran. *J Diabetes Metab Disord*. 2009;**8**:47-56.
7. Chen L, Shi L, Zhang D, Chao SM. Influence of Acculturation on Risk for Gestational Diabetes Among Asian Women. *Prev Chronic Dis*. 2019;**16**:E158. doi: [10.5888/pcd16.190212](https://doi.org/10.5888/pcd16.190212). [PubMed: [31808419](https://pubmed.ncbi.nlm.nih.gov/31808419/)]. [PubMed Central: [PMC6896832](https://pubmed.ncbi.nlm.nih.gov/PMC6896832/)].
8. Xu T, Dainelli L, Yu K, Ma L, Silva Zolezzi I, Detzel P, et al. The short-term health and economic burden of gestational diabetes mellitus in China: a modelling study. *BMJ Open*. 2017;**7**(12):e018893. doi: [10.1136/bmjopen-2017-018893](https://doi.org/10.1136/bmjopen-2017-018893). [PubMed: [29203507](https://pubmed.ncbi.nlm.nih.gov/29203507/)]. [PubMed Central: [PMC5736026](https://pubmed.ncbi.nlm.nih.gov/PMC5736026/)].
9. Srichumchit S, Luewan S, Tongsong T. Outcomes of pregnancy with gestational diabetes mellitus. *Int J Gynaecol Obstet*. 2015;**131**(3):251-4. doi: [10.1016/j.ijgo.2015.05.033](https://doi.org/10.1016/j.ijgo.2015.05.033). [PubMed: [26372349](https://pubmed.ncbi.nlm.nih.gov/26372349/)].
10. Sudasinghe BH, Wijeyaratne CN, Ginige PS. Long and short-term outcomes of Gestational Diabetes Mellitus (GDM) among South Asian women - A community-based study. *Diabetes Res Clin Pract*. 2018;**145**:93-101. doi: [10.1016/j.diabres.2018.04.013](https://doi.org/10.1016/j.diabres.2018.04.013). [PubMed: [29679628](https://pubmed.ncbi.nlm.nih.gov/29679628/)].
11. Dabelea D, Crume T. Maternal environment and the transgenerational cycle of obesity and diabetes. *Diabetes*. 2011;**60**(7):1849-55. doi: [10.2337/db11-0400](https://doi.org/10.2337/db11-0400). [PubMed: [21709280](https://pubmed.ncbi.nlm.nih.gov/21709280/)]. [PubMed Central: [PMC3121421](https://pubmed.ncbi.nlm.nih.gov/PMC3121421/)].
12. Rashidi H, Rezaianzadeh A, Tabatabaee SHR, Mousavi SA. Prevalence and Risk Factors of Gestational Diabetes Mellitus in Yazd Province, Iran. *J Midwifery Reproductive Health*. 2021;**9**(3):1-6. doi: [10.22038/JMRH.2021.56835.1696](https://doi.org/10.22038/JMRH.2021.56835.1696).
13. Grigoryan OR, Mikheev RK, Kurinova AN, Chernova MO, Sazonova DV, Akhmatova RR, et al. [Comparative impact analysis of risk factors on the course and outcomes of pregnancy with gestational diabetes mellitus]. *Probl Endokrinol (Mosk)*. 2021;**67**(3):78-86. doi: [10.14341/probli2756](https://doi.org/10.14341/probli2756). [PubMed: [34297506](https://pubmed.ncbi.nlm.nih.gov/34297506/)].
14. Zhuang W, Lv J, Liang Q, Chen W, Zhang S, Sun X. Adverse effects of gestational diabetes-related risk factors on pregnancy outcomes and intervention measures. *Exp Ther Med*. 2020;**20**(4):3361-7. doi: [10.3892/etm.2020.9050](https://doi.org/10.3892/etm.2020.9050).
15. Mirabelli M, Chiefari E, Tocci V, Greco E, Foti D, Brunetti A. Gestational diabetes: Implications for fetal growth, intervention timing, and treatment options. *Curr Opin Pharmacol*. 2021;**60**:1-10. doi: [10.1016/j.coph.2021.06.003](https://doi.org/10.1016/j.coph.2021.06.003). [PubMed: [34280703](https://pubmed.ncbi.nlm.nih.gov/34280703/)].
16. Etminan-Bakhsh M, Tadi S, Hatami M, Darabi R. Prevalence of Gestational Diabetes Mellitus and Its Associated Risk Factors in Boo-Ali Hospital, Tehran. *Galen Med J*. 2020;**9**:e1642. doi: [10.31661/gmj.v9i0.1642](https://doi.org/10.31661/gmj.v9i0.1642). [PubMed: [34466559](https://pubmed.ncbi.nlm.nih.gov/34466559/)]. [PubMed Central: [PMC8344143](https://pubmed.ncbi.nlm.nih.gov/PMC8344143/)].
17. Rashidi H, Rezaianzadeh A, Tabatabaee SHR, Mousavi SA. Prevalence and Risk Factors of Gestational Diabetes Mellitus in Yazd Province, Iran. *J Midwifery Reproductive Health*. 2021;**9**(3):1-6. doi: [10.22038/JMRH.2021.56835.1696](https://doi.org/10.22038/JMRH.2021.56835.1696).
18. Newman C, Egan AM, Ahern T, Al-Kiyumi M, Balan G, Brassill MJ, et al. Diabetes care and pregnancy outcomes for women with pregestational diabetes in Ireland. *Diabetes Res Clin Pract*. 2021;**173**:108685. doi: [10.1016/j.diabres.2021.108685](https://doi.org/10.1016/j.diabres.2021.108685). [PubMed: [33548336](https://pubmed.ncbi.nlm.nih.gov/33548336/)].
19. Ehrlich SF, Hedderson MM, Feng J, Davenport ER, Gunderson EP, Ferrara A. Change in body mass index between pregnancies and the risk of gestational diabetes in a second pregnancy. *Obstet Gynecol*. 2011;**117**(6):1323-30. doi: [10.1097/AOG.0b013e31821aa358](https://doi.org/10.1097/AOG.0b013e31821aa358). [PubMed: [21606742](https://pubmed.ncbi.nlm.nih.gov/21606742/)]. [PubMed Central: [PMC3222684](https://pubmed.ncbi.nlm.nih.gov/PMC3222684/)].

20. Najafi F, Hasani J, Izadi N, Hashemi-Nazari SS, Namvar Z, Shamsi H, et al. Risk of gestational diabetes mellitus by pre-pregnancy body mass index: A systematic review and meta-analysis. *Diabetes Metab Syndr*. 2021;**15**(4):102181. doi: [10.1016/j.dsx.2021.06.018](https://doi.org/10.1016/j.dsx.2021.06.018). [PubMed: [34214900](https://pubmed.ncbi.nlm.nih.gov/34214900/)].
21. Mohammadzadeh F, Eshghinia S, Vakili MA. The prevalence of gestational diabetes mellitus and its related risk factors in Gorgan, north of Iran. Selective or universal screening test is cost-effective? *Int J Diabetes Develop Countr*. 2015;**35**(3):225-9. doi: [10.1007/s13410-014-0209-8](https://doi.org/10.1007/s13410-014-0209-8).
22. Mocarski M, Savitz DA. Ethnic differences in the association between gestational diabetes and pregnancy outcome. *Matern Child Health J*. 2012;**16**(2):364-73. doi: [10.1007/s10995-011-0760-6](https://doi.org/10.1007/s10995-011-0760-6). [PubMed: [21365298](https://pubmed.ncbi.nlm.nih.gov/21365298/)].
23. Keshavarz M, Babaei GH. Comparison of pregnancy complications between gestational diabetes mellitus and normal group in Iran: A Cohort study. *Iran J Endocrinol Metab*. 2003;**5**(4):325-31.
24. Duman NB. Frequency of gestational diabetes mellitus and the associated risk factors. *Pak J Med Sci*. 2015;**31**(1):194-7. doi: [10.12669/pjms.311.5617](https://doi.org/10.12669/pjms.311.5617). [PubMed: [25878642](https://pubmed.ncbi.nlm.nih.gov/25878642/)]. [PubMed Central: [PMC4386185](https://pubmed.ncbi.nlm.nih.gov/PMC4386185/)].
25. Gracelyn L, Saranya N. Prevalence of gestational diabetes mellitus in antenatal women and its associated risk factors. *Int J Reprod Contracept Obstet Gynecol*. 2016;**5**(2):285-92.
26. Hussain T, Das S, Parveen F, Samanta P, Bal M, Yadav VS, et al. Prevalence, risk factors and morbidities of gestational diabetes among pregnant women attending a hospital in an urban area of Bhubaneswar, Odisha. *J Family Med Prim Care*. 2020;**9**(10):5327-33. doi: [10.4103/jfmpc.jfmpc_869_20](https://doi.org/10.4103/jfmpc.jfmpc_869_20). [PubMed: [33409210](https://pubmed.ncbi.nlm.nih.gov/33409210/)]. [PubMed Central: [PMC7773065](https://pubmed.ncbi.nlm.nih.gov/PMC7773065/)].
27. Tieu J, McPhee AJ, Crowther CA, Middleton P, Shepherd E. Screening for gestational diabetes mellitus based on different risk profiles and settings for improving maternal and infant health. *Cochrane Database Syst Rev*. 2017;**8**. CD007222. doi: [10.1002/14651858.CD007222.pub4](https://doi.org/10.1002/14651858.CD007222.pub4). [PubMed: [28771289](https://pubmed.ncbi.nlm.nih.gov/28771289/)]. [PubMed Central: [PMC6483271](https://pubmed.ncbi.nlm.nih.gov/PMC6483271/)].