



Detectable Antibodies in Newborn Cord Blood Following Maternal COVID-19 Infection Before Pregnancy: A Case Report

Adnan Batuhan Coşkun ^{1,*} and Selda Yüzer Alsaç ²

¹Department of Pediatric Nursing, Hasan Kalyoncu University, Gaziantep, Turkey

²Department of Pediatric Nursing, Yozgat Bozok University, Yozgat, Turkey

*Corresponding author: PhD, Department of Pediatric Nursing, Hasan Kalyoncu University, Gaziantep, Turkey. Email: a.batuhancoskun@gmail.com

Received 2022 June 06; Revised 2022 October 31; Accepted 2022 November 09.

Abstract

In this report, we present a case of an infant with detectable antibodies in the umbilical cord blood who was born to a healthcare professional mother diagnosed with coronavirus disease 2019 (COVID-19) before pregnancy and recovered, and also discuss the antibody level in light of relevant literature. Data consisted of severe acute respiratory syndrome coronavirus-2 (SARS-CoV-2) polymerase chain reaction (PCR) test and negative pre-pregnancy repeat real-time polymerase chain reaction (RT-PCR) test of a 29-year-old medical doctor mother, follow-up antibody test results in pregnancy, and antibodies detected in the cord blood of the newborn after birth. The study data were collected through face-to-face interviews with the mother. In conclusion, this case shows that SARS-CoV-2 immunoglobulin G (IgG) antibodies can be detected in the baby's cord blood immediately after birth from a mother who was diagnosed with COVID-19 before becoming pregnant and subsequently recovered with pharmacological treatment.

Keywords: SARS-CoV-2, Newborn, Cord Blood, Antibody Test, PCR

1. Introduction

The novel coronavirus disease 2019 (COVID-19) caused by severe acute respiratory syndrome coronavirus-2 (SARS-CoV-2) is a highly contagious disease which was first detected in Wuhan City, Hubei Province in China (1). The World Health Organization (WHO) China country office reported cases of pneumonia of unknown etiology in December 2019 (2). An analysis of all cases reported until January 2020 revealed that it was a viral disease that was not seen in humans before (3). Although the virus responsible for the disease was designated as 2019 novel coronavirus (2019-nCoV) on January 7, 2020, it was later named as SARS-CoV-2 (COVID-19) due to its similarity to SARS-CoV (4).

The incubation period for COVID-19 is considered to extend to 14 days, with a median time of 5 - 6 days from exposure to symptoms. Severe acute respiratory syndrome coronavirus-2 shares many similarities with SARS-CoV and Middle East respiratory syndrome (MERS)-CoV. Severe acute respiratory syndrome coronavirus-2 and MERS-CoV outbreaks have caused significant morbidity and mortality between 2002 and 2003 and in 2012, respectively (4, 5). Severe acute respiratory syndrome and MERS have significantly higher case fatality rates than COVID-19. However, COVID-19 is more infectious. The disease which has af-

ected nearly 213 countries to date has been declared a pandemic by the WHO on March 11, 2020 (6).

Severe acute respiratory syndrome coronavirus-2 Spike Antibody Test aids in identifying individuals with an adaptive immune response to the spike (S) protein of SARS-CoV-2. This assay allows detection of antibodies produced against the virus. Antibodies are immune system-related proteins known as immunoglobulins, which help the immune system recognize and eliminate foreign (non-self) threats to the body. Our immune system also produces some antibodies to protect our body against SARS-CoV-2 (COVID-19). In general, antiviral immunoglobulin M (IgM) antibodies appear about 10-15 days after the onset of symptoms and antiviral immunoglobulin G (IgG) antibodies become detectable 10 - 20 days after infection in patients. Quantitative measurement of the IgG antibody against the S protein of the virus is more valuable in assessing an individual's immune response. The antibody testing is performed on blood samples and an anti-SARS-CoV-2 IgG Antibody value ≥ 50 AU/mL is considered positive (7, 8).

Studies have shown that while the age group most commonly affected by this infection is the 30 to 79 age group, the mortality rate is much higher in individuals over 80 years of age. Limited information is available on the disease course in pregnant women and the risks to

the fetus and neonate; however, it is known that pregnant women constitute a vulnerable population who can be more severely affected by all viral infections compared to non-pregnant women (9). The Centers for Disease Control and Prevention (CDC) report states that women diagnosed with COVID-19 during pregnancy should be isolated under appropriate conditions and supported by local healthcare teams in order to reduce the possibility of transmission (10). In this case report, we present the results obtained from the cord blood of a baby born from a mother diagnosed with COVID-19 before getting pregnant who did not receive any vaccination during pregnancy and subsequently recovered. Before collecting study data, the mother was informed about the purpose of the study and both oral and written informed consent was obtained from the mother on 01.07.2021.

2. Case Presentation

The mother is 29 years old and works as a medical doctor. She tested positive for COVID-19 on 24.07.2020 before getting pregnant and experienced symptoms of severe arthralgia, headaches and fever during the 10-day quarantine period in which she received treatment (Table 1). At the end of 10-day treatment period following the COVID-19 (+) result, the mother tested negative on a polymerase chain reaction (PCR) test on 03.08.2020. She tested negative again on a repeat PCR test on 04.08.2020 and the result was confirmed (Table 2).

The mother reported that she had her last menstrual period on 26.09.2020 and became aware of her pregnancy at the end of September 2020. On 31.03.2021 (at gestational week 27), she underwent a follow-up antibody test which showed an IgG value of 50 AU/mL (Table 3). It was learnt that the mother did not develop any illness throughout her pregnancy but experienced “severe” COVID-19 symptoms before getting pregnant.

A male baby weighing 4170 g was born on 03.07.2021 at gestational week 41. A standard umbilical cord blood sample was collected from the newborn immediately after delivery under aseptic conditions to learn his blood type and to check his SARS-CoV-2 IgG level. Laboratory workup for the newborn’s cord blood demonstrated a SARS-CoV-2 IgG level of 15.32 AU/mL (Table 4). The baby underwent detailed health checks and was discharged as a completely healthy baby.

5. Discussion

In this case report, it was demonstrated that antibodies against COVID-19 that developed in the mother before

pregnancy passed to the fetus in utero and were detected in the newborn cord blood. Studies conducted during the pandemic have focused on mothers who were diagnosed with COVID-19 during pregnancy, prophylactic administration of COVID-19 vaccines to pregnant women, and COVID-19-related symptoms and antibody formation at birth in newborns following vertical transmission.

In a case report by Dong et al. published a 29-year-old woman at gestational week 34 was reported to present with complaints of fever and nasal congestion on 28.01.2020 and RT-PCR was positive for SARS-CoV-2 (11). The newborn showed no symptoms and at 2 hours after birth, the infant’s blood sample showed a SARS-CoV-2 IgG level of 140.32 AU/mL and an IgM level of 45.83 AU/mL. On 07.03.2020, the baby’s SARS-CoV-2 IgG level was 69.94 AU/mL and IgM level was 11.75 AU/mL, and she was discharged on 18.03.2020. That is a valuable case report showing exposure of the fetus to maternal COVID-19 infection for 23 days by vertical transmission during the intrauterine period (11).

In a 2020 study, Khan et al. reported their follow-up of 3 pregnant women who were COVID-19 positive at second and third trimesters (12). All 3 pregnant women delivered via vaginal route and were healthy, with no adverse outcomes observed in the postpartum period. PCR test results were negative for all newborns (12).

In one study, Zeng et al. evaluated six COVID-19 positive pregnant women and reported that none of the infants born to these mothers showed SARS-CoV-2 positivity on RT-PCR testing of throat swab and blood samples (13). However, all newborns had detectable antibodies against the virus in their serum samples (13).

Consistently, in a study by Mo et al. including six pregnant women, 3 of whom diagnosed with COVID-19 in the second trimester and 3 in the third trimester, all of their newborn infants were reported to have antibodies at birth (14). Among infants born to mothers infected in the 3rd trimester, 2 had detectable IgG and one had detectable IgM levels, whereas only IgG was found in infants born to mothers infected in the 2nd trimester (14).

The vertical transmission potential of COVID-19 in pregnant women was discussed in a study by Chen et al. published in the Lancet (15). Emergency preventive measures were implemented in 9 pregnant women with COVID-19 who developed pneumonia. Neonatal nasopharyngeal swab, amniotic fluid, cord blood and breast milk samples were collected and SARS-CoV-2 was diagnosed in 2 newborns (15).

In a case report by Paul and Chad, a pregnant healthcare worker with no history of COVID-19 was vaccinated with an mRNA COVID-19 vaccine (Moderna) at gestational week 36 (16). In line with hospital protocol, she underwent

Table 1. Initial Real-time Polymerase Chain Reaction Result of the Mother

Analysis Worked	Method	Reference Value	Analysis Result	Material	Result Date/Time
RT-PCR 1 reaction (SARS-CoV-2 PCR)	RT-PCR		Positive	Combined nose and throat swab	24.07.2020; 14:27 pm approved

Abbreviations: RT-PCR, Real-time polymerase chain reaction; SARS-CoV-2, severe acute respiratory syndrome coronavirus-2

Table 2. Post-treatment Real-time Polymerase Chain Reaction Results of the Mother

Analysis Worked	Method	Reference Value	Analysis Result	Material	Result Date/Time
RT-PCR 1 reaction (SARS-CoV-2 PCR)	RT-PCR		Negative	Combined nose and throat swab	03.08.2020; 14:53 pm approved
RT-PCR 1 reaction (SARS-CoV-2 PCR)	RT-PCR		Negative	Combined nose and throat swab	04.08.2020; 16:38 pm approved

Abbreviations: RT-PCR, Real-time polymerase chain reaction; SARS-CoV-2, severe acute respiratory syndrome coronavirus-2

Table 3. Antibody Test Result at Gestational Week 27

Analysis Worked	Method	Reference Value	Analysis Result	Material	Result Date/Time
COVID-19 (SARS-CoV-2) IgG		4 - 6	50 < positive	Blood test	31.03.2021; 15:00 pm approved

Abbreviations: COVID-19, Coronavirus disease 2019; SARS-CoV-2, severe acute respiratory syndrome coronavirus-2; IgG, immunoglobulin G

Table 4. Severe Acute Respiratory Syndrome Coronavirus-2 Antibody Test Result of the Newborn

Analysis Worked	Method	Reference Value	Analysis Result	Material	Result Date/Time
COVID-19 (SARS-CoV-2) IgG		4 - 6	15, 32 positive	Blood test	03.07.2021; 11:27 am approved

Abbreviations: COVID-19, Coronavirus disease 2019; SARS-CoV-2, severe acute respiratory syndrome coronavirus-2; IgG, immunoglobulin G

COVID-19 testing at the time of admission to the labor and delivery ward which was negative. At 3 weeks after the first dose of Moderna, a healthy girl was born via normal, spontaneous vaginal delivery. A standard cord blood sample (0.5 mL) was collected into a red-top tube immediately after the birth to determine newborn blood type under aseptic conditions. IgG antibodies to SARS-CoV-2 were detected in the cord blood at a level of 1.31 U/mL (16).

There are several studies in the literature showing significant differences among antibody levels in infants born to SARS-CoV-2 positive mothers or mothers who were vaccinated during pregnancy (11-16). However, no study was identified in both national and international literature reporting on the antibody level of the infants born to mothers who were SARS-CoV-2 positive before pregnancy.

3.1. Conclusions

Pregnant women and newborns should be evaluated and followed attentively since they are considered to be at-risk, prioritized populations in the COVID-19 outbreak. However, the case reported here shows us that the effects of a maternal COVID-19 infection occurring before pregnancy may be seen in the newborn. For this reason, consideration of the transmission routes of COVID-19 infection and transfer of antibodies to the newborn both before and during

pregnancy will allow us to take measures to eliminate potential health risks to the mother and the baby.

Here, we presented the first known case of an infant with detectable SARS-CoV-2 IgG antibodies in the cord blood immediately after birth born to a symptomatic mother who was diagnosed with COVID-19 before getting pregnant and subsequently recovered with pharmacological treatment. We believe that this case report provides valuable data to the literature by showing the persistent nature of antibodies against SARS-CoV-2.

Footnotes

Authors' Contribution: Conceptualization, A. B. C.; methodology, A. B. C., writing-original draft preparation, A. B. C., S. Y. A.; writing-review and editing, A. B. C., S. Y. A. All authors have read and agreed to the published version of the manuscript.

Conflict of Interests: The authors declare no conflict of interests.

Funding/Support: This research received no external funding.

Informed Consent: Written informed consent was obtained from the patient.

References

1. Rasmussen SA, Smulian JC, Lednický JA, Wen TS, Jamieson DJ. Coronavirus Disease 2019 (COVID-19) and pregnancy: what obstetricians need to know. *Am J Obstet Gynecol.* 2020;**222**(5):415–26. [PubMed: [32105680](#)]. [PubMed Central: [PMC7093856](#)]. <https://doi.org/10.1016/j.ajog.2020.02.017>.
2. Coronaviridae Study Group of the International Committee on Taxonomy of Viruses. The species Severe acute respiratory syndrome-related coronavirus: classifying 2019-nCoV and naming it SARS-CoV-2. *Nat Microbiol.* 2020;**5**(4):536–44. [PubMed: [32123347](#)]. [PubMed Central: [PMC7095448](#)]. <https://doi.org/10.1038/s41564-020-0695-z>.
3. Ministry of Health Republic of Turkey. *COVID-19 Infection Guideline General Information Epidemiology and Diagnosis*. Antalya, Turkey: Ministry of Health Republic of Turkey; 2021.
4. Coşkun AB, Çiğdem Z, Polat S. [Investigation of the Indirect (Secondary) Effects of COVID-19 on Children, Traditional Compilation]. *Turkiye Klinikleri J Pediatr.* 2021;**30**(2):145–52. Turkish. <https://doi.org/10.5336/pediatr.2020-79685>.
5. Chowell G, Abdirizak F, Lee S, Lee J, Jung E, Nishiura H, et al. Transmission characteristics of MERS and SARS in the healthcare setting: a comparative study. *BMC Med.* 2015;**13**:210. [PubMed: [26336062](#)]. [PubMed Central: [PMC4558759](#)]. <https://doi.org/10.1186/s12916-015-0450-0>.
6. World Health Organization. *Transmission of SARS-CoV-2: implications for infection prevention precautions*. Geneva, Switzerland: World Health Organization; 2020, [cited 10 Aug 2021]. Available from: <https://www.who.int/news-room/commentaries/detail/transmission-of-sars-cov-2-implications-for-infection-prevention-precautions>.
7. Mavi D, İnkaya AÇ. [COVID-19: Immunopathogenesis]. *Flora.* 2020;**25**(2):121–31. Turkish. <https://doi.org/10.5578/flora.69606>.
8. Çakal B. [Antibody Responses Against SARS-Cov-2 in COVID-19 and Serological Assays]. *Turkiye Klinikleri J Health Sci.* 2020;**5**(2):394–400. Turkish. <https://doi.org/10.5336/healthsci.2020-76285>.
9. Desdicioğlu R, Yavuz AF. [COVID-19 and pregnancy]. *Ankara Med J.* 2020;**20**(2):482–7. Turkish. <https://doi.org/10.5505/amj.2020.74318>.
10. Centers for Disease Control and Prevention (CDC). *Breastfeeding and Caring for Newborns if You Have COVID-19*. Atlanta, USA: Centers for Disease Control and Prevention (CDC); 2020, [updated 20 Jan 2022; cited 10 Aug 2021]. Available from: www.cdc.gov/coronavirus/2019-ncov/if-you-are-sick/pregnancy-breastfeeding.html.
11. Dong L, Tian J, He S, Zhu C, Wang J, Liu C, et al. Possible Vertical Transmission of SARS-CoV-2 From an Infected Mother to Her Newborn. *JAMA.* 2020;**323**(18):1846–8. [PubMed: [32215581](#)]. [PubMed Central: [PMC7099527](#)]. <https://doi.org/10.1001/jama.2020.4621>.
12. Khan S, Peng L, Siddique R, Nabi G, Xue M; Nawsherwan, et al. Impact of COVID-19 infection on pregnancy outcomes and the risk of maternal-to-neonatal intrapartum transmission of COVID-19 during natural birth. *Infect Control Hosp Epidemiol.* 2020;**41**(6):748–50. [PubMed: [32279693](#)]. [PubMed Central: [PMC7156579](#)]. <https://doi.org/10.1017/ice.2020.84>.
13. Zeng H, Xu C, Fan J, Tang Y, Deng Q, Zhang W, et al. Antibodies in Infants Born to Mothers With COVID-19 Pneumonia. *JAMA.* 2020;**323**(18):1848–9. [PubMed: [32215589](#)]. [PubMed Central: [PMC7099444](#)]. <https://doi.org/10.1001/jama.2020.4861>.
14. Mo H, Wang M, Wang M, Han Y, Zhang Y, Hu K. Detectable antibodies against SARS-CoV-2 in newborns from mothers infected with COVID-19 at different gestational ages. *Pediatr Neonatol.* 2021;**62**(3):321–3. [PubMed: [33836982](#)]. [PubMed Central: [PMC7997304](#)]. <https://doi.org/10.1016/j.pedneo.2021.03.011>.
15. Chen H, Guo J, Wang C, Luo F, Yu X, Zhang W, et al. Clinical characteristics and intrauterine vertical transmission potential of COVID-19 infection in nine pregnant women: a retrospective review of medical records. *Lancet.* 2020;**395**(10226):809–15. [PubMed: [32151335](#)]. [PubMed Central: [PMC7159281](#)]. [https://doi.org/10.1016/S0140-6736\(20\)30360-3](https://doi.org/10.1016/S0140-6736(20)30360-3).
16. Paul G, Chad R. Newborn antibodies to SARS-CoV-2 detected in cord blood after maternal vaccination - a case report. *BMC Pediatr.* 2021;**21**(1):138. [PubMed: [33752624](#)]. [PubMed Central: [PMC7982334](#)]. <https://doi.org/10.1186/s12887-021-02618-y>.