



The Relationship between Seborrheic Dermatitis and Androgenetic Alopecia in Patients Referred to a Skin Clinic in Tehran, Iran: A Retrospective Study

Amirhosein Faghihkhorsani ^{1,*}, Afsaneh Sadeghzadeh ², Azadeh Goodarzi ³ and Masoumeh Rohaninasab ²

¹Medical Student, Iran University of Medical Sciences (IUMS), Tehran, Iran

²Department of Dermatology, School of Medicine Hazrat-e Rasool General Hospital Iran University of Medical Sciences, Tehran, Iran

³Department of Dermatology, School of Medicine Iran University of Medical Sciences, Tehran, Iran

*Corresponding author: Medical Student, Iran University of Medical Sciences (IUMS), Tehran, Iran Email: amirkhorasany1612@gmail.com

Received 2023 December 18; Revised 2023 December 23; Accepted 2023 December 25.

Abstract

Background: Seborrheic dermatitis (SD) is a common skin problem that generally affects the scalp. SD causes scale-like parts, skin redness, and persistent dandruff, which can also affect the fatty parts of the body. Androgenic alopecia (AGA) is an androgen-dependent hereditary disorder characterized by progressive thinning of hair with various patterns.

Objectives: Considering that AGA is probably related to race, genetics, and geographical conditions, and timely treatment of SD and AGA is critical, the main objective of this study was to evaluate the relationship between SD and AGA.

Methods: This descriptive-analytical study was conducted on patients with AGA referred to the skin clinic of Rasoul-Akram Hospital in Tehran, Iran, in 2021. Among the mentioned patients, 250 were randomly selected by easy sampling. The patient's age, gender, occupation, marital status, history of skin, hormonal, or systemic disease, medical history, medication history, smoking history, alcohol intake, and family history of hair loss are all required information after obtaining informed consent from the patient. Skin color and face and skin type were recorded by a researcher-made checklist. Then, the patients were examined in terms of SD, and finally, the collected data were analyzed using SPSS software Version 24 and relevant statistical tests.

Results: The average age of the studied patients was 35.64 ± 8.11 (67.2%) were women, and 81 (32.8%) were men. The highest severity of SD was grade 1 in 116 people (46.6%). There was a significant relationship between the severity of SD and the history of hair loss in the family ($P = 0.03$). In addition, there was a significant correlation between the pattern of AGA in men and the severity of SD ($P < 0.05$). The study showed a significant relationship between dandruff symptoms and AGA patterns ($P < 0.05$).

Conclusions: Based on the results, this issue should be evaluated with more precision and follow-up due to the relationship between the severity of SD and the AGA pattern. Moreover, eliminating the symptoms of dandruff can help in improving hair loss. In addition, evaluating a person's medical history and family history is essential to determining the pattern of hair loss, preventing its progression, and timely treatment of this problem.

Keywords: Seborrheic Dermatitis, Androgenic Alopecia, Hair Loss, Scalp, Dandruff

1. Background

Hair manifests youth and health psychologically and based on interpersonal relationships. Hair loss has an adverse effect on a person's social status, self-confidence, and psychological dimension (1, 2). Androgenic alopecia (AGA) is an androgen-dependent hereditary disorder, the progressive thinning of hair with various patterns. The pattern of hair loss in women is seen centrally (based on Ludwig's classification) without temporal-frontal or

whole-head progression. However, in men, bilateral hair loss in the temporal region and vertex region (based on Hamilton's classification) is the predominant pattern (3, 4).

AGA is characterized by hair follicle shrinkage manifesting as hair loss due to systemic androgens and genetic factors (4). Most changes are in the third and fourth decades of life, but hair loss begins immediately after puberty and continues gradually (5). AGA is the most common cause of hair loss in men and women (6).

Female pattern hair loss (FPHL) has replaced AGA in women due to the unclear relationship between androgens and their incidence (7). FPHL is the most common hair loss disorder in women. Initial symptoms may appear in the teenage years, leading to progressive and extensive hair loss with a specific pattern (8).

Clinical evaluation in individuals with AGA should include examination of the scalp, facial hair, body, and all skin and nails. Although examining the scalp is usually normal in AGA, seborrheic dermatitis (SD) can be considered a worsening factor of AGA (9). When AGA is chronic, the scalp becomes atrophic, and contact with sunlight can aggravate AGA (3, 10, 11).

SD has a wide range of mild (in the form of dandruff), moderate (in the form of redness and scaling), and severe (in the form of secondary infections), and its initial symptoms are redness and peeling around the hair follicle (12, 13).

SD has a wide range of mild (in the form of dandruff), moderate (in the form of redness and scaling), and severe (in the form of secondary infections), and initial symptoms are redness and peeling around the hair follicle (12, 13). Although hair loss is not always related to SD, some studies state that hair loss occasionally occurs with increased telogens in people with chronic SD, which will be reversible when the inflammation is removed (14, 15). Follicular involvement in SD with extensive and progressive hair loss has not been mentioned in studies (16). Men are affected more often than women (3.0% vs. 2.6%) in all age groups, suggesting that SD may be associated with sex hormones such as androgens (12-14).

Since androgenetic alopecia is affected by race, genetics, and geographical conditions, it is necessary to investigate the prevalence of these two diseases together in the Iranian population. When these two complications are accompanied by each other, a more appropriate treatment process should be followed because their treatment affects each other.

2. Objectives

When SD and AGA appear in the same person, appropriate and timely treatment for those cases is essential. Therefore, the main aim of the present study was to evaluate the association of SD with AGA in patients referred to the skin clinic located at Rasoul Akram Hospital in Tehran in 2021.

3. Methods

This descriptive-analytical study was conducted on patients with AGA referred to the Dermatology Clinic of

Rasoul Akram Hospital in Tehran, Iran. Sampling was done sequentially and stopped when the number of samples reached the required size. The relevant sample size equation was used to determine the sample size, which is estimated to be 200 patients. Considering the possible 20% dropout of patients, the sample size was finally increased to 250. The number of effective parameters in determining the sample size for the present study was received from Kibar et al. (15).

Inclusion criteria were a clinical diagnosis of AGA, as the predominant diagnosis by a dermatologist, and an age between 18 and 60 years. Exclusion criteria included receiving previous treatment for hair loss, previous treatment for neurological disease, a history of immunodeficiency disease, and an unwillingness to participate in the study.

Once informed consent has been obtained from the patients, the required information of the patients is obtained, such as their age, gender, occupation, marital status, skin, hormonal or systemic diseases, medications taken by the patient, smoking and alcohol consumption, and family history of hair loss. Skin color and face and skin type were recorded by a researcher-made checklist. Then, the patients were examined regarding SD, and the presence or absence of seborrheic dermatitis and disease symptoms, including erythema, dandruff, and the spread of the lesion, were checked. In addition, the severity of dermatitis was examined, and its classification includes "no involvement"="level 0", involvement of less than half of the head="level 1", involvement of half of the head="level 2" and involvement in more than half of the head="level 3" (12-14). In the following, information about AGA includes the location of hair loss on the head, the male pattern hair loss according to the Hamilton-Norwood classification (HNC) (Table 1). The female pattern hair loss was determined according to the Ludwig classification pattern (LC) (Table 1) (15).

Finally, the collected data were analyzed using SPSS software Version 24 and relevant statistical tests at a significance level of $\alpha = 0.05$. The descriptive results obtained for quantitative variables were expressed as mean and standard deviation and for categorical qualitative variables as frequency and frequency percentage. Kolmogorov-Smirnov test was used to evaluate the normal distribution in different parameters. The Paired sample *t*-test was utilized to compare the quantitative variables in two different modes, but the Wilcoxon test was used when the distribution was abnormal. The Chi-squared test was used to compare the two qualitative variables. The Mann-Whitney U test was used for two quality and quantitative variables, and the Pearson correlation coefficient (*r*) test was applied to

Table 1. Hamilton-Norwood Classification and Ludwig Classification Patterns

Type/Category	Description
Hamilton-Norwood Classification	
Type 1	Absence of bilateral regression in the frontoparietal line
Type 2	Triangular hair loss on both sides of the frontoparietal line
Type 3	Patients not falling into the previous groups (types 1 and 2) have non-symmetrical hair loss in the frontoparietal line.
Type 4	Perceptible hair loss in the frontoparietal line is known as baldness.
Type 5	The addition of a bald head
Type 6	Connecting hair loss on the frontal line to the frontoparietal line
Types 7 and 8	View of horseshoe without hair
Ludwig Classification	
Grade 1	Thinning of hair about 1-3 cm behind the hairline
Grade 2	Perceptible hair loss compared to grade 1
Grade 3	Complete baldness

compare two quantitative variables.

4. Results

The average age was 35.64 ± 8.11 ; 146 people (58.6%) were 36 to 60 years old, and 104 (41.4%) were 18 to 35. The results showed that 169 (67.2%) women and 81 (32.8%) men participated in the study. The most common occupation of the patients was housewife, which was observed in 78 people (31%), and 164 people (65.5%) were married. According to the findings, 47 people (19%) had a history of hormonal diseases, 43 people (17.2%) had a history of skin diseases, 56 people (22.4%) had a history of systemic diseases, and 194 people (77.6%) had a history of hair loss in the family. According to the answers of the participants, 65 people (25.9%) were taking medicine, and 73 people (29.3%) had a history of smoking and alcohol consumption. The most common type of skin related to oily skin was observed in 112 (48.3%) patients. Further, the most significant number of patients, 138 (55.2%), were white, and the most alopecia was in the frontal area in 164 (65.5%) patients. The study showed that only 78 people (31%) had erythema and dandruff, and 142 people (56.9%) had lesions below 1%.

About 142 out of 250 (56.9%) patients had SD, and the highest severity of SD was grade 1 (involvement of less than half of the head) in 116 out of 250 people (46.6%).

The frequency of classification of hair loss patterns for men and women is presented in Table 2. Based on the

findings mentioned above, 90 people were classified in the Hamilton-Norwood classification (HNC) and 164 in the Ludwig classification (LC) (Table 2).

Table 2. Frequency of Participants in the Study Based on the Classification of Male and Female Hair Loss Patterns

Type/Category	No. (%)
Hamilton-Norwood Classification (HNC) of Male Pattern Hair Loss	
Type 1	14 (15.6)
Type 2	27 (30)
Type 3	23 (25.6)
Type 4	6 (6.6)
Type 5	10 (11.1)
Type 6	10 (11.1)
Total	90 (100)
Ludwig Classification (LC) of Female Pattern Hair Loss	
Grade 1	142 (86.6)
Grade 2	22 (13.4)
Total	164 (100)

The results of the relationship between the severity of SD and the study variables, including age, gender, history of drugs, smoking, and alcohol consumption, history of previous diseases, history of hair loss in the family, skin color, skin type, and location of facial involvement are shown in Table 3. The results of the analysis using the chi-squared test showed a significant relationship only between the severity of SD and "history of hair loss in the family" ($P < 0.05$), while no significant correlation between the severity of SD and other variables ($P > 0.05$).

The relationship between the severity of SD and the pattern of AGA in women and men showed that based on the pattern of AGA in men, the highest severity of SD (3rd degree=involvement of more than half of the head) corresponds to the pattern of type "2" hair loss (triangular hair loss in both sides of the front parietal line) in four patients. In addition, this result was observed in women with 2nd-degree dermatitis (involvement of half of the head) related to type 1 hair loss pattern (thinning of the head about 1-3 cm behind the hairline) in nine patients. Further, there was a significant relationship between the parameters of the AGA pattern in men and the severity of SD ($P < 0.05$) (Table 4).

Table 3. Correlation Between Seborrheic Dermatitis Severity and Study Variables

Variables	Severity of Seborrheic Dermatitis				P-Value
	Grade 0	Grade 1	Grade 2	Grade 3	
Age					0.09
18 - 35	47	34	13	9	
90 - 36	61	82	4	0	
Sex					0.106
Male	39	26	9	9	
Female	69	90	9	0	
History of drug use	26	35	4	0	0.81
History of smoking and alcohol consumption	26	39	0	6	0.071
History of skin disease	22	17	4	0	0.8
History of hormonal disease	17	26	4	0	0.82
History of systemic disease	22	34	0	0	0.45
Family history of hair loss	91	86	17	0	0.031
Facial skin color					0.52
White	56	73	4	4	
Wheat color	52	43	13	4	
Skin type					0.38
Greasy	56	47	13	4	
Dry	9	26	4	4	
Normal	24	19	0	0	
Scalp involvement					
Frontal	73	82	9	0	0.2
Temporal	35	30	4	9	0.18
Parietal	56	52	9	0	0.55
Occipital	39	22	9	0	0.28

Table 4. The Relationship Between the Severity of Seborrheic Dermatitis and the Pattern of Androgenetic Alopecia in Women and Men

Variable and Type/Category	Severity of Seborrheic Dermatitis				P-Value
	Grade 0	Grade 1	Grade 2	Grade 3	
Hamilton-Norwood classification (HNC) of male pattern hair loss					0.021
Type 1	4	0	4	0	
Type 2	13	9	0	4	
Type 3	9	13	0	0	
Type 4	0	4	0	0	
Type 5	5	0	5	0	
Type 6	9	0	0	0	
Ludwig classification (LC) of female pattern hair loss					0.42
Grade 1	56	77	9	0	
Grade 2	13	9	0	0	

5. Discussion

The results of the present study showed that 142 out of 250 patients (56.9%) had seborrheic dermatitis (SD). The highest severity of SD was grade "1", observed in almost half of the patients. The severity of SD with involvement of more than half of the head (grade "3") was only in the age group of 18 to 35 years and men. However, there was no significant relationship between the severity of SD and age and sex parameters. Based on the androgenic alopecia (AGA) pattern in men, the highest severity of SD is related to the pattern of hair loss type "2" (triangular hair loss on both sides of the frontoparietal line). The highest severity of SD in women was grade "2", which is observed in the pattern of hair loss type "1" (thinning of hair about 1 - 3 cm behind the hairline). Only the parameter of the AGA pattern in men had a significant relationship with the severity of SD. In a similar study by Sarlak et al., the level of seborrheic dermatitis was evaluated in 60 out of 160 patients (37.5%) (17). Moreover, in another study by Jang et al., the most common pattern of AGA in men was HNC-3 grade and LC-1 grade in women. SD was the most common comorbidity in affected women and men (18). In the study by Ummiti et al., 66 men and 25 women with AGA were included. The most common finding was the heterogeneity of hair shaft thickness in all male and female patients (19). Hu et al. included 750 men and 200 women as the patient group (case) and 100 men and 50 women as the healthy group (control). The mentioned study showed that the heterogeneity of hair shaft thickness was observed in more than 20% of men (20). Generally, the results of previous similar studies, in line with the results of the present study, have emphasized that higher degrees of hair loss due to thinner and more vulnerable form are prone to the occurrence of severe dermatitis and the hair of these patients should be strengthened in a better way.

Another important finding of the present study was that half of the patients had oily skin, and most were white. The most common location of alopecia was the frontal area of the head in 164 patients (65.5%). In addition, the most pattern of male hair loss is related to type 2-HNC (triangular hair loss on both sides of the frontoparietal line) in 27 patients (10.7%), and the most pattern of female hair loss is related to grade "1" (thinning of hair about 1 - 3 cm behind the hairline) in 149 (56.9%) patients. The highest intensity of dermatitis was related to grade 1-LC in people with white and wheatish skin color. Based on the skin type, only in dry and oily skins was the highest intensity of SD, i.e., intensity of grade "3". The highest severity of SD, i.e., grade "3", also occurred in the temporal region. The present study showed no meaningful relationship between facial skin color, skin type, and scalp involvement

parameters. Kibar et al. observed that multi-hair follicular and honeycomb pigmentation is probably correlated with androgenetic hair loss (15). Ummiti et al. found that honeycomb pigmentation was observed in the scalp of most patients, but it was not related to the severity of hair loss (19). Hu et al. reported that yellow spots, white spots, and scalp pigmentation were associated with the severity of hair loss (20). Therefore, the results of previous similar studies and the current research confirm that the presence of different skin pigments and dryness or having too much fat on the skin causes hair loss to become more severe and dermatitis to appear more severe. Therefore, solving these problems can prevent dermatitis or its progression (if it occurs).

5.1. Limitations

Several study participants were unfamiliar with the specialized terms related to dermatitis and hair loss, which was a critical limitation of the present study. This limitation in the time of questioning by the researcher caused problems in recording the correct and accurate answers. The questioner tried to familiarize the patients with specialized terms by providing appropriate explanations so that the patients could give the most accurate and realistic answers.

5.2. Conclusions

Considering the relationship between the severity of seborrheic dermatitis and male hair loss patterns, doctors and experts on skin diseases should evaluate this issue with more care and follow-up. Removing dandruff symptoms can help improve hair loss. Based on the results, evaluating a person's disease history and family history is critical in determining the pattern of hair loss, preventing the progression of this problem, and on-time treatment.

Footnotes

Authors' Contribution: A.F: Ideation in the presentation of the study topic, designing the study and presenting its methodology, writing of initial draft and revision of the revised manuscript; A.S.B., A.G., M.R.: Collect and record results by checklist, participation in writing and revising the manuscript.

Conflict of Interests: The authors confirm that there are no competing interests.

Ethical Approval: This study was approved by the Ethics Committee of the Iran University of Medical Science ([IR.IUMS.FMD.REC.1399.332](https://doi.org/10.30471/IR.IUMS.FMD.REC.1399.332)).

Funding/Support: This study, conducted as a student thesis in general medicine, was financially supported by

the Vice-Chancellor of Research and Technology of Iran University of Medical Sciences.

Informed Consent: Consent to participate in the study was obtained from all patients verbally and in written form.

References

- Alfonso M, Richter-Appelt H, Tosti A, Viera MS, Garcia M. The psychosocial impact of hair loss among men: a multinational European study. *Curr Med Res Opin.* 2005;21(11):1829-36. [PubMed ID: 16307704]. <https://doi.org/10.1185/030079905X61820>.
- Schmidt S, Fischer TW, Chren MM, Strauss BM, Elsner P. Strategies of coping and quality of life in women with alopecia. *Br J Dermatol.* 2001;144(5):1038-43. [PubMed ID: 11359394]. <https://doi.org/10.1046/j.1365-2133.2001.04195.x>.
- Lolli F, Pallotti F, Rossi A, Fortuna MC, Caro G, Lenzi A, et al. Androgenetic alopecia: a review. *Endocrine.* 2017;57(1):9-17. [PubMed ID: 28349362]. <https://doi.org/10.1007/s12020-017-1280-y>.
- Trueb RM. Molecular mechanisms of androgenetic alopecia. *Exp Gerontol.* 2002;37(8-9):981-90. [PubMed ID: 12213548]. [https://doi.org/10.1016/s0531-5565\(02\)00093-1](https://doi.org/10.1016/s0531-5565(02)00093-1).
- Wang TL, Zhou C, Shen YW, Wang XY, Ding XL, Tian S, et al. Prevalence of androgenetic alopecia in China: a community-based study in six cities. *Br J Dermatol.* 2010;162(4):843-7. [PubMed ID: 20105167]. <https://doi.org/10.1111/j.1365-2133.2010.09640.x>.
- Rathnayake D, Sinclair R. Male androgenetic alopecia. *Expert Opin Pharmacother.* 2010;11(8):1295-304. [PubMed ID: 20426708]. <https://doi.org/10.1517/14656561003752730>.
- Olsen EA. Female pattern hair loss. *J Am Acad Dermatol.* 2001;45(3 Suppl):S70-80. [PubMed ID: 11511856]. <https://doi.org/10.1067/mjd.2001.117426>.
- Vujovic A, Del Marmol V. The female pattern hair loss: review of etiopathogenesis and diagnosis. *Biomed Res Int.* 2014;2014:767628. [PubMed ID: 24812631]. [PubMed Central ID: PMC400932]. <https://doi.org/10.1155/2014/767628>.
- Karadag Kose O, Gulec AT. Clinical evaluation of alopecias using a handheld dermatoscope. *J Am Acad Dermatol.* 2012;67(2):206-14. [PubMed ID: 22024772]. <https://doi.org/10.1016/j.jaad.2011.08.019>.
- Severi G, Sinclair R, Hopper JL, English DR, McCredie MR, Boyle P, et al. Androgenetic alopecia in men aged 40-69 years: prevalence and risk factors. *Br J Dermatol.* 2003;149(6):1207-13. [PubMed ID: 14674898]. <https://doi.org/10.1111/j.1365-2133.2003.05565.x>.
- Trueb RM. Is androgenetic alopecia a photoaggravated dermatosis? *Dermatology.* 2003;207(4):343-8. [PubMed ID: 14657623]. <https://doi.org/10.1159/000074111>.
- Gupta AK, Bluhm R. Seborrheic dermatitis. *J Eur Acad Dermatol Venereol.* 2004;18(1):13-26. quiz 19-20. [PubMed ID: 14678527]. <https://doi.org/10.1111/j.1468-3083.2004.00693.x>.
- Gupta AK, Bluhm R, Cooper EA, Summerbell RC, Batra R. Seborrheic dermatitis. *Dermatol Clin.* 2003;21(3):401-12. [PubMed ID: 12956195]. [https://doi.org/10.1016/s0733-8635\(03\)00028-7](https://doi.org/10.1016/s0733-8635(03)00028-7).
- Sampaio AL, Mameri AC, Vargas TJ, Ramos-e-Silva M, Nunes AP, Carneiro SC. Seborrheic dermatitis. *An Bras Dermatol.* 2011;86(6):1061-71. quiz 1072-4. [PubMed ID: 22281892]. <https://doi.org/10.1590/s0365-05962011000600002>.
- Kibar M, Aktan S, Bilgin M. Scalp dermatoscopic findings in androgenetic alopecia and their relations with disease severity. *Ann Dermatol.* 2014;26(4):478-84. [PubMed ID: 25143677]. [PubMed Central ID: PMC4135103]. <https://doi.org/10.5021/ad.2014.26.4.478>.
- Pitney L, Weedon D, Pitney M. Is seborrheic dermatitis associated with a diffuse, low-grade folliculitis and progressive cicatricial alopecia? *Australas J Dermatol.* 2016;57(3):e105-7. [PubMed ID: 25753934]. <https://doi.org/10.1111/ajd.12289>.
- Sarlak M, Shakib P, Zia MA, Madani M. The Prevalence of Malassezia yeasts in patients with seborrheic dermatitis by PCR-RFLP method in Isfahan, Iran. *Int J Med Invest.* 2020;9(4):21-8.
- Jang WS, Son IP, Yeo IK, Park KY, Li K, Kim BJ, et al. The annual changes of clinical manifestation of androgenetic alopecia clinic in Korean males and females: a outpatient-based study. *Ann Dermatol.* 2013;25(2):181-8. [PubMed ID: 23717009]. [PubMed Central ID: PMC3662911]. <https://doi.org/10.5021/ad.2013.25.2.181>.
- Ummi A, Priya PS, Chandravathi PL, Kumar CS. Correlation of Trichoscopic Findings in Androgenetic Alopecia and the Disease Severity. *Int J Trichology.* 2019;11(3):118-22. [PubMed ID: 31360040]. [PubMed Central ID: PMC6580806]. <https://doi.org/10.4103/ij.t.103-17>.
- Hu R, Xu F, Han Y, Sheng Y, Qi S, Miao Y, et al. Trichoscopic findings of androgenetic alopecia and their association with disease severity. *J Dermatol.* 2015;42(6):602-7. [PubMed ID: 25810236]. <https://doi.org/10.1111/1346-8138.12857>.