



# Effectiveness of the Unified Protocol for Transdiagnostic Treatment in Reducing Depression Associated with Marital Problems

Mahsa Bameshgi <sup>1</sup>, Seyed Ali Kimiaei <sup>2, \*</sup> and Ali Mashhadi <sup>1</sup>

<sup>1</sup>Department of Clinical Psychology, Faculty of Education and Psychology, Ferdowsi University of Mashhad, Mashhad, Iran

<sup>2</sup>Department of Counseling and Education Psychology, Faculty of Education and Psychology, Ferdowsi University of Mashhad, Mashhad, Iran

\*Corresponding author: Department of Counseling and Education, Faculty of Education and Psychology, Ferdowsi University of Mashhad, Mashhad, Iran. Tel: +98-9155008538, Email: kimiaee@um.ac.ir

Received 2019 May 10; Revised 2019 June 14; Accepted 2019 June 16.

## Abstract

**Background:** Different methods have been used to treat depression caused by marital relationships.

**Objectives:** The present study was conducted to determine the effectiveness of the unified protocol for transdiagnostic treatment in decreasing depression associated with marital problems.

**Methods:** The present controlled clinical trial was conducted on a statistical population comprising all women with depression associated with marital problems presenting to psychological clinics in Mashhad, Iran. Thirty five of these patients were randomly selected and assigned to an intervention group and a control group. Twelve sessions of group therapy were held in the intervention group based on the unified protocol for transdiagnostic disorders, whereas members of the control group participated in a language course. The structured clinical interview for DSM-IV (SCID), the Beck Depression Inventory-II and the communication patterns questionnaire were used to collect the data.

**Results:** The pre-treatment and post-treatment scores obtained from ANCOVA showed significant differences in depression ( $P < 0.05$ ), the mutual avoidance of communication subscale ( $P < 0.05$ ), the mutual constructive communication subscale ( $P < 0.05$ ) and the demand/withdraw subscale ( $P < 0.05$ ).

**Conclusions:** According to the obtained findings, the unified protocol for transdiagnostic treatment can be considered effective in improving the symptoms of depression associated with marital relationships.

**Keywords:** Transdiagnostic Treatment, Depression, Marital Problems

## 1. Background

Communication patterns refer to different coupling methods that are commonly used in the family, including (1) Mutual constructive communication, i.e. men and women try to talk about their problems, show their feelings and find solutions to their problems, (2) Mutual avoidance of communication, i.e. both men and women try to avoid talking about their problems, and (3) demand/withdraw, i.e. the woman or man discusses their problems, criticizes and nag while the other side avoids discussing. Different studies have shown that marital conflicts and communication problems can play a key role in the incidence and exacerbation of depression symptoms (1, 2). Depression is an emotional disorder, and emotional regulation is an important mechanism contributing to the formation of emotional disturbances (3). Emotional regulation also affects interpersonal relationships, especially marital relationships (4). A two-way relationship therefore exists between marital communication and

depression symptoms, i.e. their emergence and exacerbation. Whisman and Steven found depressed married participants to exhibit a poorer marital adjustment than the non-depressed ones (5). Another study discovered negative correlations between depression and marital adjustment, and found the solving styles of marital conflict to be the predictors of depression in depressed participants (6). These studies suggest the importance of further examining depressive disorders in terms of interpersonal relationships and communication patterns, although they have failed to propose any appropriate interventions.

Although depression is a major debilitating disorder, it can be satisfactorily treated using proper treatments. Different treatments proposed so far for this disorder have resulted in different outcomes (7). A study examining the effectiveness of cognitive therapy in depressed people found this treatment in combination with antidepressants to be useful for chronic depression (8). Forman et al. showed that acceptance and commitment therapy as a third-wave

therapy and cognitive therapy are effective in patients with anxiety and depression symptoms, and their only difference to lie in their mechanism of action (9). A review study recommended that certain treatments be designed to help couples support each other in their adaptation and coping with the symptoms of depression experienced during marital relationships (10). Studies conducted so far to evaluate the effectiveness of different treatments have usually proposed a different protocol for every disorder, which optimistically caused negligible to moderate effects, while rarely addressing comorbid symptoms. Some therapeutic protocols proposed in evidence-based interventions were based on cognitive-behavioral approaches as third-wave therapies (11), including the unified protocol for transdiagnostic treatment.

This protocol was introduced in response to the limitations of specific cognitive-behavioral therapies for disorders, in which the principles and treatment protocols were identical for different emotional disorders (3). This protocol, which is designed for people with emotional disorders, especially mood and anxiety disorders (12), is focused on emotions and not dedicated to a special disorder and encompasses common factors, including negative emotions (13). A combination of depression with other psychiatric disorders and symptoms (14) can encourage the use of this protocol. Group therapy being found to be as beneficial as individual treatments justifies the use of these therapies as cost-effective methods. A study by de Ornelas Maia et al. (15) on the unified protocol for transdiagnostic group treatment in people with depression and comorbid anxiety disorder demonstrated the effectiveness of the protocol in improving the symptoms of depression, anxiety and sexual dysfunction and quality of life. Bullis et al. also found the protocol effective in depression, anxiety, performance mal-function and avoidance experiences (16). An intervention conducted in Japan caused the reduction in anxiety and depression symptoms to remain stable for three months (17). These studies prove the applicability of this protocol to emotional disturbances, and further studies are required for obtaining more concrete evidence.

## 2. Objectives

Given a lack of studies on the effectiveness of the unified protocol for transdiagnostic treatment, the present research was performed to investigate this effectiveness in the reduction of depression associated with marital problems.


## 3. Methods

The present applied controlled clinical trial was conducted on a statistical population comprising all the mar-

ried women referred by counseling centers in Mashhad to the Polyclinic Psychological and Counseling Services Center at Ferdowsi University of Mashhad. The eligible candidates were randomly assigned to an intervention group and a control group. The questionnaires were distributed among 41 participants who were initially interviewed. Seventeen of 35 participants selected according to the interviews and the scores obtained from the questionnaires were randomly assigned to the intervention group and 18 to the control group (Figure 1). The inclusion criteria consisted of having marital problems based on the interviews, their communication satisfying the demand/withdraw or mutual avoidance models, having depression diagnosed based on the therapist's assessment through a semi-structured DSM-IV clinical interview, receiving a score of over 19 from the Beck depression inventory, the absence of premarital depression, an education level of at least high school diploma and no history of using psychiatric medications. Both groups completed the Beck depression inventory and the communication patterns questionnaire. The intervention group participated in twelve 60-minute sessions of intervention based on the unified protocol for transdiagnostic treatment of emotional disorders, and the control group received an intervention, namely English language training as the placebo. After the treatment, both groups completed the questionnaires again.

The structured clinical interview for DSM-IV (SCID) is a semi-structured interview based on DSM-IV in two versions: SCID-I for axial I disorders and SCID-II for axial II disorders. This questionnaire was developed by First, Spitzer, Gibbon and Williams in 1997 (18). The reliability and validity of this questionnaire have been confirmed in literature (18), and its kappa coefficient investigated in Iran and reported as approximately 0.7 (19). Sharifi et al. reported a diagnostic consensus on most of the specific and general diagnoses ranging from moderate to good, i.e. Kappa over 0.60, an overall agreement, i.e. total Kappa of 0.52, for current diagnoses, and 0.55 for total lifetime diagnoses (19). Moreover, many of the interviewees and interviewers reported the feasibility of implementing the Persian version of this interview (19).

The 35-item communication patterns questionnaire was designed as a self-assessment tool by Christensen and Sullaway to measure marital patterns (20), and estimate couples' behaviors at three stages of marital conflict, including (a) Upon the emergence of a problem in the relationship of the couples (b) During the discussion of the communication problem (c) After discussing the communication problem. Subjects rank each behavior on a nine-point Likert scale ranging from 1: totally impossible to 9: quite likely. This questionnaire consists of three sub-scales,



**Figure 1.** The research flowchart

namely demand/withdraw, mutual constructive communication and mutual avoidance of communication (20). The correlation coefficients for the subscales were all significant, and obtained as follows: mutual constructive communication (five items): 0.85, mutual avoidance of communication: -0.85 and demand/withdraw: -0.35 (21).

The 21-item Beck depression inventory-second edition (BDI-II) is a widely-used self-reporting instrument for depression, and is rated on a four-point scale ranging from 0 to 3. This inventory is normally used to evaluate the emotional, cognitive, motivational and behavioral symptoms of depression. A test-retest reliability coefficient of 0.48 - 0.86 (22), an internal consistency of 0.87 and a high credibility coefficient of 0.74 were reported for this questionnaire (23).

The unified protocol for transdiagnostic treatment was adopted from the protocol proposed by Barlow et al. (3) comprising 12 sessions based on six components (Table 1). While emphasizing the fundamental principles of cognitive-behavioral therapy and integrating new advances in emotional regulation research, the therapist uses this approach to treat patients with comorbid emotional disorders by making efforts to develop cognitive-behavioral strategies, including increasing awareness of emotions, teaching emotional and behavioral avoidance techniques, and also helping with the identification and correction of non-adaptive cognition (3).

#### 4. Results

Out of the 35 participants, 2 in the intervention group and 3 controls withdrew before the end of the sessions. The data of 15 participants in each group were therefore evaluated. The mean age of the intervention group was  $32.66 \pm 9.39$  years and that of the control group  $24.4 \pm 1.04$ . The mean duration of marriage was  $7.33 \pm 8.8$  years in the intervention group and  $3.93 \pm 3.12$  in the control group. Twelve (80%) subjects in the intervention group and 13 (86.6%) in the control group had an undergraduate level of education, which was the most frequent level in both groups. Table 2 presents the descriptive indices associated with the pre-test and post-test scores of depression and the components of couples' communication patterns in the intervention and control groups. The two groups were significantly different in terms of all the study variables before the intervention. The status of the intervention group was generally worse than that of the controls according to the target variables. According to Table 2, the mean post-intervention scores of depression and marital communication in the intervention group were significantly different from the baseline (pre-intervention) measurements, while the mean scores were not significantly different in the control group at either stages.

ANCOVA was used to determine the potentially-significant differences between the groups in terms of the variables. The hypotheses of ANCOVA were examined to

**Table 1.** Sessions, Modules and Core Content of Sessions

Session	Content
1	Increasing motivation, motivational interviewing for participation and involvement of patients, presentation of treatment rationale and determination of treatment goals (module 1).
2	Presentation of psychoeducation, recognition of emotions and tracking emotional experiences, teaching the main components of emotional experience (module 2).
3	Emotional awareness training, learning to view emotional experiences (emotion and reaction to emotions) especially using mindfulness techniques (module 3).
4	Cognitive appraisal and reappraisal, making awareness of the impact and interaction between thoughts and emotions, identifying autonomic maladaptive appraisal, common thinking traps and increasing the flexibility of thinking (module 4).
5	Identifying emotional avoidance patterns, familiarizing with different strategies for avoiding emotions and their impact on emotional experiences and knowing the contradictory effects of emotional avoidance (module 5).
6	Emotion-driven behaviors study (EDBS), familiarity and identification of emotion-driven behaviors, understanding their Effects on emotional experiences, identifying maladaptive EDBSs (module 5).
7	Knowledge and tolerance of physical senses, increase awareness of the role of emotional feelings in emotional experiences, practice exercises or visceral confrontation in order to be aware of physical sensations and increase the tolerance of these symptoms (module 6).
8 - 11	Visceral confrontation and confrontation with situational emotions, awareness of the rationale of emotional confrontation, teaching how to prepare a fear and avoidance hierarchy, and designing repeated and effective emotional exercises (module 7).
12	Prevention of relapse, overview of the treatment content and patient progress, identify the ways in which treatment advantages maintain and predict future difficulties (module 8).

**Table 2.** Mean and Standard Deviation of Depression Scores and Communication Patterns of Two Groups in Pre-Test and Post-Test<sup>a</sup>

Variable/Stage	Intervention Group	Control Group	P Value
<b>Depression</b>			0.02
Pre-test	30.4 ± 5.44	25.6 ± 5.67	
Post-test	15.53 ± 6.64	24.4 ± 5.2	
<b>Mutual avoidance</b>			0.04
Pre-test	17.73 ± 5.95	14.33 ± 3.28	
Post-test	10.4 ± 5.52	15.06 ± 2.78	
<b>Mutual constructive</b>			0.02
Pre-test	19.86 ± 5.75	24.33 ± 5.28	
Post-test	27.6 ± 7.46	22.13 ± 5.23	
<b>Demand/withdraw</b>			0.01
Pre-test	34 ± 6.46	29.06 ± 11.09	
Post-test	26.26 ± 8.44	29.86 ± 11.26	

<sup>a</sup>Values are expressed as mean ± SD.

compare the intervention and control groups in terms of depression scores before performing this test. The results of the Kolmogorov-Smirnov test ( $P > 0.05$ ) confirmed the distribution normality of this variable. The results of the Leven's test also confirmed the homogeneity of variances ( $P = 0.19$ ,  $F = 2.67$ ). The third hypothesis regarding the homogeneity of the regression slope was also confirmed ( $P = 0.43$ ,  $F = 0.62$ ). ANCOVA was therefore applicable given the confirmation of all the three hypotheses. Table 3 suggests significant differences between the participants in the intervention and control group in terms of depression, as the severity of depression in the intervention group

significantly decreased compared to in the control group in the post-test. ANCOVA was used to investigate the effectiveness of the treatment whose effect size was found to be 0.49. The relevant hypotheses were examined before performing this test.

Table 4 suggests significant differences in communication patterns between the participants receiving the unified protocol for transdiagnostic treatment and those in the control group receiving no treatments (mutual avoidance of communication:  $P = 0.001$ ,  $F = 21.35$ ; mutual constructive communication:  $P = 0.001$ ,  $F = 22.78$ ; demand/withdraw:  $P = 0.001$ ,  $F = 24.6$ ). In other words, mu-

**Table 3.** The Results of Covariance Analysis for Comparing Intervention and Control Groups in the Depression Scale

Variable	Sum of Squares	Df	Mean of Squares	F	P Value	Effect Size
Pre-test	194.03	1	194.03	6.52	0.01	0.19
Group	775.7	1	775.7	26.07	0.001	0.49
Error	803.3	27	29.75			
Total	13547	30				

tual constructive communication pattern was found to significantly increase in the posttest in the intervention compared to control group, and mutual avoidance and demand/withdraw to significantly decrease. The unified protocol for transdiagnostic treatment was therefore effective in improving the communication patterns of the participants in the post-test. The size of this effect was 44% for mutual avoidance, 51% for mutual constructive and 47% for demand/withdraw.

## 5. Discussion

The present study was conducted to investigate the effectiveness of the unified protocol for transdiagnostic treatment. The results obtained provide empirical evidence for the efficacy of an intervention derived from cognitive-behavioral therapies, which can be applied at clinics.

The present findings suggested significant differences between the mean pre-test and the mean post-test in the intervention group. In other words, the unified protocol for transdiagnostic treatment reduced the depression associated with marital problems in the married women. The clinical improvements observed were consistent with studies by de Ornelas Maia et al. (15), Ito et al. (17) and Bullis et al. (16). These findings can be explained by the beneficial effects associated with many advances in the conceptualization and treatment of mood disorders, especially as the recently-proposed cognitive-behavioral theory, on behavioral patterns and maladaptive thinking as important features of emotional disturbances (24). Cognitive therapy is the standard treatment for depression, as patients with depression often tend to attribute negative events to internal self-conscious sustained and inclusive causes. Having such a documentary style results in reinforcing the notion that negative events are likely to return in the future following different issues, which can cause a lot of frustration (25). In other words, it is not the situations, events or triggers that directly causes an emotional response; rather, the cognitive assessment of the positions, events or triggers leads to this response. Identifying the thoughts associated with emotions is therefore crucial (26).

In the fourth module, the protocol is taught to the participants to recognize the role of maladaptive auto-evaluations in creating emotional experiences. The fifth module emphasizes the emotional experience of behavioral components. In this part of the treatment, the therapist helps the authorities identify emotional patterns and emotional-driven behaviors. After the cognitive authority has more to do with the effects of these behaviors on continuing discomfort, measures are taken to change the current patterns of emotional responses (3). The most recent psychopathological theories emphasize the role of emotions in developing, maintaining and continuing the symptoms of emotional disorders and many other mental disorders (24). Despite being rooted in cognitive-behavioral conventions, this protocol is unique. This treatment emphasizes the adaptive and functional nature of emotions, and its effectiveness in depressive disorders can be explained by primarily seeking to identify and correct inappropriate attempts, regulate emotional experiences, and therefore facilitate proportional processing and suppress disproportionate emotional responses to internal (visceral) and external symptoms (27).

The present study found significant differences between the mean pretest and posttest values in the intervention group. In fact, the unified protocol for transdiagnostic treatment reduced mutual avoidance of communication and the demand/withdraw pattern, and increased the mutual constructive communication model in the married women. Confirming these hypotheses is consistent with the findings of Etemadi et al. (28), who found marital satisfaction to be significantly higher in the intervention compared to the control group. The findings obtained by de Ornelas Maia et al. in 2017 are consistent with the recent hypotheses suggesting that this protocol with an improved emotional management serves as a potential source of improving the quality of life and sexual functioning (15). As the first explanation, depression appears to be positively correlated with marital conflicts and communication problems in couples (6, 29), and this relationship appears more significant in women than men (29, 30). The largest influence on interpersonal performance has been observed in women (30). Given the negative relationships of marital satisfaction and quality of life with the dura-

**Table 4.** The Results of Covariance Analysis for Comparing Intervention and Control Groups in the Components of Communication Patterns Scale

Dependent	Sum of Squares	Df	F	P Value	Effect Size
Mutual avoidance	278.01	1	21.35	0.001	0.44
Mutual constructive	556.23	1	28.77	0.001	0.51
Demand/withdraw	496.33	1	24.6	0.001	0.47

tion of untreated depression (31), treating depression in women can predict improvements in marital satisfaction and communication patterns in couples.

According to the present findings, couples who use the mutual constructive model and marital relationships struggle to maintain healthy relationships, avoid abusive behaviors, feel that they understand each other, present themselves well, negotiate to solve communication problems, and are more satisfied with their married life. On the other hand, couples who use demand/withdraw patterns present malicious behaviors, including criticizing, annoying, suggesting another change, not trying to establish and maintain a healthy relationship and avoiding discussing the problem. These couples cannot properly express their feelings, and do not negotiate a solution to their communication problems, which influences their lives and exerts damaging effects on their family (32). The desirable performance of the family and the mutual constructive communication model predict marital satisfaction (32, 33).

As a second explanation, emotional regulation is generally considered an important factor for successful interpersonal communication. Recent research suggests significant relationships between marital satisfaction and the interpersonal emotional regulation process. Emotional ordering in women during an unpleasant experience predicts more marital satisfaction in both women and their spouses, which suggests the importance of regulating negative emotions for marital satisfaction in women during conflicts (34). Given the key role of emotional regulation skills in improving couple's communication patterns, the unified protocol can be effective in improving the spouse's communication patterns through improving marital satisfaction. The main hypothesis of this treatment is that people with emotional disorders resort to procedural strategies against maladaptive emotions, and essentially make efforts to avoid or reduce unpleasant emotions.

In line with the present study, Hashemi and Kimiaei found emotion- focused cognitive therapy to reduce the depression caused by marital problems and the patterns of mutual avoidance and demand/withdraw in the intervention group (35) and Mahlabani Gorgian et al. found emotion-focused couple therapy to be effective in treating the depression caused by couples' communication prob-

lems (36).

The present study limitations included selecting the samples from the patients referred to the psychiatric and counseling clinics of Ferdowsi University of Mashhad. The obtained results should therefore be cautiously generalized to other affected individuals. Follow-ups were also not conducted due to time constraints. Given that all the study participants were female, future research is recommended to focus on both genders.

### 5.1. Conclusions

According to the present findings, the unified protocol for transdiagnostic treatment can be considered a useful short-term cost-effective treatment for improving the symptoms of depression caused by marital problems. Solving interpersonal problems can also help treat depressed patients.

### Footnotes

**Authors' Contribution:** Study concept and design: Mahsa Bameshgi and Seyed Ali Kimiaei. Acquisition of data: Mahsa Bameshgi. Analysis and interpretation of data: Mahsa Bameshgi. Drafting of the manuscript: Mahsa Bameshgi, Seyed Ali Kimiaei and Ali Mashhadi. Critical revision of the manuscript for important intellectual content: Seyed Ali Kimiaei and Ali Mashhadi. Statistical analysis: Mahsa Bameshgi. Administrative, technical, and material support: Mahsa Bameshgi, Seyed Ali Kimiaei and Ali Mashhadi. Study supervision: Seyed Ali Kimiaei and mashhadi.

**Conflict of Interests:** It is not declared by the authors.

**Ethical Considerations:** IR.UM.REC.1398.008.

**Funding/Support:** This study was supported in part by Ferdowsi University of Mashhad.

### References

1. Kiecolt-Glaser JK, Jaremka L, Andridge R, Peng J, Habash D, Fagundes CP, et al. Marital discord, past depression, and metabolic responses to high-fat meals: Interpersonal pathways to obesity. *Psychoneuroendocrinology*. 2015;52:239-50. doi: [10.1016/j.psyneuen.2014.11.018](https://doi.org/10.1016/j.psyneuen.2014.11.018). [PubMed: 25506778]. [PubMed Central: PMC4297566].

2. Fisher SD, Brock RL, O'Hara MW, Kopelman R, Stuart S. Longitudinal contribution of maternal and paternal depression to toddler behaviors: Interparental conflict and later depression as mediators. *Couple Fam Psychol*. 2015;**4**(2):61-73. doi: [10.1037/cfp0000037](https://doi.org/10.1037/cfp0000037).
3. Barlow DH, Farchione TJ, Sauer-Zavala S, Murray Latin H, Ellard KK, Bullis JR, et al. *Unified protocol for transdiagnostic treatment of emotional disorders: Therapist guide*. Oxford University Press; 2017. doi: [10.1093/med-psych/9780190685973.001.0001](https://doi.org/10.1093/med-psych/9780190685973.001.0001).
4. Frankel LA, Umemura T, Jacobvitz D, Hazen N. Marital conflict and parental responses to infant negative emotions: Relations with toddler emotional regulation. *Infant Behav Dev*. 2015;**40**:73-83. doi: [10.1016/j.infbeh.2015.03.004](https://doi.org/10.1016/j.infbeh.2015.03.004). [PubMed: [26047678](https://pubmed.ncbi.nlm.nih.gov/26047678/)].
5. Whisman MA, Beach SR. Couple therapy for depression. *J Clin Psychol*. 2012;**68**(5):526-35. doi: [10.1002/jclp.21857](https://doi.org/10.1002/jclp.21857). [PubMed: [22499085](https://pubmed.ncbi.nlm.nih.gov/22499085/)].
6. Ozguc S, Tanriverdi D. Relations between depression level and conflict resolution styles, marital adjustments of patients with major depression and their spouses. *Arch Psychiatr Nurs*. 2018;**32**(3):337-42. doi: [10.1016/j.apnu.2017.11.022](https://doi.org/10.1016/j.apnu.2017.11.022). [PubMed: [29784211](https://pubmed.ncbi.nlm.nih.gov/29784211/)].
7. Segal ZV, Teasdale J. *Mindfulness-based cognitive therapy for depression*. Guilford Publications; 2018.
8. Rupke SJ, Blecke D, Renfrow M. Cognitive therapy for depression. *Am Fam Physician*. 2006;**73**(1):83-6. [PubMed: [16417069](https://pubmed.ncbi.nlm.nih.gov/16417069/)].
9. Forman EM, Herbert JD, Moitra E, Yeomans PD, Geller PA. A randomized controlled effectiveness trial of acceptance and commitment therapy and cognitive therapy for anxiety and depression. *Behav Modif*. 2007;**31**(6):772-99. doi: [10.1177/0145445507302202](https://doi.org/10.1177/0145445507302202). [PubMed: [17932235](https://pubmed.ncbi.nlm.nih.gov/17932235/)].
10. Mead DE. Marital distress, co-occurring depression, and marital therapy: A review. *J Marital Fam Ther*. 2002;**28**(3):299-314. doi: [10.1111/j.1752-0606.2002.tb01188.x](https://doi.org/10.1111/j.1752-0606.2002.tb01188.x). [PubMed: [12197153](https://pubmed.ncbi.nlm.nih.gov/12197153/)].
11. Dobson KS, Dozois DJ. *Handbook of cognitive-behavioral therapies*. Guilford Publications; 2019.
12. Barlow DH, Farchione TJ, Bullis JR, Gallagher MW, Murray-Latin H, Sauer-Zavala S, et al. The unified protocol for transdiagnostic treatment of emotional disorders compared with diagnosis-specific protocols for anxiety disorders: A randomized clinical trial. *JAMA Psychiatry*. 2017;**74**(9):875-84. doi: [10.1001/jamapsychiatry.2017.2164](https://doi.org/10.1001/jamapsychiatry.2017.2164). [PubMed: [28768327](https://pubmed.ncbi.nlm.nih.gov/28768327/)]. [PubMed Central: [PMC5710228](https://pubmed.ncbi.nlm.nih.gov/PMC5710228/)].
13. Laposa JM, Mancuso E, Abraham G, Loli-Dano L. Unified protocol transdiagnostic treatment in group format. *Behav Modif*. 2017;**41**(2):253-68. doi: [10.1177/0145445516667664](https://doi.org/10.1177/0145445516667664). [PubMed: [27591430](https://pubmed.ncbi.nlm.nih.gov/27591430/)].
14. Kaplan H, Kaplan SB. *Sadock's synopsis of psychiatry: Behavioral sciences/clinical psychiatry*. Tehran: Arjmand pub; 2014. p. 43-447.
15. de Ornelas Maia ACC, Sanford J, Boettcher H, Nardi AE, Barlow D. Improvement in quality of life and sexual functioning in a comorbid sample after the unified protocol transdiagnostic group treatment. *J Psychiatr Res*. 2017;**93**:30-6. doi: [10.1016/j.jpsychires.2017.05.013](https://doi.org/10.1016/j.jpsychires.2017.05.013). [PubMed: [28575646](https://pubmed.ncbi.nlm.nih.gov/28575646/)].
16. Bullis JR, Fortune MR, Farchione TJ, Barlow DH. A preliminary investigation of the long-term outcome of the unified protocol for transdiagnostic treatment of emotional disorders. *Compr Psychiatry*. 2014;**55**(8):1920-7. doi: [10.1016/j.comppsy.2014.07.016](https://doi.org/10.1016/j.comppsy.2014.07.016). [PubMed: [25113056](https://pubmed.ncbi.nlm.nih.gov/25113056/)]. [PubMed Central: [PMC4252968](https://pubmed.ncbi.nlm.nih.gov/PMC4252968/)].
17. Ito M, Horikoshi M, Kato N, Oe Y, Fujisato H, Nakajima S, et al. Transdiagnostic and transcultural: Pilot study of unified protocol for depressive and anxiety disorders in Japan. *Behav Ther*. 2016;**47**(3):416-30. doi: [10.1016/j.beth.2016.02.005](https://doi.org/10.1016/j.beth.2016.02.005). [PubMed: [27157034](https://pubmed.ncbi.nlm.nih.gov/27157034/)].
18. First MB, Gibbon M. The Structured clinical interview for DSM-IV axis I disorders (SCID-I) and the structured clinical interview for DSM-IV axis II disorders (SCID-II). *Comprehensive handbook of psychological assessment, Vol. 2. Personality assessment*. 2. Hoboken, NJ, US: John Wiley & Sons Inc; 2004. p. 134-43.
19. Sharifi V, Asadi M, Mohammadi M, Kaviani H, Semanani Y, Shabani A. [Reliability and usability of Persian version of diagnosis structured interview for DSM-IV]. *J Cogn Sci Novels*. 2004;**6**:1&2. Persian.
20. Christensen A, Sullaway M. *Communication patterns questionnaire. Unpublished manuscript*. Los Angeles: University of California; 2012. doi: [10.1037/t02529-000](https://doi.org/10.1037/t02529-000).
21. Farmani Shahreza S, Rasouli M, Ghaedaniya Jahromi A. [Relationship between marital intimacy, patterns of marital relationship and adolescent companionship among married workers of Kharazmi University]. *Q Counsel Psychother Fam*. 2012;**4**(3):456-7. Persian.
22. Beck AT, Steer RA, Brown G. *Beck depression inventory-II*. San Antonio; 2011. doi: [10.1037/t00742-000](https://doi.org/10.1037/t00742-000).
23. Ghassemzadeh H, Mojtabai R, Karamghadiri N, Ebrahimkhani N. Psychometric properties of a Persian-language version of the Beck Depression inventory-Second edition: BDI-II-PERSIAN. *Depress Anxiety*. 2005;**21**(4):185-92. doi: [10.1002/da.20070](https://doi.org/10.1002/da.20070). [PubMed: [16075452](https://pubmed.ncbi.nlm.nih.gov/16075452/)].
24. Mohammadi A, Zargar F, Omidi A. *Introduction to the third generation of cognitive-behavioral therapies*. Tehran: Arjmand Publication; 2015. Persian.
25. Abramson LY, Seligman ME, Teasdale JD. Learned helplessness in humans: Critique and reformulation. *J Abnorm Psychol*. 1978;**87**(1):49-74. doi: [10.1037/0021-843X.87.1.49](https://doi.org/10.1037/0021-843X.87.1.49). [PubMed: [649856](https://pubmed.ncbi.nlm.nih.gov/649856/)].
26. Mohammadkhani P, Abasi I, Poursahbaz A, Mohammadi A, Fatehi M. The role of neuroticism and experiential avoidance in predicting anxiety and depression symptoms: Mediating effect of emotion regulation. *Iran J Psychiatry Behav Sci*. 2016;**10**(3). e5047. doi: [10.17795/ijpbs-5047](https://doi.org/10.17795/ijpbs-5047). [PubMed: [27822282](https://pubmed.ncbi.nlm.nih.gov/27822282/)]. [PubMed Central: [PMC5097448](https://pubmed.ncbi.nlm.nih.gov/PMC5097448/)].
27. Wilamowska ZA, Thompson-Hollands J, Fairholme CP, Ellard KK, Farchione TJ, Barlow DH. Conceptual background, development, and preliminary data from the unified protocol for transdiagnostic treatment of emotional disorders. *Depress Anxiety*. 2010;**27**(10):882-90. doi: [10.1002/da.20735](https://doi.org/10.1002/da.20735). [PubMed: [20886609](https://pubmed.ncbi.nlm.nih.gov/20886609/)].
28. Etemadi A, Gholizadeh H, Salimi Bajestani H, Farahbakhsh K. [The effectiveness of analytical-functional couple therapy with unified protocol for transdiagnostic treatment in depression, anxiety and marital satisfaction of women affected by marital disturbances]. *J Thought Behav*. 2017;**12**(46):47-56. Persian.
29. Ellison JK, Kouros CD, Papp LM, Cummings EM. Interplay between marital attributions and conflict behavior in predicting depressive symptoms. *J Fam Psychol*. 2016;**30**(2):286-95. doi: [10.1037/fam0000181](https://doi.org/10.1037/fam0000181). [PubMed: [26751758](https://pubmed.ncbi.nlm.nih.gov/26751758/)]. [PubMed Central: [PMC4767684](https://pubmed.ncbi.nlm.nih.gov/PMC4767684/)].
30. Hou Y, Jiang F, Wang X. Marital commitment, communication and marital satisfaction: An analysis based on actor-partner interdependence model. *Int J Psychol*. 2019;**54**(3):369-76. doi: [10.1002/ijop.12473](https://doi.org/10.1002/ijop.12473). [PubMed: [29318606](https://pubmed.ncbi.nlm.nih.gov/29318606/)].
31. Aggarwal S, Kataria D, Prasad S. A comparative study of quality of life and marital satisfaction in patients with depression and their spouses. *Asian J Psychiatr*. 2017;**30**:65-70. doi: [10.1016/j.ajp.2017.08.003](https://doi.org/10.1016/j.ajp.2017.08.003). [PubMed: [28802799](https://pubmed.ncbi.nlm.nih.gov/28802799/)].
32. Parvandi A, Arefi M, Moradi A. [The role of family function and couples' communication patterns in predicting marital satisfaction]. *J Pathol Counseling Fam Enrich*. 2016;**2**(1):54-65. Persian.
33. Madahi ME, Samadzadeh M, Javidi N. The communication patterns & satisfaction in married students. *Procedia Soc Behav Sci*. 2013;**84**:1190-3. doi: [10.1016/j.sbspro.2013.06.725](https://doi.org/10.1016/j.sbspro.2013.06.725).
34. Bloch L, Haase CM, Levenson RW. Emotion regulation predicts marital satisfaction: More than a wives' tale. *Emotion*. 2014;**14**(1):130-44. doi: [10.1037/a0034272](https://doi.org/10.1037/a0034272). [PubMed: [24188061](https://pubmed.ncbi.nlm.nih.gov/24188061/)]. [PubMed Central: [PMC4041870](https://pubmed.ncbi.nlm.nih.gov/PMC4041870/)].
35. Hashemi SF, Kimiaei SA. The effectiveness of emotion-focused cognitive therapy in decreasing depression due to marital relationship problems. *Univ J Psychol*. 2017;**5**(4):196-203. doi: [10.13189/ujp.2017.050405](https://doi.org/10.13189/ujp.2017.050405).
36. Mahlabani Gorgian H, Kimiaei AS, Ganbari Hashem Abadi B. [Effectiveness of emotional-focused couple therapy to reduce depression due to marital problems]. *Psychol appl res*. 2011;**2**(4):75-89. Persian.