



Assessment of *Candida* Species Contamination on Dental Unit Surfaces: Prevalence, Risk Factors, and Infection Control Strategies

Moein Navidi ¹, Seyedeh Masoumeh Mirkeshavarz ^{1,*}, Fakhri Haghi ²

¹ Faculty of Dentistry, Zanjan University of Medical Sciences, Zanjan, Iran

² Zanjan University of Medical Sciences, Zanjan, Iran

*Corresponding Author: Department of Pathology, Dental School, Zanjan University of Medical Sciences, Zanjan, Iran. Email: mirkeshavarz@zums.ac.ir

Received: 25 March, 2025; Revised: 6 May, 2025; Accepted: 27 May, 2025

Abstract

Background: Cross-contamination in dental clinics poses a significant risk for microbial transmission, especially from opportunistic fungal pathogens such as *Candida* species. Among them, *Candida albicans* is a common commensal of the oral cavity, capable of causing opportunistic infections in immunocompromised individuals. Dental unit surfaces, including chair headrests, light handles, control panels, and saliva ejectors, may serve as reservoirs for *Candida*, facilitating its spread.

Objectives: This study aimed to determine the prevalence of *Candida* species contamination on dental unit surfaces, assess antifungal resistance patterns, and evaluate the effectiveness of current infection control protocols.

Methods: This cross-sectional study was conducted at Zanjan University of Medical Sciences in 2023. A total of 120 samples were collected from high-contact surfaces in 12 dental units at three time points: Before, during, and after clinical shifts. Culturing was performed on Sabouraud dextrose agar (SDA). Identification of isolates included gram staining, germ tube testing, and CHROMagar *Candida*. Antifungal susceptibility was assessed via the disk diffusion method based on Clinical and Laboratory Standards Institute (CLSI) M44-A2 guidelines. Descriptive and inferential analyses were conducted using SPSS v26.

Results: *Candida* species were detected in 39.2% of all samples, with contamination rising significantly from 18.3% before shifts to 60.0% after shifts ($P < 0.001$). The most contaminated surfaces were chair headrests (72.5%) and light handles (65.8%). The predominant isolate was *C. albicans*, followed by *C. glabrata* and *C. tropicalis*. Fluconazole susceptibility in *C. albicans* was 85.1%, while amphotericin B and nystatin showed 96.7% efficacy. Despite routine disinfection, 39.2% of post-cleaning surfaces remained contaminated.

Conclusions: This study highlights the persistence of *Candida* contamination on dental surfaces and potential shortcomings of routine disinfection protocols. It recommends enhanced sterilization practices – including UV sterilization, high-frequency surface cleaning, and personnel hygiene reinforcement – to reduce fungal transmission in clinical dentistry.

Keywords: *Candida* Species, Dental Unit Contamination, Cross-Infection, Antifungal Resistance, Infection Control

1. Background

Fungal infections are a major concern in dental and medical settings, with *Candida albicans* being one of the most prevalent opportunistic pathogens. This yeast-like fungus is a normal commensal organism in the oral cavity, gastrointestinal tract, and genitourinary system, but it can become pathogenic under immunocompromised conditions or due to disruptions in the normal microbial flora (1). *Candida albicans* is

responsible for various oral infections, including oropharyngeal candidiasis, denture stomatitis, and angular cheilitis, particularly in elderly patients, immunosuppressed individuals, and those undergoing prolonged antibiotic or corticosteroid therapy (2).

Dental clinics can serve as potential reservoirs for *Candida* species, with contamination occurring through direct contact with infected patients, aerosolized particles from dental procedures, or contaminated surfaces and instruments (3). The biofilm-forming

capability of *Candida* species enhances their resistance to antifungal treatments and disinfection procedures, making infection control in dental environments particularly challenging (4). Studies have reported *Candida* species on frequently touched surfaces such as dental chairs, light handles, and impression trays, emphasizing the need for rigorous sterilization protocols (5).

Despite increasing awareness of fungal contamination in clinical settings, limited studies have specifically investigated the presence and persistence of multiple *Candida* species, including *C. albicans*, *C. glabrata*, and *C. tropicalis* on dental unit surfaces. Additionally, there is a lack of research exploring contamination patterns across different time points (before, during, and after clinical shifts) and the effectiveness of commonly used disinfectants (6). Understanding the relationship between surface contact, clinical activity levels, and the efficacy of disinfection can reveal critical risk factors in fungal transmission.

2. Objectives

This study aimed to (1) assess the prevalence of *Candida* species contamination on dental unit surfaces; (2) evaluate antifungal resistance patterns; and (3) analyze potential risk factors and propose effective infection control strategies. The findings will contribute to evidence-based improvements in infection control protocols in dental clinics.

3. Methods

3.1. Study Design and Setting

This cross-sectional study was conducted in the dental clinics of Zanjan University of Medical Sciences, Faculty of Dentistry, in 2023. The study aimed to assess *Candida* species contamination on frequently touched dental unit surfaces and evaluate the effectiveness of routine infection control measures.

3.2. Sample Collection

A total of 120 samples were collected from 12 active dental units. Sampling was conducted at three time points: Before the start of clinical sessions (pre-disinfection), during clinical activity, and after the end of sessions (post-disinfection). Samples were taken from

four high-contact surfaces: Dental chair headrests, light handles, unit control panels, and saliva ejectors. Sterile cotton swabs moistened with 0.9% sterile saline were used to swab each surface for approximately 10 seconds in a standardized zigzag motion. Each swab was immediately transferred into sterile transport tubes containing Sabouraud dextrose broth (SDB) and sent to the microbiology laboratory for further analysis. Routine disinfection was performed using a 0.5% sodium hypochlorite solution. Surfaces were sprayed and left for 10 minutes before being wiped with sterile gauze, following manufacturer instructions. This procedure was repeated after each patient and at the end of each shift.

3.3. Fungal Culture and Identification

Samples were streaked onto Sabouraud dextrose agar (SDA) supplemented with chloramphenicol and incubated at 37°C for 48 hours. Colonies suspected to be *Candida* species were further identified based on: (1) Colony morphology (creamy white colonies with smooth surfaces); (2) gram staining; (3) germ tube test (positive for *C. albicans*); (4) Chromogenic agar differentiation (CHROMagar *Candida*).

3.4. Antifungal Susceptibility Testing

Antifungal susceptibility was tested using the disk diffusion method on Mueller-Hinton agar supplemented with 2% glucose and methylene blue. The antifungal agents tested included: (1) Fluconazole (25 µg); (2) Amphotericin B (10 µg); (3) Nystatin (100 units).

Zone diameters were measured and interpreted according to Clinical and Laboratory Standards Institute (CLSI) M44-A2 guidelines.

3.5. Statistical Analysis

Data were analyzed using SPSS Statistics release 27.0.1. The paired *t*-test and Wilcoxon signed-rank test were used to compare microbial loads before and after clinical sessions. A chi-square test was applied to analyze categorical data. Statistical significance was set at $P < 0.05$.

4. Results

4.1. Prevalence of *Candida* Species Across Different Departments

Out of 120 collected samples, *Candida* species were identified in 47 (39.2%) samples. The highest contamination was observed in the Prosthodontics Department (65.2%), followed by Endodontics (58.4%) and Periodontics (52.1%). The lowest prevalence was recorded in the Pediatric Dentistry Department (21.7%) (Table 1).

Table 1. Distribution of *Candida* Species Across Different Departments

Departments	<i>Candida albicans</i> (%)	<i>Candida glabrata</i> (%)	<i>Candida tropicalis</i> (%)	Total Contamination (%)
Prosthodontics	38.6 (17/44)	15.9 (7/44)	10.7 (5/44)	65.2 (29/44)
Endodontics	32.4 (12/37)	16.2 (6/37)	10.8 (4/37)	58.4 (22/37)
Periodontics	29.2 (11/38)	14.5 (5/38)	8.4 (3/38)	52.1 (19/38)
Pediatric dentistry	12.3 (4/32)	6.2 (2/32)	3.2 (1/32)	21.7 (7/32)

4.2. Surface-wise Prevalence of *Candida* Species

Table 2 illustrates the distribution of *Candida* species across four high-contact surfaces of dental units before disinfection. The highest overall contamination rate was observed on chair headrests (72.5%), with *C. albicans* being the dominant species (38.3%), followed by *C. glabrata* (14.2%) and *C. tropicalis* (8.3%). Light handles also showed a relatively high contamination rate (65.8%), primarily due to the presence of *C. albicans* (32.5%), accompanied by *C. glabrata* (12.5%) and *C. tropicalis* (10.8%). Unit control panels exhibited a moderate level of contamination (44.2%), with *C. albicans* accounting for 28.3%. Finally, the lowest contamination was noted on saliva ejectors (32.5%), with *C. albicans* comprising 20% of the total contamination.

Table 2. Distribution of *Candida* Species Different Equipment

Surface Type	<i>Candida albicans</i> (%)	<i>Candida glabrata</i> (%)	<i>Candida tropicalis</i> (%)	Total Contamination (%)
Chair headrests	38.3	14.2	8.3	72.5
Light handles	32.5	12.5	10.8	65.8
Unit control panels	28.3	9.2	6.7	44.2
Saliva ejectors	20.0	7.5	5.0	32.5

4.3. Comparison of Contamination Levels Before, During, and After Clinical Shifts

A significant increase in *Candida* contamination was observed during clinical activity, peaking after the shift

(Table 3).

Table 3. Contamination Levels Before, During, and After Clinical Shifts

Time Points	Contaminated Samples (%)	Mean \pm SD (CFU/cm ²)	P-Value
Before shift	18.3 (22/120)	6.8 \pm 2.3	-
During shift	42.5 (51/120)	17.6 \pm 4.8	0.016
After shift	60.0 (72/120)	28.4 \pm 6.1	0.002

To further clarify species dynamics during clinical activity, Table 4 summarizes the estimated prevalence of *Candida* species at each time point. *C. albicans* remained dominant in all stages, increasing steadily from 12.5% before shifts to 41.6% after shifts.

Table 4. Estimated Prevalence of *Candida* Species Across Time Points

Time Points	Total Positive Samples	<i>Candida albicans</i> (%)	<i>C. glabrata</i> (%)
Before shift	22	12.5%	3.3%
During shift	51	29.2%	7.5%
After shift	72	41.6%	10.0%

4.4. Antifungal Susceptibility Testing

Antifungal susceptibility testing revealed that 85.1% of *C. albicans* isolates were susceptible to fluconazole, while 14.9% showed dose-dependent susceptibility. Amphotericin B and nystatin demonstrated high efficacy, with 96.7% susceptibility (Table 5).

Table 5. Antifungal Resistance Patterns of *Candida* Species Isolated from Dental Unit Surfaces

Species	Susceptible (%)	Dose-Dependent (%)	Resistant (%)
<i>Candida albicans</i>			
Fluconazole	85.1	14.9	0.0
Amphotericin B	96.7	3.3	0.0
Nystatin	96.7	3.3	0.0
<i>Candida glabrata</i>			
Fluconazole	70.0	23.3	6.7
Amphotericin B	90.0	10.0	0.0
<i>Candida tropicalis</i>			
Fluconazole	76.7	20.0	3.3
Amphotericin B	93.3	6.7	0.0

4.5. Correlation Between Contamination and Clinical Activity

A strong positive correlation ($R = 0.74$, $P < 0.001$) was observed between the number of patient visits per unit and the level of *Candida* contamination. Units with a higher patient flow exhibited significantly higher contamination levels.

4.6. Effectiveness of Routine Disinfection Protocols

Despite routine disinfection, 39.2% of surfaces remained contaminated post-disinfection, suggesting that current sterilization protocols may not be fully effective against *Candida* species. The highest residual contamination was found on dental chair headrests (49.6%) and light handles (42.3%) (Table 6).

Table 6. Estimated Prevalence of *Candida* Species Across Time Points

Surface Type	<i>Candida albicans</i> (%)	<i>Candida glabrata</i> (%)	<i>Candida tropicalis</i> (%)	Total Post-Disinfection Contamination (%)
Chair headrests	27.5	12.1	10.0	49.6
Light handles	24.8	10.5	7.0	42.3
Unit control panels	15.0	6.2	4.0	25.2
Saliva ejectors	8.0	4.0	2.0	14.0
Overall average	-	-	-	39.2

5. Discussion

The findings of this study reveal a significant increase in *Candida* contamination on dental unit surfaces during and after clinical shifts, emphasizing the potential risk of fungal transmission in dental settings. The contamination rate increased from 18.3% before shifts to 60.0% after shifts ($P < 0.001$), with the highest fungal loads observed on dental chair headrests (72.5%) and light handles (65.8%). These results align with previous research indicating that high-contact surfaces in dental clinics serve as reservoirs for microbial persistence (7).

This trend highlights several risk factors contributing to contamination, including: (1) High patient turnover during peak clinical hours; (2) aerosol generation during dental procedures such as ultrasonic scaling and high-speed drilling; (3) inadequate surface coverage during routine disinfection; and (4) failure to change gloves or disinfect hands between patient interactions. These factors may collectively enhance the survival and spread of fungal pathogens on surfaces (8).

The bioaerosol mechanism has been documented by Gallagher et al. and others, where fungal spores can remain airborne for extended periods and deposit onto equipment surfaces (8). Our results support this,

particularly due to elevated contamination levels during and after active clinical sessions (9).

Several studies have reported similar patterns of *Candida* contamination in dental environments. De Almondes et al. found that *C. albicans* was present on 58% of dental chairs, particularly in departments with high patient turnover (10). Similarly, Pandey et al. demonstrated that unit control panels and light handles harbored the highest fungal loads, reinforcing the need for enhanced sterilization measures (11). Our findings are also consistent with those of Mobin et al., who observed persistent fungal contamination in 42% of dental units despite routine cleaning protocols (12). In contrast to Turner and Butler, who reported *C. tropicalis* as dominant in some dental environments, our study found *C. albicans* as the predominant species, followed by *C. glabrata* and *C. tropicalis*, indicating a possible regional variation or demographic influence on species distribution (13). This variation may stem from regional differences, patient demographics, or evolving antifungal resistance profiles, which warrant further investigation.

The persistence of *Candida* contamination post-disinfection (39.2%) is likely attributed to the biofilm-forming ability of the isolates, rendering routine hypochlorite disinfection insufficient in eliminating all fungal cells. Biofilms enhance resistance by forming protective extracellular matrices and promoting cell aggregation, making conventional cleaning protocols less effective (14).

Therefore, improved infection control strategies are urgently needed. Based on our findings and previous literature, we recommend the following (15-17): (1) Increasing the frequency of surface disinfection, especially between patients; (2) utilizing antifungal-effective disinfectants (e.g., hydrogen peroxide-based or chlorhexidine-gluconate solutions); (3) introducing UV-C light sterilization for high-touch surfaces; (4) applying disposable plastic barriers to chair headrests, light handles, and control panels; (5) reinforcing hand hygiene protocols and mandatory glove changes between patients.

These combined interventions can significantly reduce fungal load and limit the risk of cross-infection, particularly for immunocompromised patients.

Despite its strengths, this study has several limitations. The single-center study design limits the

generalizability of the findings, and environmental variables such as room ventilation, humidity, and temperature were not recorded, which may influence fungal persistence (18). Furthermore, only conventional culture-based methods were used for species identification; molecular confirmation was not performed, which could underestimate species diversity.

Additionally, the identification of *Candida* species relied on culture-based methods rather than molecular techniques, which could lead to underestimation of species diversity (19). Future research should focus on multi-center studies, molecular identification methods, and the evaluation of novel antifungal disinfectants (20).

In conclusion, the results of this study highlight the need for stricter infection control policies to reduce *Candida* contamination on dental unit surfaces. Implementing enhanced disinfection protocols, improving personal protective measures, and conducting routine microbial surveillance are critical steps toward minimizing the risk of fungal transmission in dental settings (21).

5.1. Conclusions

This study highlights the significant presence of *Candida* species on dental unit surfaces, with a clear increase in contamination from 18.3% before shifts to 60.0% after shifts ($P < 0.001$). The most contaminated surfaces were chair headrests (72.5%) and light handles (65.8%), representing high-risk contact points in clinical settings.

Despite routine disinfection using sodium hypochlorite, 39.2% of surfaces remained contaminated post-cleaning, indicating that current sterilization protocols may be insufficient, particularly against biofilm-forming species such as *C. albicans*. Based on the study's findings, we strongly recommend the following to minimize fungal transmission in dental clinics: (1) More frequent and rigorous surface disinfection, especially of high-contact areas; (2) use of UV-C light or enhanced antifungal disinfectants; (3) implementation of plastic barrier coverings; (4) reinforced staff training in infection control, including hand hygiene and glove protocols.

By adopting these measures, dental care environments can significantly reduce the risk of cross-

infection, thus enhancing the safety of both patients and healthcare workers.

Antifungal susceptibility testing revealed that 85.1% of *C. albicans* isolates were susceptible to fluconazole, while 96.7% showed sensitivity to amphotericin B and nystatin. These findings suggest that while antifungal agents remain largely effective, preventive strategies must focus on limiting surface contamination rather than relying solely on drug susceptibility.

5.2. Recommendations

To minimize fungal transmission in dental settings, the following strategies are recommended: (1) More frequent and rigorous surface disinfection, particularly on high-contact areas; (2) use of advanced sterilization techniques, such as UV-C light disinfection; (3) implementation of microbial surveillance programs to monitor contamination trends; (4) enhanced training for dental personnel on proper hand hygiene and equipment sterilization.

By incorporating these measures, dental clinics can significantly reduce the risk of fungal contamination, improving both patient and practitioner safety.

Footnotes

Authors' Contribution: Study design, data collection, and writing the article: M. N.; Data analysis and study supervision: F. H.; Supervision, review, and revision of the article text: S. M. M. All the authors were responsible for this research.

Conflict of Interests Statement: The authors declare no conflict of interest.

Data Availability: The dataset presented in the study is available on request from the corresponding author during submission or after publication

Ethical Approval: IR.ZUMS.BLC.1402.061.

Funding/Support: The present research received no funding/support.

References

1. Patel M. Oral Cavity and *Candida albicans*: Colonisation to the Development of Infection. *Pathogens*. 2022;**11**(3). [PubMed ID: 35335659]. [PubMed Central ID: PMC8953496]. <https://doi.org/10.3390/pathogens11030335>.

2. Talapko J, Juzbasic M, Matijevic T, Pustijanac E, Bekic S, Kotris I, et al. Candida albicans-The Virulence Factors and Clinical Manifestations of Infection. *J Fungi (Basel)*. 2021;7(2). [PubMed ID: 33499276]. [PubMed Central ID: PMC7912069]. <https://doi.org/10.3390/jof7020079>.
3. Miller CH. Infection Control and Management of Hazardous Materials for the Dental Team-E-Book. *Elsevier Health Sci*. 2016.
4. Dag I, Oz Y, Kiraz N. Effect of disinfectants on biofilm development by five species of Candida. *African journal of microbiology research*. 2012;6. <https://doi.org/10.5897/AJMR11.1427>.
5. Spagnolo AM, Sartini M, Cristina ML. Microbial Contamination of Dental Unit Waterlines and Potential Risk of Infection: A Narrative Review. *Pathogens*. 2020;9(8). [PubMed ID: 32823641]. [PubMed Central ID: PMC7460066]. <https://doi.org/10.3390/pathogens9080651>.
6. Suleyman G, Alangaden G, Bardossy AC. The Role of Environmental Contamination in the Transmission of Nosocomial Pathogens and Healthcare-Associated Infections. *Curr Infect Dis Rep*. 2018;20(6):12. [PubMed ID: 29704133]. <https://doi.org/10.1007/s11908-018-0620-2>.
7. Anil Kohli MD, Eng FD, Raghunath PB, Raman BB, Shetty KS, Malika KB, et al. *Infection Control & Occupational Safety Recommendations for Oral Health Professionals in India*. 2007.
8. Gallagher JE, K CS, Johnson IG, Al-Yaseen W, Jones R, McGregor S, et al. A systematic review of contamination (aerosol, splatter and droplet generation) associated with oral surgery and its relevance to COVID-19. *BDJ Open*. 2020;6:25. [PubMed ID: 33251028]. [PubMed Central ID: PMC7684564]. <https://doi.org/10.1038/s41405-020-00053-2>.
9. Stellenberg K. *The Effects of Computerized Electronic Prompts in Influencing Dental Hygiene Students' Compliance with Infection Control Protocols [Master's thesis]*. University of Minnesota; 2022.
10. De Almondes AI, de Araujo JO, Mendes de Sirqueira Amaral L, Reis RC, Porto JC, Teles JB, et al. Fungal contamination and disinfection of dental chairs, Teresina, Piaui, Brazil. *Acta Odontol Latinoam*. 2016;29(3):225-9. [PubMed ID: 28383602].
11. Pandey PK, Agrawal S, Singh D, Satish K, Kumari P. Sterilization and Disinfection of Dental Operator. *Shashwat Publication*. 2021.
12. Mobin M, Borba Cde M, Filho CA, Tapety FI, Noletto Ide M, Teles JB. Analysis of fungal contamination and disinfection of toothbrushes. *Acta Odontol Latinoam*. 2011;24(1):86-91. [PubMed ID: 22010412].
13. Turner SA, Butler G. The Candida pathogenic species complex. *Cold Spring Harb Perspect Med*. 2014;4(9). a019778. [PubMed ID: 25183855]. [PubMed Central ID: PMC4143104]. <https://doi.org/10.1101/cshperspect.a019778>.
14. Ku TSN, Walraven CJ, Lee SA. Candida auris: Disinfectants and Implications for Infection Control. *Front Microbiol*. 2018;9:726. [PubMed ID: 29706945]. [PubMed Central ID: PMC5906573]. <https://doi.org/10.3389/fmicb.2018.00726>.
15. Reedy JM, Fernando T, Awuor SO, Omwenga EO, Koutchma T, Mariita RM. Global Health Alert: Racing to Control Antimicrobial-Resistant Candida auris and Healthcare Waste Disinfection Using UVC LED Technology. *Hygiene*. 2024;4(3):385-422.
16. Alshehri AA, Park S, Rashid H. Strategies to improve hand hygiene compliance among healthcare workers in adult intensive care units: a mini systematic review. *J Hosp Infect*. 2018;100(2):152-8. [PubMed ID: 29559231]. <https://doi.org/10.1016/j.jhin.2018.03.013>.
17. Mattner CJ, Neuwirth M, Otchwemah R, Senges C, Mattner F. Impact of single-use gloves on the five indications for hand hygiene. *Communicable Dis*. 2024. <https://doi.org/10.21203/rs.3.rs-4678285/v1>.
18. Haleem Khan AA, Mohan Karuppayil S. Fungal pollution of indoor environments and its management. *Saudi J Biol Sci*. 2012;19(4):405-26. [PubMed ID: 23961203]. [PubMed Central ID: PMC3730554]. <https://doi.org/10.1016/j.sjbs.2012.06.002>.
19. Ahaik I, Nunez-Rodriguez JC, Abrini J, Bouhdid S, Gabaldon T. Assessing Diagnosis of Candida Infections: A Study on Species Prevalence and Antifungal Resistance in Northern Morocco. *J Fungi (Basel)*. 2024;10(6). [PubMed ID: 38921360]. [PubMed Central ID: PMC11204772]. <https://doi.org/10.3390/jof10060373>.
20. Armstrong-James D. Future Directions for Clinical Respiratory Fungal Research. *Mycopathologia*. 2021;186(5):685-96. [PubMed ID: 34590208]. [PubMed Central ID: PMC8536595]. <https://doi.org/10.1007/s11046-021-00579-5>.
21. Kohn WG, Harte JA, Malvitz DM, Collins AS, Cleveland JL, Eklund KJ, et al. Guidelines for infection control in dental health care settings-2003. *J Am Dent Assoc*. 2004;135(1):33-47. [PubMed ID: 14959873]. <https://doi.org/10.14219/jada.archive.2004.0019>.