





Survival Rate of Hemodialysis Patients and Its Predictors: A Multicentric Retrospective Cohort Study

Sahar Ahmadi Valiabadi¹, Seyedeh Zahra Hosseinigolafshani ^{2,*}, Mehdi Ranjbaran ³

¹ Student Research Committee, Qazvin University of Medical Sciences, Qazvin, Iran

² Social Determinants of Health Research Center, Research Institute for Prevention of Non-communicable Diseases, Qazvin University of Medical Sciences, Qazvin, Iran

³ Metabolic Diseases Research Center, Research Institute for Prevention of Non-communicable Diseases, Qazvin University of Medical Sciences, Qazvin, Iran

* Corresponding author: Social Determinants of Health Research Center, Research Institute for Prevention of Non-Communicable Diseases, Qazvin University of Medical Sciences, Qazvin, Iran. Email: z.hoseinigolafshani@qums.ac.ir

Received 2023 November 4; Revised 2024 January 13; Accepted 2024 March 8.

Abstract

Background: Despite advancements in medical and nursing care, the mortality rate among hemodialysis patients remains unexpectedly high.

Objectives: This study aimed to evaluate the survival rates and identify mortality predictors among hemodialysis patients in Qazvin Province, Iran, over a five-year period from March 2015 to March 2020.

Methods: This historical cohort study involved 378 hemodialysis patients from seven hospitals in Qazvin province, including Bu Ali Sina, Velayat, Razi, Alvand, Valiasr, Bueen Zahra, and Avaj, covering the period from March 21, 2015, to March 20, 2020. The study was based on data extracted from the patient's medical records, offering a detailed overview of their medical history and treatments during this timeframe.

Results: The study revealed a median survival time of 38.5 months, with survival rates at 1, 2, 3, 4, and 5 years being 84.3%, 68.1%, 53.2%, 39.8%, and 18.0%, respectively. Cox regression analysis identified several key predictors of mortality. Mortality rates increased with age, higher levels of blood urea nitrogen, and when hemoglobin levels fell below 10 g/dL or calcium levels exceeded 9.5 mg/dL. Conversely, patients who underwent hemodialysis three times a week had higher levels of education, used fistula access, and experienced significantly lower mortality rates.

Conclusions: The study highlighted the need for targeted interventions to improve both the survival rates and quality of life for hemodialysis patients in Qazvin Province. It emphasized the importance of careful monitoring and effective management of factors such as age, blood urea nitrogen, hemoglobin, and calcium levels, as well as the patients' educational background, frequency of dialysis sessions, and type of vascular access. By focusing on these aspects, healthcare professionals in the region could significantly enhance patient outcomes and prognoses in hemodialysis care.

Keywords: Survival Rate, Hemodialysis, Chronic Kidney Disease, Iran

1. Background

Chronic kidney disease involves the bilateral, progressive, chronic deterioration of nephrons and becomes apparent when 50 - 75% of approximately two million nephrons have ceased their physiological function. In such cases, death is inevitable unless renal replacement therapies are undertaken (1, 2). This disease poses a significant health problem due to its substantial financial burden, and the absence of early and

appropriate management can lead to end-stage renal disease (3).

The World Health Organization (WHO) identifies chronic kidney disease as a global health issue, estimating that around 850 million people worldwide suffer from various degrees of renal failure. This condition, with an 11% prevalence rate, accounts for approximately 4.5 million deaths annually (4). In Iran, the prevalence and incidence rates of end-stage renal disease (ESRD) are 435.8 and 63.8 per million people, respectively (5). The United States reports the highest

prevalence of renal failure, though there has been a decreasing trend, with the number of patients reducing from 125,408 in 2016 to 124,500 in 2017 (6). Conversely, Western Europe, developed countries, and developing nations like Iran have seen an increasing trend in recent decades (7).

In addition to metabolic disorders and an elevated mortality risk from cardiovascular diseases, chronic kidney disease significantly affects patients' physical and mental health, diminishes their quality of life, and places a considerable financial strain on the healthcare system (8).

Patients undergoing hemodialysis face not only the physical and mental challenges associated with the disease but also significant economic pressures (9). The costs of managing this disease were estimated at \$4.1 billion in Australia in 2012, \$55 billion in the U.S. in 2010, and \$1.45 billion in England in 2010 (10).

Early diagnosis and the timely, effective management of chronic kidney disease can significantly reduce its physical and mental impact and markedly decrease its mortality rate (11). Dialysis and kidney transplantation are two renal replacement methods that prevent the accumulation of waste materials in the body. Peritoneal dialysis and hemodialysis represent two forms of dialysis (12), with hemodialysis being the most common method (70%) worldwide for several reasons (13). The annual growth rate of hemodialysis patients is between 10 - 12% globally (14).

Opting for hemodialysis comes with a range of complications during and after the procedure, such as hypotension (20 - 30%), muscle cramps (5 - 20%), nausea and vomiting (5 - 15%), headaches (5%), pruritus (5%), chest and back pain (2 - 5%), and fever and chills (1%) (15). Approximately four million people globally must connect to a dialysis machine 2 - 3 times a week for several hours, significantly impacting their daily lives (16). Despite considerable advancements in medical and nursing care, the mortality rate among hemodialysis patients remains alarmingly high (17, 18). Research indicates that the average life expectancy of patients on dialysis is under three years (19). Survival analysis, a statistical model widely endorsed by many experts, employs a retrospective methodology to monitor a group of subjects over a specific period until an event of interest (e.g., death) occurs. This study concludes by evaluating subjects who did not experience the event as censored data (20). Numerous global studies utilizing this approach have informed various aspects of patient management and treatment. For example, Park et al. reported that the mortality rate for hemodialysis patients after one year of treatment was 20 - 25%, with a

five-year survival rate of 35% in the United States (21). In Iran, Habibi et al. estimated the survival rates at 1, 2, 3, 4, and 5 years for these patients to be 84%, 60%, 49%, 25%, and 10%, respectively (15).

Therefore, this study specifically focuses on hemodialysis, a prevalent renal replacement therapy for ESRD. Hemodialysis is vital for managing chronic kidney disease (CKD), especially in light of its increasing prevalence and the high mortality rates associated with it. The therapy necessitates significant lifestyle adjustments and is fraught with various complications, highlighting the need to comprehend its long-term outcomes. Despite advances in healthcare, the survival rates of patients undergoing hemodialysis present a significant concern, with many patients experiencing a shortened life expectancy.

2. Objectives

Given these challenges, this research seeks to offer detailed insights into the survival rates and determinants of mortality among hemodialysis patients in Qazvin Province, Iran. This is crucial for enhancing patient management and treatment strategies, tailoring them to meet the specific needs of the Iranian populace, and enriching the global understanding of CKD and ESRD management.

3. Methods

3.1. Study Design and Participants

This historical cohort study involved 378 hemodialysis patients from seven hospitals in Qazvin province, including Bu Ali Sina, Velayat, Razi, Alvand, Valiasr, Bueen Zahra, and Avaj. The sample size was estimated using the formula related to the Cox proportional hazards regression model, with considerations for $\alpha = 0.05$ and $1 - \beta = 0.80$:

$$\text{Number of events} = \frac{\left(Z_{1-\frac{\alpha}{2}} + Z_{1-\beta} \right)^2}{\sigma^2(\log HR)^2} \quad (1)$$

This calculation was based on the variable of serum albumin, identified as an important and significant variable in previous studies (15, 22), with a standard deviation of 0.82 and an HR = 0.55 for the 173 samples that resulted in death. Accounting for approximately 45% death during the follow-up period, at least 375 samples of hemodialysis patients were needed.

The study spanned from March 21, 2015, to March 20, 2020, and analyzed data extracted from the medical

records of these patients, offering a detailed overview of their medical history and treatment during this timeframe. All patients who met the inclusion and exclusion criteria during the specified period were included in the study through the census method. The eligibility criteria were as follows:

- Willingness to participate: Participants must voluntarily agree to take part in the study, emphasizing the significance of consent given the study's focus on survival rates among hemodialysis patients.

- Minimum dialysis duration: Only individuals who had undergone dialysis for at least three months were considered to ensure the sample represented those actively managing end-stage renal disease.

- Age requirement: Participants needed to be 18 years or older, which corresponds with the study's aim to assess adult hemodialysis outcomes within the Iranian setting.

The exclusion criteria were as follows:

- Refusal to continue participation: Participants unwilling to continue in the study were excluded to maintain accurate survival rate analysis through consistent participation.

- Loss to follow-up: Participants who were untraceable or whose data could not be reliably followed through the study duration were excluded, safeguarding data integrity.

- Incomplete hospital records: Due to the reliance on medical history to understand dialysis and health outcomes, participants with incomplete medical records were excluded.

- History of kidney transplantation: To keep the study focused on hemodialysis survival rates, those with a history of kidney transplantation were not included, considering their distinct health profiles.

These criteria were critical for assembling a focused and relevant group of participants to study hemodialysis survival rates within the Iranian population, particularly in Qazvin Province. The objective was to deliver comprehensive insights into the factors that influence survival rates among this specific patient demographic. Data collection involves using a checklist to gather information from patient's medical records or through telephone or face-to-face meetings. The variables evaluated in this study included age, sex, educational level, occupation, marital status, frequency of dialysis per week, type of vascular access, blood urea nitrogen (BUN) level, hemoglobin level, serum albumin, serum calcium, and the adequacy of dialysis.

The follow-up period in this study commenced three months after the initiation of the first dialysis session,

with the endpoint being the time of death. Censored subjects at the study's conclusion included those still alive, those who died from causes unrelated to their condition, and participants lost to follow-up.

Bias was minimized through stringent inclusion and exclusion criteria, the use of the census method for thorough patient inclusion, and standardized data collection from medical records. Conducted over five years, the study aimed to ensure a representative sample and reliable data collection across multiple hospital settings, thus enhancing the validity of the findings.

3.2. Statistical Analysis

Qualitative variables were presented as numbers and percentages, while quantitative variables were reported using means and standard deviations. The Kaplan-Meier curve was utilized to assess the survival rate of hemodialysis patients, and Cox regression analysis was employed to estimate the hazard ratios of factors affecting survival. To adjust for confounders, variables with P-values ≤ 0.2 in univariable Cox regression analyses were included in the multivariable Cox regression model. The proportional hazards assumption was tested and confirmed using the estate test method. Data analysis was performed using SPSS software version 25 and STATA version 12, with a significance level set at 0.05.

3.3. Ethical Considerations

This study received approval from the ethics committee of Qazvin University of Medical Sciences (Ethical Code: [IR.QUMS.REC.1400.017](#)). Before providing informed consent, all participants were thoroughly informed about the study's objectives and methods. It was clearly communicated that they had the option to withdraw from the study at any time.

4. Results

The average age of the participants was 61.65 years. The majority of the patients were male (58.2%), married (86%), and illiterate (39.9%), with all of them having underlying diseases. Most participants (90.21%) underwent hemodialysis three times a week and utilized dialysis fistulas (66.9%) for vascular access. Other qualitative and clinical characteristics are detailed in [Table 1](#).

4.1. Survival Rate

The average follow-up duration was 22.82 ± 16.65 months. Out of 378 patients, 150 patients (39.7%)

Table 1. Demographic and Clinical Characteristics of Hemodialysis Patients ^a

Variables and categories	Total	Censored	Deceased	P-Value ^b
Sex				0.258
Male	220 (58.2)	138 (62.7)	82 (37.3)	
Female	158 (41.8)	90 (57.0)	68 (43.0)	
Marital status				0.256
Single	36 (9.5)	22 (61.1)	14 (38.9)	
Married	325 (86.0)	199 (61.2)	126 (38.8)	
Divorced/Widowed	17 (4.5)	7 (41.2)	10 (58.8)	
Education level				< 0.001
Illiterate	151 (39.9)	84 (55.6)	67 (44.4)	
Primary education	76 (20.1)	31 (40.8)	45 (59.2)	
High school education	87 (23.0)	63 (72.4)	24 (27.6)	
Academic education	64 (16.9)	50 (78.1)	14 (21.9)	
Occupation				0.165
Jobless	190 (50.3)	82 (43.2)	108 (56.8)	
Employed	188 (49.7)	120 (63.8)	68 (36.2)	
Frequency of dialysis per week				0.687
Once a or Twice	35 (9.3)	20 (57.1)	15 (42.9)	
Three or Four times	343 (90.7)	208 (60.6)	135 (39.4)	
Vascular access				< 0.001
Shaldon	73 (19.3)	13 (17.8)	60 (82.2)	
Fistula	253 (66.9)	171 (67.6)	82 (32.4)	
Permcath	48 (12.7)	41 (85.4)	7 (14.6)	
Graft	4 (1.1)	3 (75.0)	1 (25.0)	
Hb, g/dL				< 0.001
<10	138 (36.5)	61 (42.2)	77 (55.8)	
10 - 12	144 (38.1)	98 (68.1)	46 (31.9)	
> 12	96 (25.4)	69 (71.9)	27 (28.1)	
Calcium, mg/dL				0.002
< 8.4	154 (40.7)	97 (63.0)	57 (37.0)	
8.4 - 9.5	188 (49.7)	119 (63.3)	69 (36.7)	
> 9.5	36 (9.5)	12 (33.3)	24 (66.7)	
Total	378 (100)	228 (60.3)	150 (39.7)	
Age, y	61.65 ± 14.94	59.57 ± 14.11	64.81 ± 15.63	0.001
Adequacy of dialysis	1.22 ± 0.30	1.25 ± 0.34	1.16 ± 0.23	0.002
Albumin, g/dL	4.09 ± 0.73	4.09 ± 0.57	4.08 ± 0.92	0.895
BUN, mg/dL	26.54 ± 16.54	22.81 ± 12.69	32.20 ± 19.83	< 0.001

^a Values are expressed as mean ± SD or No. (%).

^b Chi-square test for qualitative variables and independent t-test for quantitative variables.

experienced the event (death). The median survival time for the patients was 38.5 months (95% CI: 31.35 - 45.65 months), indicating that half of the patients survived for at least 38.5 months. According to the Kaplan-Meier method, the 1, 2, 3, 4, and 5-year cumulative survival probabilities for the hemodialysis patients, along with their standard errors, were 84.3 ± 2.0%, 68.1 ± 2.7%, 53.2 ± 3.2%, 39.8 ± 3.9%, and 18.0 ± 7.1%, respectively (Figure 1).

4.2. Univariable Cox Regression Analysis

Cox regression analysis was utilized to explore the association between various variables and the mortality of hemodialysis patients (Table 2). The analysis revealed that for each one-year increase in age and each one-mg/dL increase in BUN, there was a significant 2% increase in mortality risk. Conversely, mortality significantly decreased with higher levels of education,




Figure 1. The survival curve of hemodialysis patients in Qazvin Province from 2015 - 2020

an increased number of dialysis sessions per week, and greater adequacy of dialysis. Regarding vascular access, mortality was significantly lower in patients with fistulas and Permcath compared to those with Shaldon catheters. A decrease in hemoglobin levels and an increase in calcium levels, compared to their normal ranges, were also significantly associated with increased mortality ($P < 0.05$).

4.3. Multivariable Cox Regression Analysis

Multivariable Cox regression analysis was utilized to identify variables associated with the mortality of hemodialysis patients, adjusted for confounding factors. Variables with P -values ≤ 0.2 in univariable Cox regression were included in the multivariable Cox regression model. The analysis revealed that age, BUN levels, education level, frequency of dialysis per week, type of vascular access, average hemoglobin, and average calcium levels were significant predictors of mortality in hemodialysis patients (Table 3). Mortality risk significantly increased by 2% with each one-year increase in age (HR = 1.02, 95% CI: 1.001 to 1.03) and by 1% with each one-mg/dL increase in BUN levels (HR = 1.01, 95% CI: 1.004 to 1.02). Patients undergoing dialysis 3 - 4 times a week experienced a significant 54% reduction in mortality compared to those dialyzing 1 - 2 times a week (HR = 0.46, 95% CI: 0.25 to 0.82). Regarding vascular access, mortality was significantly lower with a fistula compared to a Shaldon catheter (HR = 0.20, 95% CI: 0.13 to 0.28) and a Permcath (HR = 0.13, 95% CI: 0.06 to 0.31). Mortality significantly increased when hemoglobin levels fell below 10 g/dL (HR = 1.77, 95% CI: 1.18 to 2.64) and

calcium levels rose above 9.5 mg/dL (HR = 1.86, 95% CI: 1.13 to 3.04) compared to their normal ranges. Additionally, hemodialysis patients with higher education levels experienced significantly lower mortality (HR = 0.58, 95% CI: 0.34 to 0.99).

5. Discussion

This study aimed to determine the 1, 2, 3, 4, and 5-year survival rates of hemodialysis patients and their predictors among 378 patients undergoing hemodialysis in various dialysis departments within Qazvin Province, Iran. Cox regression analysis was utilized to identify mortality predictors in hemodialysis patients. The findings revealed that age, education level, type of vascular access, frequency of dialysis per week, BUN, hemoglobin level, and serum calcium were significant predictors of mortality among patients attending the dialysis wards in Qazvin Province. Previous research has emphasized the importance of estimating the survival rates of hemodialysis patients for decision-making regarding hemodialysis initiation and identifying patients at a high risk of mortality for appropriate management and follow-up (23, 24). The median 5-year survival time for patients was 38.5 months, suggesting that at least 50% of the patients survived for 38.5 months following the commencement of hemodialysis. Habibi et al. reported a median 5-year survival of 34 months, Montaseri et al. reported 37 months, and Ebrahimi et al. reported 20.9 months (15, 22, 25). In contrast, Shabankhani et al. found a median survival rate of 108 months, and Ferreira et al. reported 59 months for Brazilian hemodialysis patients (26, 27).

Table 2. Univariable Cox Regression Model for Variables Affecting the Survival Time of Hemodialysis Patients

Variable	Median of Survival, Month	Hazard Ratio (HR)	95% Confidence Interval for HR		P-Value
			Lower Bound	Upper Bound	
Gender					0.350
Male	38.77	Ref	-	-	
Female	38.50	1.17	0.85	1.61	
Marital status					
Single	44.53	Ref	-	-	
Married	38.50	1.21	0.70	2.11	0.49
Widow or divorced	21.50	2.08	0.92	4.69	0.08
Educational status					
Illiterate	29.50	Ref	-	-	
Elementary	29.97	1.15	0.79	1.68	0.476
High school education	52.93	0.38	.23	0.60	< 0.001
High school diploma or academic education	55.47	0.33	0.19	0.59	< 0.001
Occupational status					0.283
Unemployed	37.97	Ref	-	-	
Employed	39.03	0.84	0.61	1.16	
Dialysis per week					0.002
1 or 2 times	23.47	Ref	-	-	
3 or 4 times	38.77	0.42	0.24	0.72	
Vascular access					
Shaldon	10.93	Ref	-	-	
Fistula	46.27	0.17	0.12	0.23	< 0.001
Permcath	.	0.09	0.04	0.20	< 0.001
Hb, g/dL					
<10	29.50	2.24	1.56	3.24	< 0.001
10 - 12	47.87	Ref	-	-	
>12 l	52.93	0.85	0.53	1.37	0.501
Calcium, mg/dL					
< 8.4	44.53	1.01	0.71	1.43	0.970
8.4 - 9.5	45.60	Ref	-	-	
> 9.5	23.67	2.25	1.41	3.59	0.001
Age, y	-	1.02	1.01	1.03	0.001
Adequacy of dialysis	-	0.50	0.27	0.91	0.023
Albumin, g/dL	-	0.92	0.74	1.14	0.443
BUN, mg/dL	-	1.02	1.01	1.03	< 0.001

The significant variation in median survival rates across these studies may be attributed to differences in countries' development levels, healthcare access, genetics, lifestyle, and nutrition, as detailed in the study by Kazemini et al. (28).

In the present historical study, the 1, 2, 3, 4, and 5-year cumulative survival rate of the hemodialysis patients and their standard errors were $84.3 \pm 0.020\%$, $68.1 \pm 0.027\%$, $53.2 \pm 0.032\%$, $39.8 \pm 0.039\%$, and $18.0 \pm 0.071\%$, respectively. The 1, 2, 3, 4, and 5-year survival rates of the patients was 84%, 60%, 49%, 25%, and 10% in a study by Habibi et al (15). These rates were 75%, 63%, 50%, 41%, and

23% in a study by Montaseri et al. respectively, indicating different values with a similar decreasing trend (25). Ferreira et al. reported 1, 5, 10, and 20-year survival rates for Brazilian hemodialysis patients as 82.3%, 49.1%, 22.5%, and 13.3%, respectively (27). Msaad noted 1 and 3-year survival rates of 95% and 87%, and Sun et al. reported 1 and 5-year survival rates of 94% and 59%, respectively (19, 29).

Comparison of the 1, 3, and 5-year survival rates among the aforementioned studies reveals similar 1-year survival rates for patients both within and outside of Iran. However, the 3-year and 5-year survival rates are

Table 3. Multivariable Cox Regression Model for Prediction of Mortality in Hemodialysis Patients

Variable	Hazard Ratio (HR)	95% Confidence Interval for HR		P-Value
		Lower Bound	Upper Bound	
Marital status				
Single	Ref	-	-	
Married	0.55	0.30	1.03	0.060
Widow or divorced	0.68	0.26	1.77	0.431
Educational status				
Illiterate	Ref	-	-	
Elementary	1.07	0.70	1.62	0.766
High School	0.58	0.34	0.99	0.045
High school diploma or academic education	0.55	0.29	1.05	0.071
Dialysis per week				
1 or 2 times	Ref	-	-	
3 or 4 times	0.46	0.25	0.82	0.008
Vascular access				
Shaldon	Ref	-	-	
Fistula	0.20	0.13	0.28	< 0.001
Permcath	0.13	.06	0.31	< 0.001
Hb, g/dL				
< 10	1.77	1.18	2.64	0.006
10 - 12	Ref	-	-	
> 12	1.04	0.64	1.69	0.881
Calcium, mg/dL				
< 8.4	0.96	0.66	1.38	0.819
8.4 - 9.5	Ref	-	-	
> 9.5	1.86	1.13	3.04	0.014
Age, y	1.02	1.001	1.03	0.031
Adequacy of dialysis	0.82	0.38	1.74	0.598
BUN, mg/dL	1.01	1.004	1.02	0.006

notably lower for Iranian patients, necessitating further investigation into the influence of various individual and societal factors, as well as the quality of care.

Regarding the relationship between different variables and the mortality of hemodialysis patients, this study found that each one-year increase in age and each one-mg/dL increase in BUN significantly raised mortality by 2%. Msaad et al. identified age as an important predictor of survival for hemodialysis patients, with those above 65 years experiencing a 1.26 times higher risk of mortality compared to those under 65 years (19). Similarly, Montaseri et al. reported that each additional year of age increased the risk of death by 1.88 times in hemodialysis patients (25). In line with this, Bal et al. (30) and Ferreira et al. (27) also highlighted age as a crucial predictor of survival, noting that older patients with higher BUN levels had a lower chance of survival. These findings align with the current study, suggesting that older age and higher BUN levels decrease the survival rate of hemodialysis patients (19,

30), potentially due to the better physical condition and lesser progression of the underlying disease in younger patients, as well as lower BUN levels.

This study also found a significant decrease in mortality with an increase in education level, frequency of dialysis per week, and adequacy of dialysis. Shabankhani et al. (26) discovered that patients with higher education levels had a 36% lower risk of mortality compared to those with lower education levels. A study examining the relationship between dialysis adequacy and mortality in 18 242 hemodialysis patients by Hong and Lee (31) showed that patients with dialysis adequacy below 1.2 faced a much higher mortality risk than those with dialysis adequacy between 1.2 and 1.4. Similarly, Hekmat et al. (32) found dialysis adequacy to be a significant determinant of survival rate in hemodialysis patients, with an OR = 1.193.

In this research, the survival rates for patients with fistulas and Shaldons were 46.27 and 10.93 months, respectively. This contrasts with a study by Torreggiani

et al. (33), which found survival rates for patients with fistulas and Shaldons to be 31 ± 26 and 14.5 ± 14 months, respectively, and a study by Kim et al. (34), which reported 5-year survival rates for patients with fistulas and Shaldons as 27% and 12%, respectively.

According to Torreggiani et al., patients with fistulas have a survival chance that is twice as high compared to those without (33, 34). Similarly, do Sameiro-Faria et al. (35) noted that the risk of mortality was higher in patients using a Shaldon catheter as their vascular access compared to other methods, with a hazard ratio (HR) of 3.03 (95%CI = 1.49 - 6.13) (35). In the current study, the hazard ratio for patients with a fistula was estimated at 0.17, whereas Ko et al. reported a hazard ratio of 1.13. This indicates a significantly high survival rate for patients with fistulas (36). These studies suggest that patients with fistulas are typically younger and have fewer comorbidities, indicating their better physical condition. Often, in older patients with a lower life expectancy, physicians may delay placing a fistula, opting instead for alternative access methods at the initiation of dialysis. Furthermore, vascular catheters are associated with a higher risk of infection and longer hospital stays compared to fistulas, contributing to the superior survival rates of patients with fistulas over other vascular access methods (29, 33, 37).

Hemoglobin level was identified as an important mortality predictor in hemodialysis patients. A reduction in hemoglobin levels below 10 g/dL and an increase in calcium levels above 9.5 mg/dL significantly elevated mortality risks compared to their normal ranges. Ferreira et al. (27) also found that ferritin levels below the normal range and calcium levels above 11.01 mg/dL significantly increased mortality risks (HR = 4.102, 95%CI: 1.35 - 12.46).

This study investigated the survival rate and predictors of hemodialysis patients over 1 - 5 years, identifying variables such as age, education, and type of vascular access as significant factors. While its findings align with some international studies, observed variations in survival rates point to regional disparities and nuances in healthcare practices. Fistulas are shown to enhance patient survival, emphasizing the critical role of access type. Nonetheless, the presence of incomplete records and potential confounders necessitates a cautious interpretation of the results and calls for further comprehensive research. Such studies are essential for a broader validation and understanding of hemodialysis outcomes.

5.1. Limitations of the Study and Efforts Made to Address Them

1. Incomplete records: Rigorous data verification methods were implemented to mitigate the impact of incomplete records.

2. Regional specificity: By including multiple hospitals across Qazvin province, the study aimed to diversify the patient population, reducing regional bias. However, it's recognized that this does not fully represent global demographics.

3. Confounding factors: Advanced statistical models, such as Cox regression analysis, were employed to adjust for known confounders, minimizing their impact on the findings.

4. Scope and sample size: Despite being limited to 378 patients, the study used a census method for inclusion to ensure comprehensive coverage of all eligible patients during the study period, enhancing the representativeness of the sample.

5. Need for further research: Recognizing the necessity for more extensive research, this study establishes a detailed analytical framework, laying the groundwork for future studies to expand upon these findings in various or larger populations.

5.2. Conclusions

The median survival rate of hemodialysis patients in Qazvin Province, Iran, was 38.5 months (95% CI: 31.35 - 45.65 months), indicating that half of the patients survived for at least 38.5 months post-initiation of hemodialysis. The 1, 2, 3, 4, and 5-year cumulative survival rates of the hemodialysis patients, along with their standard errors, were $84.3 \pm 2.0\%$, $68.1 \pm 2.7\%$, $53.2 \pm 3.2\%$, $39.8 \pm 3.9\%$, and $18.0 \pm 7.1\%$, respectively. Regarding the relationship between mortality and various variables, mortality increased significantly by 2% with each one-year increase in age and by 1% with each one-mg/dL increase in BUN levels. Mortality significantly rose with decreases in hemoglobin levels below 10 g/dL and increases in calcium levels above 9.5 mg/dL. Mortality decreased by 54% in patients undergoing dialysis 3 - 4 times a week compared to those dialyzing 1 - 2 times a week. Regarding the type of vascular access, mortality was significantly lower with a fistula compared to Shaldon and Permcath. Additionally, patients with higher education levels and greater adequacy of dialysis experienced significantly lower mortality rates.

In summary, while the findings offer valuable insights for healthcare professionals in Qazvin Province, their applicability in other regions or countries might require further validation or adaptation due to differences in demographics, healthcare systems, and

cultural aspects. These factors should be considered before directly implementing the study's recommendations elsewhere.

Acknowledgements

In the end, it is necessary to thank the officials of the Faculty of Nursing and Midwifery and the patients who participated in this study.

Footnotes

Authors' Contribution: Study concept and design: Seyedehzahra Hosseinigolafshani; acquisition of data: Sahar Ahmadi Valiabadi; Analysis and interpretation of data: Mehdi Ranjbaran and Seyedehzahra Hosseinigolafshani; Drafting of the manuscript: Seyedehzahra Hosseinigolafshani, Mehdi Ranjbaran, Sahar Ahmadi Valiabadi; Critical revision of the manuscript for important intellectual content: Seyedehzahra Hosseinigolafshani; Statistical analysis: Mehdi Ranjbaran; Administrative, technical, and material support: Seyedehzahra Hosseinigolafshani. Study supervision.

Conflict of Interests: The authors of the article hereby officially declare that they have no conflicts of interest.

Data Availability: The dataset presented in the study is available on request from the corresponding author during submission or after publication. The data are not publicly available due to privacy.

Ethical Approval: The study received approval from the Ethics Committee of Qazvin University of Medical Science under the Ethical Code: [IR.QUMS.REC.1400.017](#). All procedures were conducted in accordance with the relevant guidelines and regulations.

Funding/Support: This study was supported in part by grant 1400.017 from Qazvin University of Medical Sciences.

Informed Consent: Written informed consent was obtained from all participants before they completed the questionnaires.

References

- Jobby S, Tan RJ. Nitrolipids in kidney physiology and disease. *Nitric Oxide*. 2018;**78**:121-6. [PubMed ID: 29605557]. [PubMed Central ID: PMC6163094]. <https://doi.org/10.1016/j.niox.2018.03.021>.
- Laheij A, Rooijers W, Bidar L, Haidari L, Neradova A, de Vries R, et al. Oral health in patients with end-stage renal disease: A scoping review. *Clin Exp Dent Res*. 2022;**8**(1):54-67. [PubMed ID: 34459147]. [PubMed Central ID: PMC8874082]. <https://doi.org/10.1002/cre2.479>.
- Askarishahi M, Atapoor A, Hemayati R, Shahidi S, Zeynali S. [Comparison of the Cox Semi-Parametric Model and Parametric Models in Analyzing the Effective Diagnostic Factors in Kidney Transplantation Survival]. *J Toloobehdasht*. 2019. Persian. <https://doi.org/10.18502/tbj.v17i5.311>.
- Peng S, He J, Huang J, Tan J, Liu M, Liu X, et al. A chronic kidney disease patient awareness questionnaire: Development and validation. *PLoS One*. 2019;**14**(5). e0216391. [PubMed ID: 31050683]. [PubMed Central ID: PMC6499466]. <https://doi.org/10.1371/journal.pone.0216391>.
- Jebrailey M, Rahimi B, Makhdoomi KF, Khorami F. Patient education: a study on the present situation and factors affecting success of patient education in hemodialysis. *Hormozgan Med J*. 2016;**20**(4):250-6.
- Saran R, Robinson B, Abbott KC, Bragg-Gresham J, Chen X, Gipson D, et al. US Renal Data System 2019 Annual Data Report: Epidemiology of Kidney Disease in the United States. *Am J Kidney Dis*. 2020;**75**(1 Suppl 1):A6-7. [PubMed ID: 31704083]. <https://doi.org/10.1053/j.ajkd.2019.09.003>.
- Khazaei S, Yaseri M, Nematollahi S, Zobdeh Z, Sheikh V, Mansournia MA. Survival Rate and Predictors of Mortality among Hemodialysis Patients in West of Iran, 1996-2015. *Int J Prev Med*. 2018;**9**:113. [PubMed ID: 30687463]. [PubMed Central ID: PMC6326016]. https://doi.org/10.4103/ijpvm.IJPVM_399_16.
- Hawamdeh S, Almari AM, Almutairi AS, Dator WLT. Determinants and prevalence of depression in patients with chronic renal disease, and their caregivers. *Int J Nephrol Renovasc Dis*. 2017;**10**:183-9. [PubMed ID: 28740416]. [PubMed Central ID: PMC5503667]. <https://doi.org/10.2147/IJNRD.S139652>.
- Lee MC, Wu SV, Hsieh NC, Tsai JM. Self-Management Programs on eGFR, Depression, and Quality of Life among Patients with Chronic Kidney Disease: A Meta-Analysis. *Asian Nurs Res (Korean Soc Nurs Sci)*. 2016;**10**(4):255-62. [PubMed ID: 28057311]. <https://doi.org/10.1016/j.anr.2016.04.002>.
- Wang Q, Liu H, Ren Z, Xiong W, He M, Li N, et al. The Associations of Family Functioning, General Well-Being, and Exercise with Mental Health among End-Stage Renal Disease Patients. *Psychiatry Investig*. 2020;**17**(4):356-65. [PubMed ID: 32252511]. [PubMed Central ID: PMC7176562]. <https://doi.org/10.30773/pi.2019.0204>.
- Pascoe MC, Thompson DR, Castle DJ, McEvedy SM, Ski CF. Psychosocial Interventions for Depressive and Anxiety Symptoms in Individuals with Chronic Kidney Disease: Systematic Review and Meta-Analysis. *Front Psychol*. 2017;**8**:992. [PubMed ID: 28659852]. [PubMed Central ID: PMC5468538]. <https://doi.org/10.3389/fpsyg.2017.00992>.
- Srinivasan R, Sangeetha R, Naik HK, Varghese CS. A Study on Medication Adherence and Quality of Life in Patients with Chronic Kidney Disease. *J Drug Deliv Therap*. 2020;**10**(3-s):52-60. <https://doi.org/10.22270/jddt.v10i3-s.4083>.
- Emami-Naini A, Mortazavi M, Azarbajani A, Hoseinzadeh B, Hosseini SM, Adibi P. [Comparison of Endoscopic Findings in Patients on Hemodialysis (HD) and Peritoneal Dialysis (PD) Waiting for Kidney Transplantation]. *J Isfahan Med Sch*. 2016;**34**(398):1066-70. Persian.
- Iran Dialysis Consortium. *Statistics of the Iranian dialysis community*. 2011, [cited 2023]. Available from: <https://icdgroup.org/>.
- Habibi D, Usefi S, Moslemi A. Five-year survival rate and factors affecting the survival of hemodialysis patients. *Kooshesh J*. 2019;**21**(2):271-5.
- Hamidi M, Roshangar F, Ghafourifard M, Hassankhani H, Sarbakhsh P. [A comparative study of the prevalence rate of common complication in routine hemodialysis and liner and stepwise sodium -ultra filtration profile in patients with chronic kidney failure under hemodialysis]. *J Urmia Nurs Midwifery*. 2019;**17**(6). Persian.

17. D'Onofrio G, Simeoni M, Rizza P, Caroleo M, Capria M, Mazzitello G, et al. Quality of life, clinical outcome, personality and coping in chronic hemodialysis patients. *Ren Fail*. 2017;**39**(1):45-53. [PubMed ID: 27778533]. [PubMed Central ID: PMC6014518]. <https://doi.org/10.1080/0886022X.2016.1244077>.
18. Garg AX, Suri RS, Eggers P, Finkelstein FO, Greene T, Kimmel PL, et al. Patients receiving frequent hemodialysis have better health-related quality of life compared to patients receiving conventional hemodialysis. *Kidney Int*. 2017;**91**(3):746-54. [PubMed ID: 28094031]. [PubMed Central ID: PMC5333984]. <https://doi.org/10.1016/j.kint.2016.10.033>.
19. Msaad R, Essadik R, Mohtadi K, Meftah H, Lebrazi H, Taki H, et al. Predictors of mortality in hemodialysis patients. *Pan Afr Med J*. 2019;**33**:61. [PubMed ID: 31448023]. [PubMed Central ID: PMC6689835]. <https://doi.org/10.11604/pamj.2019.33.61.18083>.
20. Tatari M, Rahgozar M, Amir Khanloo S, Hosseinzadeh S. [The Relationship between Blood Creatinine Levels and Survival of Patients with Kidney Disease Using Joint Longitudinal and Survival Model]. *J Health Promot Managment*. 2017;**7**(1):12-9. Persian. <https://doi.org/10.21859/jhpm-07012>.
21. Park J, Ahmadi SF, Streja E, Molnar MZ, Flegal KM, Gillen D, et al. Obesity paradox in end-stage kidney disease patients. *Prog Cardiovasc Dis*. 2014;**56**(4):415-25. [PubMed ID: 24438733]. [PubMed Central ID: PMC4733536]. <https://doi.org/10.1016/j.pcad.2013.10.005>.
22. Ebrahimi V, Khademian MH, Masoumi SJ, Morvaridi MR, Ezzatzadegan Jahromi S. Factors influencing survival time of hemodialysis patients; time to event analysis using parametric models: a cohort study. *BMC Nephrol*. 2019;**20**(1):215. [PubMed ID: 31185926]. [PubMed Central ID: PMC6558674]. <https://doi.org/10.1186/s12882-019-1382-2>.
23. Anderson RT, Cleek H, Pajouhi AS, Bellolio MF, Mayulka A, Hart A, et al. Prediction of Risk of Death for Patients Starting Dialysis: A Systematic Review and Meta-Analysis. *Clin J Am Soc Nephrol*. 2019;**14**(8):1213-27. [PubMed ID: 31362990]. [PubMed Central ID: PMC6682819]. <https://doi.org/10.2215/CJN.00050119>.
24. Soleymanian T, Ghaziani Z. [Charlson Comorbidity Index as a Strong Predictor of Mortality in Patients with Chronic Hemodialysis]. *Avicenna J Clin Med*. 2018;**25**(3):151-8. Persian. <https://doi.org/10.21859/ajcm.25.3.151>.
25. Montaseri M, Yazdani Cherat J, Espahbodi F, Mousavi SJ. [Five-year survival rate in hemodialysis patients Attending Sari Imam Khomeini Hospital]. *J Mazandaran Univ Med Sci*. 2013;**23**(101):78-85. Persian.
26. Shabankhani B, Kazemnejad A, Zaeri F, Espahbodi F, Haji Ahmadi M, Mirkazemi R. Survival Factors in Patients With End-stage Renal Disease in Mazandaran Province, Iran. *Iran J Kidney Dis*. 2016;**10**(2):79-84. [PubMed ID: 26921749].
27. Ferreira ES, Moreira TR, da Silva RG, da Costa GD, da Silva LS, Cavalier SBO, et al. Survival and analysis of predictors of mortality in patients undergoing replacement renal therapy: a 20-year cohort. *BMC Nephrol*. 2020;**21**(1):502. [PubMed ID: 33228547]. [PubMed Central ID: PMC7685664]. <https://doi.org/10.1186/s12882-020-02135-7>.
28. Kazemini M, Karimi S, Khas Shirzadi L. Nutritional Status of Hemodialysis Patients Referred to Imam Reza Hospital of Kermanshah in 2019: A Descriptive Study. *J Rafsanjan Univ Med Sci*. 2021;**20**(8):851-64. <https://doi.org/10.52547/jrums.20.8.851>.
29. Sun Y, Wang Y, Yu W, Zhuo Y, Yuan Q, Wu X. Association of Dose and Frequency on the Survival of Patients on Maintenance of Hemodialysis in China: A Kaplan-Meier and Cox-Proportional Hazard Model Analysis. *Med Sci Monit*. 2018;**24**:5329-37. [PubMed ID: 30063696]. [PubMed Central ID: PMC6083938]. <https://doi.org/10.12659/MSM.909404>.
30. Bal Z, Demirci BG, Karakose S, Tural E, Erkmen Uyar M, Acar NO, et al. Factors Influencing Hemoglobin Variability and Its Association with Mortality in Hemodialysis Patients. *Sci World J*. 2018;**2018**:8065691. [PubMed ID: 29805324]. [PubMed Central ID: PMC5899870]. <https://doi.org/10.1155/2018/8065691>.
31. Hong WP, Lee YJ. The association of dialysis adequacy, body mass index, and mortality among hemodialysis patients. *BMC Nephrol*. 2019;**20**(1):382. [PubMed ID: 31640580]. [PubMed Central ID: PMC6805311]. <https://doi.org/10.1186/s12882-019-1570-0>.
32. Hekmat R. Hemodialysis Adequacy and Its Impact on Long-Term Patient Survival in Demographically, Socially, and Culturally Homogeneous Patients. *Int J Nephrol*. 2020;**2020**:9857123. [PubMed ID: 32922996]. [PubMed Central ID: PMC7453257]. <https://doi.org/10.1155/2020/9857123>.
33. Torreggiani M, Bernasconi L, Colucci M, Accarino S, Pasquinucci E, Esposito V, et al. Vascular Access, Complications and Survival in Incident Hemodialysis Patients. *Kidney and Dialysis*. 2021;**1**(2):88-99. <https://doi.org/10.3390/kidneydial020013>.
34. Kim DH, Park JI, Lee JP, Kim YL, Kang SW, Yang CW, et al. The effects of vascular access types on the survival and quality of life and depression in the incident hemodialysis patients. *Ren Fail*. 2020;**42**(1):30-9. [PubMed ID: 31847666]. [PubMed Central ID: PMC6968432]. <https://doi.org/10.1080/0886022X.2019.1702558>.
35. do Sameiro-Faria M, Ribeiro S, Costa E, Mendonca D, Teixeira L, Rocha-Pereira P, et al. Risk factors for mortality in hemodialysis patients: two-year follow-up study. *Dis Markers*. 2013;**35**(6):791-8. [PubMed ID: 24347799]. [PubMed Central ID: PMC3857718]. <https://doi.org/10.1155/2013/518945>.
36. Ko GJ, Rhee CM, Obi Y, Chang TI, Soohoo M, Kim TW, et al. Vascular access placement and mortality in elderly incident hemodialysis patients. *Nephrol Dial Transplant*. 2020;**35**(3):503-11. [PubMed ID: 30107612]. [PubMed Central ID: PMC7057539]. <https://doi.org/10.1093/ndt/gfy254>.
37. Allon M. Vascular Access for Hemodialysis Patients: New Data Should Guide Decision Making. *Clin J Am Soc Nephrol*. 2019;**14**(6):954-61. [PubMed ID: 30975657]. [PubMed Central ID: PMC6556719]. <https://doi.org/10.2215/CJN.00490119>.