



Psychosocial and Demographic Factors Impacting the Mental Workload of Nurses: A Proposed Structural Equation Model

Maryam Nourollahi ¹, Davood Afshari ^{2,*}, Noorollah Noorollah Karimi ³, Maryam Seyedtabib ⁴

¹ Department of Occupational Health, School of Health, Alborz University of Medical Sciences, Alborz, Iran

² Department of Occupational Health, School of Health, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran

³ Department of Occupational Health Engineering, School of Health, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran

⁴ Department of Biostatistics and Epidemiology, School of Health, Ahvaz Jundishapur University of Medical Sciences

*Corresponding Author: Department of Occupational Health, School of Health Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran. Email: davodafi@yahoo.com

Received: 14 January, 2024; Revised: 27 September, 2024; Accepted: 21 October, 2024

Abstract

Background: Physical and mental workload (MWL) are very important factors that affect patient safety and the quality of nursing services. Psychosocial risk factors in the workplace are among the most important factors that threaten nurses' MWL.

Objectives: The study was carried out to examine MWL and predict the psychosocial and demographic factors that affect the MWL of nurses, based on the structural equation model (SEM).

Methods: This cross-sectional descriptive-analytical study was conducted in 2021. Participants were 234 nurses from a teaching hospital in Ilam, Iran. General Health Questionnaire-12 (GHQ-12), the NASA-TLX Questionnaire, Copenhagen Psychosocial Questionnaire (COPSOQ), and a demographic questionnaire were used. The overall relationship between the variables was analyzed using the SEM. LISREL 8.8 software was used in the study to estimate the model and test the fit of the model.

Results: Based on the NASA-TLX Index, the overall MWL score was 70.09 ± 9.51 . Examining the MWL subscales showed that the mean effort score (71.41 ± 8.93), frustration (71.38 ± 5.50), and physical demand (70.74 ± 7.00) were higher than other dimensions of MWL. In the presented SEM, there was a significant relationship between individual and psychosocial factors with the MWL, and these factors, in interaction with each other, could predict MWL by 28%. Among the psychosocial factors examined, factor1 ($\beta = 0.88, P < 0.01$), (quality of leadership, social support from supervisors, recognition, justice, trust regarding management, predictability), and factor3 ($\beta = 0.95, P < 0.01$) (meaning of work, commitment to the workplace, influence, and role clarity) were among the factors affecting MWL.

Conclusions: The study results indicated the significance of psychosocial factors in nurses' MWL. Hence, controlling and managing effective psychosocial factors could help reduce MWL. Given that nurses are the main element of care in the treatment team, the results of the current study may have several implications for policymakers and nursing managers. Since nurses' MWL was in the range of high workload, workload management programs can help reduce MWL. By considering the prediction of nurses' MWL by psychosocial factors, some strategies should be considered by managers, including improving the quality of leadership, social support from supervisors, and also improving influence and role clarity.

Keywords: Mental Workload, Nurses, Psychosocial Factors, NASA-TLX

1. Background

In health systems, nurses are the main element of care in the treatment team. Nurses have always faced a high workload, given the nature of the job and the pressures caused by the high job demands. Workload includes both physical and mental elements. A systematic review has already shown that nurses

present high levels of mental workload (MWL) (1, 2). In this regard, the results of the meta-analysis study have shown that the highest MWL was reported in China (82.26 ± 14.82), Iran (75.18 ± 20.01), and America (71.40 ± 13.62), respectively (3).

An increase in MWL occurs when a person has many tasks to process and carry out in a certain period. If the

job demands exceed the individual's capacity and ability and the individual cannot face the new conditions with proper strategies, their performance and efficiency could decrease (4). Indeed, a high MWL can affect people's attention and physical skills and increase errors in performing tasks. Fatigue and excessive workload could be the potential reasons for medical errors (5). Thus, one of the major factors significantly increasing human error is the high MWL, which finally leaves adverse effects on the efficiency of people in the workplace (6, 7). Moreover, according to the study on nurses, people's work engagement negatively correlates with excessive MWL (8). Hence, MWL is a significant variable for understanding people's performance in work environments (5). Moreover, some studies have indicated that a high MWL in people results in physical fatigue and emotional exhaustion and could increase stress and burnout (9-11). Therefore, in sensitive working conditions, especially when there is time pressure and stress, such as the working conditions of nurses, paying attention to factors affecting MWL can play an important role in reducing errors and MWL.

The MWL is a multidimensional concept and is associated with the nature of a task, the abilities of the individual, and the characteristics of the environment. Hence, the MWL is affected by the cognitive demands of tasks and other factors like working conditions and motivation levels (8, 9). Epidemiological studies have shown that variables such as age, sex, and work experience are demographic risk factors associated with MWL (4, 5). A review of demographic risk factors in this occupation specifies that age, work experience, and having no children are some of the individual factors that affect the MWL (1). Therefore, for this study, we hypothesized that demographic characteristics affect MWL (Hypothesis 1).

One of the most important factors affecting MWL is the psychosocial factors of the work environment. Psychosocial factors are actually employees' perceptual experiences of the quality of work environments. Factors such as stress, emotional needs, burnout, job satisfaction, and the quality of leadership are known as the most important psychosocial risk factors (10, 11). The results of a study conducted on nurses in intensive care units have shown that 57% of the participants perceived high levels of exposure to psychosocial risk factors, and psychosocial factors explained 10.36% of MWL (12). Also,

the results of a study conducted on emergency nurses showed that job satisfaction was significantly related to mental demand and frustration. The results of this study showed that the mean scores for mental demand and frustration increased significantly in participants with low job satisfaction (13). In this regard, the results of the study conducted on healthcare workers have shown that MWL is inversely related to job satisfaction and job control (11). Also, the results of a study conducted on MWL among nurses have shown that social support was significantly negatively associated with the MWL in nurses (12). Psychosocial factors at work are among the main triggers of MWL; however, studies carried out in this field of health and also in the field of nursing on working conditions are scarce and focused on some psychosocial factors. Given this evidence, we examined the relationship between all psychosocial factors and MWL (Hypothesis 2).

2. Objectives

The study aims to (1) determine the MWL of nurses; (2) evaluate and identify psychosocial and demographic factors that affect MWL; and (3) present a conceptual model to predict the factors that affect the MWL (Figure 1).

3. Methods

3.1. Participants

This cross-sectional descriptive-analytical study was conducted in 2021. Participants were 234 nurses from a teaching hospital in Ilam, Iran. The sample size was calculated by considering a power of 95% and a probability of type I error of 0.05. According to a previous study and Equation 1, the required sample size was 234 people (13). The criteria for inclusion were: a) work experience of more than 1 year, and b) not suffering from chronic diseases or psychological problems.

$$n = \frac{\left(z_{1-\frac{\alpha}{2}} + z_{1-\beta}\right)^2 (\sigma_1^2 + \sigma_2^2)}{(\mu_1 - \mu_2)^2} \quad (1)$$

3.2. Measurements

3.2.1. NASA Task Load Index




Figure 1. The conceptual model of mental workload (MWL)

Table 1. NASA-TLX Scales

Dimension	Description
Mental demand	How much mental and perceptual activity was required (e.g., thinking, deciding, calculating, remembering, looking, searching, etc.)? Was the task easy or demanding, simple or complex, exacting, or forgiving?
Physical demand	How much physical activity was required (e.g., turning, controlling, activating, etc.)? Was the task easy or demanding, slow or brisk, slack or strenuous, restful or laborious?
Temporal demand	How much time pressure did you feel due to the rate or pace at which the tasks or task elements occurred? Was the pace slow and leisurely or rapid and frantic?
Performance	How successful do you think you were in accomplishing the goals of the task set by the experimenter (or yourself)? How satisfied were you with your performance in accomplishing these goals?
Effort	How hard did you have to work (mentally and physically) to accomplish your level of performance?
Frustration	Level How insecure, discouraged, irritated, stressed, and annoyed versus secure, gratified, content, relaxed, and complacent did you feel during the task?

NASA Task Load Index (NASA-TLX) is a multidimensional subjective MWL evaluation method. In this method, the overall workload score is obtained according to the weighted average of 6 subscales of workload. The subscales examined in this method were mental demand, physical demand, temporal demand, performance, effort, and frustration (Table 1). In this questionnaire, each field of activity is divided into 100 points with 5-point steps. The overall workload score was calculated according to the weighted average of ratings in 6 dimensions. The NASA TLX interpretation categorizes workload score values of 0 - 9 as low, 10 - 29 as medium, 30 - 49 as somewhat high, 50 - 79 as high, and 80 - 100 as very high (14). The validity and reliability

of the Persian version of NASA-TLX have been examined in a previous study (13).

3.2.2. Copenhagen Psychosocial Questionnaire

Copenhagen Psychosocial Questionnaire (COPSOQ) is one of the most complete and reliable questionnaires that examine psychosocial factors. In this study, the second version of COPSOQ with 4 main factors and 16 subfactors was used. The respondents to the COPSOQ are asked to respond to the questions using items on a 5-point scale, most of which are as follows: A = Always, B = Often, C = Sometimes, D = Seldom, and E = Never/Hardly, or as A = To a very large extent, B = To a large extent, C = Somewhat, D = To a small extent, and E = To a very small

Table 2. Demographic Information of the Participants (n = 234)^a

Variables	Values
Age (y)	36.62 ± 8.25
Work experience (y)	8.86 ± 4.89
Height (cm)	169.70 ± 10.35
Weight (kg)	69.99 ± 11.12
Gender	
Female	114 (48.7)
Male	120 (51.3)
Marital status	
Single	57 (24.4)
Married	172 (73.5)
Divorced	5 (2.1)
Educational level	
Bachelor degree	177 (75.6)
Master degree	47 (20.1)
PhD	10 (4.3)

^a Values are expressed as No. (%) or mean ± SD.

extent. Based on the subjects' ratings, each item (scores A to E) scored 0, 25, 50, 75, and 100, respectively, and the average score of the items on each of the factors determined the score of that factor. Giving a lower rate to each of the factors indicates better and more ergonomic psychosocial conditions of that work environment (15). In this study, the Persian version of the COPSOQ was used (16).

3.2.3. Demographic Form

Demographic information of the individuals, such as age, gender, marital status, work history, work department, degree, and field of study, was collected through a form.

3.2.4. General Health Questionnaire-12

The questionnaire is the short version of Goldberg's 60-question general health questionnaire, measuring a person's mental health in 12 items. Each item is scored on a four-point scale (less than usual, no more than usual, rather more than usual, or much more than usual). The questionnaire questions examine the mental state of the person in the last four weeks (one month). The positive items scored from 0 to 3, and the negative ones from 3 to 0. Total scores range from 0 to 36, with a score of 11 or 12 considered typical, scores > 15 suggesting evidence of distress, and scores > 20 considered severe problems with psychological distress

(17). In this study, the Persian version of the General Health Questionnaire-12 (GHQ-12) was used (18).

3.3. Data Analysis

The data were analyzed using SPSS version 23 and LISREL 8.8. The correlations between demographic factors, psychosocial factors, and MWL were explored using correlation analysis. Structural equation modeling (SEM) was used to identify causal relationships between variables. Psychosocial and demographic factors were considered independent variables, and MWL was considered dependent in this study. In SEM, the fit of the model was examined after estimating the model parameters and prior to interpreting them. The goodness-of-fit index, $\chi^2/df < 2$, GFI, CFI, NNFI, NFI > 0.90, and RMSEA < 0.08 were considered to indicate an adequate model fit.

4. Results

The mean general health score of the participants in the study was 1.36 ± 1.02 , and the maximum GHQ score was 3. Thus, considering the cut-off point of the GHQ questionnaire, all the participants in the study met the inclusion criteria. The mean age of the nurses participating in the study was 36.62 ± 8.25 . The mean work experience of the participants was 8.86 ± 4.89 (Table 2).

Table 3. Mean and Standard Deviation of Psychosocial Factors (COPSQ) (n = 234)

Factors	Mean ± SD
Factor 1 (quality of leadership, social support from supervisor, recognition, justice, trust regarding management, predictability)	1.76 ± 0.44
Factor 2 (general health, burnout, work-family conflict, emotional demands, exhaustion, stress)	1.85 ± 0.27
Factor 3 (meaning of work, commitment to the workplace, influence, role clarity)	1.57 ± 0.27
Factor 4 (offensive behavior)	2.54 ± 0.2
Total score	1.87 ± 0.34

Table 4. Mean and Standard Deviation of Mental Workload Subscales (n = 234)

NASA-TLX	Mean ± SD
Mental demand	69.68 ± 6.24
Physical demand	70.74 ± 7.00
Temporal demand	68.95 ± 7.17
Effort	71.41 ± 5.93
Performance	69.70 ± 7.35
Frustration	71.38 ± 8.50
Overall work load	70.09 ± 2.51

4.1. Psychosocial Factors

The mean score of the total psychosocial factors of the subjects examined was specified as 1.87. Psychosocial factors 2 (general health, burnout, work-family conflict, emotional demands, stress) and 1 [quality of leadership, social support from supervisor, recognition (reward), justice, trust regarding management, predictability] had the highest scores, respectively (Table 3).

4.2. Mental Workload

The mean score of the overall MWL of the subjects examined was determined as 70.09 ± 9.51. The analysis of the dimensions of MWL indicated that the mean scores of efforts (71.41 ± 8.93), frustration level (71.38 ± 5.50), and physical demand (70.74 ± 7.00), respectively, in the group of nurses examined were higher than other dimensions of MWL (Table 4).

4.3. Correlation Analysis

The results of the correlation of MWL with psychosocial and demographic factors showed that MWL has a positive correlation with factor 3 (meaning of work, Commitment to the workplace, Influence, Role clarity), factor 1 (quality of leadership, Social support from supervisor, Recognition (reward), Justice, Trust regarding management, Predictability), factor 4

(offensive behavior), and the total score of psychosocial factors ($P < 0.05$) (Table 5).

4.4. Structural Model

Figure 2 shows the path diagram with standardized factor loadings (numbers outside the parentheses between the latent and visible variables and endogenous and exogenous latent variables), errors (numbers outside the parentheses of the visible variables), and t values (numbers in parentheses) of the model. As Figure 2 shows, some variables show a relatively high correlation. $R^2 = 0.28$ is the relative value of the dependent or endogenous variable variance explained by the exogenous variable. The t values indicate that some loaded factors are significant at the 95% confidence interval (the absolute value of t values is greater than 1.96) (Figure 2). There was a significant relationship between individual and psychosocial factors with the MWL, and these factors in interaction had a 28% ability to predict MWL in the presented SEM (Figure 2).

4.5. Validation of the Research Model

As Table 6 shows, most of the fit indices for measuring the fit of the whole model show a good model fit, compatibility, and agreement of the whole model with experimental data.

Table 5. Mental Workload (NASA-TLX) Subscales and Psychosocial Factors (n = 234)^a

Variables	Factor 1	Factor 2	Factor 3	Factor 4	Total Score (COPSOQ)
Mental demand	0.311*	0.180**	0.248*	0.143	0.318*
Physical demand	0.164	0.264*	0.232*	0.083	0.110
Temporal demand	0.180	0.028	0.194**	0.167	0.906
Effort	0.257*	0.030	0.188**	-0.146	0.206**
Performance	0.198**	-0.250*	0.324*	0.283*	0.227*
Frustration	0.268*	0.158	0.269*	0.153	0.313*
Overall work load	0.303*	0.034	0.334*	0.299*	0.312*

Abbreviation: COPSOQ, Copenhagen Psychosocial Questionnaire.

^a Pearson correlation test: *significant < 0.01, ** significant < 0.05.

5. Discussion

The study was carried out to predict the psychosocial factors affecting the MWL of nurses based on a structural model. The study revealed that psychosocial factors, as well as demographic variables in interaction with each other, can predict 28% of the MWL of nurses. The study results showed that the mean NASA-TLX score of the nurses who participated was 70.09 ± 9.51 . Similarly, the results of a study conducted by Nasirizad Moghadam et al. among ICU nurses in Iran showed that the mean nurses' MWL was 70.21 (19). Regarding this, the study carried out by Li et al. on the MWL of nurses indicated that the mean MWL in 56.96% of the nurses studied was 73.59 ± 8.86 , which was in the range of high workload (20). Moreover, the results of a similar study conducted by Shan et al. have indicated that the MWL of 62.9% of the nurses was in the high workload perception group (21). The results of the systematic review study conducted by Yuan et al. have also highlighted that nurses experience high MWL and need urgent interventions to reduce MWL (2). Overall, studies have revealed that nurses experience more workload than other healthcare workers, including the time needed to carry out direct and indirect care (22, 23).

Studying the NASA-TLX dimensions in the present study indicated that the mean score of the effort dimension was higher and more challenging than other dimensions. Moreover, the mean score of frustration level and physical demand were higher than other dimensions of MWL, respectively. Consistent with the present study, Li et al. reported that a relatively high score was obtained in the subscales of "effort," "physical demand," and "mental demand" among the nurses examined (20). In this line, Sonmez et al. also reported

that the highest mean score of the MWL subscales were in mental demand, performance, and effort, respectively (24). In the same way, the results of a study conducted by Bazazan et al. also showed that the effort is more problematic than other dimensions (13). The results of the current study indicated that the level of frustration is another important dimension of MWL. Regarding this, the findings of a study conducted by Ebrahimi et al. have also shown that frustration is the second subscale of MWL with the highest score in evaluating nurses' MWL (25). In this regard, the results of a study conducted by Bazazan et al. also showed that frustration is one of the important dimensions of workload and the mean scores for frustration increased significantly in participants with low job satisfaction scores (13).

The present study results indicated that MWL significantly correlates with psychosocial factors. Thus, the MWL could be reduced by enhancing the identified factors such as quality of leadership, social support from supervisors, recognition (reward), justice, predictability, role clarity, general health, and reducing burnout, stress, work-family conflict, and offensive behavior. The results of SEM analysis in the study indicated that the key psychosocial factors affecting MWL were factor 1 (quality of leadership, social support from supervisors, recognition (reward), justice, trust regarding management, predictability) and factor 3 (meaning of work, commitment to the workplace, influence, role clarity). In this regard, Ceballos-Vásquez et al. examined psychosocial factors and MWL among nurses and reported a negative correlation between psychosocial factors and MWL (26). Studies have shown that nurses with higher levels of coworker and manager support report less daily workload (26, 27). In this line, Al-Hakim




Figure 2. Structural equation model (SEM) of mental workload (MWL) with standard loads, errors, t values and latent variable variance

Table 6. The Model Fit Indices for the Effect of Psychosocial and Personal Factors on Mental Work Load

Model Fit Indices	χ^2	χ^2/df	AGFI	GFI	NFI	NNFI	CFI	RMSEA
Cutoff values	> 0.05	< 3 or 2 <	> 0.9 or 0.8	> 0.9	> 0.9	> 0.9	> 0.9	< 0.08
SEM	0.08	2.88	0.86	0.85	0.95	0.93	0.94	0.06

Abbreviations: χ^2 , chi-square statistic; χ^2/df , normed chi-square; AGFI, adjusted goodness of fit index; GFI, Goodness of Fit Index; NFI, Normed Fit Index; NNFI, Non-normed Fit Index; CFI, Comparative Fit Index; RMSEA, root mean squared error of approximation; SEM, structural equation model.

et al. also illustrate the critical role of perceived organizational support on nurses' workload and job satisfaction (28). In this regard, in Afshari et al.'s study, the quality of leadership and social support have been mentioned as important psychosocial risk factors identified from the point of view of nurses (29). Moreover, the results of García-Iglesias et al. indicated that social support and quality of leadership are important psychosocial factors (30). Concerning this, Álvarez et al. revealed that social support and quality of leadership are among the psychosocial factors that affect the workload (31). Indeed, the presence of role clarity, autonomy, and supervisor support in the work environment of the nurses could help nurses control and manage workload and thus reduce workload. The results of Cai et al. showed that the leadership style and the logical arrangement of tasks could moderate nurses' workload and prevent the negative effects of high workload (32). Changes in leadership style, influence,

and reduction of concentration are the factors that reduce workload and increase motivation and performance from the point of view of nurses (33). In this regard, the results of a study conducted by Cho et al. showed that changing the leadership style and increasing nursing teamwork is important to monitor and modify high workload to protect nurses from elevated fatigue (34). Hence, identifying the psychosocial factors affecting the MWL of nurses could be useful in determining strategies and planning to reduce the MWL and increase productivity.

Besides psychosocial factors, the model presented in the study revealed that gender is one of the demographic factors that affect the MWL. In this regard, studies have shown a significant relationship between gender and MWL (1, 35). Similarly, Bagheri Hosseinabadi et al. also concluded that gender may be effective in increasing the MWL of nurses (4). Hence, identifying

individual factors that affect MWL could control and reduce the MWL of the nurses.

5.1. Strengths and Limitations

The present study examined MWL and psychosocial factors using a self-reporting questionnaire. Thus, there could be bias in the study results, and efforts were made to solve this limitation by fully explaining the questions of the questionnaires and the objectives of the study to the participants, which increased the accuracy of answering the questions. Among the other limitations of the study is its being conducted in a teaching hospital. As working conditions in private hospitals vary from teaching hospitals, it is suggested that a study should be conducted to examine psychosocial factors that affect MWL in private hospitals to generalize the results to all hospitals.

5.2. Conclusions

The study findings indicated that nurses' workload was relatively high. Studying the mean scores of the workload scales showed that the effort, frustration, and physical demand scores, respectively, have the highest values relative to other dimensions of MWL. According to the study results, psychosocial factors, including the quality of leadership, social support from supervisors, recognition (reward), justice, trust regarding management, predictability, meaning of work, commitment to the workplace, influence, and role clarity, as well as the demographic factor of gender, were among the factors affecting the MWL of nurses. Thus, the MWL of nurses could be reduced, especially by managing and enhancing the effective psychosocial factors.

Footnotes

Authors' Contribution: All authors made contributions to the study conception and design, made contributions to the interpretation of data. In addition, they also contributed to the drafting and revision of the article as well as approval and, submission of the final version.

Conflict of Interests Statement: The authors declare no conflict of interest.

Data Availability: Data of this study will be available upon the reasonable request from corresponding author.

Ethical Approval: This study was approved by the Ethics Review Committee of the Ahvaz Jundishapur University of Medical Sciences (IR.TUMS.CHMC.REC.1399.166).

Funding/Support: This study was funded by Ahvaz Jundishapur University of Medical Sciences (grant No. U-00069).

Informed Consent: All participants provided written informed consent prior to data collection.

References

- Jin M, Qian R, Wang J, Long J, Yuan Z, Zeng L, et al. Influencing factors associated with mental workload among nurses: A latent profile analysis. *Int J Nurs Sci*. 2024;**11**(3):330-7. [PubMed ID: 39156683]. [PubMed Central ID: PMC11329065]. <https://doi.org/10.1016/j.ijnss.2024.04.002>.
- Yuan Z, Wang J, Feng F, Jin M, Xie W, He H, et al. The levels and related factors of mental workload among nurses: A systematic review and meta-analysis. *Int J Nurs Pract*. 2023;**29**(5). e13148. [PubMed ID: 36950781]. <https://doi.org/10.1111/ijn.13148>.
- Weber J, Catchpole K, Becker AJ, Schlenker B, Weigl M. Effects of Flow Disruptions on Mental Workload and Surgical Performance in Robotic-Assisted Surgery. *World J Surg*. 2018;**42**(11):3599-607. [PubMed ID: 29845381]. <https://doi.org/10.1007/s00268-018-4689-4>.
- Bagheri Hosseinabadi M, Khanjani N, Etemadinezhad S, Samaei SE, Raadabadi M, Mostafae M. The associations of workload, individual and organisational factors on nurses' occupational injuries. *J Clin Nurs*. 2019;**28**(5-6):902-11. [PubMed ID: 30357967]. <https://doi.org/10.1111/jocn.14699>.
- Schroers G, Ross JG, Moriarty H. Nurses' Perceived Causes of Medication Administration Errors: A Qualitative Systematic Review. *Jt Comm J Qual Patient Saf*. 2020. [PubMed ID: 33153914]. <https://doi.org/10.1016/j.jcjq.2020.09.010>.
- Al Ma'mari Q, Sharour LA, Al Omari O. Fatigue, burnout, work environment, workload and perceived patient safety culture among critical care nurses. *Br J Nurs*. 2020;**29**(1):28-34. [PubMed ID: 31917951]. <https://doi.org/10.12968/bjon.2020.29.1.28>.
- Zahednezhad H, Shokrollahi N, Gheshlagh RG, Afshar PF. Does heavy mental workload affect moral sensitivity among critical care unit nursing professionals? a cross-sectional study. *BMC Nurs*. 2021;**20**(1):140. [PubMed ID: 34376186]. [PubMed Central ID: PMC8353778]. <https://doi.org/10.1186/s12912-021-00662-8>.
- Longo L, Wickens CD, Hancock G, Hancock PA. Human Mental Workload: A Survey and a Novel Inclusive Definition. *Front Psychol*. 2022;**13**:883321. [PubMed ID: 35719509]. [PubMed Central ID: PMC9201728]. <https://doi.org/10.3389/fpsyg.2022.883321>.
- Galy E, Paxion J, Berthelon C. Measuring mental workload with the NASA-TLX needs to examine each dimension rather than relying on

- the global score: an example with driving. *Ergonomics*. 2018;**61**(4):517-27. [PubMed ID: 28817353]. <https://doi.org/10.1080/00140139.2017.1369583>.
10. Rosyidawati D, Noor NB, Zulkifli A. The Influence of Workload, Job Satisfaction and Work Motivation on Nurse Performance in Hospital Inpatient Installation. *J Asian Multicultural Res Soc Sci Study*. 2020;**1**(2):56-63. <https://doi.org/10.47616/jamrsss.vii2.60>.
 11. Hellin Gil MF, Ruiz Hernandez JA, Ibanez-Lopez FJ, Seva Llor AM, Roldan Valcarcel MD, Mikla M, et al. Relationship between Job Satisfaction and Workload of Nurses in Adult Inpatient Units. *Int J Environ Res Public Health*. 2022;**19**(18). [PubMed ID: 36141970]. [PubMed Central ID: PMC9517381]. <https://doi.org/10.3390/ijerph19181701>.
 12. Teng M, Yuan Z, He H, Wang J. Levels and influencing factors of mental workload among intensive care unit nurses: A systematic review and meta-analysis. *Int J Nurs Pract*. 2024;**30**(4). e13167. [PubMed ID: 37259643]. <https://doi.org/10.1111/ijn.13167>.
 13. Bazazan A, Dianat I, Bahrapour S, Talebian A, Zandi H, Sharafkhaneh A, et al. Association of musculoskeletal disorders and workload with work schedule and job satisfaction among emergency nurses. *Int Emerg Nurs*. 2019;**44**:8-13. [PubMed ID: 30902617]. <https://doi.org/10.1016/j.ienj.2019.02.004>.
 14. Hart SG, Staveland LE. Development of NASA-TLX (Task Load Index): Results of Empirical and Theoretical Research. *Human Mental Workload*. 1988. p. 139-83. [https://doi.org/10.1016/s0166-4115\(08\)62386-9](https://doi.org/10.1016/s0166-4115(08)62386-9).
 15. Pejtersen JH, Kristensen TS, Borg V, Bjorner JB. The second version of the Copenhagen Psychosocial Questionnaire. *Scand J Public Health*. 2010;**38**(3 Suppl):8-24. [PubMed ID: 21172767]. <https://doi.org/10.1177/1403494809349858>.
 16. Aminian M, Dianat I, Miri A, Asghari-Jafarabadi M. The Iranian version of the Copenhagen Psychosocial Questionnaire (COPSOQ) for assessment of psychological risk factors at work. *Health Promot Perspect*. 2017;**7**(1):7-13. [PubMed ID: 28058236]. [PubMed Central ID: PMC5209652]. <https://doi.org/10.15171/hpp.2017.03>.
 17. Goldberg D, Williams P. *A user's guide to the General Health Questionnaire*. Windsor, UK: nferNelson;1988.
 18. Montazeri A, Vahdaninia M, Ebrahimi M, Jarvandi S. The Hospital Anxiety and Depression Scale (HADS): translation and validation study of the Iranian version. *Health Qual Life Outcomes*. 2003;**1**:14. [PubMed ID: 12816545]. [PubMed Central ID: PMC161819]. <https://doi.org/10.1186/1477-7525-1-14>.
 19. Nasirizad Moghadam K, Chehrzad MM, Reza Masouleh S, Maleki M, Mardani A, Atharyan S, et al. Nursing physical workload and mental workload in intensive care units: Are they related? *Nurs Open*. 2021;**8**(4):1625-33. [PubMed ID: 33596333]. [PubMed Central ID: PMC8186696]. <https://doi.org/10.1002/nop2.785>.
 20. Li D, Hu Y, Chen H, Zhu X, Wu X, Li J, et al. Identifying the Subtypes and Characteristics of Mental Workload Among Chinese Physicians in Outpatient Practice: A Latent Profile Analysis. *Front Public Health*. 2021;**9**:779262. [PubMed ID: 34900919]. [PubMed Central ID: PMC8653799]. <https://doi.org/10.3389/fpubh.2021.779262>.
 21. Shan Y, Shang J, Yan Y, Lu G, Hu D, Ye X. Mental workload of frontline nurses aiding in the COVID-19 pandemic: A latent profile analysis. *J Adv Nurs*. 2021;**77**(5):2374-85. [PubMed ID: 33594687]. [PubMed Central ID: PMC8014576]. <https://doi.org/10.1111/jan.14769>.
 22. Rostami F, Babaei-Pouya A, Teimori-Boghsani G, Jahangirimehr A, Mehri Z, Feiz-Arefi M. Mental Workload and Job Satisfaction in Healthcare Workers: The Moderating Role of Job Control. *Front Public Health*. 2021;**9**:683388. [PubMed ID: 34540781]. [PubMed Central ID: PMC8446529]. <https://doi.org/10.3389/fpubh.2021.683388>.
 23. Sardo PMG, Macedo RPA, Alvarelhao JIM, Simoes JFL, Guedes JAD, Simoes CJ, et al. Nursing workload assessment in an intensive care unit: A retrospective observational study using the Nursing Activities Score. *Nurs Crit Care*. 2023;**28**(2):288-97. [PubMed ID: 36336353]. <https://doi.org/10.1111/nicc.12854>.
 24. Sonmez B, Oguz Z, Kutlu L, Yildirim A. Determination of nurses' mental workloads using subjective methods. *J Clin Nurs*. 2017;**26**(3-4):514-23. [PubMed ID: 27434614]. <https://doi.org/10.1111/jocn.13476>.
 25. Ebrahimi H, Jafarjalal E, Lotfolahzadeh A, Kharghani Moghadam SM. The effect of workload on nurses' quality of life with moderating perceived social support during the COVID-19 pandemic. *Work*. 2021;**70**(2):347-54. [PubMed ID: 34633341]. <https://doi.org/10.3233/WOR-210559>.
 26. Ceballos-Vasquez P, Rolo-Gonzalez G, Hernandez-Fernaund E, Diaz-Cabrera D, Paravic-Klijin T, Burgos-Moreno M. Psychosocial factors and mental work load: a reality perceived by nurses in intensive care units. *Rev Lat Am Enfermagem*. 2015;**23**(2):315-22. [PubMed ID: 26039303]. [PubMed Central ID: PMC4459006]. <https://doi.org/10.1590/0104-1169.0044.2557>.
 27. Trottier M, Bentein K. Coworker support as a moderator in the relationship between daily experience of workload and an individual's experience of same-day WFC: a buffer or an intensifier? *Commun, Work Family*. 2018;**22**(5):569-88. <https://doi.org/10.1080/13668803.2018.1434484>.
 28. Al-Hakim L, Zhang Y, Jin J, Sevdalis N. The effect of psychological meaningfulness and perceived organisational support on the relationship between nursing workload and job satisfaction: A prospective, cross-sectional investigation. *Int J Nurs Stud*. 2022;**133**:104274. [PubMed ID: 35759809]. <https://doi.org/10.1016/j.ijnurstu.2022.104274>.
 29. Afshari D, Nourollahi-Darabad M, Chinisaz N. Psychosocial Factors Associated With Resilience Among Iranian Nurses During COVID-19 Outbreak. *Front Public Health*. 2021;**9**:714971. [PubMed ID: 34422753]. [PubMed Central ID: PMC8373242]. <https://doi.org/10.3389/fpubh.2021.714971>.
 30. Garcia-Iglesias JJ, Gomez-Salgado J, Ortega-Moreno M, Navarro-Abal Y. Relationship Between Work Engagement, Psychosocial Risks, and Mental Health Among Spanish Nurses: A Cross-Sectional Study. *Front Public Health*. 2020;**8**:627472. [PubMed ID: 33575245]. [PubMed Central ID: PMC7870998]. <https://doi.org/10.3389/fpubh.2020.627472>.
 31. Álvarez AKG, Almaguer AYC, Ramírez GR, López AMM, Godales TDJG, Ruiz AL. Capacitación en seguridad psicológica a la alta dirección hospitalaria en el afrontamiento a la COVID-19. *Educación Médica Superior*. 2021;**35**.
 32. Cai Y, Li Q, Cao T, Wan Q. Nurses' work engagement: The influences of ambidextrous leadership, clinical nurse leadership and workload. *J Adv Nurs*. 2023;**79**(3):1152-61. [PubMed ID: 34723406]. <https://doi.org/10.1111/jan.15086>.
 33. Morsiani G, Bagnasco A, Sasso L. How staff nurses perceive the impact of nurse managers' leadership style in terms of job satisfaction: a mixed method study. *J Nurs Manag*. 2017;**25**(2):119-28. [PubMed ID: 27917561]. <https://doi.org/10.1111/jonm.12448>.
 34. Cho H, Sagherian K, Scott LD, Steege LM. Occupational fatigue, workload and nursing teamwork in hospital nurses. *J Adv Nurs*. 2022;**78**(8):2313-26. [PubMed ID: 35396873]. <https://doi.org/10.1111/jan.15246>.

35. Mohammadi F, Hanifi N, Bahraminegad N. [Investigating the Relationship between Nurses' Mental Workload and the Quality of Care Services in Intensive Care Unit]. *Critical Care Nurs.* 2021;**14**(2):38-47. FA.