



The Relationship Between Perceived Burdensomeness and Psychosocial Adjustment to Illness in Patients with Chronic Pain: A Mediating Role of Pain Self-Efficacy

Esmail Sadri Damirchi ^{1,*}, Shahriar Dargahi ¹, Nader Ayadi ² and Davod Fathi ¹

¹Faculty of Educational Sciences and Psychology, University of Mohaghegh Ardabili, Ardabil, Iran

²Faculty of Educational Sciences and Psychology, University of Isfahan, Isfahan, Iran

*Corresponding author: Associate Professor, Faculty of Educational Sciences and Psychology, Mohaghegh Ardabili University, Ardabil, Iran. Email: e.sadri@uma.ac.ir

Received 2019 October 22; Revised 2020 February 26; Accepted 2020 February 27.

Abstract

Background: The majority of patients living with chronic pain faces several challenges; therefore, factors affecting their adaptation to the disease to provide appropriate therapeutic methods and meditation and relaxation therapy should be identified.

Objectives: The present study aimed at investigating the relationship between perceived burdensomeness and psychosocial adjustment, and the mediating role of pain self-efficacy.

Methods: This descriptive correlational study was conducted on 240 patients with chronic pain referred to the orthopedic centers in Ardabil in 2019 selected by convenience sampling. Psychosocial Adjustment to Illness Scale, Pain Self-efficacy Questionnaire, and Interpersonal Needs Questionnaire were used for data collection. Also, Pearson correlation and multiple regressions were used for data analysis. Data were analyzed using SPSS V.19 and AMOS V.21.

Results: The findings confirmed the negative and direct effect of perceived burdensomeness ($\beta = -0.11$ and $P < 0.01$), as well as the positive and direct effect of pain self-efficacy ($\beta = 0.86$ and $P < 0.01$) on psychosocial adjustment. On the other hand, perceived burdensomeness ($\beta = -0.32$, $P < 0.01$) had a negative and direct effect on pain self-efficacy. The indirect effect of perceived burdensomeness and pain self-efficacy on psychosocial adjustment to illness was also confirmed.

Conclusions: According to the findings of the present study, it can be concluded that the feelings of pain self-efficacy mediate the relationship between perceived burdensomeness and psychosocial adjustment in patients with chronic pain. That can also be important for the patient and their caregivers in relation to other chronic illnesses. Chronic diseases can also affect patients and their caregivers.

Keywords: Perceived Burdensomeness, Psychosocial Adjustment, Pain Self-Efficacy, Chronic Pain Patients

1. Background

The pain may be moderate to severe with serious negative consequences for the working or social life of the individual and can be changed over time. The perception of pain is highly subjective and unique, and its threshold is developed in individuals through the experiences related to injury in early life (1). Chronic pain, meanwhile, is a common disease that can change affected people's lives, and experiencing chronic pain is one of the most common reasons to seek medical care (2). Chronic pain is a depressive condition, in which the person experiences not only the mental stress caused by the pain but also many other stressors that affect different parts of his life (2). Based on the existing research literature, long-term painful physical symptoms can lead to greater comorbidity of physical and

mental illnesses, in addition to the quality of life deterioration (3). Chronic pain is one of the most common reasons adults seek medical care that has been linked to restrictions in mobility and daily activities, drug dependence, anxiety, and depression (4). The prevalence of chronic pain among adults in Tehran, Iran has been reported 24%, of which chronic low back pain (12.4%) and chronic knee pain (11.2%) were the most common complaints (5).

Despite many physical and psychological problems faced by chronic pain patients, adaptation to the disease has been reported as one of the factors contributing to disease acceptance and as an important part of participation in the treatment process, which can affect the social and physical functioning of individuals (6). Adaptation to illness is the ability to maintain a positive attitude toward oneself and the world despite having physical prob-

lems (7). Psychosocial adjustment to the disease consists of six subscales: attitude toward illness, work environment, family environment, sexual relations, development of family relationships, social environment, and psychological disorders (8). Adaptation to the process of treatment of chronic disease is an important factor in attracting a patient's participation in treatment and assisting in the decision-making process (9). Studies have shown that psychosocial adjustment to illness is the strongest predictor of health care utilization compared with other disease variables (10). The importance of psychosocial factors as significant predictors of pain and functioning in different persons has been reported (11).

The process of coping with a chronic illness is a dynamic process that is constantly affected by psychological and social stimuli. In this process, the affected person must face individual and environmental challenges to reach an acceptable level of physical, mental and social wellbeing leading to achieve successful adaptation (12). One of the psychological challenges is being overwhelmed by the illness and its associated care. According to the interpersonal theory, when the person is overbearing, incapable of providing safety and well-being, and he does not help the group can cause the feeling of being overburdened (13). Perceived burdensomeness refers to the view that being overbearing is about friends, family, or the community comprising two dimensions: (a) selfishness means that I hate myself, and (b) I feel responsible; for example, my death will be worth more than my life to others (14). Perceived burdensomeness is also seen as a dynamic cognitive state that can be developed by risk factors, such as homelessness, unemployment, physical well-being, and low self-esteem (15). Various studies have investigated the relationship between feeling overwhelmed and different factors and have focused the effect of these variables on psychological trauma and self-harm behaviors indicating their importance in mental and physical health (16). It has shown that living with chronic pain is associated with significant emotional stress. Moreover, pain can decrease the patients' emotional abilities and also his continuous desire to get rid of pain will often be unattainable. This ultimately weakens the patient's mood, hopelessness, and depression (17).

Pain self-efficacy can enhance coping skills and thereby increase patients' adaptation to chronic pain by enhancing patients' cognitive beliefs about their ability to solve problems. It has defined as patients' beliefs in their abilities to successfully organize and control health habits, and also achieve valuable health outcomes in different situations and contexts (18). Pain self-efficacy beliefs are an important determinant of pain behaviors and indicate how much a person will try to endure distressing problems and experiences (19). Pain self-efficacy moderates the effects of

hopelessness in patients. Therefore, it acts against hopelessness as a bumper and causes resistance to hopelessness and fatigue caused by illness (20-22). It has also reported that pain self-efficacy improves functional ability and reduces pain-related fear. Promoting pain self-efficacy beliefs can be a key mechanism for improving behavioral outcomes (23).

Consequences of chronic pain on the different aspects of patients' lives, such as their psychological, social, and occupational functioning (2), the importance of psychosocial adjustment of patients with chronic pain, and assessment and identification of psychological adjustment affecting the social functioning of these patients have yet been addressed in previous studies and should be more considered. Therefore, this study aimed at investigating the role of burdensomeness on psychosocial adjustment in patients with chronic pain and whether pain self-efficacy can mediate the relationship between burdensomeness and psychosocial adjustment of patients with chronic pain.

2. Objectives

The present study aimed at investigating the relationship between perceived burdensomeness and psychosocial adjustment, and the mediating role of pain self-efficacy

3. Methods

The present descriptive correlational study was conducted on all patients with chronic pain referred to the orthopedic centers in Ardebil. The minimum sample size of 200 subjects was estimated for a structural equation modeling and path analysis (24). Using a purposive sampling method, 240 patients with chronic pain were selected. Inclusion criteria were having at least two years of chronic pain, no psychiatric problems, informed consent, having at least a diploma (including admission requirements). The exclusion criteria were being illiterate, unwillingness to continue the research, and having psychiatric problems.

3.1. Data Collection Tools

Psychosocial Adjustment to Illness Scale (PAIS): This tool was designed in 1990 by Leonardo Derogatis to assess the quality of a patient's psychosocial adjustment to a chronic illness. It is a 46-item scale including 7 subscales: attitude toward disease (8 questions), work environment (6 questions), family environment (8 questions), sexual relationships (6 questions), family relationship development (5 questions), social environment (6 questions),

and psychological disorders (7 questions). In this study, each item was scored on a 4-point Likert scale (range 0 - 3) with choices varying from "not at all" to "severe difficulties with household duties, with higher scores indicating greater impairment. The sum scores of each component were divided by the number of questions in that component and its mean was considered as the adjustment score of the component, whereas the sum of the total score of all questions was divided by the number of all questions and its mean was considered as the total adjustment score. The questionnaire was first translated by Feghhi et al. (8). It was then reviewed by 3 people fluent in English and finally reviewed and approved by an expert. Some changes were made on the scale based on the Iranian culture and it was localized. Since the majority of the subjects were housewives and retired men, the question related to absenteeism was omitted and the number of questions was reduced to 45 questions. The content validity of the questionnaire was confirmed by 10 university professors and its reliability was calculated to be 0.94 using Cronbach's alpha in 20 patients with type 2 diabetes (8).

Pain Self-efficacy Questionnaire (PSEQ): the PSEQ is a 10-item questionnaire, developed by Bandura in 1977 to assess the confidence people with ongoing pain have in performing activities while in pain, which is scored on a 7-point Likert scale (range 0 - 6). A Higher score indicates greater self-efficacy in performing and functioning despite ongoing pain. The Persian version of PSEQ has reported with acceptable reliability. The validity of this questionnaire by confirmatory factor analysis supported a one-factor solution (25).

Interpersonal Needs Questionnaire (INQ): INQ has multiple versions, of which the version 15 (INQ-15) is a self-report measure of perceived burdensomeness and thwarted belongingness that participants are asked to indicate the degree to which each item had been true for them recently. The first six statements measure perceived burdensomeness (e.g., "These days, I think I am a burden on society"), and the last nine statements measure thwarted belongingness (e.g., "These days, I feel like I belong"). INQ-15 is scored on a 7-point scale (range 1 - 7) with higher scores indicating greater thwarted belongingness and perceived burdensomeness. Internal consistency ($\alpha = 90\%$) and good reliability for this scale have been reported (26, 27). Also, in another study, confirmatory factor analysis of the two-factor model of perceived burdensomeness and thwarted belongingness provided an adequate fit for the data (28).

In this research, the ethical considerations, such as informed consent, willingness to participate in research, and confidentiality of participants' data were observed. The obtained data were analyzed by Pearson correlation and

structural equation modeling using SPSS V. 19 and AMOS V. 21.

4. Results

Of the participants, 61% were female and 39% were male. Also, 33% of them had a history of surgery, 20% physiotherapy, 17% were taking and injecting drugs, and 30% of them had performed physical exercise for recovery. Table 1 represents the mean, standard deviation, and correlation matrix of the variables.

The results of model fit indices (after correction) are provided in Table 2. Based on the obtained results, it can be concluded that the model fits the data well and both absolute and comparative indices reached the desired level.

Regarding the question "Does pain self-efficacy play a mediating role in the relationship between perceived burdensomeness and psychosocial adjustment?", the results indicated that pain self-efficacy had a mediating role in the research model. The detailed results of the research hypothesis are presented in Figure 1.

Based on the results of Figure 1 and Table 3, the negative and direct effect of perceived burdensomeness ($\beta = -0.11$, $P < 0.01$) on psychosocial adjustment, as well as the positive and direct effect of pain self-efficacy ($\beta = 0.86$, $P < 0.001$) on psychosocial adjustment were also confirmed. On the other hand, perceived burdensomeness ($\beta = -0.32$, $P < 0.01$) had a negative and direct effect on pain self-efficacy.

To investigate the mediating role of pain self-efficacy in the relationship between perceived burdensomeness and psychosocial adjustment to the disease, Bootstrap was used and the results are summarized in Table 4.

According to the results of Table 4, the indirect effect of exogenous variable perceived burdensomeness on pain self-efficacy and psychosocial adjustment to illness was confirmed.

5. Discussion

The aim of the present study was to investigate the relationship between perceived burdensomeness and psychosocial adjustment, and the mediating role of pain self-efficacy in chronic pain patients. The result of this study showed that the feeling of being overburdened had a negative and significant relationship with psychosocial adjustment in patients with chronic pain. The findings are consistent with the results of other relevant studies, such as Vahidi and Shaker Dioulagh study (29), in which stress had a significant and negative relationship with mental adjustment to the disease. In line with the results of the present study, it has found that the caregiver's perception

Table 1. Correlation Matrix Among the Research Variables

	M	SD	1	2	3
Perceived burdensomeness	38.57	5.69	1		
Pain self-efficacy	34.32	4.04	-0.47 ^a	1	
Psychosocial adjustment to illness	78.81	12.79	-0.19 ^b	0.89 ^a	1

^aP < 0.01.
^bP < 0.05.

Table 2. Model Fit Indices

Index	X ² /df	Df	GFI	NFI	IFI	CFI	RMSEA
Values	2.03	249	0.98	0.98	0.99	0.99	0.07




Figure 1. Standard coefficients of the research conceptual model. INQ, perceived burdensomeness; PAIS, psychosocial adjustment to illness; PSEQ, pain self-efficacy.

Table 3. Path Coefficients for Direct Effects in the Model

Direct Effects	Estimate d Value	Standard Value	S.E.	C.R.	P Value
Perceived burdensomeness on psychosocial adjustment	-1.64	-0.11	0.59	-2.77	0.001
Perceived burdensomeness on pain self-efficacy	-0.29	-0.32	0.06	-4.76	0.001
Pain self-efficacy on psychosocial adjustment	13.78	0.86	1.31	10.49	0.001

of the burden makes them susceptible to some personal and marital problems (30).

Since patients' attitudes, beliefs, and expectations about themselves and their problems affect the level of

pain, disability, and response to treatment (31), they indicate the relationship between perceived burdensomeness and low psychosocial adjustment in patients with chronic pain. It can be said that when a patient with chronic pain

Table 4. Estimation of Indirect Paths in the Model Using the Bootstrap Method

Variable			Upper Value	Lower Line	Significant Line	Estimation Level
Predictor Variable	Intermediate Variable	Criterion Variable				
Perceived burdensomeness	Pain self-efficacy	Psychosocial adjustment to illness	-0.27	-0.16	-0.38	0.01

develops a negative attitude toward himself and his illness and blames himself for it and suffers from his family members, a cycle of negative thoughts can be formed that motivates and frustrates the patient by providing the backgrounds for the development of depression. It also leads to separation from the job and social position in the patient, which gradually results in psychosocial adjustment. This cycle has defined as a process of maintaining a positive attitude toward yourself and the world despite having physical problems (32). Therefore, it is clear that feelings of perceived burdensomeness by developing a negative attitude toward yourself will decrease the psychosocial adjustment in patients with chronic pain.

In addition, the findings of the present study showed that pain self-efficacy -despite having a positive relationship with psychosocial adjustment in patients with chronic pain- can mediate the negative relationship between perceived burdensomeness and psychosocial adjustment. This finding is in line with the results of a relevant study indicating a positive and significant relationship between self-efficacy and psychosocial adjustment (33). It has also shown that self-efficacy plays a mediating role in the relationship between happiness and pain self-management in patients (22).

According to the social cognitive theory in explaining the mediating role of pain self-efficacy in the relationship between perceived burdensomeness and psychosocial adjustment in patients with chronic pain, it can be said that people with high levels of pain self-efficacy utilize desirable levels of pain relief and discomfort management, and consequently pain control (24), while reducing emotional distress and disability (34). They can not only increase their ability to cope with the disease by reducing their negative emotions and anxiety, but also improve their overall functioning levels. Such factors can reduce the feeling of helplessness and overburdened in these people. In this regard, it has believed that people who are more adaptable than their less-adapted counterparts are more optimistic about themselves and others focusing more on their life priorities, and consequently on intimate family relationships. They are more likely to cope easily with their physical conditions and increase their physical activities and are optimistic about their diseases leading to faster recovery (35). Also, regarding the relationship

between perceived burdensomeness and psychosocial adjustment of patients with chronic pain, it has reported that perceived burdensomeness and the resulted negative attitude of the patients towards themselves can decrease their mental functioning by creating negative and depressive thoughts. To explain the mediating role of pain self-efficacy in the relationship between these two variables, it is important to conduct a study on the mediating role of pain self-efficacy on the relationship between depression and suicidal ideation in patients with chronic pain indicating a significant relationship between depression and suicidal ideation through pain self-efficacy (36). The present study had some limitations. The used subjects were selected from the patients with chronic pain in Ardabil city, which can affect the generalizability of the results and was the most important limitation of this research.

5.1. Conclusions

The purpose of this study was to investigate the mediating role of pain self-efficacy in the relationship between perceived burdensomeness and psychosocial adjustment in patients with chronic pain. The findings of the present study showed that despite the negative and significant relationship between perceived burdensomeness and psychosocial adjustment in patients with chronic pain, pain self-efficacy can play a mediating role in the relationship between these two variables so that pain self-efficacy can mediate the negative relationship between perceived burdensomeness and psychosocial adjustment in patients with chronic pain.

Acknowledgments

Hereby, we express our appreciation to the Research Administration of the University of Mohaghegh Ardabili and those who helped us to conduct the study.

Footnotes

Authors' Contribution: Drafting of the manuscript: Esmaeil Sadri; study concept and design: Shahriar Dargahi; analysis and interpretation of data: Nader Ayadi and Davod Fathi; all authors read and approved the final manuscript.

Conflict of Interests: The authors have no competing interests to declare.

Ethical Approval: This article was extracted from the approved research by the Vice-Chancellor of Research of the University of Mohaghegh Ardabili (approval code: 10788-98/5/23).

Funding/Support: This study was supported by the University of Mohaghegh Ardabili.

References

- Ratter J, Radlinger L, Lucas C. Several submaximal exercise tests are reliable, valid and acceptable in people with chronic pain, fibromyalgia or chronic fatigue: A systematic review. *J Physiother.* 2014;**60**(3):144-50. doi: [10.1016/j.jphys.2014.06.011](https://doi.org/10.1016/j.jphys.2014.06.011). [PubMed: [25084634](https://pubmed.ncbi.nlm.nih.gov/25084634/)].
- Kapur BM, Lala PK, Shaw JL. Pharmacogenetics of chronic pain management. *Clin Biochem.* 2014;**47**(13-14):1169-87. doi: [10.1016/j.clinbiochem.2014.05.065](https://doi.org/10.1016/j.clinbiochem.2014.05.065). [PubMed: [24912048](https://pubmed.ncbi.nlm.nih.gov/24912048/)].
- Li CT, Su TP, Hsieh JC, Ho ST. Efficacy and practical issues of repetitive transcranial magnetic stimulation on chronic medically unexplained symptoms of pain. *Acta Anaesthesiol Taiwan.* 2013;**51**(2):81-7. doi: [10.1016/j.aat.2013.06.003](https://doi.org/10.1016/j.aat.2013.06.003). [PubMed: [23968659](https://pubmed.ncbi.nlm.nih.gov/23968659/)].
- Dahlhamer J, Lucas J, Zelaya C, Nahin R, Mackey S, DeBar L, et al. Prevalence of chronic pain and high-impact chronic pain among adults - United States, 2016. *MMWR Morb Mortal Wkly Rep.* 2018;**67**(36):1001-6. doi: [10.15585/mmwr.mm6736a2](https://doi.org/10.15585/mmwr.mm6736a2). [PubMed: [30212442](https://pubmed.ncbi.nlm.nih.gov/30212442/)]. [PubMed Central: [PMC6146950](https://pubmed.ncbi.nlm.nih.gov/PMC6146950/)].
- Mohammadzadeh F, Faghihzadeh S, Asadi Lari M, Vaez Mahdavi MR, Arab Kheradmand J, Noorbala AA, et al. A fairly comprehensive survey of chronic pain in Iranian population: Prevalence, risk factors, and impact on daily life. *Health Scope.* 2015;**4**(3). e25467. doi: [10.17795/jhealthscope-25467](https://doi.org/10.17795/jhealthscope-25467).
- Chabowski M, Junke M, Juzwisyj N, Milan M, Malinowski M, Janczak D. Adaptation to illness in relation to pain perceived by patients after surgery. *J Pain Res.* 2017;**10**:1447-52. doi: [10.2147/JPR.S129936](https://doi.org/10.2147/JPR.S129936). [PubMed: [28721086](https://pubmed.ncbi.nlm.nih.gov/28721086/)]. [PubMed Central: [PMC5499957](https://pubmed.ncbi.nlm.nih.gov/PMC5499957/)].
- Sharpe L, Curran L. Understanding the process of adjustment to illness. *Soc Sci Med.* 2006;**62**(5):1153-66. doi: [10.1016/j.socscimed.2005.07.010](https://doi.org/10.1016/j.socscimed.2005.07.010). [PubMed: [16112783](https://pubmed.ncbi.nlm.nih.gov/16112783/)].
- Feghhi H, Saadatjoo SA, Dastjerdi R. [Psychosocial adaptation in patients with type 2 diabetes referring to Diabetes Research Center of Birjand in 2013]. *Modern Care J.* 2014;**10**(4):249-56. Persian.
- Drummond-Young M, LeGris J, Browne GB, Pallister R, Roberts J. Interactional styles of out-patients with poor adjustment to chronic illness receiving problem-solving counselling. *Health Soc Care Community.* 1996;**4**(6):317-29. doi: [10.1111/j.1365-2524.1996.tb00078.x](https://doi.org/10.1111/j.1365-2524.1996.tb00078.x).
- Browne GB, Arpin K, Corey P, Fitch M, Gafni A. Individual correlates of health service utilization and the cost of poor adjustment to chronic illness. *Med Care.* 1990;**28**(1):43-58. doi: [10.1097/00005650-199001000-00006](https://doi.org/10.1097/00005650-199001000-00006). [PubMed: [2136926](https://pubmed.ncbi.nlm.nih.gov/2136926/)].
- Jensen MP, Moore MR, Bockow TB, Ehde DM, Engel JM. Psychosocial factors and adjustment to chronic pain in persons with physical disabilities: A systematic review. *Arch Phys Med Rehabil.* 2011;**92**(1):146-60. doi: [10.1016/j.apmr.2010.09.021](https://doi.org/10.1016/j.apmr.2010.09.021). [PubMed: [21187217](https://pubmed.ncbi.nlm.nih.gov/21187217/)]. [PubMed Central: [PMC3028590](https://pubmed.ncbi.nlm.nih.gov/PMC3028590/)].
- Khalilzadeh H, Nanbakhsh J, Yaghoobi M. [Evaluation of compatibility of heart failure patients with problems caused by disease in Taleghani educational center of Urmia]. *J Urmia Nurs Midwifery Facul.* 2006;**4**(4):1-6. Persian.
- Bryan CJ. The clinical utility of a brief measure of perceived burdensomeness and thwarted belongingness for the detection of suicidal military personnel. *J Clin Psychol.* 2011;**67**(10):981-92. doi: [10.1002/jclp.20726](https://doi.org/10.1002/jclp.20726). [PubMed: [20715022](https://pubmed.ncbi.nlm.nih.gov/20715022/)].
- Ma J, Batterham PJ, Calear AL, Han J. A systematic review of the predictions of the Interpersonal-Psychological Theory of Suicidal Behavior. *Clin Psychol Rev.* 2016;**46**:34-45. doi: [10.1016/j.cpr.2016.04.008](https://doi.org/10.1016/j.cpr.2016.04.008). [PubMed: [27155061](https://pubmed.ncbi.nlm.nih.gov/27155061/)].
- Van Orden KA, Lynam ME, Hollar D, Joiner TE. Perceived burdensomeness as an indicator of suicidal symptoms. *Cognit Ther Res.* 2006;**30**(4):457-67. doi: [10.1007/s10608-006-9057-2](https://doi.org/10.1007/s10608-006-9057-2).
- Silva C, Ribeiro JD, Joiner TE. Mental disorders and thwarted belongingness, perceived burdensomeness, and acquired capability for suicide. *Psychiatry Res.* 2015;**226**(1):316-27. doi: [10.1016/j.psychres.2015.01.008](https://doi.org/10.1016/j.psychres.2015.01.008). [PubMed: [25650048](https://pubmed.ncbi.nlm.nih.gov/25650048/)].
- Rosenzweig S, Greeson JM, Reibel DK, Green JS, Jasser SA, Beasley D. Mindfulness-based stress reduction for chronic pain conditions: Variation in treatment outcomes and role of home meditation practice. *J Psychosom Res.* 2010;**68**(1):29-36. doi: [10.1016/j.jpsychores.2009.03.010](https://doi.org/10.1016/j.jpsychores.2009.03.010). [PubMed: [20004298](https://pubmed.ncbi.nlm.nih.gov/20004298/)].
- Marks R, Allegrante JP, Lorig K. A review and synthesis of research evidence for self-efficacy-enhancing interventions for reducing chronic disability: Implications for health education practice (part II). *Health Promot Pract.* 2005;**6**(2):148-56. doi: [10.1177/1524839904266792](https://doi.org/10.1177/1524839904266792). [PubMed: [15855284](https://pubmed.ncbi.nlm.nih.gov/15855284/)].
- Bazaryi Meimand M, Alipour A, Poladi Reishahri A, Habibi Asgarabadi M. [Effect of chronic and recurrent pain on the self-management of chronic spinal pain: Moderator role of pain self-efficacy and catastrophic thoughts]. *J Anesth Pain.* 2017;**8**(1):60-77. Persian.
- Johnson J, Gooding PA, Wood AM, Tarrier N. Resilience as positive coping appraisals: Testing the schematic appraisals model of suicide (SAMS). *Behav Res Ther.* 2010;**48**(3):179-86. doi: [10.1016/j.brat.2009.10.007](https://doi.org/10.1016/j.brat.2009.10.007). [PubMed: [19906364](https://pubmed.ncbi.nlm.nih.gov/19906364/)].
- Knittle KP, De Gucht V, Hurkmans EJ, Vlieland TP, Peeters AJ, Ronda HK, et al. Effect of self-efficacy and physical activity goal achievement on arthritis pain and quality of life in patients with rheumatoid arthritis. *Arthritis Care Res (Hoboken).* 2011;**63**(11):1613-9. doi: [10.1002/acr.20587](https://doi.org/10.1002/acr.20587). [PubMed: [21898861](https://pubmed.ncbi.nlm.nih.gov/21898861/)].
- Meredith P, Strong J, Feeney JA. Adult attachment, anxiety, and pain self-efficacy as predictors of pain intensity and disability. *Pain.* 2006;**123**(1-2):146-54. doi: [10.1016/j.pain.2006.02.025](https://doi.org/10.1016/j.pain.2006.02.025). [PubMed: [16644132](https://pubmed.ncbi.nlm.nih.gov/16644132/)].
- Saberi A, Rezaei S, Shabani F, Ghorbani Shirkouhi S. The role of self-efficacy beliefs in psychological symptoms and clinical outcomes of primary chronic headache. *J Pract Clin Psychol.* 2019;**7**(3):175-86. doi: [10.32598/jpcp.7.3.175](https://doi.org/10.32598/jpcp.7.3.175).
- Dworkin RH, Breitbart WS. *Psychosocial aspects of pain: A handbook for health care providers (progress in pain research and management, v. 27)*. International Assn for the Study of Pain (IASP) Press; 2004.
- Siyami M, Rostami S. [Path Analysis of the Effect of mobile phone usage on academic engagement: the Mediating Role of attention]. *Educ Res J.* 2017;**4**(35):72-92. Persian.
- Boroumand A, Asghari Moghaddam MA, Shaeeri MR, Mesgarian F. [Chronic pain, pain self-efficacy and suicidal ideation: The moderating role of pain self-efficacy on relation between depression and suicidal ideation in chronic pain patients]. *J Fund Ment Health.* 2012;**14**(2):152-63. Persian.
- Hawkins KA, Hames JL, Ribeiro JD, Silva C, Joiner TE, Cogle JR. An examination of the relationship between anger and suicide risk through the lens of the interpersonal theory of suicide. *J Psychiatry.* 2014;**50**:59-65. doi: [10.1016/j.jpsychires.2013.12.005](https://doi.org/10.1016/j.jpsychires.2013.12.005). [PubMed: [24388767](https://pubmed.ncbi.nlm.nih.gov/24388767/)].
- Teo DCH, Suarez L, Oei TPS. Validation of the interpersonal needs questionnaire of young male adults in Singapore. *PLoS One.* 2018;**13**(6). e0198839. doi: [10.1371/journal.pone.0198839](https://doi.org/10.1371/journal.pone.0198839). [PubMed: [29879192](https://pubmed.ncbi.nlm.nih.gov/29879192/)]. [PubMed Central: [PMC5991743](https://pubmed.ncbi.nlm.nih.gov/PMC5991743/)].
- Van Orden KA, Cukrowicz KC, Witte TK, Joiner TE. Thwarted belongingness and perceived burdensomeness: Construct validity and psychometric properties of the Interpersonal Needs Questionnaire. *Psychol*

- Assess. 2012;**24**(1):197-215. doi: [10.1037/a0025358](https://doi.org/10.1037/a0025358). [PubMed: [21928908](https://pubmed.ncbi.nlm.nih.gov/21928908/)]. [PubMed Central: [PMC3377972](https://pubmed.ncbi.nlm.nih.gov/PMC3377972/)].
30. Vahidi A, Shaker Dioulagh A. [The role of caregiver burden, marital stress, and mindfulness in predicting psychosocial adjustment to illness in people with coronary artery disease]. *J Cardiovasc Nurs*. 2018;**7**(1):50-9. Persian.
 31. Showani E, Zaharakar K, Rasooli M. [Effectiveness of acceptance and commitment therapy on reducing the components of couple burnout in veterans spouses]. *Iran J War Public Health*. 2016;**8**(2):111-8. Persian.
 32. Main CJ, Sullivan MJ, Watson PJ. *Pain management: practical applications of the biopsychosocial perspective in clinical and occupational settings*. Elsevier Health Sciences; 2007.
 33. Kolokotroni P, Anagnostopoulos F, Missitzis I. Psychosocial Adjustment to Illness Scale: Factor structure, reliability, and validity assessment in a sample of Greek breast cancer patients. *Women Health*. 2017;**57**(6):705-22. doi: [10.1080/03630242.2016.1186780](https://doi.org/10.1080/03630242.2016.1186780). [PubMed: [27158891](https://pubmed.ncbi.nlm.nih.gov/27158891/)].
 34. Yusoff YM. Self-Efficacy, perceived social support, and psychological adjustment in international undergraduate students in a public higher education institution in Malaysia. *J Stud Int Educ*. 2012;**16**(4):353-71. doi: [10.1177/1028315311408914](https://doi.org/10.1177/1028315311408914).
 35. Wells-Federman C, Arnstein P, Caudill M. Nurse-led pain management program: effect on self-efficacy, pain intensity, pain-related disability, and depressive symptoms in chronic pain patients. *Pain Manag Nurs*. 2002;**3**(4):131-40. doi: [10.1053/jpmn.2002.127178](https://doi.org/10.1053/jpmn.2002.127178). [PubMed: [12454805](https://pubmed.ncbi.nlm.nih.gov/12454805/)].
 36. Mehnert A, Koch U. Prevalence of acute and post-traumatic stress disorder and comorbid mental disorders in breast cancer patients during primary cancer care: A prospective study. *Psychooncology*. 2007;**16**(3):181-8. doi: [10.1002/pon.1057](https://doi.org/10.1002/pon.1057). [PubMed: [16856147](https://pubmed.ncbi.nlm.nih.gov/16856147/)].