



Effect of Fibroblast Growth Factor 21 on the Expression of *TLR4* and *BAMBI* Genes in Cholesterol-Treated Human Hepatic Stellate Cells

Elham Shakerian ¹, Narges Mohammadtaghvaei ², Zohre Askari¹ and Reza Afarin ^{1,*}

¹Department of Clinical Biochemistry, School of Medicine, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran

²Hyperlipidemia Research Center, Department of Laboratory Sciences, School of Paramedical Sciences, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran

*Corresponding author: Student Research committee, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran. Email: r.afarin1988@gmail.com

Received 2021 February 08; Revised 2021 September 07; Accepted 2021 September 07.

Abstract

Background: Activated hepatic stellate cells (HSCs) are the primary mediators in the progression of hepatic fibrosis. The activation of toll-like receptor 4 (*TLR4*) signaling leads to the downregulation of the transmembrane inhibitory transforming growth factor-beta (*TGF-β*) pseudoreceptor BMP and activin membrane-bound inhibitor (*BAMBI*) on HSCs. Fibroblast growth factor 21 (FGF21) is a natural secretory protein in the body with effects, such as the reduction of fat accumulation and oxidation of lipids; however, no study has investigated FGF21 ability to prevent the progression of liver fibrosis.

Objectives: This study aimed to examine the beneficial effects of FGF21 to reduce cholesterol-activated human HSCs.

Methods: The human HSCs were incubated in media containing different concentrations of cholesterol, including 25, 50, 75, 100, 125, and 150 μ M, for 24 h and then incubated with FGF21 for 24 h. Total ribonucleic acids were extracted and reversely transcribed into complementary deoxyribonucleic acid. A quantitative real-time polymerase chain reaction was performed in this study.

Results: The results showed that the messenger ribonucleic acid (mRNA) expression of *TGF-β*, collagen, type I, alpha 1 (collagen1 α), and *TLR4* genes increased significantly in the presence of cholesterol (75 and 100 μ M), compared to that of the control group (* $P < 0.05$, ** $P < 0.01$, and *** $P < 0.001$); nevertheless, the mRNA expression of the *BAMBI* gene significantly reduced, compared to that of the control group (* $P < 0.05$). The FGF21 significantly reduced the mRNA expression of *TGF-β*, collagen1 α , and *TLR4* genes (# $P < 0.05$). The mRNA expression of the *BAMBI* gene significantly increased with FGF21 (# $P < 0.05$).

Conclusions: It was concluded that the treatment with FGF21 reduces the cholesterol-activated HSCs by decreasing the mRNA expression of the *TLR4*, *TGF-β*, and collagen1 α genes and increasing the mRNA expression of the *BAMBI* gene.

1. Background

Liver fibrosis as a metabolic disorder is currently a global problem. The transplantation of all or part of the liver tissue is the only solution to treat patients with acute liver fibrosis; therefore, it is important to find other treatments. Hepatic stellate cells (HSCs) are the main mediators of liver fibrosis. When HSCs activate, they produce the extracellular matrix (ECM) and collagen, type I, alpha 1 (collagen1 α) in response to damage to hepatocytes (1, 2). The damage can be caused by various factors, including severe viral hepatitis and non-alcoholic steatohepatitis (NASH) (3). If left untreated, the excessive accumulation of the ECM proteins ultimately leads to liver failure and hepatocellular carcinoma (4).

Among the factors that can increase the activity of star-shaped cells, it can be said that the transforming growth factor-beta (*TGF-β*) plays the most important role. According to previous studies on liver fibrosis pathways, the increased activation of the *TGF-β* signaling pathway could

eventually lead to the production of more proteins, such as collagen1 α , thereby accumulating the ECM (5). The primary activation of the *TGF-β* signaling pathway is through well-known receptors called serine/threonine kinase at the cell surface (6). In some studies, it has been suggested that by inhibiting the activation of the *TGF-β* and its downstream pathways, it is possible to prevent the activation of as many star-shaped cells as possible. Therefore, by inhibiting the effects of *TGF-β*, fibrogenic conditions in the liver can be improved (7, 8).

The activation of toll-like receptor 4 (*TLR4*) signaling leads to the downregulation of the transmembrane inhibitory *TGF-β* pseudoreceptor BMP and activin membrane-bound inhibitor (*BAMBI*) on HSCs, a critical fibrogenic cell type in the liver. As a result, net *TGF-β*1 activity increases, enhancing the ECM production (9, 10). Fibroblast growth factor 21 (FGF21) is a natural secretory protein in the body with effects, such as the reduction of fat accumulation and oxidation of lipids (11). Animal

experiments showed that the administration of recombinant FGF21 has beneficial effects on fatty liver and hepatic steatosis and improves fibrotic conditions and liver toxicity (12). However, no study has investigated the ability of FGF21 to protect against hepatic fibrosis in cholesterol-treated human HSCs.

2. Objectives

The present study aimed to investigate the effect of FGF21 to reduce cholesterol-treated activated human HSCs. The obtained results demonstrated that FGF21 inhibited *TLR4* signaling, thereby inhibiting the activation of HSCs; therefore, FGF21 can act as an influential factor in preventing liver fibrosis.

3. Methods

3.1. Materials

Cell culture grade cholesterol and FGF21 were provided from Sigma-Aldrich (USA) and Abnova (Taiwan), respectively. Fetal bovine serum (FBS) from Gibco (USA), Dulbecco's modified Eagle's Medium (DMEM) Low Glucose, and Pen/Strep from Bio-Idea (Iran) were purchased. Next, FGF21 (final concentration of 1.0 μ M) was dissolved in a culture medium and added to the cells for 24 h. The present study was designed and conducted with the permission of the Ethics Committee of Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran (IR.AJUMS.REC.1399.649).

3.2. Cell Culture

The LX-2 cell line (i.e., an immortalized human HSC line) as an HSC cell line was kindly provided by Scott L. Friedman (Mount Sinai School of Medicine, USA). The cells were seeded in DMEM containing 10% FBS and penicillin and streptomycin antibiotics (100 mg/mL each) in a 5% CO₂ humidified incubator at 37°C.

3.3. Treatment of LX-2 Cells with Different Cholesterol Concentrations

For cholesterol treatment, the cells were plated in 6-well plates at the density of 10⁵ cells per well and grown in the 5% CO₂ incubator at 37°C until 80% confluency. The cells were serum-starved for 24 h before treatments and then treated with different concentrations of cholesterol, including 25, 50, 75, 100, 125, and 150 μ M, for 24 h; then, the cells were subject to serum starvation for 24 h.

3.4. Treatment of Cholesterol-Activated HSCs with FGF21

In the following step, for the inspection of the inhibitory effects of FGF21 on activated HSCs, cholesterol-activated HSCs (75 and 100 μ M) were treated by FGF21 (1 μ M) for 24 h. Real-time polymerase chain reaction (PCR) was performed to evaluate the messenger ribonucleic acid (mRNA) expression of genes involved in liver fibrogenesis.

3.5. MTT Assay Technique

MTT (3-(4,5-Dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide) assay was performed to evaluate cell survival in the presence of cholesterol compound. In each well of 96 cells, about 5000 cells were cultured and incubated for 24 h. Then, the cells of each well (13) were treated with various cholesterol concentrations of 25, 50, 75, 100, 125, and 150 μ M for 24 h in an incubator at 37°C and 5% CO₂. Then, the medium was changed, and the MTT compound (0.5 mg/mL) was added to the cells and incubated for 4 h. The MTT-containing medium was then removed, and 100 μ L of dimethyl sulfoxide was added to each well to dissolve formazan crystals. Finally, the absorbance of the samples at 595 nm was determined.

3.6. Quantitative Real-time PCR

Ribonucleic acid (RNA) was extracted by a kit of Qiagen Company (Germany) according to the manufacturer's instructions. The optical density wavelength of 260 to 280 readings and the number 2.0 were obtained using the nanodrop device. Complementary deoxyribonucleic acid synthesis was then performed on total RNA. For the quantification of the mRNA expression, real-time PCR was performed using RealQ Plus 2x Master Mix Green "Low Rox" kit (Ampliqon, Denmark) and QuantStudio™ 3 Real-time PCR System (ABI Applied Biosystems, USA) consisted of a hot start by heating the PCR solution to 95°C for 10 min, followed by 38 cycles of denaturation at 95°C for 10 sec and annealing/extension at 58°C for 1 min. The total time of running this protocol on the system of the current study was 90 min with the following specific primers:

TGF- β : forward 5'-GTGGACATCAACGGGTTCAC-3', reverse: 5'-CTCCGTGGAGCTGAAGCAATA-3';

TLR4: forward 5'-GCCAGGATGATGCTGCCTC-3', reverse 5'-TTAGGAACCACTCCACGC-3';

BAMBI: forward 5'-GACAGACATCTGCCAAGCCA-3', reverse 5'-TGATACCTGTTTCTTGCTCTGA-3';

Collagen1 α : forward 5'-GGAATGAAGGGACACAGAGGTT-3', reverse 5'-AGTAGCACCATCATTTCCACGA-3';

Glyceraldehyde 3-phosphate dehydrogenase: forward 5'-GACAGTCAGCCGATCTTCT-3', reverse 5'-GCCCAATACGACAAATCCGT-3'

3.7. Statistical Analysis

All the experiments were performed in triplicate. The data were analyzed as mean \pm standard error of the mean using analysis of variance and Tukey's posthoc test using GraphPad Prism software (version 9.0.1). The significance level was considered less than 0.05.

4. Results

4.1. Effect of Different Cholesterol Concentrations on the Survival of HSCs

The MTT assay was performed to achieve appropriate concentrations of cholesterol for cell treatment. For this purpose, the cells were first treated with various cholesterol concentrations of 25, 50, 75, 100, 125, and 150 μ M for 24 h. The control group was not treated with cholesterol. At a concentration of 125 μ M cholesterol, the percentage of star-shaped cells was significantly reduced, compared to that of the control group ($P < 0.001$; Figure 1). Therefore, concentrations lower than the half-maximal inhibitory concentration were used for the experiment.

4.2. mRNA Expression of TLR4 and BAMBI Genes in the Presence of Cholesterol and FGF21

Qualitative real-time PCR was used for the evaluation of the mRNA expression of *TLR4* and *BAMBI* genes in the presence of different cholesterol concentrations. The results showed that the mRNA expression of these genes did not significantly change in the presence of 25 and 50 μ M cholesterol; however, in the presence of 75 and 100 μ M concentrations, the mRNA expression of the *TLR4* gene significantly increased, and the mRNA expression of the *BAMBI* gene significantly reduced, compared to those reported for the control group (* $P < 0.05$, ** $P < 0.01$, and *** $P < 0.001$; Figure 2A). The modulation of *TLR4* and *BAMBI* genes by 1 μ M FGF21 was evaluated following their induction with 75 and 100 μ M cholesterol. Qualitative real-time PCR results showed that the mRNA expression of the *TLR4* gene significantly reduced and the *BAMBI* gene significantly increased in response to FGF21, compared to those reported for the cells treated with 75 and 100 μ M cholesterol (# $P < 0.05$; Figure 2B).

4.3. mRNA Expression of TGF- β and Collagen1 α Genes in the Presence of Cholesterol and FGF21

Qualitative real-time PCR was used for the evaluation of the mRNA expression of the TGF- β and collagen1 α genes in the presence of different cholesterol concentrations. The obtained results showed that the mRNA expression of these genes did not significantly change in the presence of 25 and 50 μ M cholesterol; nevertheless, in the presence of

75 and 100 μ M cholesterol, the mRNA expression of TGF- β and collagen1 α genes significantly increased, compared to that reported for the control group (* $P < 0.05$ and ** $P < 0.01$; Figure 3A). The modulation of TGF- β and collagen1 α genes expression by 1 μ M FGF21 was evaluated following their induction with 75 and 100 μ M cholesterol. Real-time PCR results showed that the mRNA expression of TGF- β and collagen1 α genes significantly reduced in response to FGF21, compared to that reported for the cells treated with 75 and 100 μ M cholesterol (# $P < 0.05$; Figure 3B).

5. Discussion

Hepatic fibrosis is a significant health problem worldwide for which there is no acceptable therapy. The most important reason for hepatic fibrosis is the excess deposition of collagen1 α . These activated HSCs secrete the ECM in hepatic fibrogenesis. Therefore, collagen1 α is a helpful marker for the assessment of hepatic fibrosis (14). Several studies have been performed to understand the molecular mechanisms that lead to the expansion of hepatic fibrosis (15).

It has been previously reported that *TLR4* enhances TGF- β signaling in HSCs, HSC activation, and hepatic fibrosis (16, 17). Recognizing the factors involved in the onset and progression of liver fibrosis are important for the prevention of this disease. The HSCs are central contributors to the initiation and progression of liver fibrosis (18). Therefore, it is important to determine the factors that cause HSCs activation and develop hepatic fibrosis.

In this study, the effects of several cholesterol concentrations were studied on the activation of LX-2 cells and changes in the mRNA expression of fibrogenic genes. The present study showed that at high concentrations, cholesterol could activate human HSCs and increase the mRNA expression of fibrogenic genes; nevertheless, no such changes were observed at lower concentrations.

To date, numerous efforts have been made to find effective drugs to prevent the progression of liver fibrosis; however, it is still a great problem to find drugs that can control the progression of liver fibrosis. The FGF21 is a natural secretory protein in the body with effects, such as the reduction of fat accumulation and oxidation of lipids. Recently, the administration of FGF21 has been shown to have several metabolic benefits on insulin sensitivity and reduce the accumulation of fat in the liver in obese rats without specific side effects (19, 20).

Studies on humans have shown high levels of circulating FGF21 in patients with type 2 diabetes and non-alcoholic fatty liver disease/NASH (21). On the other hand, previous studies showed that the administration of recombinant FGF21 has beneficial effects on fatty liver and hepatic




Figure 1. The 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyl-2H-tetrazolium bromide assay results showing cell viability under different cholesterol concentrations over 24 h. Results are shown as mean \pm standard error of the mean. Statistical analyses were performed using GraphPad Prism software (version 9.0.1), one-way analysis of variance, and Tukey's test. Statistical significance was considered < 0.05 (** $P < 0.01$ and *** $P < 0.001$).

steatosis and improves fibrotic conditions and liver toxicity (22, 23). The current study investigated whether FGF21 can reduce activated human HSCs treated with cholesterol. The result of the present study showed that FGF21 reduced activated human HSCs treated with cholesterol. In this study, FGF21 significantly reduced the mRNA expression of $TGF-\beta$, collagen α , and *TLR4* genes in cholesterol-treated HSCs.

In summary, FGF21 can reduce the activation of LX-2 cells treated with cholesterol in vitro. The relevant mechanism is probably through the inhibition of the *TLR4*/ $TGF-\beta$ signaling pathway. However, the present study was subject to some limitations, and the results have to be observed in light of these limitations. All the experiments were carried out on cells cultured and treated in a specific cell line in a laboratory. It is advised to test on living organisms,

including mice or humans. Hepatic fibrosis is a dynamic process resulting from the cooperation of HSCs with other liver cell types that warrants further examination on the effect of cholesterol and FGF21 on all hepatic cells' interactions. It is necessary to perform further studies to evaluate the cholesterol effect on mice to measure liver damage and inflammation by measuring liver enzymes and staining liver tissue.

5.1. Conclusions

This study showed that high cholesterol treatment activates human HSCs; nevertheless, cholesterol has no such effect at lower concentrations. The FGF21 is effective in reducing the activation of HSCs treated with cholesterol in vitro. Therefore, FGF21 has significant antifibrotic properties in liver fibrosis.




Figure 2. Toll-like receptor 4 (*TLR4*) and BMP and activin membrane-bound inhibitor (*BAMBI*) messenger ribonucleic acid (mRNA) expression in the presence of cholesterol and fibroblast growth factor 21 (FGF21) in LX-2 cell line. A, *TLR4* mRNA expression in the presence of cholesterol and FGF21 in LX-2 cells. B, *BAMBI* mRNA expression in the presence of cholesterol and FGF21 in LX-2 cells. The results of three replications (mean \pm standard error of the mean) and changes in *TLR4* and *BAMBI* mRNA expression were compared to those of the control group. The significance level was considered < 0.05 . Glyceraldehyde 3-phosphate dehydrogenase was used as the reference gene (* $P < 0.05$, ** $P < 0.01$, and *** $P < 0.001$; # $P < 0.05$).




Figure 3. Transforming growth factor-beta (TGF-β) and collagen, type I, alpha 1 (collagen1α) messenger ribonucleic acid (mRNA) expression in the presence of cholesterol and fibroblast growth factor 21 (FGF21) in LX-2 cell line. A, TGF-β mRNA expression in the presence of cholesterol and FGF21 in LX-2 cells. B, collagen1α mRNA expression in the presence of cholesterol and FGF21 in LX-2 cells. The results of three replications (mean \pm standard error of the mean) and changes in TGF-β and collagen1α mRNA expression were compared to those of the control group. The significance level was considered < 0.05 . Glyceraldehyde 3-phosphate dehydrogenase was used as the reference gene (* $P < 0.05$ and ** $P < 0.01$; # $P < 0.05$).

Footnotes

Authors' Contribution: Study concept and design, R. A. and N. M.; Analysis and interpretation of the data, E. SH. and R. A.; Drafting of the manuscript, Z. A. All the authors read and approved the final manuscript.

Conflict of Interests: All the authors affirm that there is no conflict of interest.

Ethical Approval: The present study was designed and conducted with the permission of the Ethics Committee of Ahvaz Jundishapur University of Medical Sciences (IR.AJUMS.REC.1399.649).

Funding/Support: This study was financially supported by Ahvaz Jundishapur University of Medical Sciences (grant no.: 99s66).

References

- Abbas N, Getachew A, You K, Shah Z, Chen Y, Tao J, et al. Kupffer cells mediate the recruitment of hepatic stellate cells into the localized liver damage. *Biochem Biophys Res Commun.* 2020;**529**(2):474–9. doi: [10.1016/j.bbrc.2020.06.041](https://doi.org/10.1016/j.bbrc.2020.06.041). [PubMed: [32703454](https://pubmed.ncbi.nlm.nih.gov/32703454/)].
- Cai X, Wang J, Wang J, Zhou Q, Yang B, He Q, et al. Intercellular crosstalk of hepatic stellate cells in liver fibrosis: New insights into therapy. *Pharmacol Res.* 2020;**155**:104720. doi: [10.1016/j.phrs.2020.104720](https://doi.org/10.1016/j.phrs.2020.104720). [PubMed: [32092405](https://pubmed.ncbi.nlm.nih.gov/32092405/)].
- Higashi T, Friedman SL, Hoshida Y. Hepatic stellate cells as key target in liver fibrosis. *Adv Drug Deliv Rev.* 2017;**121**:27–42. doi: [10.1016/j.addr.2017.05.007](https://doi.org/10.1016/j.addr.2017.05.007). [PubMed: [28506744](https://pubmed.ncbi.nlm.nih.gov/28506744/)]. [PubMed Central: [PMC5682243](https://pubmed.ncbi.nlm.nih.gov/PMC5682243/)].
- Saber S. Angiotensin II: A key mediator in the development of liver fibrosis and cancer. *Bull Natl Res Cent.* 2018;**42**(1). doi: [10.1186/s42269-018-0020-7](https://doi.org/10.1186/s42269-018-0020-7).
- Yoshida K, Murata M, Yamaguchi T, Matsuzaki K. TGF-beta/Smad signaling during hepatic fibro-carcinogenesis (review). *Int J Oncol.* 2014;**45**(4):1363–71. doi: [10.3892/ijo.2014.2552](https://doi.org/10.3892/ijo.2014.2552). [PubMed: [25050845](https://pubmed.ncbi.nlm.nih.gov/25050845/)]. [PubMed Central: [PMC4151811](https://pubmed.ncbi.nlm.nih.gov/PMC4151811/)].
- Mu M, Zuo S, Wu RM, Deng KS, Lu S, Zhu JJ, et al. Ferulic acid attenuates liver fibrosis and hepatic stellate cell activation via inhibition of TGF-beta/Smad signaling pathway. *Drug Des Devel Ther.* 2018;**12**:4107–15. doi: [10.2147/DDDT.S186726](https://doi.org/10.2147/DDDT.S186726). [PubMed: [30584275](https://pubmed.ncbi.nlm.nih.gov/30584275/)]. [PubMed Central: [PMC6284527](https://pubmed.ncbi.nlm.nih.gov/PMC6284527/)].
- Xiang D, Zou J, Zhu X, Chen X, Luo J, Kong L, et al. Physalin D attenuates hepatic stellate cell activation and liver fibrosis by blocking TGF-beta/Smad and YAP signaling. *Phytomedicine.* 2020;**78**:153294. doi: [10.1016/j.phymed.2020.153294](https://doi.org/10.1016/j.phymed.2020.153294). [PubMed: [32771890](https://pubmed.ncbi.nlm.nih.gov/32771890/)].
- Yang X, Dan X, Men R, Ma L, Wen M, Peng Y, et al. MiR-142-3p blocks TGF-beta-induced activation of hepatic stellate cells through targeting TGFbetaRI. *Life Sci.* 2017;**187**:22–30. doi: [10.1016/j.lfs.2017.08.017](https://doi.org/10.1016/j.lfs.2017.08.017). [PubMed: [28823564](https://pubmed.ncbi.nlm.nih.gov/28823564/)].
- Zhao HW, Zhang ZF, Chai X, Li GQ, Cui HR, Wang HB, et al. Oxy-matine attenuates CCl4-induced hepatic fibrosis via modulation of TLR4-dependent inflammatory and TGF-beta1 signaling pathways. *Int Immunopharmacol.* 2016;**36**:249–55. doi: [10.1016/j.intimp.2016.04.040](https://doi.org/10.1016/j.intimp.2016.04.040). [PubMed: [27179304](https://pubmed.ncbi.nlm.nih.gov/27179304/)].
- Friedman SL. A deer in the headlights: BAMBI meets liver fibrosis. *Nat Med.* 2007;**13**(11):1281–2. doi: [10.1038/nm1107-1281](https://doi.org/10.1038/nm1107-1281). [PubMed: [17987019](https://pubmed.ncbi.nlm.nih.gov/17987019/)].
- Lin Z, Tian H, Lam KS, Lin S, Hoo RC, Konishi M, et al. Adiponectin mediates the metabolic effects of FGF21 on glucose homeostasis and insulin sensitivity in mice. *Cell Metab.* 2013;**17**(5):779–89. doi: [10.1016/j.cmet.2013.04.005](https://doi.org/10.1016/j.cmet.2013.04.005). [PubMed: [23663741](https://pubmed.ncbi.nlm.nih.gov/23663741/)].
- Ye D, Wang Y, Li H, Jia W, Man K, Lo CM, et al. Fibroblast growth factor 21 protects against acetaminophen-induced hepatotoxicity by potentiating peroxisome proliferator-activated receptor coactivator protein-1alpha-mediated antioxidant capacity in mice. *Hepatology.* 2014;**60**(3):977–89. doi: [10.1002/hep.27060](https://doi.org/10.1002/hep.27060). [PubMed: [24590984](https://pubmed.ncbi.nlm.nih.gov/24590984/)].
- Lim SC, Parajuli KR, Duong HQ, Choi JE, Han SI. Cholesterol induces autophagic and apoptotic death in gastric carcinoma cells. *Int J Oncol.* 2014;**44**(3):805–11. doi: [10.3892/ijo.2014.2246](https://doi.org/10.3892/ijo.2014.2246). [PubMed: [24399152](https://pubmed.ncbi.nlm.nih.gov/24399152/)].
- Kundu A, Dey P, Park JH, Kim IS, Kwack SJ, Kim HS. EX-527 prevents the progression of high-fat diet-induced hepatic steatosis and fibrosis by upregulating SIRT4 in Zucker rats. *Cells.* 2020;**9**(5). doi: [10.3390/cells9051101](https://doi.org/10.3390/cells9051101). [PubMed: [32365537](https://pubmed.ncbi.nlm.nih.gov/32365537/)]. [PubMed Central: [PMC7290750](https://pubmed.ncbi.nlm.nih.gov/PMC7290750/)].
- Distler JHW, Gyorfi AH, Ramanujam M, Whitfield ML, Konigshoff M, Lafyatis R. Shared and distinct mechanisms of fibrosis. *Nat Rev Rheumatol.* 2019;**15**(12):705–30. doi: [10.1038/s41584-019-0322-7](https://doi.org/10.1038/s41584-019-0322-7). [PubMed: [31712723](https://pubmed.ncbi.nlm.nih.gov/31712723/)].
- Liao X, Zhan W, Tian T, Yu L, Li R, Yang Q. MicroRNA-326 attenuates hepatic stellate cell activation and liver fibrosis by inhibiting TLR4 signaling. *J Cell Biochem.* 2019. doi: [10.1002/jcb.29520](https://doi.org/10.1002/jcb.29520). [PubMed: [31692098](https://pubmed.ncbi.nlm.nih.gov/31692098/)].
- Liu L, Zhou Y, Dai D, Xia H, Zhao K, Zhang J. Antifibrotic effects of kangxian ruangan capsule on rats with nonalcoholic fatty liver fibrosis and hepatic stellate cells through regulation of TGF-beta and TLR4 signaling pathways. *Evid Based Complement Alternat Med.* 2021;**2021**:5649575. doi: [10.1155/2021/5649575](https://doi.org/10.1155/2021/5649575). [PubMed: [34422075](https://pubmed.ncbi.nlm.nih.gov/34422075/)]. [PubMed Central: [PMC8371615](https://pubmed.ncbi.nlm.nih.gov/PMC8371615/)].
- Liu L, You Z, Yu H, Zhou L, Zhao H, Yan X, et al. Mechanotransduction-modulated fibrotic microniches reveal the contribution of angiogenesis in liver fibrosis. *Nat Mater.* 2017;**16**(12):1252–61. doi: [10.1038/nmat5024](https://doi.org/10.1038/nmat5024). [PubMed: [29170554](https://pubmed.ncbi.nlm.nih.gov/29170554/)].
- Zhang J, Li Y. Fibroblast growth factor 21, the endocrine FGF pathway and novel treatments for metabolic syndrome. *Drug Discov Today.* 2014;**19**(5):579–89. doi: [10.1016/j.drudis.2013.10.021](https://doi.org/10.1016/j.drudis.2013.10.021). [PubMed: [24189035](https://pubmed.ncbi.nlm.nih.gov/24189035/)].
- Woo YC, Xu A, Wang Y, Lam KS. Fibroblast growth factor 21 as an emerging metabolic regulator: Clinical perspectives. *Clin Endocrinol (Oxf).* 2013;**78**(4):489–96. doi: [10.1111/cen.12095](https://doi.org/10.1111/cen.12095). [PubMed: [23134073](https://pubmed.ncbi.nlm.nih.gov/23134073/)].
- Li X, Fan X, Ren F, Zhang Y, Shen C, Ren G, et al. Serum FGF21 levels are increased in newly diagnosed type 2 diabetes with nonalcoholic fatty liver disease and associated with hsCRP levels independently. *Diabetes Res Clin Pract.* 2011;**93**(1):10–6. doi: [10.1016/j.diabres.2011.02.034](https://doi.org/10.1016/j.diabres.2011.02.034). [PubMed: [21596453](https://pubmed.ncbi.nlm.nih.gov/21596453/)].
- Tucker B, Li H, Long X, Rye KA, Ong KL. Fibroblast growth factor 21 in non-alcoholic fatty liver disease. *Metabolism.* 2019;**101**:153994. doi: [10.1016/j.metabol.2019.153994](https://doi.org/10.1016/j.metabol.2019.153994). [PubMed: [31672443](https://pubmed.ncbi.nlm.nih.gov/31672443/)].
- Sanchez-Valle V, Chavez-Tapia NC, Uribe M, Mendez-Sanchez N. Role of oxidative stress and molecular changes in liver fibrosis: A review. *Curr Med Chem.* 2012;**19**(28):4850–60. doi: [10.2174/092986712803341520](https://doi.org/10.2174/092986712803341520). [PubMed: [22709007](https://pubmed.ncbi.nlm.nih.gov/22709007/)].