



Factors Influencing the Cerebroplacental Ratio in Non-Severe Small for Gestational Age Fetuses at the Age of 28 - 38 Weeks of Gestation

Mojgan Barati¹, Razeih Mohammad Jafarei¹, Sareh Aberoumand^{1,*}, Kobra Shojaei¹, Yalda Jefride¹, Sara Masihi¹ and Mahintaj Najafian¹

¹Fertility, Infertility and Perinatology Research Center, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran

*Corresponding author: Obstetrics and Gynecology Resident, Fertility, Infertility and Perinatology Research Center, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran. Email: saraaberumand@yahoo.com

Received 2019 January 13; Accepted 2019 June 23.

Abstract

Background: Cerebroplacental ratio (CPR) is an important indicator for predicting adverse outcomes during pregnancy. It is derived from the division of the PI Doppler index from the middle cerebral artery (MCA) into the umbilical cord vessels.

Objectives: To characterize the relationship between CPR and some factors such as diabetes, gestational hypertension, body mass index (BMI), PAPP-A, and drug use.

Methods: This descriptive-analytic study was conducted in Ahvaz in 2016. In this study, the CPR was evaluated in 230 pregnant women who were at 28 - 38 weeks of gestational age, except for pregnant women with severe Small for gestational age (SGA) and multiple pregnancies. In this study, a number of factors such as diabetes mellitus, gestational hypertension, BMI, PAPP-A, and drug use were investigated. Data were analyzed using SPSS software, version 19.

Results: In this study, there is a relationship between CPR and diabetes mellitus, gestational hypertension, and maternal drug use. There is not a relationship between CPR and BMI and PAPP-A.

Conclusions: Diabetes and gestational hypertension are two factors that influence the cerebroplacental ratio.

Keywords: CPR, Diabetes, SGA, Hypertension

1. Background

Increasing the health of mothers and children was one of the 10 major achievements in public health in the United States from 1900 to 1999 (1). In the United Kingdom and the United States, it is recommended that fetal Doppler should not be used as a screening tool for placental failure unless the fetus is known as small for gestational age (SGA) (2-4). Late placenta failure may occur in fetuses that are appropriate for gestational age (AGA). However, findings from hemodynamic changes in these fetuses indicate that the fetus suffers from fetal growth restriction (FGR); however, their weight remains higher than 10%. As a result, a small portion of the AGA fetuses in each percentile weight is at risk for stillbirth. Nevertheless, the prenatal diagnosis of IUGR by clinical examination is less than the optimal condition. Babies with birth weights less than 10 percentiles of weight for the gestational are called IUGR (5). Doppler ultrasonography is used to investigate the incidence of uterine growth retardation, fetal distress, asphyxia, anemia, pregnancy, twin to twin transfusion syndrome. Uterine

placental blood flow is detectable by Doppler ultrasonography, as a non-invasive method (6, 7). This sonography is very important in high-risk pregnancies. Ultrasound Doppler sonography completes the gray scale ultrasound. The blood vessels, blood flow, obstruction, and stenosis of the vessels, blood flow to organs, and also the dynamics of blood flow in organs can be evaluated for the diagnosis of physiological disorders by it. The aim of this study was to determine the relationship between the cerebroplacental ratio (CPR) in fetuses with normal weight percentile at 28 - 38 weeks of gestational age. So far, no study has been conducted to evaluate the statistical significance of uterine artery Doppler parameters in high-risk pregnancies, and most studies have studied some of the Doppler indices alone, such as PI (Pulsatility Index) or RI (Resistive Index) or early diastolic notching (8-11). One of the parameters for the evaluation of the middle cerebral artery is the following: Peak systolic velocity (PSV) (S), end-diastolic velocity (EDV) (D), and meantime span, systolic/diastolic ratio (S/D) RI, PI and comparison of arterial status of arteries (12, 13). CPR is an appropriate ratio as an important indi-

cator for predicting adverse outcomes during pregnancy, and this includes implications for the proper evaluation of the SGA and AGA fetuses. CPR is obtained by dividing the Doppler index from the middle cerebral artery (MCA) into the umbilical cord vessels. When these changes occur, diastolic flow increases from the MCA with S/D, RI, and PI (14-17). However, when calculating the CPR, it is chosen more often than the recently calculated PI (18, 19). Abnormal CPR may lead to three types of Doppler measurement patterns, including first, when UA PI and MCA PI are in the upper and lower distribution curve, respectively. Second, when UA PI is normal, but the MCA PI is reduced and the third, the UA PI is abnormally high and the MCA PI abnormally decreases and is therefore abnormal in all three CPR patterns (20-24).

2. Methods

This descriptive-analytic study was conducted in Ahvaz from January 2016 until January 2017. This research was approved by the department of obstetrics and gynecology of the Jundishapur, Ahvaz University of Medical Sciences and Ahvaz University Research Committee and Imam Khomeini Hospital. The sample consisted of 230 pregnant women referred to the Prenatal and Emergency Health Clinic of Imam Khomeini Hospital. CPR was evaluated in the pregnant women at 28 to 38 weeks of gestational age, except for women with multiple pregnancies and women who had severe SGA fetuses. The relationship between CPR and some maternal-fetal factors such as diabetes, gestational hypertension, body mass index (BMI), PAPP-A, and drug use were investigated. The ultrasound device was V10, V20 Medison. Data were put in Astraia software of obstetrics. All measurements were done in the prenatal clinic.

2.1. Ethical Considerations

The verbal consent for performing the research was obtained through the direct reference of professional assistant of Obstetrics and Gynecology to each patient. Every patient was informed that all tests were non-invasive and non-risky. Total information of the patients was encoded and preserved in the checklist and they were not available for real and legal persons. The study was approved by the Research Committee of Ahvaz University of Medical Sciences and ethically by the Ethics Committee of the University. Before starting the study, all participants were given informed consent.

3. Results

According to Table 1 among 230 pregnant women entered this study, 31 women had diabetes. Of these 31 women

with diabetes, 22 persons have normal CPR but, 9 of them have low CPR, which has significant analytic value. Of 230 pregnant women entered this study, 9 women had gestational hypertension, which all of them had Low CPR (Table 2).

Table 1. The Frequency of CPR Versus Maternal Diabetes in the Study Population (N = 230)

Maternal Diabetes	CPR Classification			Total
	> 5	2.5 - 5	< 2.5	
No	186	9	4	199
Yes	22	8	1	31
Total	208	17	5	230

Table 2. The Frequency of Gestational Hypertension in the Sample with CPR

CPR	Gestational Hypertension		Total
	Negative	Positive	
> 5	208	0	208
2.5 - 5	10	7	17
< 2.5	3	2	5
Total	221	9	230

Of total 7 women with diabetes, which used insulin, three women had significantly lower CPR. Of the total 4 women, which used Methyldopa all of them had low CPR. Other drugs had no influence on CPR (Table 3). According to Table 4, there was no significant relationship between PAPP-A and CPR in our study. There was no significant relationship between BMI and CPR in our study. According to the Figure 1, distribution of low CPR in different EFW percentiles, 3 - 10, 10 - 25, 25 - 50, 50 - 75, 75 - 95 were 14%, 13%, 10%, 6%, 6%, respectively, showing the presence of low CPR even in high EFW percentiles, but low CPR were more common in 3 - 10 EFW percentile.

4. Discussion

In this study, 230 women were studied and the cerebrospinal ratio (CPR) was measured. Diabetes and gestational hypertension are two factors that influence the cerebroplacental ratio.

Gibbons et al. (25) in their study examined the relationship between cerebrospinal CPR and the outcome of delivery and birth in complicated pregnancies, including insulin-dependent diabetes, non-insulin-dependent diabetes, and gestational diabetes, and concluded that regardless of the type of diabetes, there is a significant relationship between low CPR and diabetes. In another study con-




Figure 1. The distribution of CPR in different EFW percentiles is shown

ducted in 2016 by Gibbons et al., there was a correlation between CPR and gestational diabetes (26).

Studies have shown that diabetes can affect the outcome of pregnancy and childbirth. This effect includes fetal complications, such as macrosomia, hypoglycemia, and hypoglycemia in newborns. In a recent study, the effect of diabetes and gestational diabetes on CPR was studied. The results of this study showed that there is a relationship between blood flow to the brain in the fetus and the mother's diabetes. Gibbons et al. (25) generally examined diabetes and found it related to CPR, the results of this study are consistent with diabetes. On the contrary, the results of this

study did not match the results of Gibbons et al. (26).

Gaikwad et al. (27) showed that there was a relationship between CPR and GHTN. Shahinaj et al. (28) concluded in their research that the CPR was a predictor of the adverse outcomes in preeclampsia and GHTN. Parshuram et al. in 2014 showed that CPR is very useful for prenatal monitoring of mothers with gestational hypertension (29).

4.1. Conclusions

The results of this study indicated a relationship between CPR and maternal diabetes, GHTN and drug use. However, there was no relationship between CPR and PAPP-

Table 3. The Frequency of CPR Versus Drug Use

Drug Use	CPR Classification			Total
	> 5	2.5 - 5	< 2.5	
No use	183	10	3	196
ASA	1	1	0	2
Aspirin	1	0	0	1
Insulin	4	3	0	7
Levothyroxine	15	0	1	16
Prednisolone	1	0	0	1
Metformin	1	0	0	1
Carbamazepine	1	0	0	1
Clonazepam	1	0	0	1
Methyl dopa	0	3	1	4
Total	208	17	5	230

Table 4. The Frequency of CPR Versus PAPP-A Values

CPR	PAPP-A			Total
	< 0.4	0.4 - 2.6	> 2.6	
> 5	13	190	5	208
2.5 - 5	4	12	1	17
< 2.5	0	5	0	5
Total	17	207	6	230

Table 5. Chi-Square Test for Analyzing Research Hypotheses

Items	P Value (2-Sided)
Maternal diabetes	0.00
Maternal hypertension	0.642
Taking medication	0.00

A and BMI. To conduct multicenter studies for study generalization, further studies with a larger sample size to increase the validity of the results, and further studies on the association of CPR with PI in the uterine artery are recommended. It is hoped that more comprehensive results could be presented via further studies with a larger sample size.

4.2. Limitations

This study has some limitations that are worth mentioning. Unfortunately, the duration of study was relatively short; therefore, we suggest that for future studies other factors such as neonatal-maternal complications etc. should be investigated. We also suggest that a study with the same purpose should be conducted with a greater sample size in patients with diabetes and hypertension.

Acknowledgments

This paper is extracted from the thesis of Dr. Sara Abroumand. Thesis was supported by the Ahvaz Jundishapur University of Medical Science (AJUMS), Ahvaz, Iran. Thereby, we wish to express our profound gratitude to study participants.

Footnotes

Authors' Contribution: All authors contributed equally in planning and carrying out of this study.

Conflict of Interests: There is no conflict of interest in this article.

Ethical Approval: The study was approved scientifically by Research Committee of Ahvaz University of Medical Sciences and ethically by the Ethic Committee of the university before starting participant recruitment.

Funding/Support: This study was funded and supported by Ahvaz Jundishapur University of Medical Sciences.

Patient Consent: All participants gave informed consent.

References

- Cunningham FG. *Williams obstetrics*. 22nd ed. New York: Mc Grow-Hill; 2005.
- Partodel M. *Summery obstetrics and gynecology*. Tehran: Yad; 1994. p. 14-5.
- American College of Obstetricians and Gynecologists. ACOG practice bulletin. Intrauterine growth restriction. *Int J Gynecol Obstet*. 2000;**72**(2001):85-96. doi: [10.1016/S0020-7292\(00\)90000-6](https://doi.org/10.1016/S0020-7292(00)90000-6).
- Royal College of Obstetricians and Gynaecologists. *The investigation and management of the small-for-gestational-age fetus*. London: Green-top Guideline No. 31; 2002.
- Morales-Rosello J, Khalil A. Fetal cerebral redistribution: a marker of compromise regardless of fetal size. *Ultrasound Obstet Gynecol*. 2015;**46**(4):385-8. doi: [10.1002/uog.15664](https://doi.org/10.1002/uog.15664). [PubMed: [26224656](https://pubmed.ncbi.nlm.nih.gov/26224656/)].
- Kurmanavicius J, Florio I, Wisser J, Hebisch G, Zimmermann R, Muller R, et al. Reference resistance indices of the umbilical, fetal middle cerebral and uterine arteries at 24-42 weeks of gestation. *Ultrasound Obstet Gynecol*. 1997;**10**(2):112-20. doi: [10.1046/j.1469-0705.1997.10020112.x](https://doi.org/10.1046/j.1469-0705.1997.10020112.x). [PubMed: [9286020](https://pubmed.ncbi.nlm.nih.gov/9286020/)].
- Pijnenborg R, Dixon G, Robertson WB, Brosens I. Trophoblastic invasion of human decidua from 8 to 18 weeks of pregnancy. *Placenta*. 1980;**1**(1):3-19. [PubMed: [7443635](https://pubmed.ncbi.nlm.nih.gov/7443635/)].
- Deurloo KL, Spreuwenberg MD, Bolte AC, Van Vugt JM. Color Doppler ultrasound of spiral artery blood flow for prediction of hypertensive disorders and intra uterine growth restriction: A longitudinal study. *Prenat Diagn*. 2007;**27**(11):1011-6. doi: [10.1002/pd.1822](https://doi.org/10.1002/pd.1822). [PubMed: [17721908](https://pubmed.ncbi.nlm.nih.gov/17721908/)].
- Fratelli N, Rampello S, Guala M, Platto C, Frusca T. Transabdominal uterine artery Doppler between 11 and 14 weeks of gestation for the prediction of outcome in high-risk pregnancies. *J Matern Fetal Neonatal Med*. 2008;**21**(6):403-6. doi: [10.1080/14767050802053073](https://doi.org/10.1080/14767050802053073). [PubMed: [18570118](https://pubmed.ncbi.nlm.nih.gov/18570118/)].

10. Meler E, Figueras F, Mula R, Crispi F, Benassar M, Gomez O, et al. Prognostic role of uterine artery Doppler in patients with preeclampsia. *Fetal Diagn Ther.* 2010;**27**(1):8–13. doi: [10.1159/000258048](https://doi.org/10.1159/000258048). [PubMed: [19907134](https://pubmed.ncbi.nlm.nih.gov/19907134/)].
11. Anastasakis E, Papantoniou N, Daskalakis G, Mesogitis S, Antsaklis A. Screening for pre-eclampsia by oxidative stress markers and uteroplacental blood flow. *J Obstet Gynaecol.* 2008;**28**(3):285–9. doi: [10.1080/01443610802042852](https://doi.org/10.1080/01443610802042852). [PubMed: [18569469](https://pubmed.ncbi.nlm.nih.gov/18569469/)].
12. Cunningham G, Leveno K, Bloom S, Hauth JC. Chronic hypertension. In: Cunningham G, editor. *Williams obstetrics*. 22nd ed. USA, Philadelphia: McGraw-Hill; 2005. 1044 p.
13. Yoshimura S, Masuzaki H, Gotoh H, Ishimaru T. [The relationship between blood flow redistribution in umbilical artery and middle cerebral artery and fetal growth in intrauterine growth retardation]. *Nihon Sanka Fujinka Gakkai Zasshi.* 1995;**47**(12):1352–8. Japanese. [PubMed: [8568354](https://pubmed.ncbi.nlm.nih.gov/8568354/)].
14. Baschat AA, Gembruch U. The cerebroplacental Doppler ratio revisited. *Ultrasound Obstet Gynecol.* 2003;**21**(2):124–7. doi: [10.1002/uog.20](https://doi.org/10.1002/uog.20). [PubMed: [12601831](https://pubmed.ncbi.nlm.nih.gov/12601831/)].
15. Odibo AO, Riddick C, Pare E, Stamilio DM, Macones GA. Cerebroplacental Doppler ratio and adverse perinatal outcomes in intrauterine growth restriction: Evaluating the impact of using gestational age-specific reference values. *J Ultrasound Med.* 2005;**24**(9):1223–8. [PubMed: [16123182](https://pubmed.ncbi.nlm.nih.gov/16123182/)].
16. Ebbing C, Rasmussen S, Kiserud T. Middle cerebral artery blood flow velocities and pulsatility index and the cerebroplacental pulsatility ratio: Longitudinal reference ranges and terms for serial measurements. *Ultrasound Obstet Gynecol.* 2007;**30**(3):287–96. doi: [10.1002/uog.4088](https://doi.org/10.1002/uog.4088). [PubMed: [17721916](https://pubmed.ncbi.nlm.nih.gov/17721916/)].
17. Morales-Rosello J, Khalil A, Morlando M, Papageorgiou A, Bhide A, Thilaganathan B. Changes in fetal Doppler indices as a marker of failure to reach growth potential at term. *Ultrasound Obstet Gynecol.* 2014;**43**(3):303–10. doi: [10.1002/uog.13319](https://doi.org/10.1002/uog.13319). [PubMed: [24488879](https://pubmed.ncbi.nlm.nih.gov/24488879/)].
18. Williams KP, Farquharson DF, Bebbington M, Dansereau J, Galerneau F, Wilson RD, et al. Screening for fetal well-being in a high-risk pregnant population comparing the nonstress test with umbilical artery Doppler velocimetry: A randomized controlled clinical trial. *Am J Obstet Gynecol.* 2003;**188**(5):1366–71. doi: [10.1067/mob.2003.305](https://doi.org/10.1067/mob.2003.305). [PubMed: [12748513](https://pubmed.ncbi.nlm.nih.gov/12748513/)].
19. Mari G, Hanif F. Fetal Doppler: Umbilical artery, middle cerebral artery, and venous system. *Semin Perinatol.* 2008;**32**(4):253–7. doi: [10.1053/j.semperi.2008.04.007](https://doi.org/10.1053/j.semperi.2008.04.007). [PubMed: [18652923](https://pubmed.ncbi.nlm.nih.gov/18652923/)].
20. Jugovic D, Tumbri J, Medic M, Jukic MK, Kurjak A, Arbeille P, et al. New Doppler index for prediction of perinatal brain damage in growth-restricted and hypoxic fetuses. *Ultrasound Obstet Gynecol.* 2007;**30**(3):303–11. doi: [10.1002/uog.4094](https://doi.org/10.1002/uog.4094). [PubMed: [17721870](https://pubmed.ncbi.nlm.nih.gov/17721870/)].
21. Mula R, Savchev S, Parra M, Arranz A, Botet F, Costas-Moragas C, et al. Increased fetal brain perfusion and neonatal neurobehavioral performance in normally grown fetuses. *Fetal Diagn Ther.* 2013;**33**(3):182–8. doi: [10.1159/000350699](https://doi.org/10.1159/000350699). [PubMed: [23594501](https://pubmed.ncbi.nlm.nih.gov/23594501/)].
22. Roza SJ, Steegers EA, Verburg BO, Jaddoe VW, Moll HA, Hofman A, et al. What is spared by fetal brain-sparing? Fetal circulatory redistribution and behavioral problems in the general population. *Am J Epidemiol.* 2008;**168**(10):1145–52. doi: [10.1093/aje/kwn233](https://doi.org/10.1093/aje/kwn233). [PubMed: [18826969](https://pubmed.ncbi.nlm.nih.gov/18826969/)].
23. Khalil AA, Morales-Rosello J, Elsaddig M, Khan N, Papageorgiou A, Bhide A, et al. The association between fetal Doppler and admission to neonatal unit at term. *Am J Obstet Gynecol.* 2015;**213**(1):57 e1–7. doi: [10.1016/j.ajog.2014.10.013](https://doi.org/10.1016/j.ajog.2014.10.013). [PubMed: [25447961](https://pubmed.ncbi.nlm.nih.gov/25447961/)].
24. Khalil AA, Morales-Rosello J, Morlando M, Hannan H, Bhide A, Papageorgiou A, et al. Is fetal cerebroplacental ratio an independent predictor of intrapartum fetal compromise and neonatal unit admission? *Am J Obstet Gynecol.* 2015;**213**(1):54 e1–54 e10. doi: [10.1016/j.ajog.2014.10.024](https://doi.org/10.1016/j.ajog.2014.10.024). [PubMed: [25446667](https://pubmed.ncbi.nlm.nih.gov/25446667/)].
25. Gibbons A, Flatley C, Kumar S. The fetal cerebro-placental ratio in diabetic pregnancies is influenced more by the umbilical artery rather than middle cerebral artery pulsatility index. *Eur J Obstet Gynecol Reprod Biol.* 2017;**211**:56–61. doi: [10.1016/j.ejogrb.2017.02.001](https://doi.org/10.1016/j.ejogrb.2017.02.001). [PubMed: [28189724](https://pubmed.ncbi.nlm.nih.gov/28189724/)].
26. Gibbons A, Flatley C, Kumar S. Cerebroplacental ratio in pregnancies complicated by gestational diabetes mellitus. *Ultrasound Obstet Gynecol.* 2017;**50**(2):200–6. doi: [10.1002/uog.17242](https://doi.org/10.1002/uog.17242).
27. Gaikwad PR, Gandhewar MR, Rose N, Suryakar V. Significance of obstetric Doppler studies in prediction of perinatal outcome in pregnancy induced hypertension. *Int J Reprod Contraception Obstetr Gynecol.* 2017;**6**(6):2354. doi: [10.18203/2320-1770.ijrcog20172312](https://doi.org/10.18203/2320-1770.ijrcog20172312).
28. Shahinaj R, Manoku N, Kroj E, Tasha I. The value of the middle cerebral to umbilical artery Doppler ratio in the prediction of neonatal outcome in patient with preeclampsia and gestational hypertension. *J Prenat Med.* 2010;**4**(2):17–21. [PubMed: [22439055](https://pubmed.ncbi.nlm.nih.gov/22439055/)]. [PubMed Central: [PMC3279170](https://pubmed.ncbi.nlm.nih.gov/PMC3279170/)].
29. Parshuram PL, Mwangi G, Wambugu M, Ong'ech J, Mwachaka P. The cerebro-placental ratio as a prognostic factor of foetal outcome in patients with third trimester hypertension. *East Cent Afr J Surg.* 2014;**19**(1):41–51.