Research Article



Balance Changes Following a Controlled Whole-Body Vibration Training: A Pilot Analysis Addressing Balance Control in Adult Women with Obesity

Manijeh Soleimanifar¹, Masoumeh Abedi², Mostafa Sarabzadeh ib^{3,*}, Afsaneh Dadarkhah⁴, Zahra Sadat Asadi ib⁵

¹Assistant Professor, Department of Lifestyle Medicine , Medical Laser Research Center, ACECR, Tehran, Iran

² School of Health and Rehabilitation Science, University of Queensland, Brisbane, Australia

³ Research Associate in Exercise Physiology & NeuroPhysiotherapy, National Elites Foundation, Tehran, Iran

⁴ Instructor of Physiotherapy, School of Medicine, Research Center of Clinical Biomechanics and Ergonomics, Imam Reza Hospital, AJA University of Medical Sciences, Tehran, Iran

⁵ Department of Community Medicine, School of Medicine, AJA University of Medical Sciences, Tehran, Iran

^{*}Corresponding author: Research Associate in Exercise Physiology & NeuroPhysiotherapy, National Elites Foundation, Tehran, Iran. Email: m.sarab68@yahoo.com

Received 2023 December 29; Revised 2024 March 14; Accepted 2024 April 2.

Abstract

Background: The prevalence of obesity among older individuals has significantly increased in recent years, potentially doubling the risk of postural imbalance and falls. Controlled whole-body vibration (CWBV) training is considered an alternative/complementary therapy for fall prevention, but its mechanisms still require further investigation.

Objectives: This study aimed to investigate the effects of CWBV on balance and fall risk among older adult women with obesity. **Methods:** Fifteen adult women over the age of 60 (mean age 68.55 \pm 4.86, BMI: 30.57 \pm 2.97) participated in this quasiexperimental study. They underwent CWBV training three times weekly for six weeks. Gait stability and balance were assessed using a gait analysis system before and after the intervention.

Results: The results indicated a significant decrease in both static and dynamic balance parameters, except for the variability of velocity index (P = 0.9) and fear of falling (FES-I score, P = 0.56), after the intervention.

Conclusions: Controlled whole-body vibration training appears to be an effective method for improving static and dynamic balance parameters in older adult women. However, further research is necessary to explore different dosages and protocols of vibration training, incorporating more challenging/specialized proprioceptive exercises to potentially improve the fear of falling in this population.

Keywords: Vibration, Obesity, Postural Balance, Gait Analysis, Frail Elderly

1. Background

The aging community is growing worldwide. Population growth has increased geometrically, rising by 5.2% from 2009 to 2015, resulting in 40,000 middleaged and 250,000 elderly individuals (1). Additionally, the prevalence of obesity in these age categories increased from 16% to 18%, with women at a higher risk than men across all continents, both in developed and developing countries (2). Alarmingly, a recent study revealed a decline in public awareness and healthseeking behaviors regarding obesity from 2004 to 2021 (3). If this trend continues, a majority of the world's adult population will be either overweight or obese by 2030 (4). However, these figures, derived from BMI-based estimates, are not highly sensitive tests for determining obesity (5), and the true prevalence is likely much higher than what these tests regularly report (6). Given the significant underestimates of obesity prevalence and its association with multiple comorbidities (7), as well as being a gateway to many non-communicable diseases (8) leading to a high risk of hospitalizations and intubation (9), different preventive and therapeutic measures to assist the daily living

Copyright © 2024, Jundishapur Journal of Health Sciences. This open-access article is available under the Creative Commons Attribution-NonCommercial 4.0 (CC BY-NC 4.0) International License (https://creativecommons.org/licenses/by-nc/4.0/), which allows for the copying and redistribution of the material only for noncommercial purposes, provided that the original work is properly cited.

activities of the elderly, most directly impacted by obesity, would be beneficial. This need is more pressing than ever, especially in light of the world's new COVID-19-related challenges and other global crises predicted in the coming years (10). Furthermore, the role of obesity as a risk factor has been emphasized by data collected across all ethnicities during the current SARS-CoV-2 pandemic (11, 12).

The impact of obesity on falls has not been thoroughly examined. Rosenblatt and Madigan and Choi et al. identified a strong positive correlation between fat mass index and trip-induced fall risk among older adults (13, 14). This finding is recently supported by Thiamwong et al., who demonstrated that body fat mass (BFM), percent body fat (PBF), and Body Mass Index (BMI) are correlated with fear of falling (FoF) and balance impairment (15). Furthermore, a distinct pattern of force application during walking has been observed between obese individuals and weight-matched control groups (14), suggesting that obesity may contribute to fall risk, although it is not the sole factor (16). Other independent factors such as disability, history of falls, hand grip strength (HGS), number of medications used, and pain have also been shown to influence FoF (17). The complexity of these factors, combined with the use of different stability measures, may explain why some studies have found no effect of obesity on gait stability in this population (18). Greater muscle weakness in the lower limbs and 34% balance disorders are common in older adults, which may be exacerbated by a variety of biological and social factors, including physical disability, chronic illnesses, incontinence, inactivity, and obesity (19). Regarding the role of gender, both younger women under 50 years of age (20, 21) and older women aged 65 to 75 years (22) exhibit lower stability limits, making them more susceptible to falls. However, irrespective of gender, obesity alone leads to altered postural control, associated with increased activity of plantar flexion. This underscores the importance of focusing on proprioception and neuromuscular control in fall prevention rehabilitation protocols (23). This is why individuals with obesity are often twice as likely to fall as those who are not obese (24).

In recent years, there has been growing interest in using controlled whole-body vibration (CWBV) training not only as an alternative rehabilitation method for fall prevention but also across various fields exploring a wide range of demand-based applications. These include improving overall fatigue (25) and athletic performance (26), and adopting recovery-oriented approaches for the rehabilitation of neurological (27), musculoskeletal (28), and metabolic conditions by inducing biological responses that enhance functional parameters in these patients (29). However, despite promising results regarding the efficacy of WBV or CWBV combined with exercise in improving balance and proprioception, a recent systematic review indicated that no definitive conclusions could be drawn due to the heterogeneity of articles, duration of exposure, parameters characterizing mechanical vibration, and the lack of clinical trials exploring different frequencies and amplitudes (30). Therefore, identifying these contributing factors through various assessment models should be among the first steps toward developing appropriate intervention strategies to reduce the fall rate.

The neuromuscular adaptations acquired through CWBV appear to be an important mechanism for improving basic living skills in older adults by enhancing proprioceptive sense. As a complementary therapy, this technique, alongside unstable shoes, could be proposed as a beneficial intervention with relatively long-term effects on balance measures in older people (31). A recent study concluded that this model may effectively improve neuromuscular performance and balance in healthy individuals (28), as well as in patients using CWBV for neurological recovery. Chang et al. reported altered neuromuscular control in patients with Parkinson's disease through a pilot analysis (27), Ruhde and Hulla observed improvements in physical function in individuals with cerebral palsy (CP) (32), Li et al. described this technique as a passive and safe clinical intervention as an alternative treatment for individuals with motor impairments or poor balance function (33), and Guadarrama-Molina et al. recently proposed this therapy as a co-adjuvant to conventional therapy for improving functional balance status (34). However, some studies report the opposite, with no motor/balance symptom-reducing effect of CWBV (35), and the reasons behind this should not be underestimated. Implementation also contributes to these discrepancies, such as the different stances on a force plate—semi-squat position or deep-squat position (36)—and performing dynamic exercises, such as squats or leg lunges, while on the plate. The frequencies of vibrations used in studies ranged from 10 to 50 Hz, with amplitudes from 1 to 5 mm, and some studies progressively increased vibration frequencies and amplitudes from session to session; however, using a 20 Hz frequency and 2 mm amplitude applied in 30 to 60second intervals (36) and a 30 or 60-second interval (37) has been reported to be most beneficial in improving balance, supported by an earlier study that reported improved functional gait efficiency, dynamic balance, functional strength of lower body muscles, and exercise

tolerance in women over 60 at increased risk of falling (38).

The complexity and multifactorial nature of falls, along with challenges in measuring proprioception in older adults with obesity, have led to conflicting findings and hindered our understanding of the neurophysiological mechanisms of vibration. This understanding is key to prescribing a safe and effective CWBV protocol aimed at reducing the aging effects on the musculoskeletal structures of obese individuals prone to falling. To date, studies investigating the health-enhancing effects of CWBV-either alone or in conjunction with other therapies—specifically in obese individuals at risk of falling, have not reached a consensus to ascertain its potential impact on balance parameters and home-related falls to aid these individuals in maintaining the physical activity they need. Thus, this should also be considered in addition to addressing the obesity-associated metabolic parameters that conventional therapeutic protocols have focused on.

2. Objectives

To advance our understanding, as prompted by other researchers' recent endeavors, this pilot study aims to investigate the effects of CWBV on balance parameters and fall risk among obese adult women. The hypotheses are as follows: First, the therapeutic effect of CWBV on balance parameters and FoF is significantly different from that of conventional therapies routinely used in older adult women with obesity. This is because CWBV offers a more versatile and patient-centered rehabilitation protocol, meeting users' diverse physical fitness needs and specific requirements through adjustable vibration setups and tailored degrees of muscle activation. Secondly, CWBV training may provide a quicker additional modality to enhance the active lifestyles of this population by improving stability and addressing personal concerns about falling.

3. Methods

3.1. Participants

Fifteen adult women with obesity, who are employees at a university's biomechanical research center and aged over 60 (68.55 ± 4.86) with a BMI of 30.57 ± 2.97 , were voluntarily recruited and participated in this study. The study used a one-group pretest-posttest design (Figure 1 and Table 1). The exercise and measurement protocols were conducted in accordance with the Declaration of Helsinki-Tokyo and approved by the local ethics committee.



After obtaining the necessary permissions for research, candidates completed a medical history form and a consent letter. BMI, a reliable indicator of body fatness with high sensitivity and specificity for individuals over 60 (29), was used to assess obesity and to include only those with a BMI over 30. Our inclusion criteria were as follows: (a) Having normal vision, hearing, and dexterity; (b) no personal history of cancer, diabetes, mental or neurological disorders, or other significant diseases that could bias the results or contraindicate vibration exercise; (c) not having taken any medications that could affect balance or movement smoothness (either increasing or reducing movement speed, whether voluntary or involuntary) in the last six months; and (d) having physical and cognitive abilities confirmed through medical records and being

interested in participating in the intervention program. Participants who did not meet these criteria were excluded from the study. A table of random numbers was used to create our final sample. Initially, each participant was assigned a number between 1 and 16. Then, a team member blindfolded herself and randomly pointed to a spot on the chart containing three-digit numbers, selecting the first two digits one time and the first digit another time to match one of our subjects until we identified our final fifteen participants for the intervention.

3.2. Intervention

After identifying participants who met the inclusion criteria, they first received a detailed explanation of how to use the device effectively while adhering to all safety precautions during a session of familiarization trials. Subsequently, all participants were assigned to CWBV training three times weekly for six weeks under the supervision of an experienced physiotherapist and exercise physiologist. Each session lasted on average 10 minutes, with participants standing with knees flexed at 20 degrees and their trunk in an upright vertical position on the marked footprint over the vibration platform (Extream1000; AMH International Co., Ltd., Incheon, Republic of Korea). They underwent five cycles of one-minute exposure to the vibration training protocol with a one-minute rest interval between cycles and a vibration frequency set at a maximum of 30 Hz (Figure 2). The vibration intensity was progressively increased by adjusting the frequency up to 30 Hz and the amplitude up to 1 - 2 mm(39, 40).



Figure 2. Participant's position during the controlled whole-body vibration (CWBV) experiment

3.3. Measurement

Recent studies have identified practical functional balance measures with the highest levels of ability to indicate the risk of falls among older adults, boasting a sensitivity of 92.68%, specificity of 84.09%, and an AUC of 0.931 (95% CI = 0.860 to 1.000) (41). These tests, suggested for roughly gauging individual functional levels, are simple, feasible, and practical tools that can be administered to larger populations and conducted almost anywhere. Therefore, static and dynamic balance parameters were selected for evaluation before starting CWBV training and again six weeks afterward, using the Zebris FDM system (ZEBRIS Medical GmbH, Germany)-a portable force platform that provides access to spatiotemporal gait parameters with high accuracy. The reliability of this system has been reported as acceptable to excellent in balance assessments by Van Alsenoy et al. (42).

Static balance was assessed over a 20-second period with subjects standing barefoot on the platform, feet shoulder-width apart. Dynamic balance was evaluated through a walking test where subjects walked barefoot on a 2-meter platform at their comfortable and safe speed (Figure 3).

The analysis of the force's application point between the feet and the ground was based on Romberg's test, assessing static balance during free-standing on both lower limbs with eyes alternately closed and open. Ellipse sway area (ESA) (COP displacement) and path length (the distance the COP covers) were selected to determine static balance. For dynamic balance, step width (the distance between the line of progression of the left foot and the line of progression of the right foot measured in cm), total double support (duration of the gait cycle phase when both lower limbs are in contact with the treadmill belt, recorded and expressed in percent), stride length (data from overground walking gait used to calculate stride length), stride time (duration of foot displacement between two consecutive steps on the same side, recorded and expressed in ms), cadence (number of steps per minute -sum of both feet in this study), and velocity (time taken by each subject to traverse the entire distance, measured in m/s), were all monitored according to their standard definitions and norms (43) to assess dynamic stability. The mean values of three repetitive trials for each pre and post-test were taken as the test outcomes. All data were then recorded and stored on a laptop. Testretest reliability and validity of these center of pressurebased parameters were explored and found to be acceptable for assessing balance in older adults, both



standing with eyes opened or closed, and also in seated positions (ICC = 0.99)(44).

To assess FoF, the Fall Efficacy Scale-International (FES-I) was used. The country-specific standardized version of FES-I has been found acceptable to assess FoF among older adults (45), as well as the international version (46, 47). In this test, FoF is operationalized as the level of concern about falling over 16 different physical activities. It has a four-point scale ranging from 1 (not concerned) to 4 (very concerned), resulting in a total sum-score between 16 and 64— the higher each parameter score, the higher the levels of concern/FoF.

3.4. Statistical Analysis

We first estimated the number of participants using G*Power (Version 3.1.9.4.) (48), required for a significant group by time (pre, post) interaction. This test revealed that enrolling 15 patients for the intervention phase would produce a medium effect of 0.5 (alpha: 0.05, power: 0.82). The results of the Kolmogorov–Smirnov test did not confirm normally distributed data in all variables (P-value \leq 0.05), and the Wilcoxon signed-rank test, as a non-parametric paired t-test, was used

accordingly to examine pre-post-treatment changes. Meanwhile, statistical analysis was carried out using SPSS (IBM, USA, version 17.0) with a 95% confidence interval.

4. Results

According to Table 2, a significant difference was observed in all parameters of both static and dynamic balance, including Ellipse Area (P = 0.008, ES: 1.82) and path length (P = 0.013, ES: 2.66), step width (P = 0.03, ES: 0.78), total double support (P = 0.021, ES: 0.73), stride length (P = 0.022, ES: 0.76), stride time (P = 0.003, ES: 1.18), cadence (P = 0.003, ES: 1.06), and velocity (P < 0.001, ES: 1.06), except for the variability of velocity (P = 0.9) in the pretest-posttest score comparison. The test result of the FES-I questionnaire (P = 0.56) also found no statistically significant difference in FoF following the intervention. Meanwhile, the formula Meanpost -Meanpre / Sdpre was applied for this one-group pre-post design to assess the effect size of each parameter. According to the study of Mordkoff, the test of all variables is interpreted as having a large effect size-0.01, 0.06, and 0.14 indicate small effect, medium effect, and large effect, respectively (49, 50).

Table 2. Descriptive Data and Wilcoxon Signed-Rank Test for Parameters of Static and Dynamic Balance, and FES-I Test Before and After Treatment ^a

Stability Parameters	Condition		ny t
	Pre-intervention	Post-intervention	P-value ²
Static balance			
Ellipse Area1 (mm ²)	99.04 (3.906)	91.90 (3.812)	0.008
Path length2 (mm)	147.92 (2.887)	140.24 (3.109)	0.013
Dynamic balance			
Step width	13.00 (2.8)	10.81 (2.48)	0.03
Total Double Support	36 (3.8)	33.2 (4.7)	0.021
Stride length	88.8 (10.4)	95.18 (11.01)	0.022
Stride time	1.28 (0.11)	1.15 (0.08)	0.003
Cadence	47.09 (4.78)	52.18 (3.97)	0.003
Velocity	2.56 (0.48)	3.07(0.52)	< 0.001
Variability of Velocity	6.54 (1.50)	6.54 (2.16)	0.9
FES-I3 score (range 1 to 64)	25.55 (4.19)	24.3 (5.1)	0.56
^a Values are expressed as mean (SD).			

values are expressed as mean (3D).

^b P-values are significant at P < 0.05.

5. Discussion

To begin with an overview of the findings, the improvement of static and dynamic balance reported among adult women with obesity following our CWBV training is clearly the main scientific discovery of the present study. However, this technique did not support our pre-study hypothesis regarding the efficiency of CWBV intervention on FoF in obese fallers.

Whole-body vibration has recently been found to be a potential tool in the rehabilitation of the elderly and patients with chronic diseases. However, few studies have been performed concerning whole-body vibration in these individuals who exhibit poor balance performing ability and to investigate its potential impact on proprioception to clarify the connection between proprioceptive function and improved motor function. Among all the well-documented obesity comorbidities, balance impairment has always been one of the most ignored risk factors that had to be addressed as equal to metabolic comorbidities in healthcare and rehabilitation programs for older adults with obesity. Although metabolic healthy obesity (MHO) shows reduced cardiometabolic risks and MHO individuals are closer to normal than those with unhealthy phenotypes (51), it could not vet be considered a safe condition (52). and a modest weight loss (5 - 10% of body weight) has been proven to alleviate their cardiovascular risk (53), recently supported on a broader scale by Haase et al. (54). Increasing ambulatory activity might also be one of the other clues to solving the increasing prevalence of obesity and its associated comorbidities (55). Therefore,

the present study is one of the first studies aimed to provide evidence in support of an additional effect of CWBV training on stability parameters while verifying the efficiency of its controlled exposure on the risk of fall and on the active life of our older adult women with obesity (20).

What this study found may coincide with the results of studies that reveal improvements in balance and postural control following CWBV training. Onal et al. reported significant improvements in static and dynamic balance among stroke patients but not obese individuals (56). Melo et al., through a recent systematic review, found positive results when vibration-with frequency values of 30 to 35 Hz as set in our study's protocol-was used alone or in association with other types of exercises in older adults with obesity (57). Maghbouli et al. reported that CWBV training using a vibration frequency higher than 100 Hz may improve muscle strength and open-eye mediolateral postural control, recommending this technique as a promising approach for decreasing fall risk (58). Additionally, several recent studies found this modality to be an effective stimulus to improve neuromuscular performance and balance in healthy individuals (26), as well as in elderly individuals with neurological disorders (27, 32).

Looking at these previous studies, it has been shown that most of the physical activity-induced health benefits have been acquired from the simplest, most fun, and entertaining programs, which provide a sufficient training stimulus that older adults are able to comply with and complete eagerly due to age-related physical debilities. This is partly why CWBV training seems capable of delivering the same benefits seen with conventional exercise programs, but in a simpler format that allows older adults to be mentally and physically engaged on a routine basis. However, a few studies that failed to show any positive effect of this technique (35, 59) highlight the importance of setting up an optimal amplitude and frequency of the vibration, coupled with an optimal level of muscle activity by which the vibration stimulation could be superimposed.

For instance, the study conducted by Miller et al. shows that exposure to continuous whole-body vibration may reach its peak application with the participant on the platform, allowing fatigue to set in, speculating greater change post-CWBV possibly due to the balance of potentiation and fatigue, which is missed in our vibration protocol and other intermittent wholebody vibrations potentiate performance without fatigue accumulation (59). In another study, fixed-frequency and amplitude vibration training did not result in any meaningful changes in any of the balance variables (60), where the exact mechanism behind this failure compared to stability improvements concluded by the protocol designed an individualized vibration frequency and amplitude according to their EMG data, may be attributed to the provision of stronger stimulation to skin receptors, muscle spindles, and the vestibular system. Other than that, most of the studies have focused on the leg extensors, neglecting plantar flexors, which have been shown to be up to five times more effective in increasing electromyographic activity with vibration. All these factors may explain in part why our CWBV protocol, despite the balance improvements, was not sufficient to impact the fall efficacy scale, which reflects real-life functional limitation-related concerns.

In the case of older adults with obesity specifically, FoF and its relationship with activity have not been sufficiently investigated. Rosic et al. appear to be the most recent study suggesting that FoF is associated with low levels of physical activity (20). However, no links have been found between FoF and obesity (61), except for some conflicting findings demonstrated contrariwise (62-64). Perhaps a new somato-sensory stimulation (SSS) type of exercise would improve these fall-related psychological concerns through triggering brain plasticity (65), or even historical derivatives of Chinese healing practices based on mind-body interaction protocols, as recently claimed by Nissim et al. (66), would be a better option to enhance cerebellar activation, which in turn might improve balance. This is because, compared to voluntary muscle control during conventional exercise training, these techniques rely more on contractions induced by passive stretch reflexes, leading to facilitating the excitability of the spinal reflex. After reaching this level of physiological and neuro-muscular adaptations (67), our older adults with obesity would end up with a more adaptable body, making its control easier to perform smoother movements (68)—as compensation for all possible agerelated physical debilities and inactivities they are most likely exposed to. Therefore, despite the lack of significance of FoF, our results imply clinical relevance particularly in individuals showing contraindications toward conventional exercise therapies, considering and addressing the limitations of the present study listed as follows.

5.1. Limitations

The present study has some limitations that need to be discussed and considered thoroughly by future works. First, the sample size was small, as reasons such as not exceeding the budget and not violating our intraorganizational rules and policies (not recruiting subjects outside of the department) hindered us from enrolling a higher number of participants. Secondly, there is not yet standardization of a vibration protocol for obese individuals in the literature. Therefore, we tried to apply the most commonly used protocol with the vibration setup designed by evidence-focused databases, but not with individualized vibration thatbased on others' up-to-date findings-may present stronger stimulation to the body than the fixedfrequency and amplitude vibration training being used in the present study. Thirdly, daily activity, work, smoking, physical activity, and daily energy intake could all be disturbing variables which were not controlled and may have affected the present findings. Fourthly, this study was limited to the short-term effect of CWBV training for obese individuals and the findings are not generalizable beyond this scope; thus, the long-lasting effects of this technique need to be investigated. Fifthly, this study was also restricted to older adult women as they have been shown to have higher levels of FoF in the elderly than men. Lastly, this is an exploratory study conducted without a control group, which reflects back to our criteria and the number of patients that fit it when undertaking their routine treatment in the physiotherapy department.

5.2. Conclusions

We consider CWBV training to be an effective method for improving static and dynamic balance parameters in older adult women. However, further research is warranted to explore different dosages of vibration protocols or to incorporate more challenging/specialized proprioceptive exercises into CWBV. This could help target the improvement of FoF in those who are unable to participate in other forms of conventional exercise programs for rehabilitation. Additionally, it is important for further studies, in addition to addressing the aforementioned limitations of the present study, to explore the neurophysiological mechanisms involved in muscle activation with individualized vibration. This will enable the prescription of safe and more effective CWBV programs, addressing possible age-related physical debilities and inactivities that these individuals are likely to suffer from.

Footnotes

Authors' Contribution: Study concept and design: Manijeh soleimanifar and Masoumeh Abedi; acquisition of data: Manijeh soleimanifar and Afsaneh Dadarkhah; analysis and interpretation of data: Zahra Sadat Asadi, Masoumeh Abedi and Mostafa Sarabzadeh; drafting of the manuscript: Mostafa Sarabzadeh and Manijeh soleimanifar; critical revision of the manuscript for important intellectual content: Manijeh soleimanifar and Mostafa Sarabzadeh; statistical analysis: Zahra Sadat Asadi; administrative, technical, and material support: Masoumeh Abedi and Afsaneh Dadarkhah; study supervision: Manijeh soleimanifar and Mostafa Sarabzadeh.

Clinical Trial Registration Code: UMIN000046896.

Conflict of Interests: Authors mention that there is no conflict of interest in this study.

Ethical Approval: Ethical approval for this study was obtained from the institutional ethics committee (approval ID: IR.AJAUMS.REC.1396.98).

Funding/Support: This study was supported by AJA university of Medical Sciences, Tehran, Iran.

Informed Consent: After obtaining the necessary permissions for research, candidates completed a medical history form and a consent letter.

References

- Overgaard A, Frederiksen P, Kristensen LE, Robertsson O, W. Dahl A. The implications of an aging population and increased obesity for knee arthroplasty rates in Sweden: A register-based study. *Acta Orthop.* 2020;91(6):738-42. [PubMed ID: 32895012]. [PubMed Central ID: PMC8023941]. https://doi.org/10.1080/17453674.2020.1816268.
- 2. Bluher M. Obesity: global epidemiology and pathogenesis. *Nat Rev Endocrinol.* 2019;**15**(5):288-98. [PubMed ID: 30814686].

https://doi.org/10.1038/s41574-019-0176-8.

- 3. Tantengco OAG. Decreased global online interest in obesity from 2004 to 2021: An infodemiology study. *Obesity Medicine*. 2022;**30**. https://doi.org/10.1016/j.obmed.2022.100389.
- 4. Haththotuwa RN, Wijeyaratne CN, Senarath U. Worldwide epidemic of obesity. *Obesity and Obstetrics.* 2020. p. 3-8. https://doi.org/10.1016/b978-0-12-817921-5.00001-1.
- Marzban M, Farhadi A, Asadipooya K, Jaafari Z, Ghazbani A, Husseinzadeh S, et al. Evaluation of different anthropometric indices and association with metabolic syndrome in community-dwelling older adults: Bushehr Elderly Health (BEH) program. *Obesity Medicine*. 2022;30. https://doi.org/10.1016/j.obmed.2021.100387.
- Reilly JJ, El-Hamdouchi A, Diouf A, Monyeki A, Somda SA. Determining the worldwide prevalence of obesity. *Lancet.* 2018;**391**(10132):1773-4. [PubMed ID: 29739565]. https://doi.org/10.1016/S0140-6736(18)30794-3.
- Emami A, Akbari A, Basirat A, Zare H, Javanmardi F, Falahati F, et al. The role of comorbidities on mortality of COVID-19 in patients with diabetes. *Obes Med.* 2021;25:100352. [PubMed ID: 34027220]. [PubMed Central ID: PMC8123934]. https://doi.org/10.1016/j.obmed.2021.100352.
- Censin JC, Peters SAE, Bovijn J, Ferreira T, Pulit SL, Magi R, et al. Causal relationships between obesity and the leading causes of death in women and men. *PLoS Genet.* 2019;**15**(10). e1008405. [PubMed ID: 31647808]. [PubMed Central ID: PMC6812754]. https://doi.org/10.1371/journal.pgen.1008405.
- Shrestha E, Charkviani M, Musurakis C, Kansakar AR, Devkota A, Banjade R, et al. Type 2 diabetes is associated with increased risk of critical respiratory illness in patients COVID-19 in a community hospital. *Obes Med.* 2021;22:100316. [PubMed ID: 33392411]. [PubMed Central ID: PMC7772088]. https://doi.org/10.1016/j.obmed.2020.100316.
- Mulugeta W, Hoque L. Impact of the COVID-19 lockdown on weight status and associated factors for obesity among children in Massachusetts. *Obes Med.* 2021;22:100325. [PubMed ID: 36567747]. [PubMed Central ID: PMC9764432]. https://doi.org/10.1016/j.obmed.2021.100325.
- AbdelMassih A, Yacoub E, Husseiny RJ, Kamel A, Hozaien R, El Shershaby M, et al. Hypoxia-inducible factor (HIF): The link between obesity and COVID-19. *Obes Med.* 2021;22:100317. [PubMed ID: 33521378]. [PubMed Central ID: PMC7832240]. https://doi.org/10.1016/j.obmed.2020.100317.
- Sattar N, Valabhji J. Obesity as a Risk Factor for Severe COVID-19: Summary of the Best Evidence and Implications for Health Care. *Curr Obes Rep.* 2021;10(3):282-9. [PubMed ID: 34374955]. [PubMed Central ID: PMC8353061]. https://doi.org/10.1007/s13679-021-00448-8.
- Rosenblatt NJ, Madigan ML. Exploring the Association Between Measures of Obesity and Measures of Trip-induced Fall Risk Among Older Adults. Arch Phys Med Rehabil. 2021;102(12):2362-8. [PubMed ID: 34343524]. [PubMed Central ID: PMC8648946]. https://doi.org/10.1016/j.apmr.2021.06.013.
- 14. Choi H, Lim J, Lee S. Body fat-related differences in gait parameters and physical fitness level in weight-matched male adults. *Clin Biomech (Bristol, Avon).* 2021;**81**:105243. [PubMed ID: 33309933]. https://doi.org/10.1016/j.clinbiomech.2020.105243.
- Thiamwong L, Xie R, Conner NE, Renziehausen JM, Ojo EO, Stout JR. Body composition, fear of falling and balance performance in community-dwelling older adults. *Transl Med Aging.* 2023;7:80-6. [PubMed ID: 38516177]. [PubMed Central ID: PMC10957135]. https://doi.org/10.1016/j.tma.2023.06.002.
- Dowling L, McCloskey E, Cuthbertson DJ, Walsh JS. Dynapenic Abdominal Obesity as a Risk Factor for Falls. *J Frailty Aging*. 2023;**12**(1):37-42. [PubMed ID: 36629082]. https://doi.org/10.14283/jfa.2022.18.

- Fernandes VO, Moreira BS, de Melo G, de Avelar NCP, Costa HS, Bastone AC. Factors associated with fear of falling in older women with knee osteoarthritis: A cross-sectional study. *Geriatr Nurs*. 2024;55:333-8. [PubMed ID: 38154414]. https://doi.org/10.1016/ji.gerinurse.2023.12.018.
- Gonzalez M, Gates DH, Rosenblatt NJ. The impact of obesity on gait stability in older adults. *J Biomech*. 2020;**100**:109585. [PubMed ID: 31911052]. [PubMed Central ID: PMC7061260]. https://doi.org/10.1016/j.jbiomech.2019.109585.
- Deger TB, Sarac ZF, Savas ES, Akcicek SF. The Relationship of Balance Disorders with Falling, the Effect of Health Problems, and Social Life on Postural Balance in the Elderly Living in a District in Turkey. *Geriatrics (Basel)*. 2019;4(2). [PubMed ID: 31108836]. [PubMed Central ID: PMC6630729]. https://doi.org/10.3390/geriatrics4020037.
- Rosic G, Milston AM, Richards J, Dey P. Fear of falling in obese women under 50 years of age: a cross-sectional study with exploration of the relationship with physical activity. *BMC Obes.* 2019;6:7. [PubMed ID: 30867933]. [PubMed Central ID: PMC6398230]. https://doi.org/10.1186/s40608-019-0230-z.
- Cieslinska-Swider J, Blaszczyk JW, Opala-Berdzik A. The effect of body mass reduction on functional stability in young obese women. *Sci Rep.* 2022;**12**(1):8876. [PubMed ID: 35614189]. [PubMed Central ID: PMC9132994]. https://doi.org/10.1038/s41598-022-12959-y.
- Carneiro JA, Santos-Pontelli TE, Vilaca KH, Pfrimer K, Colafemina JF, Carneiro AA, et al. Obese elderly women exhibit low postural stability: a novel three-dimensional evaluation system. *Clinics (Sao Paulo)*. 2012;**67**(5):475-81. [PubMed ID: 22666792]. [PubMed Central ID: PMC3351265]. https://doi.org/10.6061/clinics/2012(05)12.
- Maktouf W, Durand S, Boyas S, Pouliquen C, Beaune B. Combined effects of aging and obesity on postural control, muscle activity and maximal voluntary force of muscles mobilizing ankle joint. *J Biomech.* 2018;**79**:198-206. [PubMed ID: 30205976]. https://doi.org/10.1016/j.jbiomech.2018.08.017.
- A H, A E, M S, S A. Risk of Falls among Obese Elderly Females. *The Egyptian Journal of Geriatrics and Gerontology*. 2018;5(1):1-4. https://doi.org/10.21608/ejgg.2018.30880.
- 25. Wu B, Hsu C, Liu Y. [Effects of whole body vibration and electromuscular stimulation recovery methods used betweeninning on baseball pitching performance during a simulated game]. [College Physical Edu. 2020;22(3):247-59. Chinese.
- 26. Yoon JY, Kang SR, Kim HS, Won YH, Park SH, Seo JH, et al. Effects of Low-Frequency Whole-Body Vibration on Muscle Activation, Fatigue, and Oxygen Consumption in Healthy Young Adults: A Single-Group Repeated-Measures Controlled Trial. *J Sport Rehabil*. 2022;**31**(8):984-92. [PubMed ID: 35584804]. https://doi.org/10.1123/jsr.2021-0170.
- Chang CM, Tsai CH, Lu MK, Tseng HC, Lu G, Liu BL, et al. The neuromuscular responses in patients with Parkinson's disease under different conditions during whole-body vibration training. *BMC Complement Med Ther.* 2022;**22**(1):2. [PubMed ID: 34980075]. [PubMed Central ID: PMC8722001]. https://doi.org/10.1186/s12906-021-03481-1.
- 28. de Araujo AXP, de Araujo M, Mendes Fontes LA, Souto Braz RR, Sanudo B, Taiar R, et al. Do two whole-body vibration amplitudes improve postural balance, gait speed, muscle strength, and functional mobility in sedentary older women? A crossover randomized controlled trial. *J Bodyw Mov Ther.* 2022;**32**:143-8. [PubMed ID: 36180141]. https://doi.org/10.1016/j.jbmt.2022.05.010.
- de Souza HCM, Pessoa MF, Dos Santos Clemente R, da Silva AV, de Andrade EAM, Neves PAF, et al. Inspiratory muscle training in addition to whole body vibration for functional and physical outcomes in pre-frail older women: a randomized controlled trial. *Age Ageing.* 2022;**51**(4). [PubMed ID: 35397159]. https://doi.org/10.1093/ageing/afac083.

- Tariq N, Khan Z, Veqar Z. Effect of Whole-Body Vibration on Balance or Proprioception in Nonspecific Chronic Low Back Pain: A Systematic Review. J Chiropr Med. 2023;22(4):284-93. [PubMed ID: 38205229]. [PubMed Central ID: PMC10774621]. https://doi.org/10.1016/j.jcm.2023.04.006.
- Sobhani S, Sinaei E, Motealleh A, Hooshyar F, Kashkooli NS, Yoosefinejad AK. Combined effects of whole body vibration and unstable shoes on balance measures in older adults: A randomized clinical trial. *Arch Gerontol Geriatr.* 2018;78:30-7. [PubMed ID: 29883807]. https://doi.org/10.1016/j.archger.2018.05.016.
- Ruhde L, Hulla R. An overview of the effects of whole-body vibration on individuals with cerebral palsy. *J Pediatr Rehabil Med.* 2022;15(1):193-210. [PubMed ID: 35275570]. https://doi.org/10.3233/PRM-201508.
- Li KY, Cho YJ, Chen RS. The Effect of Whole-Body Vibration on Proprioception and Motor Function for Individuals with Moderate Parkinson Disease: A Single-Blind Randomized Controlled Trial. *Occup Ther Int.* 2021;2021:9441366. [PubMed ID: 34992511]. [PubMed Central ID: PMC8709745]. https://doi.org/10.1155/2021/9441366.
- Guadarrama-Molina E, Barron-Gamez CE, Estrada-Bellmann I, Melendez-Flores JD, Ramirez-Castaneda P, Hernandez-Suarez RMG, et al. Comparison of the effect of whole-body vibration therapy versus conventional therapy on functional balance of patients with Parkinson's disease: adding a mixed group. Acta Neurol Belg. 2021;121(3):721-8. [PubMed ID: 32651877]. https://doi.org/10.1007/s13760-020-01439-7.
- Chang WD, Chen S, Tsou YA. Effects of Whole-Body Vibration and Balance Training on Female Athletes with Chronic Ankle Instability. J Clin Med. 2021;10(11). [PubMed ID: 34071325]. [PubMed Central ID: PMC8198174]. https://doi.org/10.3390/jcm10112380.
- Ochi A, Abe T, Yamada K, Ibuki S, Tateuchi H, Ichihashi N. Effect of balance exercise in combination with whole-body vibration on muscle activity of the stepping limb during a forward fall in older women: a randomized controlled pilot study. *Arch Gerontol Geriatr.* 2015;60(2):244-51. [PubMed ID: 25482957]. https://doi.org/10.1016/j.archger.2014.11.011.
- Parsons J, Mathieson S, Jull A, Parsons M. Does vibration training reduce the fall risk profile of frail older people admitted to a rehabilitation facility? A randomised controlled trial. *Disabil Rehabil.* 2016;**38**(11):1082-8. [PubMed ID: 26693802]. https://doi.org/10.3109/09638288.2015.1103793.
- Nawrat-Szoltysik A, Sieradzka M, Nowacka-Chmielewska M, Piejko L, Duda J, Brachman A, et al. Effect of Whole-Body Vibration Training on Selected Intrinsic Risk Factors in Women Aged 60+ at Fall Risk: A Randomized Controlled Trial. *Int J Environ Res Public Health.* 2022;**19**(24). [PubMed ID: 36554961]. [PubMed Central ID: PMC9779491]. https://doi.org/10.3390/ijerph192417066.
- Liu Y, Fan Y, Chen X. Effects of Whole-Body Vibration Training with Different Body Positions and Amplitudes on Lower Limb Muscle Activity in Middle-Aged and Older Women. *Dose Response*. 2022;20(3):15593258221113000. [PubMed ID: 35859854]. [PubMed Central ID: PMC9289914]. https://doi.org/10.1177/15593258221112960.
- Pujari AN, Neilson RD, Cardinale M. Effects of different vibration frequencies, amplitudes and contraction levels on lower limb muscles during graded isometric contractions superimposed on whole body vibration stimulation. *J Rehabil Assist Technol Eng.* 2019;6:2055668319827470. [PubMed ID: 31245030]. [PubMed Central ID: PMC6582277]. https://doi.org/10.1177/2055668319827466.
- Poncumhak P, Srithawong A, Duangsanjun W, Amput P. Comparison of the Ability of Static and Dynamic Balance Tests to Determine the Risk of Falls among Older Community-Dwelling Individuals. *J Funct Morphol Kinesiol*. 2023;8(2). [PubMed ID: 37092375]. [PubMed Central ID: PMC10123699]. https://doi.org/10.3390/jfmk8020043.

Jundishapur J Health Sci. 2024; 16(2): e144460.

- Van Alsenoy K, Thomson A, Burnett A. Reliability and validity of the Zebris FDM-THQ instrumented treadmill during running trials. *Sports Biomech.* 2019;**18**(5):501-14. [PubMed ID: 29785869]. https://doi.org/10.1080/14763141.2018.1452966.
- Zebris Medical GmbH. *zebris FDM Software Manual*. 2022. Available from: https://www.hpcosmos.com/sites/default/files/uploads/documents/2 0150709_cos102245manen_instruction_for_use_software_zebris_fdm_1.12_r2_en_0.pdf.
- 44. Li Z, Liang YY, Wang L, Sheng J, Ma SJ. Reliability and validity of center of pressure measures for balance assessment in older adults. *J Phys Ther Sci.* 2016;28(4):1364-7. [PubMed ID: 27190484]. [PubMed Central ID: PMC4868244]. https://doi.org/10.1589/jpts.28.1364.
- 45. Baharlouei H, Salavati M, Akhbari B, Mosallanezhad Z, Mazaheri M, Negahban H. Cross-cultural validation of the Falls Efficacy Scale International (FES-I) using self-report and interview-based questionnaires among Persian-speaking elderly adults. Arch Gerontol Geriatr. 2013;57(3):339-44. [PubMed ID: 23830993]. https://doi.org/10.1016/j.archger.2013.06.005.
- Franca AB, Low G, de Souza Santos G, da Costa Serafim R, Vitorino LM. Psychometric properties of the falls efficacy scale-international and validating the short version among older Brazilians. *Geriatr Nurs.* 2021;42(2):344-50. [PubMed ID: 33556901]. https://doi.org/10.1016/j.gerinurse.2021.01.012.
- Kamide N, Shiba Y, Sakamoto M, Sato H. Reliability and validity of the Short Falls Efficacy Scale-International for Japanese older people. *Aging Clin Exp Res.* 2018;30(11):1371-7. [PubMed ID: 29594873]. https://doi.org/10.1007/s40520-018-0940-y.
- Kang H. Sample size determination and power analysis using the G*Power software. J Educ Eval Health Prof. 2021;18:17. [PubMed ID: 34325496]. [PubMed Central ID: PMC8441096]. https://doi.org/10.3352/jeehp.2021.18.17.
- Mordkoff J. A Simple Method for Removing Bias From a Popular Measure of Standardized Effect Size: Adjusted Partial Eta Squared. Advances in Methods and Practices in Psychological Science. 2019;2(3):228-32. https://doi.org/10.1177/2515245919855053.
- Rahlfs V, Zimmermann H. Effect size measures and their benchmark values for quantifying benefit or risk of medicinal products. *Biom J.* 2019;61(4):973-82. [PubMed ID: 30821037]. [PubMed Central ID: PMC6618136]. https://doi.org/10.1002/bimj.201800107.
- 51. Ahmad MN, Zawatia AA. Current prospects of metabolically healthy obesity.
 Obesity Medicine.
 2021;25.

 https://doi.org/10.1016/j.obmed.2021.100361.
 2021;25.
 2021;25.
- Bluher M. Metabolically Healthy Obesity. Endocr Rev. 2020;41(3). [PubMed ID: 32128581]. [PubMed Central ID: PMC7098708]. https://doi.org/10.1210/endrev/bnaa004.
- Yumuk V, Tsigos C, Fried M, Schindler K, Busetto L, Micic D, et al. European Guidelines for Obesity Management in Adults. *Obes Facts*. 2015;8(6):402-24. [PubMed ID: 26641646]. [PubMed Central ID: PMC5644856]. https://doi.org/10.1159/000442721.
- Haase CL, Lopes S, Olsen AH, Satylganova A, Schnecke V, McEwan P. Weight loss and risk reduction of obesity-related outcomes in 0.5 million people: evidence from a UK primary care database. *Int J Obes* (Lond). 2021;45(6):1249-58. [PubMed ID: 33658682]. [PubMed Central ID: PMC8159734]. https://doi.org/10.1038/s41366-021-00788-4.
- 55. Sakai M, Nakazawa M, Abraham D. The comparison of ambulatory activity among people living in different regions, and relationship with BMI in Pohnpei state, the Federated States of Micronesia: Crosssectional study. *Obesity Medicine*. 2020;20. https://doi.org/10.1016/j.obmed.2020.100297.
- 56. Onal B, Sertel M, Karaca G. Effect of plantar vibration on static and dynamic balance in stroke patients: a randomised controlled study.

Physiotherapy. 2022;**116**:1-8. [PubMed ID: 35462214]. https://doi.org/10.1016/j.physio.2022.02.002.

- Melo FATD, Melo GFD, Albuquerque Neto SLD, Silva RWD, França NMD, Silva AAD, et al. Whole-Body Vibration Training Protocols in Obese Individuals: A Systematic Review. *Revista Brasileira de Medicina do Esporte*. 2019;25(6):527-33. https://doi.org/10.1590/1517-869220192506211005.
- Maghbouli N, Khodadost M, Pourhassan S. The effectiveness of vibration therapy for muscle peak torque and postural control in individuals with anterior cruciate ligament reconstruction: a systematic review and meta-analysis of clinical trials. *J Orthop Traumatol*. 2021;22(1):28. [PubMed ID: 34259935]. [PubMed Central ID: PMC8280257]. https://doi.org/10.1186/s10195-021-00589-5.
- Miller RM, Heishman AD, Freitas EDS, Bemben MG. Comparing the Acute Effects of Intermittent and Continuous Whole-Body Vibration Exposure on Neuromuscular and Functional Measures in Sarcopenia and Nonsarcopenic Elderly Women. *Dose Response*. 2018;16(3):1559325818797010. [PubMed ID: 30202250]. [PubMed Central ID: PMC6125857]. https://doi.org/10.1177/1559325818797009.
- Ko MC, Wu LS, Lee S, Wang CC, Lee PF, Tseng CY, et al. Whole-body vibration training improves balance control and sit-to-stand performance among middle-aged and older adults: a pilot randomized controlled trial. *Eur Rev Aging Phys Act.* 2017;14:11. [PubMed ID: 28729887]. [PubMed Central ID: PMC5516349]. https://doi.org/10.1186/s11556-017-0180-8.
- Garcia PA, Queiroz LL, Caetano MBD, Silva K, Hamu T. Obesity is associated with postural balance on unstable surfaces but not with fear of falling in older adults. *Braz J Phys Ther.* 2021;**25**(3):311-8. [PubMed ID: 32830064]. [PubMed Central ID: PMC8134782]. https://doi.org/10.1016/j.bjpt.2020.08.003.
- 62. Neri SGR, Gadelha AB, Correia ALM, Pereira JC, Safons MP, Lima RM. Association between obesity, risk of falls and fear of falling in older women. *Revista Brasileira de Cineantropometria & Desempenho Humano*. 2017;**19**:450-8.
- 63. Neri SGR, Gadelha AB, de David AC, Ferreira AP, Safons MP, Tiedemann A, et al. The Association Between Body Adiposity Measures, Postural Balance, Fear of Falling, and Fall Risk in Older Community-Dwelling Women. *J Geriatr Phys Ther.* 2019;**42**(3):E94-E100. [PubMed ID: 29227413]. https://doi.org/10.1519/JPT.00000000000165.
- 64. Pereira JC, de Moraes Elias J, Neri SG, Gadelha AB, Lemos RR, Lima RM. Dynapenic Abdominal Obesity as a Risk Factor for Falls in Older Women. *Topics in Geriatric Rehabilitation*. 2019;**35**(2):149-55. https://doi.org/10.1097/tgr.00000000000225.
- Korman M, Herling Z, Levy I, Egbarieh N, Engel-Yeger B, Karni A. Background matters: Minor vibratory stimulation during motor skill acquisition selectively reduces off-line memory consolidation. *Neurobiol Learn Mem.* 2017;**140**:27-32. [PubMed ID: 28189551]. https://doi.org/10.1016/j.nlm.2017.02.002.
- Nissim M, Livny A, Barmatz C, Tsarfaty G, Berner Y, Sacher Y, et al. Effects of Ai-Chi Practice on Balance and Left Cerebellar Activation during High Working Memory Load Task in Older People: A Controlled Pilot Trial. *Int J Environ Res Public Health*. 2021;**18**(23). [PubMed ID: 34886482]. [PubMed Central ID: PMC8656797]. https://doi.org/10.3390/ijerph182312756.
- Alam MM, Khan AA, Farooq M. Effects of vibration therapy on neuromuscular efficiency & features of the EMG signal based on endurance test. *J Bodyw Mov Ther.* 2020;24(4):325-35. [PubMed ID: 33218530]. https://doi.org/10.1016/j.jbmt.2020.06.037.
- Blazevich AJ, Collins DF, Millet GY, Vaz MA, Maffiuletti NA. Enhancing Adaptations to Neuromuscular Electrical Stimulation Training Interventions. *Exerc Sport Sci Rev.* 2021;49(4):244-52. [PubMed ID: 34107505]. [PubMed Central ID: PMC8460078]. https://doi.org/10.1249/JES.00000000000264.