



Prevalence of SARS-CoV-2 Specific Antibodies in the Staff of a Children's Hospital, in Tehran, Iran

Shahnaz Armin¹, Fereshteh Karbasian², Seyedeh Mahsan Hoseinialfatemi¹, Roxana Mansour Ghanaie¹, Sedigheh Rafiei Tabatabaei¹, Seyed Alireza Fahimzad¹, Maryam Rajabnejad¹, Ghamartaj Khanbabaee³, Soheila Sadat Vaghefi⁴, Farid Imanzadeh⁵, Saeed Maham¹, Raheleh Razmara⁶, Abdollah Karimi^{1,*}, Peyman Eshghi^{7,8,**}, Mahtab Maghsudlu⁷ and Ali Arabkhazaeli⁷

¹Pediatric Infections Research Center, Research Institute for Children's Health, Shahid Beheshti University of Medical Sciences, Tehran, Iran

²Department of Pediatric Emergency Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran

³Department of Pediatric Pulmonology, Mofid Children's Medical Center, Shahid Beheshti University of Medical Sciences, Tehran, Iran

⁴Mofid Children's Hospital, Shahid Beheshti University of Medical Sciences, Tehran, Iran

⁵Pediatric Gastroenterology, Hepatology and Nutrition Research Center, Research Institute for Children Health, Shahid Beheshti University of Medical Sciences, Tehran, Iran

⁶Faculty of Nursing & Midwifery, Shahid Beheshti University of Medical Sciences, Tehran, Iran

⁷Blood Transfusion Research Center, High Institute for Research and Education in Transfusion Medicine, Tehran, Iran

⁸Pediatric Congenital Hematologic Disorders Research Center, Shahid Beheshti University of Medical Sciences, Tehran, Iran

*Corresponding author: Pediatric Infections Research Center, Research Institute for Children's Health, Shahid Beheshti University of Medical Sciences, Tehran, Iran. Email: dry.karimi@gmail.com

**Corresponding author: Pediatric Congenital Hematologic Disorders Research Center, Shahid Beheshti University of Medical Sciences and Iran Blood Transfusion Organization, Tehran, Iran. Email: peshghi64@gmail.com

Received 2020 August 17; Revised 2020 October 21; Accepted 2020 November 06.

Abstract

Background: In December 2019, a new coronavirus appeared in China, as a cause of acute respiratory disease. Healthcare Workers (HCWs) in children's hospitals are one of the groups who are at a high rate of exposure to COVID-19 patients. The detection of antibodies is useful and helps diagnose late or recent SARS-CoV-2 infection. Most children may present with asymptomatic or mild SARS-CoV-2 infection and can be silent sources of infection in the community and hospitals.

Objectives: This study was conducted to determine the antibodies (IgM and IgG) against SARS-CoV-2 in Mofid children's hospital staff.

Methods: This cross-sectional study evaluated 475 staff from Mofid children's hospital from April 20 to May 5, 2020, in Tehran. We collected blood samples for the antibody assay with a rapid test kit. A questionnaire was used to collect demographic and clinical data.

Results: Of 475 staff who participated in this study, 25 (5.3%) were diagnosed with COVID-19 by Polymerase Chain Reaction (PCR) and/or CT scan. Besides, 140 (29.4%) of them were positive for IgM and/or IgG SARS-CoV-2 antibodies.

Conclusions: The seropositivity of antibodies against SARS-CoV-2 among children's hospital staff was higher than expected. Approximately 5% of the participants were diagnosed as definitive positive COVID-19 cases by PCR and/or CT scan, but 29.4% of them were seropositive. The difference is a warning, as it shows seropositive people could be silent sources of transmission during asymptomatic infection.

Keywords: SARS-CoV-2, COVID-19, Antibodies, Seroprevalence, Healthcare Workers

1. Background

New unknown pneumonia was reported from Wuhan, China, at the end of 2019 (1, 2). The etiology was known to be a coronavirus family member, named Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2). It spread around the world very soon. On March 11, 2020, the World Health Organisation (WHO) declared this global health emergency a pandemic (3, 4).

The severity of COVID-19 symptoms can range from

very mild to severe, and some people may show no symptoms. Healthcare Workers (HCWs) are one of the high-risk groups who are exposed to COVID-19 (5). Many reports from different countries show the outbreak of COVID-19 among HCWs (6-8). Infection in hospital staff is a big challenge, as it reduces the capacity of the health system and increases secondary transmission (9). Thus, it seems that the early detection of infected hospital staff is crucial to prevent the transmission of the virus to other employees and

patients and maintain their stable workforce at optimum performance.

Typically, seroconversion occurs 11 to 14 days after the first symptoms (10). According to various studies, the overall seroprevalence of COVID-19 in HCWs is low. Based on data from Santa Clara state of USA (11), Germany (8), and Dutch hospitals (12), the rates of SARS-CoV-2 IgG antibody were 1.5, 4.5, and 6%, respectively. However, another study from the United Kingdom revealed that the rate was higher (18%) (13). The Real-Time Reverse Transcription Polymerase Chain Reaction (RT-PCR) in respiratory samples was introduced as the gold standard COVID-19 diagnostic test (14) and is a sensitive and accurate way to confirm the diagnosis of SARS-CoV-2 infection in practice (15). However, this test takes a long time (an average of two to three hours) to get the result and requires certified laboratories, expensive equipment, and trained technicians.

Most importantly, several false-negative results have been reported for RT-PCR of COVID-19 (14) that can lead to a major problem in the diagnosis and controlling of virus transmission (16). To identify a large number of symptomatic and asymptomatic infected patients quickly, an accurate and rapid testing method is needed (15). Serological tests can be helpful (17). These tests allow experts to know what proportion of the population has been involved, and assess their types of involvement (symptomatic or asymptomatic) and the probability of spreading the virus without being aware of it (18). Also, it helps estimate how many people are at risk. On the other hand, in workplaces, serology detection can help get a decision on who can return to work safely, especially for healthcare workers, provided that high-quality tests are used.

2. Objectives

At Mofid children's hospital, based on unpublished data, 36 of 800 employees became infected within two months of the COVID-19 epidemic. No data is available for asymptomatic cases because we did not conduct periodic surveillance by PCR in our hospital. As children can be silent carriers or mild non-specific symptom presenters, the HCWs in children's hospitals may be at higher risk of infection in contact with apparently asymptomatic cases, causing a neglected risk. This study aimed at determining the antibodies (IgM and IgG) against SARS-COV-2 in Mofid children's hospital staff to define the rate of seropositivity.

3. Methods

3.1. Specimens Collection, Preparation, and Testing

This cross-sectional study was performed in April 2020 in a referral pediatric subspecialty tertiary center in Tehran

(the capital of Iran). From Mofid children's hospital, 475 staff participated in the study. The inclusion criteria included all Mofid children's hospital staff who were presented in the period of this study. The COVID-19 IgG/IgM antibodies were detected by Rapid Test Dipstick (Hangzhou AllTest Biotech Co. INCP-401, China). Rapid serological tests provide a means to quickly diagnose COVID-19-suspected cases by detecting IgG and IgM antibodies in blood samples. The participant's finger was punctured by a lancet, and the blood was applied to the test card by using a capillary tube. Then, 20 μ l of blood and one drop of buffer were added to the test card, and we waited for the colored line(s) to appear. The result was interpreted by the company instructions based on three lines of color (Control, IgG, and IgM). All the tests were performed under strict biosafety conditions. Informed consent was obtained from all participants in this study. The sensitivity, specificity, and accuracy of COVID-19 IgG/IgM antibodies according to the manufacturer's instructions are shown in Table 1.

3.2. Statistical Analyses

Statistical analyses were performed using SPSS software, version 25. Simple and multiple logistic regressions were used to analyze the relationship between risk factors and the disease. P values of less than 0.05 were considered statistically significant.

4. Results

Of 475 staff participating in the study, 387 were female, and 97 were male. The demographic data of the participants by serological results and the relationship between individual independent variables and COVID-19 serology results are presented in Table 2. A total of 140 (29.4%) staff were infected with COVID-19 based on serologic tests, including 115/140 health care workers, 13 office workers, and 12 hospital service workers. Among 140 seropositive cases, 52 were asymptomatic, 83 were symptomatic, and five had severe symptoms who were hospitalized. Besides, 51 (10.7%) and 65 (13.7%) cases had positive IgM and IgG antibodies, respectively. Also, 24 (5%) cases had both IgG and IgM positive results (Figure 1). To investigate the effect of family crowding, we divided the sample based on the number of family members into two groups of < 4 and > 4 persons. The results showed that 66 (47.14%) cases with positive serology belonged to the group of < 4 persons, and 74 (52.86%) cases belonged to the group of > 4 persons (P value = 0.06). Besides, 25 (5.26%) staff had a history of confirmed COVID-19 by CT scan (16 cases) or PCR (nine cases), of whom 21 (84%) were positive (3 IgM+, 10 IgG+, 8 both of them) and four (16%) were seronegative.

Table 1. Sensitivity, Specificity, and Accuracy of COVID-19 IgG/IgM Rapid Test Kits

	IgG	IgM
Relative Sensitivity	> 99.9% (95%CI ^a : 82.5%-100%)	90.9% (95%CI ^a : 71.0%-98.7%)
Relative Specificity	98.0% (95%CI ^a : 92.6%-99.9%)	97.0% (95%CI ^a : 91.8%-99.4%)
Accuracy	98.4% (95%CI ^a : 93.9%-99.9%)	95.9% (95%CI ^a : 90.5%-98.5%)

^aConfidence Interval.**Table 2.** Demographic Data of Participants by Serological Results and Relationship Between Individual Independent Variables and COVID-19 Serology Results in Hospital Staff

Variable	Positive Serology		OR	95%CI	P Value
	Negative (n=335)	Positive IgM, IgG, or both (n=140)			
Education level					
Diploma	62 (71.26%)	25 (28.74%)	Ref.		
BSc	178 (71.77%)	70 (28.23%)	0.98	0.57 - 1.67	0.93
MSc	31 (62.00%)	19 (38.00%)	1.52	0.73 - 3.17	0.26
Doctorate	64 (71.11%)	26 (28.89%)	1.01	0.53 - 1.93	0.98
Age group					
< 40	238 (74.84%)	80 (25.16%)	Ref.		
40-62	97 (61.78%)	60 (38.22%)	1.84	1.22 - 2.77	0.004
Type of service					
Service	27 (69.23%)	12 (30.77%)	Ref.		
Official	16 (55.17%)	13 (44.83%)	1.83	0.67 - 4.96	0.24
Health/Medical	292 (71.74%)	115 (28.26%)	0.89	0.43 - 1.81	0.74
Family size					
1-3	190 (74.22%)	66 (25.78%)	Ref.		
≥ 4	145 (66.21%)	74 (33.79%)	1.47	0.99 - 2.18	0.06
Used masks					
N95	46 (73.02%)	17 (26.98%)	Ref.		
Surgery	218 (68.99%)	98 (31.01%)	1.22	0.66 - 2.23	0.53
Other types	67 (76.14%)	21 (23.86%)	0.85	0.40 - 1.78	0.66
None	4 (50.00%)	4 (50.00%)	2.71	0.61 - 12.05	0.19
Body mass index					
< 24	169 (76.47%)	52 (23.53%)	Ref.		
≥ 24	166 (65.35%)	88 (34.65%)	1.72	1.15 - 2.58	0.008
Contact to persons with COVID-19					
No	135 (70.31%)	57 (29.69%)	Ref.		
Yes	200 (70.67%)	83 (29.33%)	0.98	0.66 - 1.47	0.93

The results of multiple analyses showed that the two variables of BMI and family members had relationships with positive serology. The odds ratio was 1.72 times higher for health workers with a BMI ≥ 24 compared to BMI below 24 (95% confidence interval: 1.17 to 2.57; P value = 0.006) and 1.49 times higher for those with a family of four mem-

bers or more in comparison with families with three or fewer members (95% confidence interval: 1.01 to 2.20; P value = 0.04). By the elimination of the number of family members, variables that were independently related to positive serology of COVID-19 were BMI and age. It means that the odds ratio is 1.62 times higher in health care work-




Figure 1. Serological results of the participants

ers with BMI ≥ 24 in comparison with ones who had BMI below 24 (95% confidence level: 1.09 to 2.43; P value = 0.02) and 1.52 times higher in health workers ≥ 40 -years-old in comparison with ones who were < 40 years (95% confidence interval: 1.01 to 2.30; P value = 0.04). Other variables were not significant in this study.

5. Discussion

Children can be silent carriers or show non-specific mild symptoms. Children are admitted to the hospital for different reasons while they carry viruses. As all hospitalized children stay with their mothers in the wards, people who work in children's hospitals may be at higher risk of being infected by SARS CoV-2. This study aimed to investigate the effect of different variables on the seropositivity of Mofid children's hospital staff in Tehran, Iran. Based on our knowledge, this research was the first large-scale prevalence study on Iranian children's hospital staff during the COVID-19 pandemic. Our results showed that 140 (29.4%) staff were seropositive for IgG, IgM, or both. Other studies from Belgium and the USA showed rates of 6.4% and 1.5%, respectively, which were lower than our result (11, 19). In a study conducted at a pediatric and woman hospital in Germany, 1.8% were serologically positive (20). The high rate of antibodies among our staff may be due to some

reasons. First of all, the epidemic of COVID-19 started in Iran sooner than in Europe and the USA. As our study was conducted four months after the epidemic onset, our population had more time to being infected more. Second, it may be related to the lack of compulsory quarantine in our country compared to other countries.

This study was done in a children's hospital where patients were admitted because of different complaints and stayed with their parents who might be carriers of the virus. Also, at the first stages of the epidemic, according to the data from Wuhan, it seemed that the rate of infection in children is low; so HCWs in children hospitals did not follow the protocols carefully. As 51 of our cases were IgM positive, it may show that they were infected recently, and their immune system started responding to the virus. Besides, 65 cases were IgG positive; this antibody indicates that they may have acquired COVID-19 before. Similar to the results of our study, other studies have shown that age has a significant role in the incidence of the disease (21-23).

Our findings showed that the risk of seropositivity was not different between staff who were working in administrative departments of the hospital and those working as health care or service workers. In contrast, a study from Wuhan showed that the number of HCWs with COVID-19 was significantly lower in office buildings than in clinical wards (24). A higher rate of seroconversion in our study

than in Lai et al. study from Wuhan may show that our staff in different parts of the hospital did not appropriately follow protocols because the risk of contact and working with children was not clear as an important risk factor for them. Thus, we think awareness and education have important roles in better protection. The number of family members in our study did not correlate with seropositivity. Unlike our results, Song et al. reported that close contact and even casual contact with family members increased the risk of COVID-19 transmission (25).

Although most of the participants announced that they were wearing masks, positive serological tests for 140 (29.4%) cases represented that they had not used proper protection during the epidemic, and still, we need more education. Based on another study, a high BMI was a risk factor for the severity of COVID-19 (26). Although we did not compare the seropositivity rate between high and normal BMI people because of limited cases, our results showed that staff with high BMI were more seropositive than normal BMI people. Based on this data, a high BMI can be a risk factor for getting the disease. Besides, 245 participants announced contact with COVID-19 cases in the hospital, 174 (71%) of whom were seronegative. Thus, in this study, having contact with COVID-19 cases was not statistically effective. Our findings are also in line with Fusco et al.'s study in Naples and Korth et al.'s study in Germany that reported 1.6% seropositivity in HCWs (8, 27). The low rate of seropositivity in these staff may be related to some reasons. First, based on the data, only eight participants did not use masks or appropriate personal preventive equipment during the epidemic. Second, it was not confirmed which participants exactly had contact with suspected and probable COVID-19 cases.

Of 230 people who did not appear to have specific contact with patients, 57 (24.7%) had positive serology. Ningxia et al. and Gandhi et al. confirmed that asymptomatic COVID-19 infection is possible, and our data again confirmed that the sources of infection can be asymptomatic or may have mild symptoms (28, 29). This study had some limitations. Because of the pandemic situation, only 60% of the hospital staff were presented at the time of the survey. We did not conduct PCR testing in this study because of limited resources. Thus, we considered negative serology equal to not dealing with COVID-19. However, we know it may happen in the current infection, and even it may change to negative in cured cases.

5.1. Conclusions

In summary, the serological test of the staff from a tertiary children's hospital in Iran had 29.4% positivity after four months of starting the COVID-19 epidemic, which was higher than the rate in the general population at the time

of the study. It could be an alarming sign to insist on continuous education on using PPE and other necessary preventive protocols and being aware of asymptomatic carriers.

Acknowledgments

This research was partially supported by the Iranian Blood Transfusion Organization. We thank our colleagues from the Nursery Office and Laboratory Infections Research Center of Mofid children's hospital that provided insight and expertise, which greatly assisted the research.

Footnotes

Authors' Contribution: Shahnaz Armin, Saeed Maham, Abdollah Karimi, and Peyman Eshghi contributed to the study design and interpretation of data. Seyedeh Mahsan Hoseini-Alfatemi, Shahnaz Armin, Abdollah Karimi, and Fereshteh Karbasian were involved in drafting. Roxana Mansour Ghanaie, Sedigheh Rafiei Tabatabaei, Seyed Alireza Fahimzad, and Maryam Rajabnejad contributed to the critical revision of the manuscript. Seyedeh Mahsan Hoseini-Alfatemi and Raheleh Razmara took part in data acquisition. Saeed Maham, Ghamartaj Khanbabaee, Soheila Sadat Vaghefi, Farid Imanzadeh, Mahtab Maghsudlu, and Ali Arabkhazaeli contributed to the management and coordination of the research activity planning and execution.

Conflict of Interests: There is no conflict of interest.

Ethical Approval: The study protocol was approved by the Pediatric Infections Research Center (PIRC) of Shahid Beheshti University of Medical Sciences. The ethical approval code is IR.SBMU.RICH.REC.1399.031.

Funding/Support: This work was supported by the Research Institute for Children's Health, Shahid Beheshti University of Medical Sciences, Tehran, Iran.

Informed Consent: Informed consent was obtained from all participants in this study.

References

1. Wu JT, Leung K, Leung GM. Nowcasting and forecasting the potential domestic and international spread of the 2019-nCoV outbreak originating in Wuhan, China: a modelling study. *The Lancet*. 2020;395(10225):689-97. doi: 10.1016/S0140-6736(20)30260-9.
2. WHO. *Rolling updates on coronavirus disease (COVID-19)*. 2020, [cited May 25 2020]. Available from: <https://www.who.int/emergencies/diseases/novel-coronavirus-2019/events-as-they-happen>.
3. Lau SK, Woo PC, Yip CC, Tse H, Tsoi HW, Cheng VC, et al. Coronavirus HKU1 and other coronavirus infections in Hong Kong. *J Clin Microbiol*. 2006;44(6):2063-71. doi: 10.1128/JCM.02614-05. [PubMed: 16757599]. [PubMed Central: PMC1489438].

4. Singhal T. A Review of coronavirus disease-2019 (COVID-19). *Indian J Pediatr.* 2020;**87**(4):281-6. doi: [10.1007/s12098-020-03263-6](https://doi.org/10.1007/s12098-020-03263-6). [PubMed: [32166607](https://pubmed.ncbi.nlm.nih.gov/32166607/)]. [PubMed Central: [PMC7090728](https://pubmed.ncbi.nlm.nih.gov/PMC7090728/)].
5. Koh D. Occupational risks for COVID-19 infection. *Occup Med (Lond).* 2020;**70**(1):3-5. doi: [10.1093/occmed/kqaa036](https://doi.org/10.1093/occmed/kqaa036). [PubMed: [32107548](https://pubmed.ncbi.nlm.nih.gov/32107548/)]. [PubMed Central: [PMC7107962](https://pubmed.ncbi.nlm.nih.gov/PMC7107962/)].
6. Bahl P, Doolan C, de Silva C, Chughtai AA, Bourouiba L, MacIntyre CR. Airborne or droplet precautions for health workers treating COVID-19? *J Infect Dis.* 2020. doi: [10.1093/infdis/jiaa189](https://doi.org/10.1093/infdis/jiaa189). [PubMed: [32301491](https://pubmed.ncbi.nlm.nih.gov/32301491/)]. [PubMed Central: [PMC7184471](https://pubmed.ncbi.nlm.nih.gov/PMC7184471/)].
7. Chou R, Dana T, Buckley DI, Selph S, Fu R, Totten AM. Update alert 6: epidemiology of and risk factors for coronavirus infection in health care workers. *Ann Intern Med.* 2020. doi: [10.7326/L20-1323](https://doi.org/10.7326/L20-1323). [PubMed: [33226856](https://pubmed.ncbi.nlm.nih.gov/33226856/)].
8. Korth J, Wilde B, Dollf S, Anastasiou OE, Krawczyk A, Jahn M, et al. SARS-CoV-2-specific antibody detection in healthcare workers in Germany with direct contact to COVID-19 patients. *J Clin Virol.* 2020;**128**:104437. doi: [10.1016/j.jcv.2020.104437](https://doi.org/10.1016/j.jcv.2020.104437). [PubMed: [32434708](https://pubmed.ncbi.nlm.nih.gov/32434708/)]. [PubMed Central: [PMC7219425](https://pubmed.ncbi.nlm.nih.gov/PMC7219425/)].
9. Adams JG, Walls RM. Supporting the health care workforce during the COVID-19 global epidemic. *JAMA.* 2020;**323**(15):1439-40. doi: [10.1001/jama.2020.3972](https://doi.org/10.1001/jama.2020.3972). [PubMed: [32163102](https://pubmed.ncbi.nlm.nih.gov/32163102/)].
10. Petherick A. Developing antibody tests for SARS-CoV-2. *The Lancet.* 2020;**395**(10230):1101-2. doi: [10.1016/s0140-6736\(20\)30788-1](https://doi.org/10.1016/s0140-6736(20)30788-1).
11. Bendavid E, Mulaney B, Sood N, Shah S, Ling E, Bromley-Dulfano R, et al. COVID-19 antibody seroprevalence in Santa Clara County, California. *medRxiv.* 2020. doi: [10.1101/2020.04.14.20062463](https://doi.org/10.1101/2020.04.14.20062463).
12. Kluytmans M, Buiting A, Pas S, Bentvelsen R, van den Bijllaardt W, van Oudheusden A, et al. SARS-CoV-2 infection in 86 healthcare workers in two Dutch hospitals in March 2020. *medRxiv.* 2020.
13. Keeley AJ, Evans C, Colton H, Ankcorn M, Cope A, State A, et al. Roll-out of SARS-CoV-2 testing for healthcare workers at a large NHS Foundation Trust in the United Kingdom, March 2020. *Euro Surveill.* 2020;**25**(14). doi: [10.2807/1560-7917.ES.2020.25.14.2000433](https://doi.org/10.2807/1560-7917.ES.2020.25.14.2000433). [PubMed: [32290904](https://pubmed.ncbi.nlm.nih.gov/32290904/)]. [PubMed Central: [PMC7160437](https://pubmed.ncbi.nlm.nih.gov/PMC7160437/)].
14. Li Z, Yi Y, Luo X, Xiong N, Liu Y, Li S, et al. Development and clinical application of a rapid IgM-IgG combined antibody test for SARS-CoV-2 infection diagnosis. *J Med Virol.* 2020;**92**(9):1518-24. doi: [10.1002/jmv.25727](https://doi.org/10.1002/jmv.25727). [PubMed: [32104917](https://pubmed.ncbi.nlm.nih.gov/32104917/)]. [PubMed Central: [PMC7228300](https://pubmed.ncbi.nlm.nih.gov/PMC7228300/)].
15. Long Q, Deng H, Chen J, Hu J, Liu B, Liao P, et al. Antibody responses to SARS-CoV-2 in COVID-19 patients: the perspective application of serological tests in clinical practice. *medRxiv.* 2020. doi: [10.1101/2020.03.18.20038018](https://doi.org/10.1101/2020.03.18.20038018).
16. Guo L, Ren L, Yang S, Xiao M, Yang F, Chang, et al. Profiling early humoral response to diagnose novel coronavirus disease (COVID-19). *Clin Infect Dis.* 2020;**71**(15):778-85. doi: [10.1093/cid/ciaa310](https://doi.org/10.1093/cid/ciaa310). [PubMed: [32198501](https://pubmed.ncbi.nlm.nih.gov/32198501/)]. [PubMed Central: [PMC7184472](https://pubmed.ncbi.nlm.nih.gov/PMC7184472/)].
17. Cassaniti I, Novazzi F, Giardina F, Salinaro F, Sachs M, Perlini S, et al. Performance of VivaDiag COVID-19 IgM/IgG Rapid Test is inadequate for diagnosis of COVID-19 in acute patients referring to emergency room department. *J Med Virol.* 2020. doi: [10.1002/jmv.25800](https://doi.org/10.1002/jmv.25800). [PubMed: [32227490](https://pubmed.ncbi.nlm.nih.gov/32227490/)]. [PubMed Central: [PMC7228409](https://pubmed.ncbi.nlm.nih.gov/PMC7228409/)].
18. Du Z, Zhu F, Guo F, Yang B, Wang T. Detection of antibodies against SARS-CoV-2 in patients with COVID-19. *J Med Virol.* 2020. doi: [10.1002/jmv.25820](https://doi.org/10.1002/jmv.25820). [PubMed: [32243608](https://pubmed.ncbi.nlm.nih.gov/32243608/)]. [PubMed Central: [PMC7228383](https://pubmed.ncbi.nlm.nih.gov/PMC7228383/)].
19. Steensels D, Oris E, Coninx L, Nuyens D, Delforge ML, Vermeersch P, et al. Hospital-wide SARS-CoV-2 antibody screening in 3056 staff in a tertiary center in Belgium. *JAMA.* 2020;**324**(2):195-7. doi: [10.1001/jama.2020.11160](https://doi.org/10.1001/jama.2020.11160). [PubMed: [32539107](https://pubmed.ncbi.nlm.nih.gov/32539107/)]. [PubMed Central: [PMC7296458](https://pubmed.ncbi.nlm.nih.gov/PMC7296458/)].
20. Brandstetter S, Roth S, Harner S, Buntrock-Dopke H, Toncheva AA, Borchers N, et al. Symptoms and immunoglobulin development in hospital staff exposed to a SARS-CoV-2 outbreak. *Pediatr Allergy Immunol.* 2020. doi: [10.1111/pai.13278](https://doi.org/10.1111/pai.13278). [PubMed: [32413201](https://pubmed.ncbi.nlm.nih.gov/32413201/)].
21. Liu Y, Mao B, Liang S, Yang JW, Lu HW, Chai YH, et al. Association between age and clinical characteristics and outcomes of COVID-19. *Eur Respir J.* 2020;**55**(5). doi: [10.1183/13993003.01112-2020](https://doi.org/10.1183/13993003.01112-2020). [PubMed: [32312864](https://pubmed.ncbi.nlm.nih.gov/32312864/)]. [PubMed Central: [PMC7173682](https://pubmed.ncbi.nlm.nih.gov/PMC7173682/)].
22. Zhou F, Yu T, Du R, Fan G, Liu Y, Liu Z, et al. Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. *The Lancet.* 2020;**395**(10229):1054-62. doi: [10.1016/s0140-6736\(20\)30566-3](https://doi.org/10.1016/s0140-6736(20)30566-3).
23. Wang W, Tang J, Wei F. Updated understanding of the outbreak of 2019 novel coronavirus (2019-nCoV) in Wuhan, China. *J Med Virol.* 2020;**92**(4):441-7. doi: [10.1002/jmv.25689](https://doi.org/10.1002/jmv.25689). [PubMed: [31994742](https://pubmed.ncbi.nlm.nih.gov/31994742/)]. [PubMed Central: [PMC7167192](https://pubmed.ncbi.nlm.nih.gov/PMC7167192/)].
24. Lai X, Wang M, Qin C, Tan L, Ran L, Chen D, et al. Coronavirus disease 2019 (COVID-2019) infection among health care workers and implications for prevention measures in a tertiary hospital in Wuhan, China. *JAMA Netw Open.* 2020;**3**(5). e209666. doi: [10.1001/jamanetworkopen.2020.9666](https://doi.org/10.1001/jamanetworkopen.2020.9666). [PubMed: [32437575](https://pubmed.ncbi.nlm.nih.gov/32437575/)]. [PubMed Central: [PMC7243089](https://pubmed.ncbi.nlm.nih.gov/PMC7243089/)].
25. Song R, Han B, Song M, Wang L, Conlon CP, Dong T, et al. Clinical and epidemiological features of COVID-19 family clusters in Beijing, China. *J Infect.* 2020;**81**(2):e26-30. doi: [10.1016/j.jinf.2020.04.018](https://doi.org/10.1016/j.jinf.2020.04.018). [PubMed: [32335171](https://pubmed.ncbi.nlm.nih.gov/32335171/)]. [PubMed Central: [PMC7177072](https://pubmed.ncbi.nlm.nih.gov/PMC7177072/)].
26. Kass DA, Duggal P, Cingolani O. Obesity could shift severe COVID-19 disease to younger ages. *The Lancet.* 2020;**395**(10236):1544-5. doi: [10.1016/s0140-6736\(20\)31024-2](https://doi.org/10.1016/s0140-6736(20)31024-2).
27. Fusco FM, Pisaturo M, Iodice V, Bellopede R, Tambaro O, Parrella G, et al. COVID-19 among healthcare workers in a specialist infectious diseases setting in Naples, Southern Italy: results of a cross-sectional surveillance study. *J Hosp Infect.* 2020;**105**(4):596-600. doi: [10.1016/j.jhin.2020.06.021](https://doi.org/10.1016/j.jhin.2020.06.021). [PubMed: [32565367](https://pubmed.ncbi.nlm.nih.gov/32565367/)]. [PubMed Central: [PMC7301109](https://pubmed.ncbi.nlm.nih.gov/PMC7301109/)].
28. Yu X, Yang R. COVID-19 transmission through asymptomatic carriers is a challenge to containment. *Influenza Other Respir Viruses.* 2020;**14**(4):474-5. doi: [10.1111/irv.12743](https://doi.org/10.1111/irv.12743). [PubMed: [32246886](https://pubmed.ncbi.nlm.nih.gov/32246886/)]. [PubMed Central: [PMC7228388](https://pubmed.ncbi.nlm.nih.gov/PMC7228388/)].
29. Gandhi M, Yokoe DS, Havlir DV. Asymptomatic transmission, the achilles' heel of current strategies to control Covid-19. *N Engl J Med.* 2020;**382**(22):2158-60. doi: [10.1056/NEJMe2009758](https://doi.org/10.1056/NEJMe2009758). [PubMed: [32329972](https://pubmed.ncbi.nlm.nih.gov/32329972/)]. [PubMed Central: [PMC7200054](https://pubmed.ncbi.nlm.nih.gov/PMC7200054/)].