



Epidemiological Trends and Demographic Determinants of Malaria in Chabahar and Suburbs, Southeastern Iran: Insights from a Six-Year Study (2019 - 2024)

Erfan Ebrahimi Eskandar Kolaie ¹, Mahdi Rezaei Kahkha-Zhaleh ², Bahman Fouladi ³, Shaghayegh Dabirzadeh ⁴, Mansour Dabirzadeh ^{5,*}

¹ Student Research Committee, Zabol University of Medical Sciences, Zabol, Iran

² Ph.D Student, Health Promotion Research Center, Zahedan University of Medical Sciences, Zahedan, Iran

³ Assistant Professor of Mycology, Department of Parasitology and Mycology, Faculty of Medical Sciences, Zabol University of Medical Sciences, Zabol, Iran

⁴ Student of Public Health, School of Medicine, Najaf Abad (IAUN), Islamic Azad University of Medical Sciences, Najaf Abad, Iran

⁵ Associate Professor of Parasitology, Department of Parasitology and Mycology, Faculty of Medicine, Zabol University of Medical Sciences, Zabol, Iran

*Corresponding Author: Associate Professor of Parasitology, Department of Parasitology and Mycology, Faculty of Medicine, Zabol University of Medical Sciences, Zabol, Iran. Tel: +98-9131002177; Email: mdabirzadeh20002000@gmail.com

Received: 20 April, 2025; Accepted: 30 April, 2025

Abstract

Background: Malaria remains a critical public health challenge in southeastern Iran, where Chabahar's tropical climate and cross-border migration from endemic regions exacerbate transmission.

Objectives: This six-year study (2019 - 2024) investigates malaria trends, climatic drivers, and demographic risks to inform targeted control strategies.

Methods: A retrospective analysis of 3,393 confirmed malaria cases was conducted using blood smears, rapid diagnostic tests, and demographic questionnaires. Meteorological data were integrated to assess the impacts of temperature, humidity, and precipitation. Statistical analyses (SPSS version 26) included ANOVA, Spearman's correlation, and chi-square tests.

Results: Malaria incidence peaked in 2023 (1,272 cases), with *Plasmodium vivax* dominating (98.7% in 2023; 85.8% in 2024), while *P. falciparum* prevailed in 2021 (84.9%). Temperature strongly correlated with transmission, particularly for mixed infections (mean: 33.6°C; $P < 0.001$), whereas humidity and precipitation showed no significant effect. Males constituted 72.9% to 81.6% of cases, and international travel was linked to higher *P. vivax* incidence (35.8% of travel-related infections). Afghan nationals exhibited the highest *P. vivax* burden (83%), underscoring cross-border transmission risks.

Conclusions: Temperature-driven transmission and migration dynamics necessitate climate-informed strategies in Chabahar. Prioritizing male populations, travelers, and cross-border collaboration, alongside enhanced surveillance for relapsing *P. vivax*, is critical for elimination. Integrating temperature-based predictive models and addressing regional genetic factors (e.g., G6PD deficiency) could optimize interventions in this high-risk zone.

Keywords: Malaria, Epidemiology, Climate, *Plasmodium vivax*, Iran, Migration

1. Background

Malaria, a vector-borne disease caused by *Plasmodium* parasites, remains a pressing global health challenge, with approximately 247 million cases and 619,000 deaths reported in 2021 (WHO, 2022). Despite advancements in prevention and treatment, the disease

persists as a significant burden, underscoring the need for context-specific strategies to achieve control and elimination (1, 2). Climate change has amplified concerns about the spread of vector-borne diseases, with rising temperatures, humidity, and rainfall influencing mosquito breeding and malaria transmission (3). In Chabahar, southeastern Iran,

Copyright © 2025, Ebrahimi Eskandar Kolaie et al. This open-access article is available under the Creative Commons Attribution 4.0 (CC BY 4.0) International License (<https://creativecommons.org/licenses/by/4.0/>), which allows for unrestricted use, distribution, and reproduction in any medium, provided that the original work is properly cited.

How to Cite: Ebrahimi Eskandar Kolaie E, Rezaei Kahkha-Zhaleh M, Fouladi B, Dabirzadeh S, Dabirzadeh M. Epidemiological Trends and Demographic Determinants of Malaria in Chabahar and Suburbs, Southeastern Iran: Insights from a Six-Year Study (2019 - 2024). Jundishapur J Microbiol. 2025; 18 (5): e161892. <https://doi.org/10.5812/jjm-161892>.

monsoon patterns and temperatures of 25 - 35°C create ideal conditions for *Anopheles* mosquito proliferation, exacerbating malaria transmission in both urban and rural areas. Cross-border migration from malaria-endemic countries like Pakistan and Afghanistan further complicates the situation.

2. Objectives

The present study aims to analyze malaria trends in Chabahar and its surrounding areas from 2019 to 2024, focusing on the impact of climatic factors, cross-border migration, and demographic determinants. By identifying key challenges and contributing factors, this research seeks to inform targeted strategies for malaria control and elimination in southeastern Iran. The prevalence of *Plasmodium vivax* in the region, combined with genetic factors like G6PD deficiency and thalassemia, further adds to the complexity of malaria management (4, 5).

3. Methods

3.1. Study Areas

Chabahar, located in the Sistan-Baluchistan province of southeastern Iran, was selected as the study area due to its high malaria burden and unique ecological and demographic characteristics (Figure 1). The district comprises three urban centers and six rural districts (Pirsohrab, Kambad-e Soleyman, Nowbandian-e Payyan, Bahoukalat, Sand-e-Mirsuban, and Polan), encompassing 446 inhabited villages. Situated on the northern coast of the Oman Sea, Chabahar's warm and humid climate provides ideal conditions for *Anopheles* species, the primary vectors of malaria in the region (6). The majority of the population is ethnic Baloch, with Balochi and Persian being the predominant languages, reflecting the area's cultural and linguistic diversity (7).

3.2. Study Design and Data Collection

This retrospective, cross-sectional study analyzed malaria cases in Chabahar and its surrounding areas from 2019 to 2024. Blood samples were collected from individuals presenting with malaria-like symptoms at health centers across the district. Malaria diagnosis was confirmed using Giemsa-stained blood smears and rapid diagnostic tests (RDTs) targeting *Plasmodium*-specific antigens (pLDH/HRP2). A standardized questionnaire was administered to collect demographic data, including age, sex, nationality, and travel history. Comprehensive malaria diagnoses were performed in multiple laboratories across Chabahar and its suburbs,

with data gathered through a computerized system. Demographic data, including gender, age, and occupation, were recorded annually based on census data. Meteorological data, including maximum temperature, minimum temperature, average temperature, humidity, and precipitation, were sourced from local weather stations and cross-verified with national meteorological agencies to ensure reliability. All data were aligned temporally and spatially to match the study region and period, ensuring accurate analysis of the relationship between malaria incidence, weather patterns, and demographic factors. Missing data were handled using multiple imputation techniques, and quality control measures such as outlier analysis and data validation were implemented to ensure data accuracy and integrity.

3.3. Statistical Analysis

The statistical analysis explored the relationship between parasite type and various factors, including month and average temperature. Data were analyzed using the statistical package for the social sciences (SPSS), version 26.0. An analysis of variance (ANOVA) was performed to determine significant differences between groups, followed by Spearman's correlation coefficients between environmental variables and type of parasitology to identify specific group differences. Results were reported as means \pm standard deviations to accurately convey the data distribution. To compare seroprevalence rates across different groups, chi-square and Fisher's exact tests were employed. A P-value of less than 0.05 was considered statistically significant, indicating a strong likelihood that the observed differences were not due to chance. This rigorous statistical approach ensured the validity and reliability of the findings, allowing for meaningful interpretations of the data in relation to the study's objectives.

4. Results

Over the six-year study period from 2019 to 2024, a total of 3,393 malaria cases were documented in Chabahar, Iran, representing one of the highest burdens in the Sistan-Baluchistan province. The number of infections peaked significantly in 2023, with 1,272 reported cases, likely due to favorable climatic conditions for mosquito breeding and increased cross-border migration (Figure 2). Figure 2 illustrates the distribution of malaria cases by species and year, highlighting the dominance of *P. vivax* in 2023 - 2024.

4.1. Annual Trends and Demographics



Figure 1. The geographical location of Chabahar city and its suburbs (Konarak, Qasr-e Qand, Nik Shahr, Dashtiari) on the map is shown at the bottom of the map.




Figure 2. Distribution of malaria cases by species of malaria and years in Chabahar city and its suburbs

Over the six-year study period, malaria incidence varied significantly, with the lowest number of cases

(199) recorded in 2021 and a marked peak in 2023 (1,272 cases). The annual parasite incidence (API), a key metric


Table 1. Malaria Epidemiologic Indices and Demographic Variables in Chabahar City of Sistan-Baluchistan Province in the Years 2019 - 2024 ^a

Indices	2019	2020	2021	2022	2023	2024	P-Value
Age	26.45 ± 14.16	30.46 ± 14.32	25.87 ± 15.17	29.42 ± 17.46	26.77 ± 14.87	25.62 ± 14.73	< 0.001
Gender							< 0.001
Male	180 (85.7)	178 (87.3)	148 (72.9)	678 (73.2)	1038 (81.6)	437 (75.6)	
Female	30 (14.3)	26 (12.7)	55 (27.1)	248 (26.8)	234 (18.4)	141 (24.4)	
Total population	216295	222985	228703	233785	238867	241949	-
API ^b	0.97	0.91	0.87	3.96	5.32	0.72	-
Parasites							< 0.001
<i>Plasmodium vivax</i>	163 (77.6)	184 (90.2)	2 (1.0)	493 (53.2)	1256 (98.7)	496 (85.8)	
<i>Plasmodium falciparum</i>	47 (22.4)	19 (9.3)	171 (84.2)	432 (46.7)	16 (1.3)	78 (13.5)	
Mixed	0 (0)	1 (0.5)	30 (14.8)	1 (0.1)	0 (0.0)	4 (0.7)	

Abbreviation: API, the annual parasite incidence.

^a Values are expressed as No. (%) or mean ± SD.

^b The annual parasite incidence is a crucial metric that illustrates the prevalence of malaria by indicating the number of cases per 1,000 population.

**Figure 3.** Comparison of environmental factors (temperature, humidity, and precipitation) across malaria parasite groups (mix, *Plasmodium falciparum*, and *Plasmodium vivax*)

for malaria prevalence, reached its highest in 2023 at 5.32 per 1,000 population before declining to 0.72 in 2024 (Table 1). The mean age of malaria patients fluctuated significantly, with a notable increase in 2022 to 29.42 years ($P < 0.001$). Gender distribution also varied, with male patients consistently comprising the majority, peaking at 81.6% in 2023 (Table 1).

4.2. Parasite Distribution

There was a notable shift in the prevalence of *Plasmodium* species. *Plasmodium vivax* dominated in

2023 (98.7%) and 2024 (85.8%), while *P. falciparum* was predominant in 2021 (84.9%). Mixed infections were rare but showed a significant association with higher temperatures ($P < 0.001$) (Table 1).

4.3. Changes in Parasite Species

4.3.1. Species Distribution

There was a notable shift in the prevalence of *Plasmodium* species. *Plasmodium vivax* dominated in 2023 (98.7%) and 2024 (85.8%), while *P. falciparum* was

Table 2. Spearman's Correlation Coefficients Between Environmental Variables and Type of Parasite, Chabahar City

Factors	Correlation Coefficient	Significance (2-tailed)	N
Max. temperature	0.024	0.163	3393
Min. temperature	0.048 ^a	0.005	3393
Average temperature	0.039 ^b	0.024	3393
Average humidity (%)	0.024	0.156	3393
The amount of precipitation	0.025	0.139	3393

^a Correlation is significant at the 0.05 level (2-tailed).

^b Correlation is significant at the 0.01 level (2-tailed).

predominant in 2021 (84.9%). This change was statistically significant ($P < 0.001$), indicating a shifting epidemiological landscape (Table 1).

4.3.2. Temperature Effects

Analysis revealed significant correlations between temperature and malaria transmission. Higher temperatures were strongly associated with increased incidence, particularly for mixed infections ($P < 0.001$). Mean temperatures for mixed infections were notably higher (33.631°C) compared to *P. falciparum* (31.563°C) and *P. vivax* (31.723°C), suggesting a robust association between temperature fluctuations and transmission dynamics (Figure 2).

4.4. Analysis of Meteorological Variables by Malaria Type

Figure 3 provides descriptive statistics and P-values for various meteorological variables across different types of malaria, including *P. falciparum*, *P. vivax*, and mixed infections. An analysis of key climatic factors – maximum temperature, minimum temperature, average temperature, average humidity, and precipitation – is presented across four groups: Mixed infections, *P. falciparum*, *P. vivax*, and total cases. For each variable, we report the sample size ($n = 2$), mean values, standard deviations, 95% confidence intervals (CIs) for the means, and associated P-values (Table 2).

4.4.1. Humidity and Precipitation

Average humidity and precipitation showed no significant impact on malaria transmission ($P = 0.19$ and $P = 0.50$, respectively), indicating that temperature is the primary climatic driver of malaria incidence in Chabahar (Figure 3).

4.4.2. Temperature Measurements

Significant statistical differences in maximum, minimum, and average temperatures were observed in

the mixed infections group, as all reported P-values were < 0.001 . This finding underscores a robust association between higher temperature levels and increased malaria incidence. This suggests that fluctuations in temperature could critically influence malaria transmission dynamics. Conversely, the data for *P. falciparum* and *P. vivax* exhibit no significant temperature relationships, indicating potential differences in the ecological or biological responses to climatic variations.

4.5. Key Implications

4.5.1. Temperature

Higher temperatures are strongly associated with mixed malaria infections, suggesting temperature fluctuations critically influence transmission dynamics.

4.5.2. Variability

Mixed infections exhibited more consistent exposure to higher temperatures (lower standard deviations) compared to *P. vivax*, potentially affecting transmission patterns.

4.5.3. Public Health

Temperature monitoring should be prioritized in malaria control strategies, as humidity and precipitation showed minimal impact. Predictive models based on temperature data could enhance proactive interventions.

4.5.4. Research Needs

Further studies are needed to explore the ecological and biological responses of *P. falciparum* and *P. vivax* to temperature variations, particularly threshold effects on parasite lifecycles.

In conclusion, temperature is a key climatic factor driving malaria transmission in Chabahar, with mixed infections showing distinct responses. Public health strategies should focus on temperature-based predictive models and interventions, while further research is needed to understand species-specific climatic adaptations.

4.5.5. Gender

Over time, there were fluctuations in the proportion of male and female malaria patients. Male patients fluctuated between 72.9% in 2021 and 81.6% in 2023, while female patients fluctuated between 12.7% in 2020 and 27.1% in 2021. The P-value for the gender variable is $P < 0.001$, indicating a statistically significant variation in the gender distribution of malaria patients across different years (Figure 4).

4.6. Malaria Cases by Job Type and Type of Parasite

This document presents a bar chart that illustrates the distribution of malaria cases by job type and type of parasite. The data reflect different categories of employment and the prevalence of *P. falciparum*, *P. vivax*, and mixed infections. The total number of cases for each job category is also shown to provide context. Figure 5 provides information about the distribution of occupational categories among people who tested positive for the parasitic species of malaria.

Plasmodium falciparum shows the highest count in autumn, with a significant increase compared to spring and summer. The count remains high in winter (Figure 6). *Plasmodium vivax* exhibits a similar pattern to *P. falciparum*, increasing from spring to autumn. It remains consistently high in winter as well, with very similar numbers in autumn. Mixed infections occur less frequently across all seasons, peaking in summer with a noticeable drop in other seasons.

4.7. Analysis of Parasite Types by Nationality

In our study, we conducted a crosstabulation of malaria parasite types with the nationality of infected individuals. The data reveal significant differences in the prevalence of specific malaria parasites among Iranian, Pakistani, and Afghan populations.

4.7.1. Total Cases

The total number of malaria cases recorded in this study was 3,393, with the majority of infections identified as *P. vivax* (2,594 cases), followed by *P. falciparum* (763 cases) and mixed infections (36 cases).

4.7.2. Iranian Nationals

Among the Iranian population, *P. vivax* was the most prevalent type, with 1,677 cases. This indicates the enduring transmission dynamics of this species in Iran, which is often associated with relapsing malaria. *Plasmodium falciparum* cases were also significant, totaling 510, suggesting that severe malaria remains a concern in certain regions.

4.7.3. Pakistani Nationals

The data for Pakistani nationals show a considerable incidence of both *P. falciparum* (235 cases) and *P. vivax* (804 cases), indicating a diverse burden of malaria infections. The presence of mixed infections (12 cases) among this group also raises concerns regarding the complexities of treatment and potential resistance issues.

4.7.4. Afghan Nationals

The Afghan population showed the lowest total count, with 135 cases. The distribution revealed a stark predominance of *P. vivax* (113 cases) compared to *P. falciparum* (18 cases), which may reflect geographical factors influencing malaria transmission in Afghanistan.

Figure 7 shows the chart depicting the variations in malaria cases by different parasite types – trophozoite, schizont, gametocyte, and mixed infections – which is a crucial tool for understanding the epidemiological landscape of malaria in Chabahar city over five years. By categorizing the data into specific stages of the parasite lifecycle, the graph allows for a detailed analysis of the prevalence and distribution of *P. vivax* and *P. falciparum*, the two primary malaria-causing parasites in the region.

Table 3 illustrates the distribution and percentage of different parasite types (mix, *P. falciparum*, and *P. vivax*) among collected samples based on age, nationality, previous history of malaria, and travel destination (no travel, inside the country, and abroad). The data shows the highest percentage of *P. vivax* parasites among those who traveled abroad (35.8%), while the lowest is for the mixed type in domestic travelers (0.1%).

4.7.5. Inside the Country

Plasmodium vivax is predominant at 12.5%, but there is also a notable presence of *P. falciparum*. *Plasmodium vivax* remains the highest at 35.8%, with significant *P. falciparum* presence. *Plasmodium vivax* is consistently the most prevalent type across all travel categories. Traveling abroad shows the highest total count and




Figure 4. Distribution of malaria cases by genus and year




Figure 5. Distribution of job categories among individuals testing positive for malaria parasites

percentage for all parasite types, indicating perhaps a higher risk or exposure during international travel.

5. Discussion

The epidemiological trends observed in Chabahar reflect a dynamic interplay of climatic, demographic, and cross-border factors that drive malaria transmission in southeastern Iran. The resurgence of *P. vivax* in 2023-2024 (98.7% and 85.8% of cases, respectively; (Table 1) contrasts starkly with the predominance of *P. falciparum* in 2021 (84.9%), underscoring ecological and operational influences. The dominance of *P. vivax* may stem from its

ability to form hypnozoites, which enable relapses even during suboptimal transmission periods (8). This challenge is exacerbated by limited access to primaquine – a critical hypnozoitocidal drug – in Sistan-Baluchistan, where G6PD deficiency affects approximately 15% of the population (3). Conversely, the decline in *P. falciparum* after 2022 likely reflects the success of artemisinin-based combination therapies (ACTs), which disproportionately target *P. falciparum* due to its shorter extrinsic incubation period (1).

The robust association between temperature and malaria incidence, particularly for mixed infections (mean: 33.6°C; $P < 0.001$; (Figure 3), aligns with




Figure 6. Distribution of malaria cases by species of malaria and season

Paaijmans et al. (9), who demonstrated that warmer temperatures accelerate *Plasmodium* development in *Anopheles* vectors. However, the lack of humidity and precipitation effects diverges from tropical studies (10), where rainfall sustains breeding sites. In Chabahar's arid monsoon climate, ephemeral water bodies from irregular rainfall may desiccate before sustaining larvae, rendering temperature the dominant driver. The interplay between ephemeral water bodies and temperature plays a crucial role in the survival and development of aquatic larvae. Irregular rainfall patterns lead to the formation of temporary water bodies that may desiccate before larvae can complete their life cycles, making temperature a critical factor influencing their survival (11, 12).

However, the variability in rainfall patterns can lead to prolonged periods of drought, resulting in the formation of ephemeral water bodies. These water bodies are temporary habitats that form after rainfall events but are subject to rapid desiccation due to high temperatures and evaporation rates (13). Ephemeral water bodies are crucial habitats for various aquatic organisms, including insect larvae, amphibian larvae, and crustacean larvae. These habitats provide breeding grounds and resources for larval development. However,

the transient nature of these water bodies poses a significant challenge to the survival of larvae. If the water body dries up before the larvae reach a certain developmental stage, they may not survive (14).

Gender disparities, with males constituting 72.9 - 81.6% of cases (Table 1), mirror trends in agrarian communities where occupational exposure (e.g., outdoor labor, fishing) elevates bite risk. Similar patterns were observed by Raasti et al. in Baluchistan, where socio-cultural norms limit female outdoor activity (15). However, the rise in female cases in 2021 (27.1%) may reflect increased indoor transmission during COVID-19 lockdowns, a phenomenon documented in sub-Saharan Africa (16).

Cross-border migration emerged as a critical driver of transmission, with Afghan nationals disproportionately affected by *P. vivax* (83% of cases; Table 3). This aligns with Xiaohong Li et al. (14), who identified migration routes from Afghanistan as hotspots for *P. vivax* reintroduction. The high *P. falciparum* burden among Pakistani travelers (235 cases) likely reflects chloroquine resistance in Pakistan's Sindh province (8), underscoring the WHO's call for cross-border collaboration to disrupt parasite reservoirs (1).




Figure 7. Parasite stages of malaria seen in microscopic examination in Chabahar (2019 - 2024). This graph illustrates the variation in malaria cases by different parasite types (trophozoite, schizont, gametocyte, and mixed infections) observed during microscopic examinations from 2019 to 2024. The data highlights the temporal trends of *Plasmodium vivax* and *Plasmodium falciparum* infections over these six years. The abbreviations used for the stages of the parasite are T, trophozoite; V, *P. vivax*; S, schizont; G, gametocyte; F, *P. falciparum*, and mix: Infection of both *P. vivax* and *P. falciparum*; T.V: The trophozoite *P. vivax* parasitic stage.

The predominance of trophozoite-stage *P. vivax* in microscopic analyses (Figure 6) highlights challenges in diagnosing relapsing malaria. Hypnozoite activation often evades routine surveillance, perpetuating transmission (17). Despite this, the low recurrence rate (5.1% with prior malaria history; Table 3) suggests either underdiagnosis or effective radical cure adherence, warranting pharmacovigilance (18).

5.1. Conclusions

Chabahar's malaria landscape is shaped by temperature-driven transmission, gendered exposure, and cross-border migration. Elimination requires climate-informed strategies, gender-sensitive interventions, and regional partnerships.

5.2. Limitations and Future Directions

While this study provides critical insights, passive surveillance likely underestimates asymptomatic carriers, who account for approximately 80% of transmissions in endemic regions. The absence of molecular data on parasite resistance (e.g., *dhfr/dhps* mutations) limits insights into treatment efficacy. Future studies should integrate genomic surveillance, as demonstrated in Southeast Asia, to track resistance and hypnozoite reservoirs.

Acknowledgements

We want to thank the Vice Chancellor for Research and Technology at Zabol Medical University, Chabahar Medical Center, and the dedicated health professionals and support staff at Chabahar for their support and cooperation in our research on malaria. Additionally, we

Table 3. Distribution and Percentage of Different Parasite Types

Categories	Type of Parasite			Total	P-Value
	Mix	<i>Plasmodium falciparum</i>	<i>Plasmodium vivax</i>		
Travel					< 0.001
No	3 (8.3)	230 (30.1)	594 (36.8)	1187 (35.0)	
Inside the country	3 (8.3)	128 (16.8)	425 (16.4)	3 (8.3)	
Abroad	30 (83.4)	405 (53.1)	1215 (46.8)	1650 (48.6)	
Nationality					0.021
Iranian	20 (55.6)	510 (66.8)	1677 (64.6)	2207 (65.0)	
Pakistani	12 (33.3)	235 (30.8)	804 (31.0)	1051 (31.0)	
Afghan	4 (11.1)	18 (2.4)	113 (4.4)	135 (4.0)	
Previous history of malaria					0.32
No	35 (97.2)	731 (95.8)	2453 (94.6)	3219 (94.9)	
Yes	1 (2.8)	32 (4.2)	141 (5.4)	174 (5.1)	
Total	36 (100.0)	763 (100.0)	2594 (100.0)	3393 (100.0)	

want to thank the front-line workers, the entire team at Chabahar, and the community members for their contributions. We are deeply grateful for the crucial support of the regional health authorities in Chabahar city and the surrounding areas, whose endorsement and facilitation have propelled our research endeavors forward.

Footnotes

Authors' Contribution: E. E. E. K. conducted extensive field research in Chabahar city and its surroundings to gather crucial epidemiological data and drafted the introduction and results sections, thoroughly reviewing the manuscript content. M. R. K. Zh. coordinated field study operations and played a key role in formulating the research design, specifically focusing on data collection methodologies, assisted in interpreting epidemiological data and participated actively in discussions regarding regional health policies and approved the final manuscript for publication; B. F. assisted in interpreting epidemiological data and participated in discussions about regional health policies and contributed significantly to gathering essential epidemiological data. Sh. D. conducted extensive field research in Chabahar city and its environs, collecting critical epidemiological data pivotal to the study, analyzed the gathered data, employing statistical tools to discern patterns relevant to the malaria incidence over the observed period; drafted initial sections of the manuscript, particularly the introduction and results, and revised the document for critical content and provided final approval of the

version to be published. M. D. (corresponding author) led the project and supervised the study design, execution, and coordination, condensed research findings, offered critical insights for data interpretation, and led the manuscript writing process, incorporating contributions from co-authors while maintaining coherence and academic integrity; and served as the guarantor for the work, provided final approval for publication, and addressed any post-publication issues, ensuring the accuracy and integrity of the study.

Conflict of Interests Statement: The authors declare no conflict of interests.

Data Availability: Data is gathered separately as an Excel file.

Ethical Approval: Ethics approval for the study was obtained from the relevant Ethical Review Committee ([IR.ZBMU.REC.1403.077](#)). This approval was crucial in guiding our research practices and ensuring compliance with ethical guidelines.

Funding/Support: The present study received no funding/support.

Informed Consent: Consent was obtained from all participants regarding the publication of data collected during the study. Any data or images that could potentially identify participants were handled with strict confidentiality and published only after securing explicit consent. The participants were informed about the nature of the data being published and the intended audience, and they agreed to the dissemination of their anonymized information within the scientific community.

References

- World Health Organization. *World malaria report 2023*. Geneva, Switzerland: World Health Organization; 2023. Available from: <https://www.who.int/teams/global-malaria-programme/reports/world-malaria-report-2023>.
- Adam M, Nahzat S, Kakar Q, Assada M, Witkowski B, Tag Eldin Elshafie A, et al. Antimalarial drug efficacy and resistance in malaria-endemic countries in HANMAT-PIAM_net countries of the Eastern Mediterranean Region 2016-2020: Clinical and genetic studies. *Trop Med Int Health*. 2023;**28**(10):817-29. [PubMed ID: 37705047]. <https://doi.org/10.1111/tmi.13929>.
- Babaie J, Barati M, Azizi M, Ephtekharian A, Sadat SJ. A systematic evidence review of the effect of climate change on malaria in Iran. *J Parasit Dis*. 2018;**42**(3):331-40. [PubMed ID: 30166779]. [PubMed Central ID: PMC6104236]. <https://doi.org/10.1007/s12639-018-1017-8>.
- Brandt J. *Thalassemia and the protection against malaria*. 2024. Available from: https://dc.ewu.edu/srcw_2024/ps_2024/pi_2024/11/.
- Danaei M, Dabirzadeh S, Dabirzadeh M. Evaluation of G6PD deficiency in malaria patients in the south-east of Iran. *Ann Parasitol*. 2022;**68**(2):247-56. [PubMed ID: 35809480]. <https://doi.org/10.17420/ap6802.430>.
- Tavakoli M, Mokhtari Karchegani A. Analysis of the efficiency of crisis management of local communities in the face of floods due to climate change: Five selected villages of port Chabahar County, Iran. *Int J Coastal, Offshore Environ Engin (ijcoe)*. 2024;**9**(1):67-78.
- Jahani C. State control and its impact on language in Balochistan. *Role State West Asia*. 2005;**14**(1):151.
- Li X, Snow RW, Lindblade K, Noor AM, Steketee R, Rabinovich R, et al. Border malaria: Defining the problem to address the challenge of malaria elimination. *Malar J*. 2023;**22**(1):239. [PubMed ID: 37605226]. [PubMed Central ID: PMC10440889]. <https://doi.org/10.1186/s12936-023-04675-3>.
- Paaijmans KP, Blanford S, Chan BH, Thomas MB. Warmer temperatures reduce the vectorial capacity of malaria mosquitoes. *Biol Lett*. 2012;**8**(3):465-8. [PubMed ID: 22188673]. [PubMed Central ID: PMC3367745]. <https://doi.org/10.1098/rsbl.2011.1075>.
- Ghanbarnejad A, Turki H, Yaseri M, Raeisi A, Rahimi-Foroushani A. Spatial modelling of malaria in South of Iran in line with the implementation of the malaria elimination program: A bayesian poisson-gamma random field model. *J Arthropod Borne Dis*. 2021;**15**(1):108-25. [PubMed ID: 34277860]. [PubMed Central ID: PMC8271232]. <https://doi.org/10.18502/jad.v15i1.6490>.
- Bahmanzadegan Jahromi A, Ezam Mojtaba, Lari Kamran, and Ali Akbari Bidokhti AA. An investigation of the hydrography of Chabahar bay using FVCOM model and EOF analysis. *Marine Geodesy*. 2022;**45**(4):360-79. <https://doi.org/10.1080/01490419.2022.2042434>.
- Sedgh ZK, Fakhrian M. Changes in the biochemical composition of sargassum ilicifolium before and after the monsoon in chabahar bay, Gulf of Oman, Iran. *Thalassas: Int J Marine Sci*. 2022;**39**(1):1-6. <https://doi.org/10.1007/s41208-022-00495-5>.
- Liu Y, Wu C, Jia R, Huang J. An overview of the influence of atmospheric circulation on the climate in arid and semi-arid region of Central and East Asia. *Sci China Earth Sci*. 2018;**61**(9):1183-94. <https://doi.org/10.1007/s11430-017-9202-1>.
- Nag D, Phartiyal B, Agrawal S, Kumar P, Sharma R, Kumar K, et al. Westerly-monsoon variations since the last deglaciation from semi-arid Ladakh region, Trans Himalaya, India. *Palaeogeography, Palaeoclimatol, Palaeoecol*. 2023;**618**. <https://doi.org/10.1016/j.palaeo.2023.111515>.
- Raasti A, Nasir O, Khalid MA, Zafar S, Khan W, Nadeem SF. Evaluating malaria prevalence across different age and gender groups in peshawar through light microscopic analysis. *J Health Rehabil Res*. 2024;**4**(2):767-71. <https://doi.org/10.61919/jhrr.v4i2.952>.
- Ansumana R, Sankoh O, Zumla A. Effects of disruption from COVID-19 on antimalarial strategies. *Nat Med*. 2020;**26**(9):1334-6. [PubMed ID: 32807937]. <https://doi.org/10.1038/s41591-020-1047-5>.
- Anwar MN, Hickson RI, Mehra S, Price DJ, McCaw JM, Flegg MB, et al. Optimal interruption of *P. vivax* malaria transmission using mass drug administration. *Bull Math Biol*. 2023;**85**(6):43. [PubMed ID: 37076740]. [PubMed Central ID: PMC10115738]. <https://doi.org/10.1007/s11538-023-01153-4>.
- Talisuna AO, Staedke SG, D'Alessandro U. Pharmacovigilance of antimalarial treatment in Africa: Is it possible? *Malar J*. 2006;**5**:50. [PubMed ID: 16780575]. [PubMed Central ID: PMC1523354]. <https://doi.org/10.1186/1475-2875-5-50>.