

# Expression, Purification and Evaluation of Antigenicity of CagA Antigenic Fragment of *Helicobacter pylori*

Vahideh Farjadi<sup>1</sup>, Hamid Abtahi<sup>2,\*</sup>, Mohammad Reza Zolfaghari<sup>1</sup>, Safieh Soufian<sup>3</sup>, Leila Hasanzadeh<sup>4</sup>

<sup>1</sup>Department of Microbiology, Faculty of Science, Islamic Azad University, Qom Branch, Qom, IR Iran

<sup>2</sup>Molecular and Medicine Research Center, Arak University of Medical Sciences, Arak, IR Iran

<sup>3</sup>Department of Biology, Payame Noor University, Arak, IR Iran

<sup>4</sup>Department of Biotechnology and Microbiology, Arak University of Medical Sciences, Arak, IR Iran

\*Corresponding author: Hamid Abtahi, Molecular and Medicine Research Center, Arak University of Medical Sciences, Arak, IR Iran. Tel: +98-8614173502. Fax: +98-8614173526. E-mail: abtahi@arakmu.ac.ir.

Received: July 22, 2012; Revised: December 8, 2012; Accepted: January 12, 2013

**Background:** *Helicobacter pylori* is a human pathogen that causes chronic gastritis, which plays role in gastric and duodenal ulcers, is also involved in gastric carcinogenesis and may be regarded as a possible important factor in at least a subset of patients with functional dyspepsia.

**Objectives:** This study was aimed to construct a recombinant protein containing *H. pylori* antigenic CagA region and determine its antigenicity as a vaccine candidate against *H. pylori*.

**Materials and Methods:** The antigenic region of CagA gene was detected by bioinformatics techniques. In this study, the *H. pylori* antigenic CagA region was amplified by PCR and sub-cloned to prokaryotic expression vector pET32a. *Escherichia coli* BL21 (DE3) pLysS was transformed with pET32a-CagA and gene expression was induced by IPTG. The expressed protein was purified by affinity chromatography using Ni-NTA resin. The integrity of the product was confirmed by western-blot analysis using Sera of infected individual. Finally antigenicity was studied by western-blot analysis using human sera infected with *H. pylori*.

**Results:** Enzyme digestion analysis, PCR and sequencing showed that the target gene (1245 bp) was correctly inserted into the recombinant vector. The expressed protein was purified using affinity chromatography by Ni-NTA resin. The data also indicated that CagA protein from *H. pylori* detected from patients' sera.

**Conclusions:** Results indicates that antigenic region of recombinant CagA protein were recognized as an antigen, so it might be a candidate for the development of *H. pylori* vaccine, ELISA kit designs and serological diagnosis of *H. pylori* infections.

**Keywords:** Antigenic Region; Antigenicity; Cytotoxin-Associated Gene (cagA); *Helicobacter pylori*

## 1. Background

*Helicobacter pylori* is a Gram negative, microaerophilic spiral bacterium that colonizes in the human stomach and induces gastric mucosal inflammations that are linked to severe gastritis-associated diseases including atrophic gastritis, peptic ulcer and gastric cancer (1). *H. pylori* is detected in half of the world's population. In 1994, the World Health Organization and International Agency for Research on Cancer categorized *H. pylori* infection as a group I carcinogens (2). An important virulence factor of *H. pylori* is a secreted toxin known as Cytotoxin-associated gene A (CagA) which is located at the end of an approximately 40-kb cluster of genes called cag pathogenicity island (PAI).

Cag PAI encodes a type-IV secretion system (3) and

transfers CagA protein into the intracellular region of gastric epithelial cell, inducing cellular hyper proliferation, apoptosis and leading to failure of gastric epithelial cell in maintaining its normal cytoskeletal structure (4). Also increases the secretion of cytokines such as tumor necrosis factor (TNF) and interleukin (IL) 1, 6 and 8 by the epithelial cells. The IL-8 leads to intense inflammatory responses and stimulates the production of oxygen free radicals that could cause DNA damage to adjacent cells. Accumulation of oxidative DNA damage could lead to genetic modifications of gastric epithelial cells which are carcinogenic (5). A large number of studies have proven the association between CagA positive strains and gastritis and peptic ulcer. Also it may induce the persistent systemic inflammatory responses and the increased risk of recurrent atherosclerotic events (6).

### Implication for health policy/practice/research/medical education:

Currently, the main treatment to eliminate *H. pylori* is combined antibiotics therapy. However, due to incomplete eradication of *H. pylori* through antibiotic treatment, the cost of combination therapy, re-infection, patients' complaints and emerges of antibiotic resistance strains, vaccination could be suggested as a preventive approach. The CagA has been approved as an effective virulence factor involved in the pathogenesis of *H. pylori* infection. Thus, this protein can be considered as a vaccine candidate.

Copyright © 2013, Ahvaz Jundishapur University of Medical Sciences; Published by Kowsar Corp. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

## 2. Objectives

By using major epitopes of each antigen instead of the whole antigen, the specificity of fragment will increase and more specific immune responses are stimulated. Isolation, characterization and expression of *H. pylori* antigenic region of *CagA* (*arCagA*) gene in *E. coli* as a host, was presented in this paper. We also showed that recombinant *CagA* protein was recognized by infected human sera using Western blot analysis.

## 3. Materials and Methods

### 3.1. Strains, Plasmids, Media and Sera

*E. coli* strain DH5 $\alpha$  (Strata gene) (*f* - *gyr* A96 *Nalr*; *recA1 relA1 Thi-1 hsdR17 r-k m+k*) were used for initial cloning and maintenance of different DNA fragments. To produce the recombinant protein, prokaryotic expression vector pET32a (Novagene) was used. This vector can express a fusion protein with a six-histidine tag, a thrombin recognition site and a T7 tag at the N-terminus. These additional amino acids increase the size of expressed protein to about 20 kDa. The recombinant pET32a (pET32a-CagA) is transformed into *E. coli*, BL21 (DE3) pLysS (*f* - *ompHsdB*, *rB* - *mB* -, *dcm gal*, *DE3*, *pLysScmr*). The required antibiotics were added to LB media according to the reference recommendation (7). All chemicals were purchased from Merck (Germany) and all of the enzymes were provided from Fermentas (Lithuania) or Cinagen (Iran).

Fifteen patients (8 male, 7 female) referred to the Vali-Asr Hospital (a teaching university hospital in Arak, Iran) from May to October 2011 for diagnostic upper gastrointestinal (UGI) endoscopy, were prospectively enrolled in this study. Patients were considered *H. pylori*-positive when the results of RUT were positive. 5-10 cc blood sample was taken from each patient after red blood cell separation (3000 rpm for 5 minutes). Sera were collected and frozen at -20°C.

*H. pylori* was isolated from a biopsy sample of a dyspeptic patient undergoing endoscopy, who granted informed consent. Two biopsy samples were processed for cultivation on 5% packed cell *Columbia agar* containing 2-4% fetal calf serum (Oxoid-Basingstoke, UK) and Skirrow's antibiotic supplements (Oxoid, UK). The plates were incubated at 37°C for 5-7 days in microaerophilic conditions. The isolates were identified as *H. pylori* by urease, catalase and oxidase reactions and Gram staining (8). Molecular identification was confirmed by PCR amplification of the conserved *ureC* gene of *H. pylori* as described (9).

**Antigenic regions:** To find out the antigenic region of *CagA*, the sequence of *CagA* gene (NCBI GenBank, gi: 259123369) was submitted to ABCpred, Bcepred and Emboss Antigenic web servers. Two specific Primers were designed base on the sequences with OLIGO5 software. The forward primer (5' TATGATCCACAATAACGCTC 3') starts from the beginning of the gene and contain *BamHI* site.

Reverse primers (5'ATTCTCGAGTTCATCAAAAGATT 3') contain recognition site for *XhoI*. The restriction enzyme sites (underlined) were added to the primers for subsequent cloning procedure. These two primers were designed to amplify a 1245 bp fragment of the *CagA* coding region.

### 3.2. Chromosomal DNA Isolation

Genomic DNA was extracted from cultured samples according to the standard CTAB/NaCl protocol (7, 10). Then, the pellet of 1.5 mL bacterial culture was resuspended overnight in TE buffer (Tris 10 mM, EDTA 1 mM, and pH 8), the bacterial cell was lysed by sodium dodecyl sulphate (SDS 10%) and proteinase K (20 mg/mL), the chromosomal DNA was extracted by CTAB/NaCl solution (10%CTAB and 0.7 M NaCl). Cell debris and proteins were removed by mixing and genomic DNA was extracted with two times washing with phenol/chloroform/isoamylalcohol (25:24:1) mixture. The tubes were centrifuged at 10000 rpm for 10 minutes at 4°C and the supernatants were transferred to a fresh tube. In this phase, protein was deposited and RNA and DNA remained in aqueous solution. DNA was precipitated by isopropanol and washed with 70% ethanol and dissolved in TE buffer and stored at -20°C.

### 3.3. Gene Amplification

PCR was performed in a 50  $\mu$ L total volume containing 500 ng of template DNA, 1  $\mu$ M of each primer, 2.5 mM Mg<sup>2+</sup>, 200  $\mu$ M (each) deoxynucleoside triphosphates, 1 $\times$  PCR buffer and 2.5 unit of pfu DNA polymerase (Fermentas, Lithuania). The following conditions were used for amplification: hot start at 94°C for five minutes, followed by thirty cycles of denaturation at 94°C for one minute, annealing at 50°C for one minute and extension at 72°C for one minute. The program followed by a final extension at 72°C for five minutes. The PCR product was analyzed by electrophoresis in 1% agarose gel in 1 $\times$  TBE buffer and visualized by ethidiumbromide staining on UV transilluminator (11).

### 3.4. Cloning of *arCagA* Gene in Bacterial Expression Vector

The PCR product was purified from the agarose gel with high pure PCR product purification kit (Roche, Germany). The PCR product was digested with *BamHI* and *XhoI* and ligated to pET32a, which were digested by the same restriction enzymes, using T4 DNA ligase at 16°C overnight. Moreover *Escherichia coli* DH5 $\alpha$  and *E. coli* BL21 (DE3) pLysS competent cells were prepared by calcium chloride method and were used for transformation of plasmid (7).

### 3.5. Expression and Purification of Recombinant *CagA*

The pET32a-*arCagA* plasmids was transformed into *E.*

*coli* BL21 (DE3) pLYS according to the standard method (7) and grown on 2 mL Luria Bertani (LB) Broth supplemented with Ampicillin (100 µg/mL) at 37°C with agitation. A colony containing the recombinant plasmid was incubated on shaking incubator for overnight at 37°C in 2 mL LB medium containing 100 µg/mL Ampicillin. For expression of recombinant protein, 500 µL of culture was added in 50 mL LB broth (per liter contains: 10 g yeast extract (Difco, USA), 20 g Bacto tryptone broth (Difco, USA), 0.2% (mass/vol) glucose, 10 g NaCl, 1 g KCl, 0.5 g MgCl<sub>2</sub>, 0.5 g CaCl<sub>2</sub>, 100 µg/mL Ampicillin), incubated at 37°C and 200 rpm shaking with vigorous agitation until the optical density reached 0.6 at 600 nm. Expression of the CagA protein was then induced by the addition of Isopropyl thio β-D-galactosidase (IPTG) to a final concentration of 1 mM and incubation was continued for the next 4 hours.

The expressed protein was purified by Ni-NTA agarose resin affinity chromatography according to manufacture instruction (Qiagen, USA). The quality and quantity of purified recombinant arCagA protein was analyzed by SDS polyacrylamide gel electrophoresis (SDS-PAGE 12%) and Bradford methods, respectively (7). In order to analyze the cross-reaction between fused segments (20 kDa) of recombinant arCagA protein (in pET32a) with patient sera, an *E. coli* BL21 (DE3) pLYS containing pET32a vector was induced by IPTG as a control.

### 3.6. Antigenicity of Recombinant arCagA

The integrity of the product was confirmed by western blot analysis. Serum from an *H. pylori* sero-positive patient (1:100 dilution), CagA antibody and normal human

sera were used as primary antibody. Western blotting was performed according to standard protocol (12) using normal human sera and CagA antibody as negative and positive control, respectively. For western blot analyses, 0.5 µg purified recombinant arCagA protein was used per well. The gel was blotted on to poly vinylidenedifluoride (PVDF) Membrane (Roche, Germany) using transfer buffer containing 25 mM Tris (pH = 8.3), 192 mM glycine and 20% methanol at 90 V for 3 hours at 4°C. The blotted membrane was blocked with 2.5% (w/v) BSA in TBS buffer (0.5 M NaCl, 0.02 M Tris pH 8.5, 0.05% tween 20) for 1 hour at room temperature. Membranes were incubated for 2 hours at room temperature with diluted sera 1:100 from patient's sera, normal sera and CagA antibody. After reaction with the primary antibody, the membranes were washed three times with TBS and incubated with anti-human IgG (Bioscience, USA) in 1:1000 dilution of TBS.

Then washed three times with TBS and reactions were developed by Diamino Banzidine (DAB) solution (Roche, Germany). As a negative control, the bacterial lysate from induced *E. coli* BL21 (DE3) pLYS containing pET32a vector was analyzed by Western blot.

## 4. Results

### 4.1. Antigenic Regions

According to the results of three data banks which were in accordance, amino acids sequence of 886 till 2131 was selected as regions with high antigenic properties (Figure 1).

```

YGSTITLYLLCQWGVHSNIEPEKVSLLYGDNGGPEARHDWNATVGYK
DQQGSNVATLINAHLYNGSGLIAGNEDGIKNPSFYLYKEDQLTGLKQ
AMSQEEIQNKVDFLEFLAKNNAKLDNLSEKEKEKFQTEIENFQKDRKA
YLDALGNDHVAFAVSKKDQKHLALITEFGNGEVSYTLKDYGKKQDKLL
DGEVKTTLQGSLLKYDGMFVDYSNFKYTNASKSPDKGVSATNGVSHL
EANFSKVAVFNLPNLLNLAITNYIRDLKEDKLWAKGLSPQEAANKLIKD
FLNSNKELVVKVSFNKAVAEAKNTGNYDEVKKAQKDLEKSLRKREH
LEKEVAKKLESRIDNKNRMEAKAQANSQKDKIFALINKEASKEARAAA
FDPNLKGIRSELSDKLENINKNLKDFGKSFDELEN

```

**Figure 1.** The Result of Antigenic Fragments

### 4.2. DNA Amplification and Molecular Characterization of arCagA Gene


The chromosomal DNA of *H. pylori* was extracted for amplification of arCagA gene. The amplified fragment had the expected size of 1245 bp comparing to 100 bp DNA ladder (Fermentas, Lithuania).

### 4.3. PCR Product Analysis and Cloning

The recombinant plasmid (pET32a-arCagA) was sequenced by standard primers and dideoxy chain ter-

mination method (Figure 2).

The sequencing result was confirmed by comparing databases and using basic local alignment search tool (BLAST) software (data not shown). Data showed that, PCR product was identical to CagA gene (NCBI GenBank, gi: 259123369). To confirm the transformation of pET32a-arCagA into *E. coli* BL21(DE3) pLYS, PCR reaction and enzymatic digestion with *Bam*HI and *Xho*I were performed and showed that the target gene was inserted correctly into it.



**Figure 2.** Schematic Representation of the PET32a and rCagA

#### 4.4. Expression and Purification of Recombinant Protein


pET32a-arCagA which was constructed by insertion of 1245 bp fragment of *CagA* gene (encoding a ~66 kDa mature peptide) into the pET32a (expression vector) was introduced into *E. coli* BL21 (DE3) pLYsS and the expression of recombinant protein was induced by IPTG. SDS-PAGE analysis of induced cells showed a band with desired molecular weight of 65 kDa. Due to the presence of 6x His-tag at the N-terminal region of *arCagA*, we could purify the recombinant protein using Ni-NTA resin affinity chromatography (Figure 3) and we obtained ~13% purity of *arCagA* expressed protein in *E. coli*. The concentration of recombinant protein was assayed after purification and calculated to 470 mg/L of the initial culture.

#### 4.5. Immunoblotting Analysis in Infected Human

To determine the antigenicity of recombinant antigenic region of CagA protein in patients' sera, identity confirmation of antigenic region r-CagA was performed via Western blotting and the desired band was observed with an estimated molecular weight to f 65 kDa. Figure 4 shows the specific interaction between patients' serum, human normal sera and CagA antibody with purified recombinant *arCagA* protein. Human normal sera were used as a negative control and CagA antibody used as a positive control.


There was no reaction between the expressed pET32a in *E. coli* BL21 (DE3) pLYsS and patient serum (data not shown).

**Figure 3.** Expression of Recombinant Antigenic Fragment of CagA Protein and It's Purification.



Lane 1: Protein marker (Fermentas, Lithuania), Lane 2: pET32a-CagA before induction, Lane 3: pET32a-CagA after 4 hours induction, Lane 4: purification of recombinant fragment of CagA protein.

**Figure 4.** Western Blot Analysis of CagA Recombinant Antigenic Fragment Using Patient Sera.



Lane 1: Protein marker (Fermentas, Lithuania), Lane 2: Western blotting by patient's sera, Lane 3: Western blotting by Human normal sera (negative control), Lane 4: Western blotting by CagA antibody (positive control).

### 3. Discussion

In this study, we have shown that recombinant antigenic region of CagA (arCagA) protein can be produced by the means of pET32a in *E. coli*. Our findings showed that antigenic protein of CagA of *H. pylori* could be detected by human sera infected with *H. pylori*. The discovery of *H. pylori* in 1982 was the starting point of a revolution concerning the concepts and management of gastroduodenal diseases. It is now well accepted that the most common stomach diseases, peptic ulcer diseases, are infectious diseases and all consensus conferences agree that the causative agent, *H. pylori* infection, must be treated with antibiotics. Furthermore, this possibility that bacterium could be the trigger of various malignant diseases of the stomach has been emerged and it is now a model for chronic bacterial infections causing cancer (13).

*H. pylori* infection can be diagnosed by invasive and non-invasive tests. Serological testing can be performed on noninvasive collected clinical samples. Serological detection of infection caused by *H. pylori* containing a CagA by the use of anti-CagA ELISA and Western blot of CagA are the only noninvasive diagnostic tests at present available for assessing strain virulence potential and possible disease risk. The reliability of CagA serology as a predictive test for determining the CagA genotype of the infecting strain is important because various serological assays are now available (14).

Many strains of *H. pylori* produce CagA, which is an immunogenic high molecular weight protein (120- 145 kDa protein) that induces numerous alterations in cellular signaling pathways, leading to the designation of CagA as a "bacterial oncoprotein" (15). *H. pylori* CagA+ strains have been associated with gastric ulcers, duodenal ulcers and gastric cancers (16). Thus this protein can be a strong vaccine candidate against *H. pylori* infections. Production of recombinant CagA has been already done by the Tumuru (17), but due to heavy molecular weight (120 KD), the protein was degraded. Hence in this study, the part of CagA protein with suitable antigenic properties which had been determined with bioinformatics methods, was used for detecting antibody. Therefore the protein obtained in this study is much smaller than normal CagA protein, presenting the same antigenic properties. This recombinant CagA can be used for detecting CagA antibody. By expression of protein in pET32a, several amino acids such as thioredoxin protein, 6xHis tag and T7 tag are added to the C or N terminal of recombinant peptide. These additional amino acids may lead to an about 20 kDa increase of the synthesized peptide molecular weight, as shown in Figure 3, which can be interfered with the subsequent immunological analyses.

In order to investigate the effect of additional amino acids (thioredoxin protein, 6xHis tag and T7 tag) pET32a vector in *E. coli* [BL21 (DE3) pLysS] was also induced by IPTG and further Western blotting on human sera was performed. The results showed that there was no inter-

ference related to fused amino acids.

In this study, we used *E. coli* BL21 (DE3) pLysS; this strain has a deficiency in the cytoplasmic protease gene products including Lon, OmpT, DegP or HtpR, in order to produce high expression level of fusion proteins. Therefore the highest expression of recombinant CagA in *E. coli* BL21 (DE3) pLysS could be due to protease deficiency in this strain. The pET system has been recognized as one of the most powerful methods for producing recombinant proteins in *E. coli* and the significant advantages of this system have been widely discussed. All pET vectors are available in three reading frames. The plasmid contains the *ori* origin of replication and the T7 lac promoter using IPTG as the inducer (10).

Our findings also indicate that the highly regulated expression vector with powerful T7 promoter (pET32a), in conjunction with suitable host cell *E. coli*, BL21 (DE3) pLysS, could influence the expression level. The expressed protein in pET system containing 6xHis tag is linked to either C or N terminal extension of protein for purification through an immobilized metal affinity chromatography (IMAC). Another advantage of using pET32a is the presence of fusion tag at N-terminal of the multiple cloning sites (MCS) which provides stability for RNA polymerase during transcription of the cloned gene.

In this study, we have cloned and expressed antigenic region of CagA under the control of T7 promoter in optimized condition. Present data showed that recombinant antigenic region of CagA is known as an antigen by patients' immune system. Therefore, recombinant antigenic region of CagA protein has same epitopes with natural form of this antigen. Recombinant antigenic region CagA also seemed to be a promising antigen for the serologic diagnosis of *H. pylori* infections.

Data indicates that antigenic parts of *H. pylori* recombinant CagA protein were recognized by human immunity system of infected individual. It can be concluded that antigenic regions of CagA have antigenic property that can be further used for the development of *H. pylori* vaccine, ELISA kits and diagnostic purpose.

### Acknowledgements

The present study was supported by Arak University of Medical Sciences.

### Authors' Contribution

All authors had equal contribution.

### Financial Disclosure

None declared.

### Funding/Support

This study was supported by Arak University of Medical Sciences.

## References

1. Suerbaum S, Michetti P. Helicobacter pylori infection. *N Engl J Med*. 2002;**347**(15):1175-86.
2. Vaira Dino, Malfertheiner Peter, Megraud Francis, Axon Anthony TR, Deltenre Michel, Hirschl Alexander M, et al. Diagnosis of Helicobacter pylori infection with a new non-invasive antigen-based assay. *The Lancet*. 1999;**354**(9172):30-33.
3. Backert S, Selbach M. Role of type IV secretion in Helicobacter pylori pathogenesis. *Cell Microbiol*. 2008;**10**(8):1573-81.
4. Leite KR, Darini E, Canavez FC, Carvalho CM, Mitteldorf CA, Camara-Lopes LH. Helicobacter pylori and cagA gene detected by polymerase chain reaction in gastric biopsies: correlation with histological findings, proliferation and apoptosis. *Sao Paulo Med J*. 2005;**123**(3):113-8.
5. Marshall B. Helicobacter pylori: 20 years on. *Clin Med*. 2002;**2**(2):147-52.
6. Diomedes M, Pietroiusti A, Silvestrini M, Rizzato B, Cupini LM, Ferrante F, et al. CagA-positive Helicobacter pylori strains may influence the natural history of atherosclerotic stroke. *Neurology*. 2004;**63**(5):800-4.
7. Sambrook Joseph, Russell David W. *Molecular cloning: a laboratory manual (3-volume set)*. 3 ed New York: Cold Spring Harbor Laboratory Press; 2001.
8. Radcliff FJ, Hazell SL, Kolesnikow T, Doidge C, Lee A. Catalase, a novel antigen for Helicobacter pylori vaccination. *Infect Immun*. 1997;**65**(11):4668-74.
9. Labigne A, Cussac V, Courcoux P. Shuttle cloning and nucleotide sequences of Helicobacter pylori genes responsible for urease activity. *J Bacteriol*. 1991;**173**(6):1920-31.
10. Mahmoudi S, Abtahi H, Bahador A, Mosayebi G, Salmanian AH. Production of recombinant streptokinase in E. coli and reactivity with immunized mice. *Pak J Biol Sci*. 2010;**13**(8):380-4.
11. Mc Pherson MJ, Moller SG. *PCR, The basic from background to bench*. Oxford, UK: BIOS Scientific Publishers Ltd; 2000.
12. Ausubel Frederick M. *Current protocols in molecular biology*. J. Wiley; 1987.
13. Megraud F, Lehours P. Helicobacter pylori detection and antimicrobial susceptibility testing. *Clin Microbiol Rev*. 2007;**20**(2):280-322.
14. Saffari M, Abtahi H. Comparison between invasive and noninvasive tests in diagnosis of Helicobacter pylori infection. *Pak J Biol Sci*. 2010;**13**(10):509.
15. Hatakeyama M. Oncogenic mechanisms of the Helicobacter pylori CagA protein. *Nat Rev Cancer*. 2004;**4**(9):688-94.
16. Figueiredo C, Van Doorn LJ, Nogueira C, Soares JM, Pinho C, Figueira P, et al. Helicobacter pylori genotypes are associated with clinical outcome in Portuguese patients and show a high prevalence of infections with multiple strains. *Scand J Gastroenterol*. 2001;**36**(2):128-35.
17. Tummuru MK, Cover TL, Blaser MJ. Cloning and expression of a high-molecular-mass major antigen of Helicobacter pylori: evidence of linkage to cytotoxin production. *Infect Immun*. 1993;**61**(5):1799-809.