

Detection of Antiseptic Resistance Genes among *Staphylococcus aureus* Colonising Nurses and Coagulase-Negative Staphylococci Isolated from Clinical Specimens at Teaching Hospitals in Southwest of Iran

Mohammad Sadegh Damavandi,¹ Maryam Safarpour Dehkordi,² Alireza Dehghan,³ Fatemeh Heibati,³

Roohollah Taghaddosi,¹ and Abolfazl Gholipour^{1*}

¹Department of Microbiology and Immunology, Cellular and Molecular Research Center, Shahrekord University of Medical Sciences, Shahrekord, IR Iran

²Young Researchers and Elite Club, Islamic Azad university, Shahrekord Branch, Shahrekord, Iran

³Cellular and Molecular Research Center, Shahrekord University of Medical Sciences, Shahrekord, IR Iran

*Corresponding author: Abolfazl Gholipour, Department of Microbiology and Immunology, Shahrekord University of Medical Sciences, Rahmatiyeh, Shahrekord, IR Iran. Tel: +98-9137046656, E-mail: abolfazl_gholipour@yahoo.com

Received 2016 May 28; Revised 2016 October 17; Accepted 2016 December 05.

Abstract

Background: The wide application of antibiotics and antiseptics for patient therapy and medical equipment and surfaces disinfection has resulted in the emergence of resistant microorganisms. *Staphylococcus aureus* and coagulase-negative Staphylococci (CoNS) are found as a part of the normal resident flora in human so that up to two-thirds of the healthy populations are permanently or transiently colonized by *S. aureus* and CoNS. Chlorhexidine is an antiseptic agent particularly effective against Gram-positive bacteria. It is widely used for hygienic hand wash to prevent transmission of Staphylococci nosocomial infections. The plasmid-borne *qacA/B*, *qacC*, and *smr* genes confer resistance to cationic antiseptic agents in *S. aureus* and CoNS.

Objectives: The objective of the current study was to characterize the antibiotic resistance and susceptibility to quaternary ammonium compounds (QACs) in methicillin-sensitive *S. aureus* (MSSA), methicillin-resistant *S. aureus* (MRSA), methicillin-sensitive coagulase-negative staphylococci (MSCoNS), and methicillin-resistant coagulase-negative Staphylococci (MRCoNS).

Methods: In this study, the antibiotic susceptibility and resistance to Chlorhexidine in 120 Staphylococcal strains were evaluated by disc diffusion and Minimum inhibitory concentration (MIC) of Chlorhexidine gluconate (CHG) methods, respectively. The MICs of CHG were determined in triplicate by broth micro-dilution, and the presence of *mecA*, *qacA/B*, *qacC*, and *smr* genes was examined by PCR assay.

Results: Of total 60 *S. aureus* isolates, 51 (85%) were MRSA, and of 60 CoNS, 7 (11.66%) were MRCoNS. The results showed that the MIC of Chlorhexidine for all 120 isolates was 1-16 $\mu\text{g}/\text{mL}$. 15 (12.5%) isolates carried *qacA/B* gene, 26 (21.7%) carried *qacC* gene, and 38 (31.7%) carried *smr* gene.

Conclusions: Maintenance of MRSA isolates in the attendance of low amounts of antiseptics could result in the decreased susceptibility to antiseptics.

Keywords: Quaternary Ammonium Compound, Minimum Inhibitory Concentration, Methicillin-Resistant *Staphylococcus aureus*

1. Background

Staphylococcus aureus is one of the well known pathogens that can live in a wide variety of environments. It also has an inherent ability to form biofilms on biotic and a-biotic surfaces (1). Moreover, bacteremia caused by *S. aureus* and CoNS is a serious infection associated with high morbidity and mortality and refers to the development of resistance virtually to all antibiotics including vancomycin. The capacity to cause infection is probably due to the organism's capacity to colonize and survive in host during infection process (2).

Staphylococcus aureus is a major cause of illness and death and imposes serious economic costs on patients and hospitals. It can infect a wide variety of human tis-

sues, resulting in different clinical appearances that vary in severity from slight purulent infections such as impetigo to more serious conditions like infective endocarditis (3). The rates of *S. aureus* infections, both in community and hospital-acquired strains, are increasing steadily. Treatment of these infections is becoming more difficult because of the increasing prevalence of oxacillin and multidrug-resistant *S. aureus* isolates (4).

Methicillin-resistant *S. aureus* (MRSA) is proposed as a main pathogen since it is resistant to many antibiotics and responsible for nosocomial infections worldwide. It is alarming that *S. aureus* strains carrying resistance genes have been isolated from various samples (5, 6). There are rising reports regarding CoNS as a cause of disease in

immunosuppressive patients and increased prevalence of multiple drug resistant strains (7). Disinfectants and antiseptics are widely used in hospitals and other health care centers as a type of topical and hard-surface practices. Especially, they are a necessary portion of infection control programs and help prevent nosocomial infections (8, 9). The quaternary ammonium compounds (*qacs*) and small multidrug-resistance (*smr*) genes are responsible for resistance to antiseptics (10). On the other hand, one of the resistance mechanisms to disinfectants and antiseptics in *S. aureus* is mediated by two gene families named *qac* and *smr* (11, 12).

Three determinants including *qacA/B*, *qacC*, and *smr* genes cause resistance to organic cations by means of multidrug efflux pumps dependent on the proton motive force. The *qacA* gene causes resistance to some different organic cations, including benzalkonium, ethidium bromide, cetrimide, and Chlorhexidine. This gene is located both on chromosome and pSK1 plasmid family. The *qacB* gene gives resistance mostly to monovalent organic cations and to a lower extent to certain divalent compounds. It is located on pSK23 plasmid. The *qacA* gene is closely related to the *qacB* gene (5, 6). Resistance to ethidium bromide and quaternary ammonium compounds can be acquired by the *qacC* gene. It is usually located on conjugative plasmids in *S. aureus* and other Staphylococci clinical isolates (13, 14).

2. Objectives

The objective of the present study was to characterize the antibiotic resistance and also determine the prevalence of QACs genes in Methicillin-sensitive *S. aureus* (MSSA) and MRSA isolated from nares of nurses, and also in MSCoNS and MRCoNS from clinical samples. The association between the presence of *mecA* gene and antiseptic resistance genes was also investigated in Staphylococci isolates.

3. Methods

3.1. Study Design

In this cross-sectional study, samples were obtained from nurse's nose and clinical specimens. The study included a total of 120 Staphylococcal strains isolated from Kashani and Hajar teaching hospitals in the southwest of Iran during the years 2012 to 2013. Staphylococcal strains included 60 *S. aureus* strains isolated from anterior nares of the nurse's nose, and 60 CoNS strains isolated from clinical specimens (wound, soft tissues, blood cultures, intravascular catheter tips, and midstream urine).

3.2. Bacterial Identification

Each nasal swab was inoculated in a trypticase soy broth (TSB) medium (Merck, Darmstadt, Germany) and incubated for 24 hours at 37°C. TSB mediums were subcultured on to mannitol salt agar (Merck, Germany) and incubated for 24 hours. Colonies with Staphylococcal morphology were identified by biochemical tests such as catalase and coagulase and confirmed by 16S rRNA genes. Tube coagulase negative strains were reported as CoNS. Staphylococci isolates were stored at -70°C in TSB mediums supplemented with 30% glycerol.

3.3. Susceptibility Testing

Antibiotic susceptibility testing was performed by disk diffusion method according to the CLSI. All *S. aureus* isolates were tested by oxacillin (1 µg), Gentamycin (10 µg), Ticoplanin (30 µg), Linezolid (30 µg), quinopristin-dalfopristin (15 µg), Tigecycline (15 µg), Rifampin (5 µg), and Vancomycin (30 µg) and all CoNS isolates were examined by Bacitracin (0.4 U), Novobiocin (30 µg), and Vancomycin (30 µg) (MAST Diagnostics, Merseyside, UK).

Methicillin-resistant Staphylococci (MRS) strains were known by multiplication on Muller-Hinton agar and using oxacillin disc (1 µg) diffusion method. The inhibition zone size was interpreted according to the CLSI criteria (15). The MRSA strains were also confirmed through the presence of *mecA* gene by PCR technique that is so far considered as the gold standard method for the detection of methicillin resistance.

MICs of Chlorhexidine gluconate (Sigma Aldrich, St Louis, USA) (ranging from 1 to 64 µg/mL, by serial twofold dilutions) were determined in triplicate by the broth micro dilution method according to the CLSI guideline (15). The lowest concentration totally inhibiting growth after 24 hours incubation at 37°C was considered the MIC. *S. aureus* ATCC 13353 and *S. aureus* ATCC 25923 were used as positive and negative controls, respectively.

3.4. Qac genes Detection

Total genomic DNA of the microorganisms in this study was extracted using boiling method. The *qacA/B*, *qacC*, and *smr* resistance genes were tested in all isolates by multiplex PCR. The *qacA* and *qacB* genes were considered together because simple PCRs cannot discriminate between them. To detect *qacA/B*, *qacC*, and *smr* genes, the multiplex PCR conditions were as follows: 1 cycle of initial denaturation at 95°C and 35 cycles of denaturation (at 94°C for 40 seconds), annealing (at 54°C for 50 s), extension (at 72°C for 50 seconds), and one final extension (at 72°C for 5 minutes).

The detection of *qacA/B*, *qacC*, and *smr* genes was performed by multiplex PCR with the following sets of primers: 5'-GCA GAA AGT GCA GAG TTG G-3' and 5'-CCA GTC CAA TCA TGC CTG -3' for *qacA/B* (product size, 361 bp), and 5'-GGC TTT TCA AAA TTT ATA CCA TCCT -3' and 5'-ATG CGAT GTT TCC GAA AAT GT -3' for *qacC* (product size, 246 bp), and 5'-GCC ATA AGT ACT GAA GTT ATT GGA-3' and 5'-GAC TAC GGT TGT TAA GAC TAA ACC T -3' for *smr* (product size 195 bp), as previously described (16, 17).

3.5. Statistical Analysis

Statistical analyses were performed using SPSS software version 18.0 for Windows (SPSS Inc., Chicago, IL, USA). The association of categorical variables was determined by Chi Square or Fisher's exact tests. $P < 0.05$ was considered statistically significant.

4. Results

Of total 60 *S. aureus* isolates included in this study, 51 (85%) isolates were MRSA and, of 60 CoNS, 7 (11.66%) isolates were MRCoNS. The frequency of antimicrobial resistance in the isolated MRSA strains was 17 (33%) for rifampin, 15 (29.5%) for Tigecycline, 12 (23.5%) for Ticoplanin, and 7 (13.7%) for Gentamycin. All MRSA isolates showed susceptibility to vancomycin, Linezolid, and quinopristindalfopristin. As well as, only 2 isolates (1.3%) of MRCoNS strains were resistant to Vancomycin. The MICs of Chlorhexidine were 1 $\mu\text{g}/\text{mL}$ for 42 (35%) isolates, 2 $\mu\text{g}/\text{mL}$ for 22 (18.3%) isolates, 4 $\mu\text{g}/\text{mL}$ for 40 (33.3%) isolates, 8 $\mu\text{g}/\text{mL}$ for 14 (11.6%) isolates, and 16 $\mu\text{g}/\text{mL}$ for 2 (1.6%) isolates. The association between the presence of *mecA* gene and Chlorhexidine MIC is shown in Table 1.

The association between the presence of *qac* genes and a Chlorhexidine MIC of $> 4 \mu\text{g}/\text{mL}$ was statistically significant ($P < 0.0001$). It is considerable that among two strains with the highest MIC of Chlorhexidine gluconate ($\text{MIC} \geq 8 \mu\text{g}/\text{mL}$), one of them was *qac* negative and the other strain carried *qacC* and *smr* genes at the same time. Clinical strains with reduced susceptibility to Chlorhexidine ($\text{MIC} \geq 2 \mu\text{g}/\text{mL}$) had a strong association with the presence of *qacA/B* and *smr* determinants ($P < .0001$). As well as, among the tested bacteria with various Chlorhexidine MICs, there were some strains of Staphylococci isolates that carried the two genes at the same time (Table 2).

The *qacA/B* gene was identified in 15 (12.5%) isolates from which, 6 (40%) were *qacA/B* alone, 5 (33.3%) with both *smr* and *qacA/B*, and 4 (26.7%) with both *qacA/B* and *qacC* at the same time. The *qacC* was identified in 26 (21.7%) isolates from which, 8 (30.8%) were *qacC* alone, 4 (15.4%) plus *qacA/B*, and 14 (53.8%) plus *smr* at the same time. The *smr* was identified in 38 (31.7%) isolates from which, 19 (50%) *smr* alone,

5 (13.1%) plus *qacA/B*, and 14 (36.9%) plus *qacC* at the same time (Figure 1). Also, 41 out of 120 (34.1%) isolated strains were *qac*-negative. Isolates with *qac* genes ($N = 56$) had significantly a higher mean and wider range of Chlorhexidine MICs. The results of this study also showed that methicillin resistant Staphylococci were more often resistant to Chlorhexidine than methicillin sensitive isolates (Table 3).




Figure 1. Line M ladder 100 bp (Fermentas), Line 1 positive control (*S. aureus* ATCC 13353), line 2 isolates harboring *qacA/B* (361 bp) and *qacC* (247 bp), line 3 and 4 isolates harboring *qacC* (247 bp), line 5 isolates negative for *qac*s gene, line 6 isolates harboring *smr* (195 bp) and *qacC* (247 bp), line 7 isolates harboring *qacA/B* (361 bp).

5. Discussion

Staphylococcus aureus is a flexible pathogen, which has been known as a common cause of various community-acquired and nosocomial infections (18). Hospital staff nasal carriers of MRSA can increase the risk of outbreaks of MRSA at Hospitals because they are potential reservoirs for the spread of MRSA strains among patients. Nasal colonization of MRSA and MRCoNS is different all over the world. For example, a Korean study reported a rate of MRCoNS colonization of non-healthcare workers (13%) that is similar to

Table 1. Association of Chlorhexidine MIC with MRSA, MSSA, MRCoNS, and MSCoNS^a

Species (no. of Isolates)	Chlorhexidine MIC Level, $\mu\text{g/mL}$					
	MIC Range 1 - 2		MIC Range 2 - 4		MIC Range 4 - 16	
	Positive	P Value	Positive	P Value	Positive	P Value
MRSA (51)	1 (1.96)	0.15	34 (66.6)	0.002	16 (31.3)	0.05
MSSA (9)	8 (88.8)		1 (11.1)		0	
MRCoNS (7)	3 (42.8)	< 0.001	4 (57.1)	< 0.001	0	NS
MSCoNS (53)	52 (98.1)		1 (1.88)		0	
Total	64		40		16	120

Abbreviation: NS, non-significant.

^aValues are expressed as No.(%).

Table 2. Association Between the Presence of QAC Genes and Chlorhexidine MIC

Qac Genes	Frequency (%)	Chlorhexidine MIC Level, $\mu\text{g/mL}$, %					
		1	2	4	8	16	P Value
<i>qacA/B</i>	6 (5)	2 (33.3)	1 (16.7)	3 (50)	0	0	
<i>qacC</i>	8 (6.7)	1 (12.5)	0	7 (87.5)	0	0	
<i>smr</i>	19 (15.8)	1 (5.2)	1 (5.2)	17 (89.4)	0	0	
<i>qacA/B + smr</i>	5 (4.2)	0	0	5 (100)	0	0	< 0.0001
<i>qacA/B + qacC</i>	4 (3.3)	0	0	2 (50)	2 (50)	0	
<i>qacC + smr</i>	14 (11.7)	0	0	1 (7.15)	12 (85.7)	1 (7.15)	
Negative	64 (53.3)	38 (59.4)	20 (31.2)	5 (7.8)	0	1 (1.56)	
Total	120 (100)	42 (35)	22 (18.3)	40 (33.35)	14 (11.7)	2 (1.6)	

Table 3. Association Between the Presence of Antiseptic-Resistance Genes and Methicillin Resistance (MR) in Isolates of *Staphylococcus aureus* (SA) and Coagulase-Negative Staphylococci (CoNS)^a

Species (No. of Isolates)	<i>qacA/B</i>				<i>qacC</i>				<i>smr</i>			
	Positive	P Value	OR	95% CI	Positive	P Value	OR	95% CI	Positive	P Value	OR	95% CI
MRSA (51)	4 (7.8)	0.38	0.92	0.85 - 0.99	7 (13.7)	0.83	1.2	0.13 - 11	17 (33.3)	0.041	0.66	0.54 - 0.8
MSSA (9)	0				1 (11.1)				0			
MRCoNS (7)	1 (14.28)	0.086	8.6	0.47 - 157	0	NS	NS		2 (28.57)	< 0.001	0.71	0.44 - 1.1
MSCoNS (53)	1 (1.88)				0				0			

Abbreviations: CI, confidence interval; NS, non-significant; OR, odds ratio.

^aValues are expressed as No.(%).

the rate found in our study (19). The unnecessary use of disinfectants at hospitals results in the permanence of Staphylococcal isolates, as well as, the augmented prevalence of *qac* genes in such strains (20).

In order to evaluate the MIC of Chlorhexidine, determine the presence of *qac* genes, and investigate the association between the presence of *mecA* gene and Chlorhexidine MIC, 60 isolates of *S. aureus* and 60 isolates of CoNS were studied. The Chlorhexidine MIC defined in this study was equal to or higher than 4 $\mu\text{g/mL}$. According to previ-

ous studies (16-18), the Chlorhexidine MIC for MRSA strains was typically equal to or higher than 4 $\mu\text{g/mL}$. Therefore, our finding was in accordance with those of the above-mentioned studies. The correlation between the presence of *qac* genes (*qacA/B*, *qacC*, and *smr*) and a Chlorhexidine MIC of $\geq 4 \mu\text{g/mL}$ was statistically significant ($P \leq 0.0001$). It is noteworthy that 2 (33.3%) out of 6 strains with *qacA/B* gene had Chlorhexidine MIC = 1 $\mu\text{g/mL}$. As well as 1 (12.5%) out of 8 strains with *qacC* and 1 (5.2%) out of 19 strains with *smr* genes had Chlorhexidine MIC = 1 $\mu\text{g/mL}$. The presence

of *smr* gene had a strong relationship with the presence of *mecA* gene in *S. aureus* ($p \leq 0.041$) and CoNS ($P \leq 0.001$). The only one out of 14 *qacC* plus *smr*-carrying strains showed a MIC value of 16 $\mu\text{g}/\text{mL}$. It is necessary to mention that among 64 strains which did not have *qac* genes, one strain also showed an MIC value of 16 $\mu\text{g}/\text{mL}$. In fact, we observed a significant increase in Chlorhexidine MICs associated with the *qacA/B*, *qacC*, and *smr* genes. In a Canadian study on MRSA isolates, there was no significant increase in MICs with the *qac* and *smr* genes (21); while our data showed an increase in the MIC value of Chlorhexidine in the presence of *qacA/B*, *qacC*, *smr*, and *mecA* genes. The increased MIC values for Chlorhexidine (MIC of $> 2 \mu\text{g}/\text{ml}$) had a strong relationship with the presence of *mecA* in *S. aureus* ($P \leq 0.002$) and CoNS ($p \leq 0.001$). In summary, our data showed that 85% of MRSA and 11.6% of MRCoNS isolates carried *qac* genes, which were in association with reduced susceptibility to Chlorhexidine.

The QACs resistant genes such as *qacA/B*, *qacC*, and *smr* have been isolated in various environments from clinical CoNS isolates and different Staphylococcal species (13, 16). Whereas the current antiseptic MICs related to *qac* gene positive Staphylococci will not allow survival at in-use concentrations, even moderately increased resistance may allow persistence when residual disinfectants are present (16, 22). The over-use of Chlorhexidine as a decontaminant could result in the appearance of MRSA isolates carrying *qacA/B* resistance genes (23). While hospital environments act as a source of *qac* genes, long-term care facilities and nurses may play a contributory role in the transmission of antiseptic-resistant Staphylococci between hospital sections and patients (24). Many studies have been performed in order to understand the incidence and possible genetic link of antiseptic and antibiotic resistance genes in Staphylococci. Similar to the findings of other studies (20, 25), in our study the existence of *mecA* gene in these isolates was related to the presence of *qac* genes. Our study demonstrated that the raised ratio of *qac* gene positive Staphylococci strains offers the co-selection of these genes due to the increased contact with MRSA-infected patients. Decreased susceptibility to antiseptic (Chlorhexidine) could be associated with *qac* genes, which is also consistent with the reports of clinical isolates in above studies.

In this study, the frequency of *qacA/B*, *qacC*, and *smr* genes was 7.8%, 13.7%, and 33.3%, respectively while in the study of Noguchi et al. in Japan (26), the frequency of *qac* genes was 45.9% that is higher than that in our study; but the frequency of *smr* gene was 5.3% that is much lower than the value obtained in our study. In another study by Miyazaki et al. conducted on 74 Brazilian MRSA isolates (27), the rate of *qacA/B* was that higher than the ratio in our study. Lee et al. in Switzerland identified *qacA/B* in

91% of the MRSA isolates (28). In a study by Longtin et al. in Canada (21), only 2% of the MRSA strains possessed the *qacA/B* while *smr* gene was detected in 7% of strains. In Iran, a study by Hasanvand et al. (29) showed that the frequency of *qacA/B* antiseptic gene in MRSA isolates was 9% that is in agreement with our study. On the other hand, the *smr* gene in the same study was not detected in both MRSA and MSSA isolates while the frequency of the *smr* gene in this study was 17 (33.3%) in MRSA and 2 (28.57%) in MRCoNS isolates. The *mecA* gene is located on a mobile genomic island Staphylococcal cassette chromosome (SCC), which not only serves as a vehicle for the genetic exchange of genes among Staphylococci, but also as a carrier for virulence and additional drug-resistant genes (17, 30). Genetic linkage between *qac* genes and *mecA* genes conferring resistance to methicillin on the same Staphylococcal plasmids has also been reported elsewhere (31).

Application of antiseptics might be chosen for strains resistant to antibiotics and help them maintain in health-care settings. The presence of an association between *mecA* and *qacA/B*, *qacC*, and *smr* may promote survival of MRSA and MRCoNS in the hospital environments. These Staphylococci may be a hazard for infection control because of their persistence in places with low amounts of antiseptic residues. In conclusion, our study showed Chlorhexidine resistance is commonly found in MRSA isolates from nurse's nose and clinical specimens. This study had limitations such as coordination problems in nasal swab sampling from staff and clinical specimens found in different units of hospitals.

The increase of antiseptic-resistant bacteria is one of the most significant problems and a serious threat to public health. Understanding the selection of gene transfer that causes the distribution of resistance genes is very important for long-term strategies in order to treat microbial diseases.

5.1. Ethical Considerations

Ethical issues (including plagiarism, informed consent, misconduct, data fabrication and/or falsification, double publication and/or submission, redundancy, etc.) have been completely observed by the authors.

Acknowledgments

We gratefully thank the staff of the microbiology laboratory and cellular and molecular research center at the faculty of medicine, Shahrekord University of Medical Sciences, Shahrekord, Iran, for their cooperation in this study.

Footnotes

Authors' Contribution: Study concept and design: Roohollah Taghaddosi; acquisition of data: Maryam Safarpour Dehkordi; analysis and interpretation of data, Alireza Dehghan; administrative, technical, and material support: Abolfazl Gholipour; statistical analysis: Fatemeh Heibati; drafting of the manuscript: Mohammad Sadegh Damavandi; revising the manuscript: Abolfazl Gholipour.

Conflicts of Interest: The authors of the present work declare no conflict of interests.

Funding/Support: This study is a research project with grant number 1085 from deputy of research and technology of Shahrekord University of Medical Sciences.

References

- Mirani ZA, Aziz M, Khan MN, Lal I, Hassan NU, Khan SI. Biofilm formation and dispersal of *Staphylococcus aureus* under the influence of oxacillin. *Microb Pathog.* 2013;**61-62**:66-72. doi: [10.1016/j.micpath.2013.05.002](https://doi.org/10.1016/j.micpath.2013.05.002). [PubMed: [23719663](https://pubmed.ncbi.nlm.nih.gov/23719663/)].
- Liu CL, Chen ZJ, Wang F, Hou HY, Wang Y, Zhu XH, et al. The impact of *mgrA* on progression of *Staphylococcus aureus* sepsis. *Microb Pathog.* 2014;**71-72**:56-61. doi: [10.1016/j.micpath.2014.03.012](https://doi.org/10.1016/j.micpath.2014.03.012). [PubMed: [24746531](https://pubmed.ncbi.nlm.nih.gov/24746531/)].
- Abraham NM, Jefferson KK. A low molecular weight component of serum inhibits biofilm formation in *Staphylococcus aureus*. *Microb Pathog.* 2010;**49**(6):388-91. doi: [10.1016/j.micpath.2010.07.005](https://doi.org/10.1016/j.micpath.2010.07.005). [PubMed: [20673798](https://pubmed.ncbi.nlm.nih.gov/20673798/)].
- Ikonomidis A, Vasdeki A, Kristo I, Maniatis AN, Tsakris A, Malizos KN, et al. Association of biofilm formation and methicillin-resistance with accessory gene regulator (*agr*) loci in Greek *Staphylococcus aureus* clones. *Microb Pathog.* 2009;**47**(6):341-4. doi: [10.1016/j.micpath.2009.09.011](https://doi.org/10.1016/j.micpath.2009.09.011). [PubMed: [19786085](https://pubmed.ncbi.nlm.nih.gov/19786085/)].
- David MZ, Daum RS. Community-associated methicillin-resistant *Staphylococcus aureus*: epidemiology and clinical consequences of an emerging epidemic. *Clin Microbiol Rev.* 2010;**23**(3):616-87. doi: [10.1128/CMR.00081-09](https://doi.org/10.1128/CMR.00081-09). [PubMed: [20610826](https://pubmed.ncbi.nlm.nih.gov/20610826/)].
- Costa SS, Viveiros M, Amaral L, Couto I. Multidrug Efflux Pumps in *Staphylococcus aureus*: an Update. *Open Microbiol J.* 2013;**7**:59-71. doi: [10.2174/1874285801307010059](https://doi.org/10.2174/1874285801307010059). [PubMed: [23569469](https://pubmed.ncbi.nlm.nih.gov/23569469/)].
- Carbon C. MRSA and MRSE: is there an answer?. *Clin Microbiol Infect.* 2000;**6 Suppl 2**:17-22. [PubMed: [11523520](https://pubmed.ncbi.nlm.nih.gov/11523520/)].
- Gerba CP. Quaternary ammonium biocides: efficacy in application. *Appl Environ Microbiol.* 2015;**81**(2):464-9. doi: [10.1128/AEM.02633-14](https://doi.org/10.1128/AEM.02633-14). [PubMed: [25362069](https://pubmed.ncbi.nlm.nih.gov/25362069/)].
- Deress T, Girma M, Birhan W, Biadgo B, Alemu A. Isolation of Bacteria from Commonly Used Antiseptic and Disinfectant Solutions in Gondar University Hospital. North West Ethiopia. *Am J Nurs Res.* 2014;**2**(3):44-9. doi: [10.12691/ajnr-2-3-3](https://doi.org/10.12691/ajnr-2-3-3).
- Tacconelli E, Pop-Vicas AE, D'Agata EM. Increased mortality among elderly patients with methicillin-resistant *Staphylococcus aureus* bacteraemia. *J Hosp Infect.* 2006;**64**(3):251-6. doi: [10.1016/j.jhin.2006.07.001](https://doi.org/10.1016/j.jhin.2006.07.001). [PubMed: [16978733](https://pubmed.ncbi.nlm.nih.gov/16978733/)].
- Anthonsen IL, Sunde M, Steinum TM, Sidhu MS, Sorum H. Organization of the antiseptic resistance gene *qacA* and Tn552-related beta-lactamase genes in multidrug-resistant *Staphylococcus haemolyticus* strains of animal and human origins. *Antimicrob Agents Chemother.* 2002;**46**(11):3606-12. [PubMed: [12384372](https://pubmed.ncbi.nlm.nih.gov/12384372/)].
- Horner C, Mawer D, Wilcox M. Reduced susceptibility to chlorhexidine in staphylococci: is it increasing and does it matter?. *J Antimicrob Chemother.* 2012;**67**(11):2547-59. doi: [10.1093/jac/dks284](https://doi.org/10.1093/jac/dks284). [PubMed: [22833635](https://pubmed.ncbi.nlm.nih.gov/22833635/)].
- Furi L, Ciusa ML, Knight D, Di Lorenzo V, Tocci N, Cirasola D, et al. Evaluation of reduced susceptibility to quaternary ammonium compounds and bisbiguanides in clinical isolates and laboratory-generated mutants of *Staphylococcus aureus*. *Antimicrob Agents Chemother.* 2013;**57**(8):3488-97. doi: [10.1128/AAC.00498-13](https://doi.org/10.1128/AAC.00498-13). [PubMed: [23669380](https://pubmed.ncbi.nlm.nih.gov/23669380/)].
- Ortega Morente E, Fernandez-Fuentes MA, Grande Burgos MJ, Abriouel H, Perez Pulido R, Galvez A. Biocide tolerance in bacteria. *Int J Food Microbiol.* 2013;**162**(1):13-25. doi: [10.1016/j.ijfoodmicro.2012.12.028](https://doi.org/10.1016/j.ijfoodmicro.2012.12.028). [PubMed: [23340387](https://pubmed.ncbi.nlm.nih.gov/23340387/)].
- CLSI . Performance Standards for Antimicrobial Susceptibility Testing; Twenty-Fifth Informational Supplement. CLSI document M100-S25. Wayne, PA: Clinical and Laboratory Standards Institute; 2015.
- Sidhu MS, Heir E, Leegaard T, Wiger K, Holck A. Frequency of disinfectant resistance genes and genetic linkage with beta-lactamase transposon Tn552 among clinical staphylococci. *Antimicrob Agents Chemother.* 2002;**46**(9):2797-803. [PubMed: [12183231](https://pubmed.ncbi.nlm.nih.gov/12183231/)].
- Noguchi N, Suwa J, Narui K, Sasatsu M, Ito T, Hiramatsu K, et al. Susceptibilities to antiseptic agents and distribution of antiseptic-resistance genes *qacA/B* and *smr* of methicillin-resistant *Staphylococcus aureus* isolated in Asia during 1998 and 1999. *J Med Microbiol.* 2005;**54**(Pt 6):557-65. doi: [10.1099/jmm.0.45902-0](https://doi.org/10.1099/jmm.0.45902-0). [PubMed: [15888465](https://pubmed.ncbi.nlm.nih.gov/15888465/)].
- Papa R, Artini M, Cellini A, Tilotta M, Galano E, Pucci P, et al. A new anti-infective strategy to reduce the spreading of antibiotic resistance by the action on adhesion-mediated virulence factors in *Staphylococcus aureus*. *Microb Pathog.* 2013;**63**:44-53. doi: [10.1016/j.micpath.2013.05.003](https://doi.org/10.1016/j.micpath.2013.05.003). [PubMed: [23811076](https://pubmed.ncbi.nlm.nih.gov/23811076/)].
- Lee YL, Cesario T, Tran C, Stone G, Thrupp L. Nasal colonization by methicillin-resistant coagulase-negative staphylococcus in community skilled nursing facility patients. *Am J Infect Control.* 2000;**28**(3):269-72. doi: [10.1067/mic.2000.103243](https://doi.org/10.1067/mic.2000.103243). [PubMed: [10840350](https://pubmed.ncbi.nlm.nih.gov/10840350/)].
- Smith K, Gemmell CG, Hunter IS. The association between biocide tolerance and the presence or absence of *qac* genes among hospital-acquired and community-acquired MRSA isolates. *J Antimicrob Chemother.* 2008;**61**(1):78-84. doi: [10.1093/jac/dkm395](https://doi.org/10.1093/jac/dkm395). [PubMed: [17981834](https://pubmed.ncbi.nlm.nih.gov/17981834/)].
- Longtin J, Seah C, Siebert K, McGeer A, Simor A, Longtin Y, et al. Distribution of antiseptic resistance genes *qacA*, *qacB*, and *smr* in methicillin-resistant *Staphylococcus aureus* isolated in Toronto, Canada, from 2005 to 2009. *Antimicrob Agents Chemother.* 2011;**55**(6):2999-3001. doi: [10.1128/AAC.01707-10](https://doi.org/10.1128/AAC.01707-10). [PubMed: [21402848](https://pubmed.ncbi.nlm.nih.gov/21402848/)].
- Edgeworth JD. Has decolonization played a central role in the decline in UK methicillin-resistant *Staphylococcus aureus* transmission? A focus on evidence from intensive care. *J Antimicrob Chemother.* 2011;**66**(Suppl 2):ii:41-7. doi: [10.1093/jac/dkq325](https://doi.org/10.1093/jac/dkq325). [PubMed: [20852273](https://pubmed.ncbi.nlm.nih.gov/20852273/)].
- Eveillard M, Martin Y, Hidri N, Boussougant Y, Joly-Guillou ML. Carriage of methicillin-resistant *Staphylococcus aureus* among hospital employees: prevalence, duration, and transmission to households. *Infect Control Hosp Epidemiol.* 2004;**25**(2):114-20. doi: [10.1086/502360](https://doi.org/10.1086/502360). [PubMed: [14994935](https://pubmed.ncbi.nlm.nih.gov/14994935/)].
- Walsh SE, Maillard JY, Russell AD, Catrenich CE, Charbonneau DL, Bartolo RG. Development of bacterial resistance to several biocides and effects on antibiotic susceptibility. *J Hosp Infect.* 2003;**55**(2):98-107. [PubMed: [14529633](https://pubmed.ncbi.nlm.nih.gov/14529633/)].
- Hanssen AM, Ericson Sollid JU. SCCmec in staphylococci: genes on the move. *FEMS Immunol Med Microbiol.* 2006;**46**(1):8-20. doi: [10.1111/j.1574-695X.2005.00009.x](https://doi.org/10.1111/j.1574-695X.2005.00009.x). [PubMed: [16420592](https://pubmed.ncbi.nlm.nih.gov/16420592/)].
- Noguchi N, Nakaminami H, Nishijima S, Kurokawa I, So H, Sasatsu M. Antimicrobial agent of susceptibilities and antiseptic resistance gene distribution among methicillin-resistant *Staphylococcus aureus* isolates from patients with impetigo and staphylococ-

- cal scalded skin syndrome. *J Clin Microbiol.* 2006;**44**(6):2119-25. doi: [10.1128/JCM.02690-05](https://doi.org/10.1128/JCM.02690-05). [PubMed: [16757607](https://pubmed.ncbi.nlm.nih.gov/16757607/)].
27. Miyazaki NH, Abreu AO, Marin VA, Rezende CA, Moraes MT, Villas Boas MH. The presence of qacA/B gene in Brazilian methicillin-resistant *Staphylococcus aureus*. *Mem Inst Oswaldo Cruz.* 2007;**102**(4):539-40. [PubMed: [17607497](https://pubmed.ncbi.nlm.nih.gov/17607497/)].
28. Lee AS, Macedo-Vinas M, Francois P, Renzi G, Schrenzel J, Vernaz N, et al. Impact of combined low-level mupirocin and genotypic chlorhexidine resistance on persistent methicillin-resistant *Staphylococcus aureus* carriage after decolonization therapy: a case-control study. *Clin Infect Dis.* 2011;**52**(12):1422-30. doi: [10.1093/cid/cir233](https://doi.org/10.1093/cid/cir233). [PubMed: [21628482](https://pubmed.ncbi.nlm.nih.gov/21628482/)].
29. Hasanvand A, Ghafourian S, Taherikalani M, Jalilian FA, Sadeghifard N, Pakzad I. Antiseptic Resistance in Methicillin Sensitive and Methicillin Resistant *Staphylococcus aureus* Isolates from Some Major Hospitals, Iran. *Recent Pat Antiinfect Drug Discov.* 2015;**10**(2):105-12. [PubMed: [26100979](https://pubmed.ncbi.nlm.nih.gov/26100979/)].
30. Taghaddosi R, Gholipour A, Zeraatpisheh M, Safarpour-Dehkordi M, Darban-Sarokhalil D, Heibati-Goujani F. A frequency and Molecular Typing Study of Methicillin-Resistant *Staphylococcus aureus* Isolates in Teaching Hospitals in Shahrekord, SouthWestern Iran. *Jundishapur J Microbiol.* 2016;**Inpress**(Inpress) doi: [10.5812/jjm.39654](https://doi.org/10.5812/jjm.39654).
31. Sidhu MS, Heir E, Sorum H, Holck A. Genetic linkage between resistance to quaternary ammonium compounds and beta-lactam antibiotics in food-related *Staphylococcus* spp. *Microb Drug Resist.* 2001;**7**(4):363-71. doi: [10.1089/10766290152773374](https://doi.org/10.1089/10766290152773374). [PubMed: [11822776](https://pubmed.ncbi.nlm.nih.gov/11822776/)].