Published online 2021 September 13.

Research Article

Harmine Augments the Cytotoxic and Anti-invasive Potential of Temozolomide Against Glioblastoma Multiforme Cells

Mehran Kamani 😳 1, Ali Ghanbari 😳 1, Mahdi Taghadosi 2, Kamran Mansouri 3 and Cyrus Jalili 😳 3,*

¹Department of Anatomical Sciences, Medical School, Kermanshah University of Medical Sciences, Kermanshah, Iran ²Department of Immunology, School of Medicine, Kermanshah University of Medical Sciences, Kermanshah, Iran ³Medical Biology Research Center, Health Technology Institute, Kermanshah University of Medical Sciences, Kermanshah, Iran

^{*} Corresponding author: Medical Biology Research Center, Health Technology Institute, Kermanshah University of Medical Sciences, Kermanshah, Iran. Email: cjalili@yahoo.com

Received 2021 May 11; Revised 2021 July 20; Accepted 2021 August 01.

Abstract

Background: Glioblastoma multiforme (GBM) is considered the deadliest human cancer. Temozolomide is now a part of postresection standard chemotherapy for this type of cancer. Unfortunately, resistance to temozolomide is a major obstacle to treatment success. Combination therapy with natural anticancer agents increases the activity of temozolomide against cancer cells. **Objectives:** This study aimed to assess the effects of temozolomide in combination with harmine against GBM cells.

Objectives: This study affield to assess the effects of temporation and in combination with narmine against GBM cents.

Methods: Cancer cells were treated with temozolomide and/or harmine. After 24, 48, 72, and 96 h, the viability of the cells was assessed by the MTT test. The combination index and dose reduction index were determined by CompuSyn software. Tumor invasion potential was investigated by evaluating cell migration, invasion, and adhesion. The real-time PCR technique was done to study the expression pattern of two genes involved in cancer cell invasion. Statistical analysis was performed using one-way analysis of variance and Tukey's post-hoc test, and differences were considered non-significant at P > 0.05.

Results: After treatment with temozolomide, cell viability showed a concentration- and time-dependent decrease, and the cells' survival rate decreased. The combination of temozolomide and harmine had a synergistic effect. Also, temozolomide and/or harmine treatment decreased cancer cells' migration, invasion, and adhesion potentials, as well as the expression of metalloproteinases 2 and 9 in T98G cells.

Conclusions: The combination of temozolomide and harmine can be promising for the successful treatment of GBM.

Keywords: Temozolomide, Harmine, Glioblastoma Multiforme, Drug Resistance

1. Background

Glioma is the most prevalent type of the central nervous system's cancers and accounts for more than 30% of all primary malignancies of the central nervous system. This cancer originates from glial cells, and the main cause of it is not known. Hereditary genetic disorders such as neurofibromatosis types 1 and 2 and tuberous sclerosis are known to predispose to glioma (1). Another predisposing factor for this cancer is cell phone electromagnetic radiation (2). Most brain tumors of the glioma type have been associated with cytomegalovirus infection, but the direct role of viruses in triggering this type of cancer has not yet been confirmed (3). Due to the invasive behavior of this tumor, the rapid proliferation of cancerous cells, and the lack of a proper treatment, this cancer causes many problems for the patient (4). Grade IV glioma, also known as glioblastoma multiforme (GBM), is the most malignant

and progressive brain cancer and one of the deadliest human solid tumors, accounting for about 54% of malignant gliomas (5). Some GBM tumors arise from the development of pre-existing grade II and III astrocytomas (known as secondary GBM). However, about 90 to 95% of GBM cases develop de novo (i.e., primary GBM) (6). The risk of GBM increases with age, and men are about 1.6 times more likely than women to develop the disease. Also, this type of cancer is more prevalent among whites than other ethnic groups, and its rate is 3.44 cases per 100,000 people (6).

Temozolomide (TMZ), under the brand name of Temodar or temodal, is used as a standard oral chemotherapy drug to treat GBM (7). In cancer cells, DNA methylation is thought to be the main mechanism responsible for the toxicity induced by TMZ (8). Resistance to drugs is an important obstacle restricting the effectiveness of cancer drugs and is caused by some mechanisms such as drug

Copyright © 2021, Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (http://creativecommons.org/licenses/by-nc/4.0/) which permits copy and redistribute the material just in noncommercial usages, provided the original work is properly cited.

inactivation, drug expulsion from tumor cells, repair of cellular destruction through chemotherapy, activation of survival pathways, and inactivation of cellular death pathways. The main effort to overcome therapy resistance is to use combinations of several drugs with minimal toxicities from different pharmaceutical groups (9). The standard treatments used for GBM, including surgical tumor resection, chemotherapy, and radiation therapy, generally fail to extend the short survival of patients after diagnosis. In fact, long-term survival is currently not accessible due to the innate and acquired resistance of the tumor to therapy.

Over the past millions of years, many natural products have been used to treat a variety of ailments despite the lack of scientific confirmation of their effectiveness and safety. Herbal remedies have been used by experimental physicians for centuries (10). Advances in technology have allowed scientists to identify the active ingredients of herbal extracts. Thus, efforts are underway to find new drugs to either complement or replace conventional therapies (11).

Harmine, a β -carboline alkaloid, was first separated from the seeds of *Peganum harmala* and *Banisteriopsis caapi* in 1847. It is extensively present in numerous medicinal herbs and has long been utilized in folk medicine in the Middle East and Asia. Previous research has shown the anticancer activities of harmine against different cancers, including lung (12), gastric (13), breast (14), and hepatic (15) cancer.

As previously mentioned, patients with GBM have a relatively short lifespan due to the low sensitivity of tumor cells to TMZ. On the other hand, increasing the concentration of the drug to boost its effectiveness causes very severe side effects.

2. Objectives

Due to the need to identify new effective therapies and enhance the efficacy of TMZ, this study aimed to investigate the effects of the combination of TMZ and harmine on the cell viability and invasion potential of T98G cells, a GBM cell line.

3. Methods

3.1. Cell Line and Reagents

The GBM cell line (T98G) was purchased from the Pasteur Institute of Iran. Harmine, TMZ, 3-(4, 5dimethylthiazol-2-yl)-2, 5-diphenyltetrazolium bromide (MTT), ethanol, crystal violet, trypsin, and dimethyl sulfoxide (DMSO) were provided from Sigma-Aldrich Chemical Co. (St. Louis, MO, USA). Roswell Park Memorial Institute (RPMI) 1640 medium, penicillin/streptomycin solution, and fetal bovine serum (FBS) were procured from Gibco (USA). Stock solutions (200-mM) of TMZ and harmine were prepared in DMSO and diluted in the culture medium to reach final desired concentrations. The cells were seeded in the culture flasks containing RPMI 1640 supplemented with 10% FBS and 1% antibiotics (penicillin/streptomycin). The cells were maintained at 37°C in a humidified incubator under 5% CO₂.

3.2. Viability Assay

The MTT assay was used to investigate the effects of TMZ and harmine on cell viability. The cells were cultured into a 96-well plate (1.5×10^4 cells/well) and incubated overnight. Then the cells were treated with different concentrations (1.56, 3.12, 6.25, 12.5, 25, 50, 100, and $200 \ \mu$ M) of TMZ and harmine in a serum-free medium for 24, 48, 72, and 96 hours. Next, the medium was discarded, and $50 \ \mu$ L of MTT solution ($5 \ mg/mL$ in PBS) was added to the wells. After 2.5 h, to evaluate the rate of the reaction, $100 \ \mu$ L of DMSO was added. The plate was then placed on a shaker at room temperature for 20 min, and the absorbance was read by an ELISA reader at 570 and 630 nm (16). Cell survival percentage was determined according to the following formula:

$$Survival \ percentage = \frac{the \ absorption \ of \ cotrol \ cells}{adsorption \ of \ treated \ cells} \ \times 100$$

The half-maximal inhibitory concentration (IC_{50}) values of TMZ and harmine were calculated using GraphPad Prism software version 6.

3.3. Median Effect Analysis

The quantitative evaluation of the type of the interaction (i.e., synergistic, additive, or antagonistic) between TMZ and harmine was assessed using MTT viability measurement and data analysis by CompuSyn software, according to Chou and Talalay (17, 18). The combination index (CI) is an indicator of the type of the interaction between two agents. A CI value less than one is an indicator of synergism; a CI value equal to one shows an additive interaction, and a CI value greater than one reflects antagonism. Besides, the dose reduction index (DRI) was calculated. In this regard, a DRI value of less than one shows that using the combination of two drugs reduces the dose required for each when they are utilized alone. The combination of harmine and TMZ, with a constant ratio of 1: 2.66, was administered at two concentrations; above and below their IC_{50} values. The MTT test was performed again for the cotreatments. Using CompuSyn software, the plots of the effect vs. dose (Fa vs. dose), CI (Fa vs. CI), and DRI (Fa vs. DRI), as well as the normalized isobologram were drawn.

3.4. Migration Assay

The in vitro scratch test was used to evaluate the effects of TMZ and/or harmine on the cells' migration ability. The T98G cells were cultured in the logarithmic phase of growth in 6-well plates (1.2×10^6 cells/well). After the cells reached a density of 80%, a scratch was made in the middle of the wells by a sterile yellow pipette tip. The wells were washed three times with PBS, and 1.5 mL of the culture medium containing 800 μ M TMZ and/or 300 μ M harmine was added. After 24 hours, the wells were evaluated and photographed under a light microscope (19). The images were analyzed by Tathcratch software (MathWorks Inc).

3.5. Invasion Assay

To evaluate the effects of TMZ and/or harmine on the cells' invasion potential, the CytoSelect[™] 24-Well Cell Invasion Assay kit (Cell Biolabs, San Diego, CA) was used. First, the inserts were placed in a plate in the presence of 300 μ L serum-free medium in each well and incubated at 37°C for one hour. The hydrating medium was removed gently, and 500 μ L of the medium containing 10% FBS was transferred to the lower chamber of the plate. The cells (7×10^5) pretreated with 800 μ M TMZ and/or 300 μ M harmine were cultured in the upper chamber for 24 h. Then the cells adhering to the surface inside the matrigel were removed by a swab. The inserts were transferred to a well containing 400 μ L of the day solution and incubated for 10 min. Afterward, the inserts were gently rinsed with distilled water and dried. Finally, the inserts were transferred to an empty well, and 200 μ L of the extraction solution was added to each well, and the plate was placed on a rotary shaker for 10 min. The absorption of the samples was measured at 560 nm. By comparing the absorption intensity of the treated samples with that of the control, changes in the invasion ability of the cells under the influence of the treatments were evaluated.

3.6. Adhesion Assay

For this assay, the wells of a 96-well plate were coated with a matrigel solution (5 mg/mL) and allowed to dry at room temperature for an hour. The cells exposed to 800 μ M TMZ and/or 300 μ M harmine for 24 h were trypsinized and suspended in the culture medium. Five hundred cells were cultured in each well and incubated for 24 h. Then the wells were washed with PBS, and the remaining cells were fixed with 4% paraformaldehyde solution for 20 min and then stained with 5% crystal violet for 10 min. To quantify connected cells, violet crystals were dissolved in 70% ethanol, and the absorbance of each sample was measured at 570 nm (20).

3.7. Gene Expression Analysis

The effects of TMZ and/or harmine on the expression of the genes encoding matrix metalloproteinase (MMP)-2 and 9 were assessed by real-time PCR. Total RNA from both control and treated cells was extracted by the TRIzol reagent (Thermo Fisher Scientific, MA, USA), and RNA quality was tested by determining the absorbance ratios of A260/280 and A260/230, and its integrity was assessed by agarose gel electrophoresis. Complementary DNA (cDNA) synthesis was performed utilizing 1 μ g of the extracted RNA by a cDNA synthesis kit (Vivantis Technologies, Selangor DE, Malaysia) according to the manufacturer's protocol. As an internal control, β -actin was used. Thermal cycles were as follows: 15 min at 50°C for cDNA synthesis and 10 min at 95°C, followed by 40 cycles of 15 sec at 95°C to denature DNA and 45 sec at 60°C to anneal and extend the template for the PCR reaction. Real-time PCR was conducted by SYBR Premix Ex Tag Technology (Takara Bio Inc., Shiga, Japan) in the Applied Biosystems StepOne real-time PCR system. All primers were designed utilizing GeneRunner software, synthesized by CinnaGen Co. (Tehran, Iran), and checked at the NCBI Primer Blast. The primers' sequences were as follows: (1) MMP-2: (F) 5'TTGGCAGTG-CAATACCTGAA3', and (R) 5'GAGTCCGTCCTTACCGTCAA3'; (2) MMP-9: (F) 5'CATCGTCATCCAGTTTGGTG3', and (R) 5'CA-GAAGCCCCACTTCTTGTC3', and (3) β -actin: (F) 5'GTGGGCGC-CCAGGCACCA3', and (R) 5'CTCCTTAATGTCACGCACGATTT3'.

3.8. Statistical Analysis

All experimental tests were independently repeated at least three times. The data were presented as mean \pm standard deviation (SD). Statistical analysis was conducted in SPSS version 16.0 software using one-way analysis of variance and Tukey's post-hoc test, and differences were considered significant at P < 0.05.

4. Results

4.1. Cytotoxic Effects of TMZ and Harmine on T98G Cells

Since the main goal of this study was to investigate the potential of harmine in improving the toxicity of TMZ on T98G cells, first, the effect of each agent on cell viability was investigated independently. The effects of 1.56, 3.12, 6.25, 12.5, 25, 50, 100, and 200 μ M doses of TMZ and harmine were investigated on cell survival and proliferation after 24, 48, 72, and 96 h of exposure (Figure 1A and B). The results revealed that both TMZ and harmine reduced the viability of the cells in a concentration- and time-dependent manner. Table 1 shows IC₅₀ values.

4.2. Synergistic Effects of TMZ and Harmine on T98G Cells

The effect-dose plot showed that the combination of the two agents was more toxic than each of them alone (Figure 2A). The calculated CI values for all five cotreatments were less than one, indicating a synergistic interaction between TMZ and harmine in reducing cell viability (Figure 2B). The DRI values of TMZ and harmine were greater than one, indicating a decrease in the dose required to produce a specific therapeutic effect in both cases (Figure 2C). Finally, the isobologram diagram was drawn (Figure 2D), in which the concentrations of TMZ and harmine, either alone and in combination, reducing the cellular population by 50, 75, and 90% were plotted. In this diagram, the localization of compound points on the chord, its bottom, and its top represented additive, synergistic, and antagonistic interactions, respectively.

4.3. Inhibitory Effect of TMZ and Harmine on GBM cells' Migration, Invasion, and Adhesion

The analysis of the images obtained from the scratch test by Tscratch software showed that both TMZ and harmine decreased cellular migration capacity, with the combination of the two agents revealing a greater inhibitory effect than either alone (Figure 3). As shown in Figure 4, TMZ and harmine reduced cellular invasion potential, with a greater inhibitory effect in the combined compared to individual treatments. The results of the adhesion assay showed that TMZ and harmine reduced cellular adhesion, and the combination treatment had a greater inhibitory effect than either agent alone (Figure 5).

4.4. Effects of TMZ and Harmine on MMP-2 and MMP-9 Gene Expression in GBM Cells

Real-time PCR data revealed that both MMP-2 and MMP-9 were downregulated by TMZ and/or harmine (Figure 6).

5. Discussion

In this study, the effects of TMZ (an FDA-approved chemotherapy drug for GBM) and harmine (the main active ingredient of *P. harmala* seeds), both alone and in combination, were investigated on the cell viability of a GBM cell line. The results showed that at all doses of TMZ, cell viability gradually decreased in a concentration-dependent manner. After 24 h of treatment, cellular viability significantly reduced at the concentrations of 50, 100, and 200 μ M compared to the control group. After 48 and 72 h of treatment, the effects of TMZ on cellular viability were significant at the 3.12, 6.25, 12.5, 25, 50, 100, and 200 μ M concentrations, and after 96 h treatment, the cells' viability significantly reduced at all concentrations. Also, harmine gradually decreased cell viability in a dose-dependent way. After 24 h of treatment, viability significantly diminished at the concentrations of 6.25, 12.5, 25, 50, 100, and 200 μ M compared to the control. Following 48 h of treatment, the reduction in cellular viability was significant at concentrations of 3.12, 6.25, 12.5, 25, 50, 100, and 200 μ M. After 72 and 96 h, all concentrations of harmine significantly reduced viability compared to the control group.

Previous in vitro and in vivo studies have demonstrated that harmine shows significant anticancer properties, including reducing cellular proliferation (21), migration (13), and invasion (22), as well as activating apoptosis (13) and preventing tumorigenesis. The cytotoxic effects of harmine have been noted against some cancers, such as gastric (23), lung (24), breast (25), and hepatic (26). It appears that harmine arrests the cell cycle at the G0/G1 phase (27), reduces cyclin-dependent kinase activity (28), induces autophagy and apoptosis, enhances the level of pro-apoptotic factors, and decreases the expression of pro-inflammatory cytokines (21). Harmine induces autophagy via upregulating LC3-II and down-regulating P62 (29) and also suppresses the expression of pro-metastatic genes such as MMP-9 and ERK, as well as vascular endothelial growth factors to reduce cancer invasion (30).

Despite recent advances in cancer treatment, there are still no significant improvements in GBM patients' life expectancy and quality of lives, mostly because of drug resistance. Therefore, the development of new therapeutic strategies is necessary to overcome this problem. Recently, drug combinations have been widely used to treat fatal diseases such as cancer and the acquired immunodeficiency syndrome. The main goal of this strategy is to achieve a synergistic therapeutic effect, decrease drug dose and toxicity, and minimize or delay the development of drug resistance. In fact, synergistic interactions can reduce drugs' toxicities and minimize the resistance of cancer cells against them. Today, the use of several anticancer drugs from different groups is widely applicable to treat various malignancies. According to research evidence, the combination of TMZ with other anticancer agents can boost its activity. For example, clinical studies have shown the benefi-



Figure 1. A, Harmine; and B, temozolomide effects on the viability of T98G cells. Viability was evaluated by the MTT assay after 24, 48, 72, and 96 h of treatment with the 1.56, 3.12, 6.25, 12.5, 25, 50, 100, and 200 μ M doses of temozolomide and harmine. The control group received the same volume of serum-free medium (* P < 0.05; ** P < 0.01; and *** P < 0.001 compared to the control).

Table 1. The IC ₅₀ Values of Temozolomide and Harmine Against the Glioblastoma Multiforme T98MG Cell Line after 24, 48, 72, and 96 h of Treatment ^a				
Variables	24 hr	48 hr	72 hr	96 hr
Temozolomide (µM)	802.27 ± 3.85	341.11± 7.26	138.93 ± 3.71	44.75 ± 6.28
Harmine (µM)	316.74 ± 5.04	62.68 ± 4.57	9.82 ± 5.75	6.60 ± 5.10

 a The values are presented as mean \pm SD from three independent experiments.

cial outcomes of adding chloroquine to the standard GBM therapeutic regimen (31). Also, the combination of TMZ and carmustine, as a new adjunctive therapy in patients with GBM, showed satisfactory effects and tolerable toxicity (32). In vitro studies have also shown that the anticancer effects of TMZ are augmented in combination with some natural anti-tumor agents (33-42). However, no studies have been performed on the efficacy of TMZ + harmine co-treatment in suppressing the growth of GBM cells.

The results of this study showed that harmine significantly increased the cytotoxicity of TMZ, with a combination index between 0.28 and 0.84, indicating a synergistic effect between the two agents in the co-treatment state. The mean combination index for all the tests was 0.48, reflecting an overall synergistic effect for the combination of TMZ and harmine against the T98G cell line. This combination reduced the concentration of TMZ and harmine required for a specific therapeutic effect. Using this combination, the calculated IC50 values of harmine and TMZ reduced 1.22 and 2.54 times, respectively. Reducing the dose of TMZ for creating a certain effect is clinically valuable because it reduces the general side effects of chemotherapy. In this study, for the first time, the synergistic effects of TMZ and harmine on the viability of GBM cells were demonstrated after combined treatment. Our data also revealed that TMZ and/or harmine decreased the migration, invasion, and adhesion activities of T98G cells.

Cellular migration and invasion are major features of malignant tumors, especially in GBM. Although GBM cells cannot develop metastasis to other organs, their invasion and proliferation in the brain tissue are among the leading causes of death in these patients. A disseminated growth pattern is one of the characteristics of this type of tumor (43). Therefore, this ability to invade the brain, along with inherent resistance to TMZ, is the main barrier to the successful treatment of this disease, so managing this invasive behavior can be a useful strategy for effectively treating GBM.

As proteolytic enzymes, MMPs play important roles in the metastasis, migration, invasion, growth, and angiogenesis of tumors. These enzymes catalyze the decomposition of various components of the extracellular matrix. Among more than 20 members of this enzymatic family, MMP-2 and MMP-9 are of particular importance in GBM studies because their expression is directly related to the grade of the malignancy and its progression rate. The expression of both MMP-2 and MMP-9 is increased in human glioma tissues compared to the normal brain tissue, especially in GBM tumors (43). Since tumor cells' ability to migrate across the intercellular matrix primarily depends on the secretion of MMPs, suppressing these enzymes can be a suitable therapeutic option (44). The results of our study showed that TMZ and/or harmine reduced the ability of T98G cells to migrate, adhere, and invade, accompanied by



Figure 2. The plots of A, effect-dose; B, combination index; C, dose reduction index; and D, isobologram for T98G cells after 24 h of treatment with temozolomide and harmine, alone and in combination with each other. Fa represents the fraction of the cells affected, and Fu represents the fraction of non-affected cells (i.e., treated and non-treated). The data were obtained using the MTT assay.

the downregulation of the genes of MMP-2 and MMP-9.

The results of previous studies have shown that concentrations below the lethal dose of TMZ reduce the invasive and migratory properties of GBM cell lines (45, 46). Also, harmine was shown to repress cellular migration and invasion in gastric cancer via reducing cyclo-oxygenase-2 gene expression and inhibiting angiogenesis and tumor growth by activating the p53 molecule in endothelial cells (13, 22).

5.1. Conclusion

According to the results of the present study, TMZ and harmine inhibited the cellular proliferation of a GBM cancerous cell line, with the combination of the two agents exerting a more prominent synergistic inhibitory effect. Harmine and TMZ also suppressed the migration, invasion, and adhesion capacities of GBM cells, and these effects were greater in the co-treatment state.

Footnotes

Authors' Contribution: C. Jalali was responsible for overall supervision. K. Mansouri revised the manuscript. M. Taghadosi and A. Ghanbari participated in study design, data evaluation, drafting the manuscript, and statistical analysis. M. Kamani contributed to experimental tests. All authors contributed to editing and approving the final version of the manuscript for submission.

Conflict of Interests: There is no conflict of interest.

Ethical Approval: IR.KUMS.REC.1399.866

Funding/Support: This work was financially supported by the Vice-Chancellor for Research of Kermanshah University of Medical Sciences, Kermanshah, Iran (grant No. 990770).



Figure 3. The effect of temozolomide and/or harmine on the migration ability of T98G cells was measured by the scratch test. The control group A, on day zero; B, after 24 h; C, in the presence of 300 μ M harmine; D, in the presence of 800 μ M temozolomide; and E, in the presence of a combination of the two agents; F, the column diagram of the average percentage of scratch closure in the T98G cell monolayer (* P < 0.05; ** P < 0.01; and *** P < 0.001 compared to the control).



Figure 4. Temozolomide and/or harmine effect on the invasion ability of T98G cells after 24 h was measured by the invasion test. A, the control group; B, in the presence of 300 μ M harmine; C in the presence of 800 μ M temozolomide; D, in the presence of a combination of the two agents; and E, the column diagram of the average absorption at 560 nm (* P < 0.05; ** P < 0.01; and *** P < 0.001 compared to the control).



Figure 5. Temozolomide and/or harmine effect on the adhesion ability of T98G cells after 24 h was measured by the adhesion test. A, the control group; B, after treatment with 300 μ M harmine; C, after treatment with 800 μ M temozolomide; D, after treatment with a combination of the two agents; and E, the column diagram of the average absorption of violet crystals at a wavelength of 590 nm (* P < 0.05; ** P < 0.01; and *** P < 0.001 compared to the control).



Figure 6. Temozolomide and/or harmine effect on the gene expressions of MMP-2 and MMP-9 in T98G cells after 24 h was measured by real-time PCR (* P < 0.05; ** P < 0.01; and *** P < 0.001 compared to the control).

References

- Walker C, Baborie A, Crooks D, Wilkins S, Jenkinson MD. Biology, genetics and imaging of glial cell tumours. *Br J Radiol.* 2011;84 Spec No 2:S90-106. [PubMed ID: 22433833]. [PubMed Central ID: PMC3473897]. https://doi.org/10.1259/bjr/23430927.
- Benson VS, Pirie K, Schuz J, Reeves GK, Beral V, Green J, et al. Mobile phone use and risk of brain neoplasms and other cancers: Prospective study. *Int J Epidemiol*. 2013;**42**(3):792–802. [PubMed ID: 23657200]. https://doi.org/10.1093/ije/dyt072.
- Barami K. Oncomodulatory mechanisms of human cytomegalovirus in gliomas. J Clin Neurosci. 2010;17(7):819–23. [PubMed ID: 20427188]. https://doi.org/10.1016/j.jocn.2009.10.040.
- Koh I, Cha J, Park J, Choi J, Kang SG, Kim P. The mode and dynamics of glioblastoma cell invasion into a decellularized tissuederived extracellular matrix-based three-dimensional tumor model. *Sci Rep.* 2018;8(1):4608. [PubMed ID: 29545552]. [PubMed Central ID: PMC5854588]. https://doi.org/10.1038/s41598-018-22681-3.
- Hanif F, Muzaffar K, Perveen K, Malhi SM, Simjee Sh U. Glioblastoma multiforme: A review of its epidemiology and pathogenesis through clinical presentation and treatment. *Asian Pac J Cancer Prev.* 2017;18(1):3–9. [PubMed ID: 28239999]. [PubMed Central ID: PMC5563115]. https://doi.org/10.22034/APJCP.2017.18.1.3.
- Westphal M, Lamszus K. The neurobiology of gliomas: from cell biology to the development of therapeutic approaches. *Nat Rev Neurosci.* 2011;12(9):495–508. [PubMed ID: 21811295]. https://doi.org/10.1038/nrn3060.
- Goellner EM, Grimme B, Brown AR, Lin YC, Wang XH, Sugrue KF, et al. Overcoming temozolomide resistance in glioblastoma via dual inhibition of NAD+ biosynthesis and base excision repair. *Cancer*

Res. 2011;**71**(6):2308-17. [PubMed ID: 21406402]. [PubMed Central ID: PMC3077901]. https://doi.org/10.1158/0008-5472.CAN-10-3213.

- Chien CH, Hsueh WT, Chuang JY, Chang KY. Dissecting the mechanism of temozolomide resistance and its association with the regulatory roles of intracellular reactive oxygen species in glioblastoma. *J Biomed Sci.* 2021;28(1):18. [PubMed ID: 33685470]. [PubMed Central ID: PMC7938520]. https://doi.org/10.1186/s12929-021-00717-7.
- Mansoori B, Mohammadi A, Davudian S, Shirjang S, Baradaran B. The different mechanisms of cancer drug resistance: A brief review. Adv Pharm Bull. 2017;7(3):339–48. [PubMed ID: 29071215]. [PubMed Central ID: PMC5651054]. https://doi.org/10.15171/apb.2017.041.
- Ekor M. The growing use of herbal medicines: Issues relating to adverse reactions and challenges in monitoring safety. *Front Pharmacol.* 2014;4:177. [PubMed ID: 24454289]. [PubMed Central ID: PMC3887317]. https://doi.org/10.3389/fphar.2013.00177.
- Woo CC, Kumar AP, Sethi G, Tan KH. Thymoquinone: potential cure for inflammatory disorders and cancer. *Biochem Pharmacol.* 2012;83(4):443–51. [PubMed ID: 22005518]. https://doi.org/10.1016/j.bcp.2011.09.029.
- Shen J, Wang B, Zhang T, Zhu N, Wang Z, Jin J, et al. Suppression of non-small cell lung cancer growth and metastasis by a novel small molecular activator of RECK. *Cell Physiol Biochem*. 2018;**45**(5):1807-17. [PubMed ID: 29510387]. https://doi.org/10.1159/000487872.
- Zhang H, Sun K, Ding J, Xu H, Zhu L, Zhang K, et al. Harmine induces apoptosis and inhibits tumor cell proliferation, migration and invasion through down-regulation of cyclooxygenase-2 expression in gastric cancer. *Phytomedicine*. 2014;21(3):348–55. [PubMed ID: 24176842]. https://doi.org/10.1016/j.phymed.2013.09.007.
- 14. Ding Y, He J, Huang J, Yu T, Shi X, Zhang T, et al. Harmine induces anticancer activity in breast cancer cells via targeting TAZ. *Int J On*-

col. 2019;**54**(6):1995-2004. [PubMed ID: 31081045]. [PubMed Central ID: PMC6521938]. https://doi.org/10.3892/ijo.2019.4777.

- Xu B, Li M, Yu Y, He J, Hu S, Pan M, et al. Effects of harmaline on cell growth of human liver cancer through the p53/p21 and Fas/FasL signaling pathways. *Oncol Lett.* 2018;15(2):1931-6. [PubMed ID: 29434892]. [PubMed Central ID: PMC5774346]. https://doi.org/10.3892/ol.2017.7495.
- Karakas D, Ari F, Ulukaya E. The MTT viability assay yields strikingly false-positive viabilities although the cells are killed by some plant extracts. *Turk J Biol.* 2017;41(6):919–25. [PubMed ID: 30814856]. [PubMed Central ID: PMC6353273]. https://doi.org/10.3906/biy-1703-104.
- Chou TC. Drug combination studies and their synergy quantification using the Chou-Talalay method. *Cancer Res.* 2010;**70**(2):440–6. [PubMed ID: 20068163]. https://doi.org/10.1158/0008-5472.CAN-09-1947.
- Sliwka L, Wiktorska K, Suchocki P, Milczarek M, Mielczarek S, Lubelska K, et al. The comparison of MTT and CVS assays for the assessment of anticancer agent interactions. *PLoS One*. 2016;11(5). e0155772. [PubMed ID: 27196402]. [PubMed Central ID: PMC4873276]. https://doi.org/10.1371/journal.pone.0155772.
- Pinto BI, Cruz ND, Lujan OR, Propper CR, Kellar RS. In vitro scratch assay to demonstrate effects of arsenic on skin cell migration. *J Vis Exp.* 2019;(144). [PubMed ID: 30855562]. [PubMed Central ID: PMC7537821]. https://doi.org/10.3791/58838.
- Liu Y, Cao W, Zhang B, Liu YQ, Wang ZY, Wu YP, et al. The natural compound magnolol inhibits invasion and exhibits potential in human breast cancer therapy. *Sci Rep.* 2013;3:3098. [PubMed ID: 24226295].
 [PubMed Central ID: PMC3827615]. https://doi.org/10.1038/srep03098.
- Chen Q, Chao R, Chen H, Hou X, Yan H, Zhou S, et al. Antitumor and neurotoxic effects of novel harmine derivatives and structure-activity relationship analysis. *Int J Cancer.* 2005;**114**(5):675–82. [PubMed ID: 15609303]. https://doi.org/10.1002/ijc.20703.
- 22. Dai F, Chen Y, Song Y, Huang L, Zhai D, Dong Y, et al. A natural small molecule harmine inhibits angiogenesis and suppresses tumour growth through activation of p53 in endothelial cells. *PLoS One*. 2012;7(12). e52162. [PubMed ID: 23300602]. [PubMed Central ID: PMC3531399]. https://doi.org/10.1371/journal.pone.0052162.
- Li C, Wang Y, Wang C, Yi X, Li M, He X. Anticancer activities of harmine by inducing a pro-death autophagy and apoptosis in human gastric cancer cells. *Phytomedicine*. 2017;28:10–8. [PubMed ID: 28478808]. https://doi.org/10.1016/j.phymed.2017.02.008.
- Abe A, Yamada H. Harmol induces apoptosis by caspase-8 activation independently of Fas/Fas ligand interaction in human lung carcinoma H596 cells. *Anticancer Drugs*. 2009;20(5):373-81. [PubMed ID: 19318910]. https://doi.org/10.1097/CAD.0b013e32832a2dd9.
- Hashemi Sheikh Shabani S, Seyed Hasan Tehrani S, Rabiei Z, Tahmasebi Enferadi S, Vannozzi GP. Peganum harmala L.'s anti-growth effect on a breast cancer cell line. *Biotechnol Rep (Amst)*. 2015;8:138-43. [PubMed ID: 28352583]. [PubMed Central ID: PMC4980734]. https://doi.org/10.1016/j.btre.2015.08.007.
- Zhang L, Zhang F, Zhang W, Chen L, Gao N, Men Y, et al. Harmine suppresses homologous recombination repair and inhibits proliferation of hepatoma cells. *Cancer Biol Ther.* 2015;16(11):1585– 92. [PubMed ID: 26382920]. [PubMed Central ID: PMC4846143]. https://doi.org/10.1080/15384047.2015.1078021.
- Hamsa TP, Kuttan G. Harmine activates intrinsic and extrinsic pathways of apoptosis in B16F-10 melanoma. *Chin Med.* 2011;6(1):11. [PubMed ID: 21429205]. [PubMed Central ID: PMC3076298]. https://doi.org/10.1186/1749-8546-6-11.
- Liu P, Li H, Wang Y, Su X, Li Y, Yan M, et al. Harmine ameliorates cognitive impairment by inhibiting NLRP3 inflammasome activation and enhancing the BDNF/TrkB signaling pathway in STZ-induced diabetic rats. Front Pharmacol. 2020;11:535. [PubMed ID: 32425784]. [PubMed

Central ID: PMC7206617]. https://doi.org/10.3389/fphar.2020.00535.

- Zou N, Wei Y, Li F, Yang Y, Cheng X, Wang C. The inhibitory effects of compound Muniziqi granule against B16 cells and harmine induced autophagy and apoptosis by inhibiting Akt/mTOR pathway. *BMC Complement Altern Med.* 2017;**17**(1):517. [PubMed ID: 29197358]. [PubMed Central ID: PMC5712103]. https://doi.org/10.1186/s12906-017-2017-4.
- Hamsa TP, Kuttan G. Harmine inhibits tumour specific neo-vessel formation by regulating VEGF, MMP, TIMP and pro-inflammatory mediators both in vivo and in vitro. *Eur J Pharmacol.* 2010;**649**(1-3):64–73. [PubMed ID: 20858484]. https://doi.org/10.1016/j.ejphar.2010.09.010.
- Golden EB, Cho HY, Jahanian A, Hofman FM, Louie SG, Schonthal AH, et al. Chloroquine enhances temozolomide cytotoxicity in malignant gliomas by blocking autophagy. *Neurosurg Focus*. 2014;**37**(6). E12. [PubMed ID: 25434381]. https://doi.org/10.3171/2014.9.FOCUS14504.
- 32. Chang SM, Prados MD, Yung WK, Fine H, Junck L, Greenberg H, et al. Phase II study of neoadjuvant 1, 3-bis (2-chloroethyl)-1-nitrosourea and temozolomide for newly diagnosed anaplastic glioma: A North American Brain tumor consortium trial. *Cancer*. 2004;**100**(8):1712–6. [PubMed ID: 15073861]. https://doi.org/10.1002/cncr.20157.
- Torres S, Lorente M, Rodriguez-Fornes F, Hernandez-Tiedra S, Salazar M, Garcia-Taboada E, et al. A combined preclinical therapy of cannabinoids and temozolomide against glioma. *Mol Cancer Ther.* 2011;10(1):90–103. [PubMed ID: 21220494]. https://doi.org/10.1158/1535-7163.MCT-10-0688.
- Chen M, Rose AE, Doudican N, Osman I, Orlow SJ. Celastrol synergistically enhances temozolomide cytotoxicity in melanoma cells. *Mol Cancer Res.* 2009;7(12):1946–53. [PubMed ID: 19934274]. https://doi.org/10.1158/1541-7786.MCR-09-0243.
- Osmond GW, Augustine CK, Zipfel PA, Padussis J, Tyler DS. Enhancing melanoma treatment with resveratrol. J Surg Res. 2012;172(1):109–15. [PubMed ID: 20855085]. https://doi.org/10.1016/j.jss.2010.07.033.
- Qi Q, Liu X, Li S, Joshi HC, Ye K. Synergistic suppression of noscapine and conventional chemotherapeutics on human glioblastoma cell growth. *Acta Pharmacol Sin.* 2013;34(7):930– 8. [PubMed ID: 23708557]. [PubMed Central ID: PMC4002615]. https://doi.org/10.1038/aps.2013.40.
- Moskwa J, Borawska MH, Markiewicz-Zukowska R, Puscion-Jakubik A, Naliwajko SK, Socha K, et al. Polish natural bee honeys are anti-proliferative and anti-metastatic agents in human glioblastoma multiforme U87MG cell line. *PLoS One*. 2014;9(3). e90533. [PubMed ID: 24594866]. [PubMed Central ID: PMC3942434]. https://doi.org/10.1371/journal.pone.0090533.
- Markiewicz-Zukowska R, Borawska MH, Fiedorowicz A, Naliwajko SK, Sawicka D, Car H. Propolis changes the anticancer activity of temozolomide in U87MG human glioblastoma cell line. *BMC Complement Altern Med.* 2013;13:50. [PubMed ID: 23445763]. [PubMed Central ID: PMC3598711]. https://doi.org/10.1186/1472-6882-13-50.
- Jakubowicz-Gil J, Langner E, Badziul D, Wertel I, Rzeski W. Apoptosis induction in human glioblastoma multiforme T98G cells upon temozolomide and quercetin treatment. *Tumour Biol.* 2013;34(4):2367-78. [PubMed ID: 23580181]. [PubMed Central ID: PMC3713258]. https://doi.org/10.1007/s13277-013-0785-0.
- 40. Lin CJ, Lee CC, Shih YL, Lin TY, Wang SH, Lin YF, et al. Resveratrol enhances the therapeutic effect of temozolomide against malignant glioma in vitro and in vivo by inhibiting autophagy. *Free Radic Biol Med.* 2012;**52**(2):377-91. [PubMed ID: 22094224]. https://doi.org/10.1016/j.freeradbiomed.2011.10.487.
- Filippi-Chiela EC, Thome MP, Bueno e Silva MM, Pelegrini AL, Ledur PF, Garicochea B, et al. Resveratrol abrogates the temozolomideinduced G2 arrest leading to mitotic catastrophe and reinforces the temozolomide-induced senescence in glioma cells. *BMC Cancer*. 2013;13:147. [PubMed ID: 23522185]. [PubMed Central ID: PMC3635906]. https://doi.org/10.1186/1471-2407-13-147.

- Borges KS, Brassesco MS, Scrideli CA, Soares AE, Tone LG. Antiproliferative effects of Tubi-bee propolis in glioblastoma cell lines. *Genet Mol Biol.* 2011;34(2):310–4. [PubMed ID: 21734835]. [PubMed Central ID: PMC3115328]. https://doi.org/10.1590/s1415-47572011000200024.
- Konnecke H, Bechmann I. The role of microglia and matrix metalloproteinases involvement in neuroinflammation and gliomas. *Clin Dev Immunol*. 2013;2013:914104. [PubMed ID: 24023566]. [PubMed Central ID: PMC3759277]. https://doi.org/10.1155/2013/914104.
- Kessenbrock K, Plaks V, Werb Z. Matrix metalloproteinases: Regulators of the tumor microenvironment. Cell. 2010;141(1):52-

67. [PubMed ID: 20371345]. [PubMed Central ID: PMC2862057]. https://doi.org/10.1016/j.cell.2010.03.015.

- 45. Wick W, Wick A, Schulz JB, Dichgans J, Rodemann HP, Weller M. Prevention of irradiation-induced glioma cell invasion by temozolomide involves caspase 3 activity and cleavage of focal adhesion kinase. *Cancer Res.* 2002;**62**(6):1915–9. [PubMed ID: 11912174].
- Shen W, Hu JA, Zheng JS. Mechanism of temozolomide-induced antitumour effects on glioma cells. J Int Med Res. 2014;42(1):164–72. [PubMed ID: 24326954]. https://doi.org/10.1177/0300060513501753.