



Effect of Pretreatment Time with Enhancers on Caffeine Skin Permeability in Rats

Saeed Mohammad Soleymani ¹, Anayatollah Salimi ^{2,3,*}, Heibatullah Kalantari ⁴ and Adel Sheykhi ⁵

¹Department of Clinical Pharmacy, Faculty of Pharmacy, Shahid Beheshti University of Medical Sciences, Tehran, Iran

²Nanotechnology Research Center, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran

³Department of Pharmaceutics, Faculty of Pharmacy, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran

⁴Department of Toxicology, Faculty of Pharmacy, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran

⁵Student Research Committee, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran

*Corresponding author: Department of Pharmaceutics, Faculty of Pharmacy, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran. Email: anayatsalimi2003@yahoo.com

Received 2023 June 03; Revised 2023 October 04; Accepted 2023 October 13.

Abstract

Background: Caffeine is an edible chemical compound obtained from various plants, such as tea and coffee. Caffeine is an alkaloid that is highly hydrophilic and has limited skin permeability. The lipophilic nature of the stratum corneum is a major barrier to the passage of this substance through the skin. Topical drug delivery systems can effectively transfer caffeine to the skin.

Objectives: This study investigated the effect of pretreatment time with chemical enhancers on caffeine's skin permeation.

Methods: The skin was subjected to additives such as sodium lauryl sulfate, sodium lauryl ethyl sulfate, tynoline, nanoxinol, and lecithin for 5, 15, and 30 minutes. Then, the parameters of caffeine permeability and structural changes in the skin due to additive adsorption were studied using Fourier Transform Infrared (FT-IR) spectrometry.

Results: The enhancers increased the permeation of caffeine through the skin. There are different mechanisms for penetration enhancers, including lipid liquefaction, disruption of lipid bilayers, and irreversible denaturation of intracellular keratin.

Conclusions: Sodium lauryl sulfate can affect the skin permeability of caffeine.

Keywords: Enhancing Absorption, Skin Permeability, Caffeine, FT-IR, Sodium Lauryl Sulfate

1. Background

Caffeine is an edible chemical compound found in a variety of herbs, including the seed of the *Coffea* plant and *Camellia sinensis*, and is a stimulant that can prevent sleep deprivation. Caffeine is an alkaloid in the methyl xanthine tribe that has similar properties to theophylline (1). Pure caffeine powder is soft, odorless, white, shiny, and bitter. Caffeine is the most extensively used drug among people. Almost 90% of people use it daily. Increasing the body's metabolism, stimulating the central nervous system, and increasing awareness and perception of the environment are the main effects of caffeine (2).

Most drugs pass through human skin so slowly that they cannot exert a sufficient systemic effect (3). The stratum corneum is the main limiting layer for topical drug delivery. Therefore, the main challenge for topical formulation today is to achieve an increase in drug penetration into the skin (4). So far, many studies

have been carried out to identify and investigate the mechanism of various factors affecting the increase in permeability and adsorbent through the skin. These agents are used in designing various drug formulations under the name of adsorbent and in various natural and synthetic forms. Chemical enhancers are materials that increase the adsorbent of drugs from the skin layers and accelerate absorption (5).

Caffeine is a compound that is highly hydrophilic and has limited skin permeability. The lipophilic nature of the stratum corneum is a major barrier to the passage of this substance through the skin, so it is important to develop topical drug delivery systems that can effectively transport caffeine into the skin (1).

2. Objectives

Topical use of this substance can increase the effectiveness of the drug in the local area about local

treatment goals (such as hair loss) while reducing the side effects caused by oral use. Given all this, the effect of several chemical enhancers (Sodium Lauryl Sulfate [SLS], Sodium Lauryl Ethyl Sulfate [SLES], tynoline, nonoxinol, and lecithin) on the skin permeation of caffeine was investigated in this study. It is expected that the results may be effective in the skin permeability of caffeine and in designing a new topical formulation.

3. Methods

3.1. Materials

Caffeine powder, SLS, SLES, tynoline, nonoxinol, and lecithin were prepared from Behsa Daruo Pharmaceutical Company, Iran.

3.1.1. Animals

The present study was conducted on 10- to 12-week-old male Wistar rats (150 - 170 g). The Ethics Committee of Jundishapur University of Medical Sciences (IR.AJUMS.ABHC.REC.1398.053) provided ethical approval for this study. All rats involved in the study underwent anesthesia via ketamine/xylazine at a dose of 50/10 mg/kg before being relieved. The rats were humanely sacrificed. Hairs from the specimens' abdomens were then cut and collected, followed by skin removal. The skin samples were stored at a temperature of -20°C in a freezer until permeability tests (6).

3.2. Caffeine Determination

A valid measurement method is required to accurately calculate how much of an administered drug penetrates the skin. The protocol employed in the present study included UV spectroscopy. The wavelength chosen was 273 nm (per permeability and release studies), which corresponds to the caffeine's spectral absorption spectrum when integrated into a phosphate buffer solution with a pH of 7. During the first step, caffeine was added to the solution at an amount of 5 mg in 10 mL (0.5 mg/mL).

The linearity of the relationship between the concentration and the response of the device (in this study, a UV spectrophotometer) was used to measure caffeine. The response produced in UV spectroscopy is light absorption. At least four standard concentrations of 0.0025, 0.005, 0.015, and 0.05 mg/mL of the drug were prepared in phosphate buffer (pH 7) to investigate the relationship between concentration and response. Four concentrations of 0.0025, 0.005, 0.015, and 0.05 mg/mL of caffeine in phosphate buffer (pH 7) were prepared to establish a standard curve of stock solution with a concentration of 0.5 mg/mL. The absorbance of the samples was measured by UV spectrophotometry. This method was repeated for 3 consecutive days, then the calibration curve was drawn using different

concentrations in relation to atomic absorption, and the permeability of the drug was obtained by each of the enhancers (6).

3.3. Investigation of Caffeine Permeability

Rat skin was cut into pieces and placed on the Franz diffusion cell. One gram of one of the chemical enhancers was taken and poured on the skin. Then, the enhancer was removed from the skin, and the receptor phase was filled with 35 mL of phosphate buffer solution (pH 7). This process was repeated separately for each time (5, 15, or 30 minutes) and chemical enhancer. Meanwhile, 5 mL of a 0.05% drug solution was added during the donor phase.

In the next step, all Franz diffusion cells were placed on a heater (37°C) and stirred by a magnet at 200 rpm. Then, 2 mL of the receiver phase was removed after the elapse of certain predetermined periods (0.5, 1, 2, 3, 4, 5, 6, 7, 8, and 24 hours). Next, 2 mL of phosphate buffer with pH 7 was withdrawn, and the absorption of the sample was read with UV spectrophotometry. As a negative control for permeability, the skin was placed on the bulb. Only distilled water was used in the donor phase, while phosphate buffer (pH 7) was used in the receptor chamber to set the instrument to zero. Then, the water on the donor phase was removed, and the drug was placed instead, and the permeability factors of the drug were measured for the negative control (7).

3.4. Statistical Analysis and Permeability Calculation

Each experiment was repeated three times, and the data were presented as a mean \pm standard deviation (SD). The results were then analyzed statistically using a two-sample *t*-test and analysis of variance. Furthermore, a full factorial test was devised using Minitab 17 software (8).

This study was designed to measure caffeine's permeability from rat skin. Additional parameters of interest included steady-state flux (J_{ss}), permeability coefficient (P), incubation time (T_{lag}), and apparent diffusion coefficient (D_{app}). Findings comparing the drug-containing microemulsions' ER_{flux} , ER_D , and ER_p levels to those of the drug-saturated control samples are shown in Tables 1 and 2. The total amounts of the drug that passed through the skin surface were plotted against time to determine the values for the permeability parameters mentioned above.

Equation was employed to calculate the permeability coefficient (p) (7),

$$J_{ss} = P.C \quad (1)$$

Where C represents the concentration of the drug during the donor phase.

Moreover, Equation was utilized to calculate D (diffusion coefficient) (2, 7):

Table 1. Effects of Enhancers' Adsorption on Caffeine Permeability Parameters of Whole Rat Skin (n = 3)^a

Parameter	Flux (mg.cm ⁻² .h ⁻¹)	T _{lag} (h)	D (cm ² .h ⁻¹)	P (cm/h)
After 5 Minutes of Skin Contact				
Control (water)	0.001 ± 0.00015	5.71 ± 0.07	0.01865 ± 0.0002	0.00018 ± 0.00002
SLS	0.007 ± 0.00060	4.35 ± 0.09	0.02450 ± 0.0005	0.00076 ± 0.00006
SLES	0.001 ± 0.00006	3.03 ± 0.03	0.03514 ± 0.001	0.00017 ± 0.00001
Tynolin	0.018 ± 0.00046	5.71 ± 0.01	0.01868 ± 0.0004	0.00181 ± 0.00005
Nanoxinol	0.010 ± 0.00040	5.94 ± 0.11	0.01796 ± 0.00028	0.00103 ± 0.00004
Lecithin	0.006 ± 0.00021	1.57 ± 0.001	0.06796 ± 0.003	0.00065 ± 0.00002
After 15 Minutes of Skin Contact				
Control (water)	0.001 ± 0.00015	5.71 ± 0.07	0.01865 ± 0.0002	0.00018 ± 0.00002
SLS	0.007 ± 0.00060	4.35 ± 0.09	0.02450 ± 0.0005	0.00076 ± 0.00006
SLES	0.001 ± 0.00006	3.03 ± 0.03	0.03514 ± 0.001	0.00017 ± 0.00001
Tynolin	0.018 ± 0.00046	5.71 ± 0.01	0.01868 ± 0.0004	0.00181 ± 0.00005
Nanoxinol	0.010 ± 0.00040	5.94 ± 0.11	0.01796 ± 0.00028	0.00103 ± 0.00004
Lecithin	0.006 ± 0.00021	1.57 ± 0.001	0.06796 ± 0.003	0.00065 ± 0.00002
After 30 Minutes of Skin Contact				
Control (water)	0.001 ± 0.0001	5.71 ± 0.07	0.018 ± 0.0002	0.00018 ± 0.00002
SLS	0.009 ± 0.0006	2.005 ± 0.14	0.053 ± 0.0033	0.0009 ± 0.00006
SLES	0.003 ± 0.0002	0.05 ± 0.0008	1.843 ± 0.0849	0.0003 ± 0.00002
Tynolin	0.017 ± 0.0004	5.44 ± 0.05	0.019 ± 0.0002	0.0017 ± 0.00005
Nanoxinol	0.014 ± 0.001	4.70 ± 0.20	0.022 ± 0.001	0.0014 ± 0.00011
Lecithin	0.003 ± 0.0003	1.588 ± 0.19	0.068 ± 0.009	0.0003 ± 0.00003

^a Values are expressed as mean ± SD.

$$D = \frac{h^2}{6T_{lag}} \quad (2)$$

In the above equation, T_{lag} denotes the commune time (per the drug curve) from the skin along the equilibrium line to the time axis, while his skin diameter.

Note that D, as determined by Equation equals the apparent D. This is because h is not representative of the true drug passage length. Because the calculations made using the above equations depend on the steady-state range of the cumulative drug passage chart, the parameters are not valid unless the appropriate synchronous conditions are met. During the receptor phase, concentrations did not reach 10% of the drug saturation solution.

4. Results

The permeability of caffeine was studied after skin contact with absorption enhancers for 5, 15, and 30 minutes to evaluate the effect of absorption enhancers on skin permeability. The parameters of caffeine permeability for each component are given in Tables 1 and 2. The

cumulative graph of caffeine passing through the surface unit before contact for 5, 15, and 30 minutes is shown in Figures 1 - 3. Phosphate buffer was used as a control in the phase chamber.

The FT-IR spectrum analysis (before adsorption) evaluated chemical adsorbent-stratum corneum interactions in the skin by observing absorption bands at various wavelengths. The bands observed via this analysis indicate lipid and protein molecule vibrations within the stratum corneum (9). Tables 3 - 5 show the FT-IR data depicting how adsorption affected the rat skin samples.

5. Discussion

The impact of additive uptake on drug passage through rat skin compared to control was determined by calculating ER_{flux}, ER_p, and ER_D. The results showed that the adsorption of SLS, SLES, tynoline, nanoxinol, and lecithin caused a significant increase after ER_{flux}, ER_p, and ER_D at 5-, 15- and 30-minutes' contact with skin (P < 0.05). While lecithin was not significantly different from the control at five-minute pre-exposure, 15- and 30-minute

Table 2. Results for ER_{flux}, ER_D and ER_P, and Uptake (n = 3)^{a, b}

Parameter	ER _{flux}	ER _D	ER _P
After 5 Minutes of Skin Contact			
SLS	4.18 ± 0.18	1.24 ± 0.02	4.18 ± 0.18
SLES	1.52 ± 0.28	1.35 ± 0.02	1.52 ± 0.28
Tynolin	5.16 ± 0.27	2.86 ± 0.21	5.16 ± 0.27
Nanoxinol	5.48 ± 0.56	0.82 ± 0.007	5.48 ± 0.56
Lecithin	1.30 ± 0.03	1.07 ± 0.05	1.30 ± 0.03
After 15 Minutes of Skin Contact			
SLS	4.31 ± 0.43	1.31 ± 0.04	4.31 ± 0.43
SLES	0.98 ± 0.08	1.88 ± 0.06	0.98 ± 0.08
Tynolin	10.31 ± 1.18	1.001 ± 0.01	10.31 ± 1.18
Nanoxinol	5.85 ± 0.70	0.964 ± 0.01	5.85 ± 0.70
Lecithin	3.67 ± 0.34	3.64 ± 0.18	3.67 ± 0.34
After 30 Minutes of Skin Contact			
SLS	5.16 ± 0.27	2.86 ± 0.21	5.16 ± 0.27
SLES	1.78 ± 0.17	98.87 ± 5.37	1.78 ± 0.17
Tynolin	9.82 ± 0.89	1.05 ± 0.008	9.82 ± 0.89
Nanoxinol	8.24 ± 1.30	1.21 ± 0.08	8.24 ± 1.30
Lecithin	2.15 ± 0.06	3.65 ± 0.54	2.15 ± 0.06

^a Values are expressed as mean ± SD.

^b ER = Caffeine permeability parameter from treated skin/Caffeine permeability parameter from hydrated skin).




Figure 1. Caffeine cumulative diagram crossing the surface unit after 5 minutes of skin contact with enhancers from whole rat skin (SLS: sodium lauryl sulfate, TAN: tynolin, NANO: nanoxinol, SLES: sodium lauryl ethyl sulfate, LES: lecithin).




Figure 2. Caffeine cumulative diagram crossing the surface unit after 15 minutes of skin contact with enhancers from whole rat skin (SLS: sodium lauryl sulfate, TAN: tynolin, NANO: nanoxinol, SLES: sodium lauryl ethyl sulfate, LES: lecithin).




Figure 3. Caffeine cumulative diagram crossing the surface unit after 30 minutes of skin contact with enhancers from whole rat skin (SLS: sodium lauryl sulfate, TAN: tynolin, NANO: nanoxinol, SLES: sodium lauryl ethyl sulfate, LES: lecithin).

pre-exposure to this combination resulted in a significant increase in caffeine skin passage ($P < 0.05$).

The results similarly demonstrated a significant relationship between J_{ss} and the adsorption of the sample after 5, 15, and 30 minutes compared to the control sample, indicating an increase in permeability with the adsorption studied ($P < 0.05$). According to a study by Tabosa et al., high J_{ss} can lead to higher concentrations of the drug in the upper layers of the skin. It may also enhance the penetration of the components of the formulation into the bloodstream through the skin (10, 11) because, in any formulation, all components can act as penetration components with different mechanisms (12).

The results revealed that adsorption of all enhancers had a greater increase in drug delivery rate (flux) than in drug release (D_{app}) (except for nanoxinol, there was no significant difference in the permeability increase at 5 and 15 minutes, but permeability increased at 30 minutes). Among them, tynolin and lecithin showed the greatest effect on increasing the permeability of rat skin at 5 and 15 minutes, respectively.

Also, the results showed that after 30 minutes of using all the enhancers, J_{ss} had a significant increase compared to the control group, which indicates the increase in drug permeability by using the enhancers after 30 minutes ($P < 0.05$).

Table 3. Post-component Effect Peak Heights (and Peak Height Percentage Reductions) for Symmetric and Asymmetric, as Well as -CH and C=O Tensile Groups (n = 3)^{a, b}

Enhancer	Asymmetric C-H Stretching		Symmetric C-H Stretching		C=O stretching of Lipid Ester	
	Peak Height	D%	Peak Height	D%	Peak Height	D%
Control (Water)	0.395	-	0.242	-	0.015	-
SLS	2.916	N. S	0.807	N. S	1.164	N. S
SLES	1.531	N. S	1.121	N. S	3.747	N. S
Tynolin	1.717	N. S	0.989	N. S	1.280	N. S
Nanoxinol	0.745	N. S	0.538	N. S	0.850	N. S
Lecithin	0.450	N. S	0.590	N. S	1.450	N. S

Abbreviation: N. S: Not seen.

^a Values are expressed as mean ± SD.

^b %Decrease in peak height (%D) = ((Peak height from untreated whole skin - Peak height from solvent-treated whole skin)/[Peak height from untreated whole skin] × 100)

Table 4. Post-component Effect Peak Heights (and Peak Height Percentage Reductions) for C=O Tensile and C-N Tensile Groups (n = 3)^a

Enhancer	C=O Stretching of Keratin		C-N Stretching of Keratin	
	Peak Height	% D	Peak Height	% D
Control (Water)	0.012	-	0.414	-
SLS	2.490	N. S	0.708	N. S
SLES	1.444	N. S	1.438	N. S
Tynolin	1.555	N. S	1.096	N. S
Nanoxinol	1.610	N. S	1.118	N. S
Lecithin	0.650	N. S	0.770	N. S

Abbreviation: N. S: Not seen.

^a Values are expressed as mean ± SD.

Table 5. Wavelengths for -CH and C=O groups in Hydrated Skin and Pre-contacted Skin (n = 3)^a

Enhancer	C-H Stretching Asy	C-H Stretching Sym	C=O Stretching of Lipid Ester	Amid I	Amid II
Control (Water)	2915.13 ± 0.2	2840.74 ± 0.4	1690.2 ± 1	1642.58 ± 0.9	1535.9 ± 0.2
SLS	2999.98 ± 0.15	2863.51 ± 0.11	1760.89 ± 0.24	1640.87 ± 0.21	1580.65 ± 0.17
SLES	2946.49 ± 0.1	2845.88 ± 0.1	1725.45 ± 0.1	1676.76 ± 0.9	1537.27 ± 0.2
Tynolin	2960.66 ± 0.15	2880.40 ± 0.11	1769.663 ± 0.24	1599.42 ± 0.21	1502.55 ± 0.17
Nanoxinol	2917 ± 0.382	2855.32 ± 0.229	1760.89 ± 0.19	1640.87 ± 0.404	1629.58 ± 0.258
Lecithin	2916.92 ± 0.15	2808.49 ± 0.11	1758.58 ± 0.24	1672.06 ± 0.21	1672.06 ± 0.17

^a Values are expressed as mean ± SD.

The highest and lowest absorption efficiencies of the drug were related to the increase in dermal absorption of the drug, respectively.

In order from the highest to the lowest increase in permeability belongs to the use of lecithin, SLES, SLS, nanoxinol, and tynolin (5 minutes); SLES, tynolin, nanoxinol, lecithin, and SLS (15 minutes); SLES, lecithin, tynolin, nanoxinol, and SLS (30 minutes).

According to the results of FT-IR from pre-contact skin, the absorption wavelength within the asymmetric CH region was improved after the application of sodium

lauryl sulfate. Also, the absorption wavelength's position advanced in the symmetric CH. These findings signify that liquefaction (and subsequent impairments to barrier properties) occurred in the stratum corneum membrane. It is also possible that increasing the passage of the drug through the skin's surface caused the C=O wavelength of stress to increase, which indicates that the hydrogen bonds weakened between lipid molecules. It also reduces amide I's wavelength number while increasing that of amide II; as a result, the related dam effects are increased and decreased, respectively. These results indicate that this

compound affects both the lipid bilayers and protein parts of horny tissues. Nevertheless, this adsorption process's effect on the lipid parts of the stratum corneum exceeds its effect on the protein part, and it seems that this compound is disturbed in the group. Also, it seems that the presence of horny lipids enhances the drug's permeability through the skin.

The results related to sodium lauryl ethyl sulfate's FT-IR skin adsorption (before contact) indicate an increase in the absorption wavelength within both the asymmetric and symmetrical CH region. Such a finding signifies that the bilayers of the stratum corneum undergo a blue shift in liquefaction. This finding also suggests an increase in the extent to which dam properties are influenced. The drug passes through. The adsorption of the drug as it permeates the stratum corneum causes the C=O stress wavelength to increase, which, in turn, causes the bonds between the hydrogen molecules located between the lipid molecules to weaken. Furthermore, the presence of sodium lauryl ethyl sulfate mitigates the barrier's effects by causing amide I's and amide II's wavelength numbers to increase.

The results of FT-IR of pre-contact skin with tynolin demonstrate that C-H is asymmetric and C-H is more symmetrical towards the absorption band, indicating an aqueous shift. Tynolin indicates the orientation of lipid bilayer groups, impairs barrier properties and possibly increases drug passage through the stratum corneum. The results also showed that tynolin shifted the absorption band in the amide regions I and II to a lower wavelength. Such a state shows the orientation of protein groups in the bilayer of the stratum corneum.

The FT-IR results from the skin of pre-contact rats with nanoxinol show that this compound caused a very small increase in the wavelength of asymmetric CH peak absorption and an increase in the absorption wavelength in the CH symmetric region, indicating liquefaction of bilayers in the stratum corneum membrane. Moreover, there is a disruption in the skin's barrier properties to this drug. This attraction decreased the absorption wavelength in the amide region 1 and increased the absorption wavelength in the amide region 2, which indicates the increase and decrease of the skin barrier effects, respectively.

5.1. Conclusions

Transdermal drug delivery never occurs without some problems. This study represents an attempt to overcome challenges in drug delivery caused by the barrier formed by the cutaneous stratum corneum. The utilized method was intended to aid transcutaneous drug delivery. The use of chemical and herbal additives changed the skin structure, thus leading to improvements in drug delivery and absorption into the skin. The future use of the

adsorbents considered in this study could significantly improve the dermal delivery of caffeine.

Acknowledgments

This article was extracted from the pharm.D thesis (Sheykhi) and sponsored by Ahwaz Jundishapur University of Medical Sciences, Ahwaz, Iran.

Footnotes

Authors' Contribution: A. S. conceived and designed the evaluation and drafted the manuscript. S. M. S. participated in designing the evaluation, performed parts of the statistical analysis, and helped to draft the manuscript. H. K. re-analyzed statistical data and revised the manuscript. A. S., S. M. S., H. K., and A. SH. approved the final version to be published.

Conflict of Interests: One author is the editorial board member or a reviewer of this journal.

Ethical Approval: This study is approved under the ethical approval code of IR.AJUMS.ABHC.REC.1398.053.

Funding/Support: This study was extracted from Doctor of Pharmacy thesis of Adel Sheykhi and was supported by a grant from Ahwaz Jundishapur University of Medical Sciences (grant number: B-98070).

References

- Ma H, Yu M, Lei M, Tan F, Li N. A novel topical targeting system of caffeine microemulsion for inhibiting UVB-induced skin tumor: characterization, optimization, and evaluation. *AAPS PharmSciTech*. 2015;**16**(4):905-13. [PubMed ID: 25591953]. [PubMed Central ID: PMC4508291]. <https://doi.org/10.1208/s12249-014-0278-5>.
- Nawrot J, Jordan S, Eastwood J, Rotstein J, Hugenholtz A, Feeley M. Effects of caffeine on human health. *Food Addit Contam*. 2003;**20**(1):1-30. [PubMed ID: 12519715]. <https://doi.org/10.1080/0265203021000007840>.
- Trommer H, Neubert RH. Overcoming the stratum corneum: the modulation of skin penetration. A review. *Skin Pharmacol Physiol*. 2006;**19**(2):106-21. [PubMed ID: 16685150]. <https://doi.org/10.1159/000091978>.
- Abdul Mudalip SK, Abu Bakar MR, Jamal P, Adam F. Solubility and Dissolution Thermodynamic Data of Mefenamic Acid Crystals in Different Classes of Organic Solvents. *J Chem Eng Data*. 2013;**58**(12):3447-52. <https://doi.org/10.1021/jje400714f>.
- Florence AT, Attwood D. *Physicochemical principles of pharmacy: In manufacture, formulation and clinical use*. London, UK: Pharmaceutical Press; 2015.
- Salimi A, Moghimipour E, Kogani P, Mohammad Soleymani S. The effect of various penetration enhancers on the octyl methoxycinnamate permeability: mechanisms of action study. *Iran J Pharm Sci*. 2020;**16**(2):87-104.
- Hussain A, Khan GM, Jan SU, Shah S, Shah K, Akhlaq M, et al. Effect of olive oil on transdermal penetration of flurbiprofen from topical gel as enhancer. *Pak J Pharm Sci*. 2012;**25**(2):365-9. [PubMed ID: 22459463].
- Salimi A, Emam M, Mohammad Soleymani S. Increase adapalene delivery using chemical and herbal enhancers. *J Cosmet Dermatol*. 2021;**20**(9):3011-7. [PubMed ID: 33528091]. <https://doi.org/10.1111/jocd.13960>.

9. Constantinides P. Enhanced intestinal absorption of an RGD peptide from water-in-oil microemulsions of different composition and particle size. *J Control Release*. 1995;**34**(2):109-16. [https://doi.org/10.1016/0168-3659\(94\)00129-i](https://doi.org/10.1016/0168-3659(94)00129-i).
10. Saadatzadeh A, Salimi A, Zarooni M. Influence of Permeation Enhancers on the in Vitro Skin Permeation of Ketorolac Tromethamine through Excised Rat Skin: A Mechanistic Study. *Asian J Pharm Clin Res*. 2018;**11**(7):242. <https://doi.org/10.22159/ajpcr.2018.viii7.24162>.
11. Tabosa MAM, de Andrade ARB, Lira AAM, Sarmento VHV, de Santana DP, Leal LB. Microemulsion Formulations for the Transdermal Delivery of Lapachol. *AAPS PharmSciTech*. 2018;**19**(4):1837-46. [PubMed ID: 29637497]. <https://doi.org/10.1208/s12249-018-0995-2>.
12. Chaiyana W, Rades T, Okonogi S. Characterization and in vitro permeation study of microemulsions and liquid crystalline systems containing the anticholinesterase alkaloidal extract from *Tabernaemontana divaricata*. *Int J Pharm*. 2013;**452**(1-2):201-10. [PubMed ID: 23680734]. <https://doi.org/10.1016/j.ijpharm.2013.05.005>.