Published online: 2024 April 8.

Research Article

In Vivo Wound-Healing and Anti-Inflammatory Activities of Honey Produced by *Melipona beecheii* Bees

Ina Ramírez Miranda (b)¹, Juan Acevedo-Fernandez (b)², Elizabeth Negrete-Leon (b)², David Abram Betancur-Ancona (b)³, Yolanda Beatriz Moguel-Ordoñez (b)^{4,*}

¹ Posgrado Institucional en Ciencias Agropecuarias y Manejo de Recursos Naturales Tropicales, Universidad Autonoma de Yucatan, Merida, Yucatan, Mexico

² Facultad de Medicina, Universidad Autonoma del Estado de Morelos, Cuernavaca, Morelos, Mexico

³ Facultad de Ingenieria Quimica, Universidad Autonoma de Yucatan, Merida, Yucatan, Mexico

⁴ C.E. Mococha, Instituto Nacional de Investigaciones Forestales, Agricolas y Pecuarias, Mexico

Corresponding author: C.E. Mococha, Instituto Nacional de Investigaciones Forestales, Agricolas y Pecuarias, Mexico. Email: ybmo2009@gmail.com

Received 2023 December 4; Revised 2024 February 28; Accepted 2024 March 7.

Abstract

Background: Honey produced by the *Melipona beecheii* bees has been highly prized in Mexico since pre-Hispanic times for its therapeutic properties, particularly in wound healing.

Objectives: The aim of this study was to assess the in vivo wound-healing and anti-inflammatory effects of *M. beecheii* honey using an animal model.

Methods: Honey samples were collected from the state of Yucatan, Mexico. Male albino mice were utilized to test the woundhealing and anti-inflammatory properties of the honey, with pirfenidone (KitosCell®) and indomethacin (1 and 30 mg/mL) serving as positive controls, respectively.

Results: From day one, enhancements in wound healing, granulation tissue formation, and continuous contraction of the wound edges were observed. In terms of anti-inflammatory activity, the honey dose was capable of reducing edema comparably to the control group receiving 1 mg/mL of indomethacin. However, differences were noted when compared to the 30 mg/mL indomethacin control group in three of the groups.

Conclusions: It can be concluded that *M. beecheii* honey exhibits wound-healing and anti-inflammatory effects comparable to those of the positive controls used in this study. Therefore, it offers a viable alternative for the topical treatment of wounds.

Keywords: Honey, Stingless Bees, Therapeutic Properties, Animal Model

1. Background

Wound healing is a remarkable biological process essential for restoring skin integrity after an injury (1). Presently, wound care represents a significant health concern due to the challenges posed by chronic wounds in comorbid conditions, including prolonged hospitalization, loss of mobility, decreased quality of life, and increased risk of nosocomial infections, among other economic burdens (2).

Bee honey has been utilized for medicinal purposes across various cultures since ancient times. Evidence from archaeological findings and literature reveals that ancient Egyptians, Greeks, and Romans used honey for wound treatment (3). Despite its longstanding use, the advent of antibiotics led to a decline in honey's medicinal application (4). However, with the increasing prevalence of antibiotic-resistant bacteria, honey has reemerged as a viable alternative in clinical medicine for wound treatment (5).

The effectiveness of honey in treating wounds, including those resulting from skin ulcers of various etiologies, is well-documented (6). A review by Jull et al. indicated that, in over 470 cases where honey was applied as a therapeutic agent, only five cases did not achieve satisfactory healing (7).

Copyright © 2024, Miranda et al. This open-access article is available under the Creative Commons Attribution 4.0 (CC BY 4.0) International License (https://creativecommons.org/licenses/by/4.0/), which allows for unrestricted use, distribution, and reproduction in any medium, provided that the original work is properly cited.

Honey stands as a potential alternative to current wound care products due to its antibacterial and antiinflammatory properties and its ability to promote autolytic debridement, deodorization, tissue growth stimulation, pain control, and minimal scarring (8). These benefits are attributed to multiple factors, including moisture, pH level, sugar concentration, the generation of reactive oxygen species (ROS), and the presence of peptides, which collectively aid the four phases of wound healing: Hemostasis, inflammation, proliferation/epithelialization, and tissue remodeling (3).

Inflammation is the immune system's biological response to various stimuli, such as pathogens, damaged cells, and toxic compounds. It is clinically characterized at the tissue level by redness, heat, pain, swelling, and loss of function (9). Normally, the inflammatory response eliminates infectious organisms and facilitates tissue repair. However, prolonged inflammation can alter the cellular composition at the site of injury and potentially hinder the wound healing process (10). Inflammation is a dynamic response that occurs following tissue damage, involving the activation of resident cells, production of pro-inflammatory mediators and cytokines, and recruitment of blood cells to initiate phagocytosis for clearing debris and foreign cells (11).

Despite the availability of numerous antiinflammatory agents, many are not conducive to wound healing; non-steroidal anti-inflammatory drugs (NSAIDs) exhibit cytotoxicity towards tissues, while corticosteroids can impede epithelial tissue development (12).

The anti-inflammatory activity of honey is linked to phenolic compounds known to suppress the overproduction of inflammatory mediators (13). This effect, combined with honey's antioxidant activity and its capacity to reduce edema, shortens the inflammation phase during wound healing (14).

While all honey possesses wound-healing properties, honey from stingless bees, sourced from a variety of native, often endemic, plant species, stands out (14). The unique practice of these bees storing honey in cerumen pots (a blend of wax and plant resins) enables the infusion of phytochemicals into the honey. This attribute is believed to enhance the therapeutic value of honey from stingless bees over that produced by *Apis mellifera*(15).

Although numerous reports highlight the therapeutic effects of *A. mellifera* honey, there has been limited research on the in vivo anti-inflammatory and healing efficacy of *Melipona beecheii* honey, particularly that produced in the Yucatan Peninsula of Mexico.

In the Yucatan Peninsula, two species of melipona bees, *M. beecheii* and *Melipona yucatanica*, are found. The latter is rare, with minimal records of its use in meliponiculture (16) compared to *M. beecheii*. The cultivation and production of *M. beecheii* bee honey, known locally as "xunaan-kab," "colel-kab," or "poolkab," was a widespread practice among the Maya of the Yucatan Peninsula well before the arrival of the Spaniards, marking it as a significant cultural and economic endeavor in the region (17). Its honey has long been a component of traditional Maya medicine (18). Over the past 15 years, there has been a resurgence (19) in interest in this honey due to the increasing demand for natural products with functional properties.

2. Objectives

The aim of this study was to assess the in vivo woundhealing and anti-inflammatory properties of honey produced by *M. beecheii* bees collected in Yucatan, Mexico.

3. Methods

3.1. Sample Collection

Twelve honey samples were collected from a traditional meliponiculturist who produces honey from *M. beecheii* bees. These samples came from meliponaries located in various municipalities across Yucatan, Mexico, to ensure a broad representation. The meliponaries were situated in a low deciduous forest, home to native and endemic species of the Yucatan Peninsula, from which the bees gather nectar and pollen.

The collection period spanned from July to October 2020. The honey was transported to the laboratory in polyethylene containers, shielded from light in an isothermal box. It was then transferred into 50 mL Falcon tubes and stored at 4°C until analyzed.

3.2. Animal Model

The study of wound-healing and anti-inflammatory kinetics was conducted on CD1 mice (*Mus musculus*), male, albino variety, aged 2 - 3 months and weighing 30 - 40 g, sourced from the Biotherium of Universidad Autonoma de Morelos (UAEM). The experimental protocols received prior approval from the Institutional Animal Care Committee of the Faculty of Medicine at UAEM (approval number: 005/2016), adhering to the NOM-062-ZOO-1999 Technical Specifications for the Care and Use of Laboratory Animals and the European Community Council Directive 86/609/EEC guidelines for experimental animal care and use.

3.3. Evaluation of Wound-Healing Activity

The model employed was adapted from Haryanto et al. (20) with modifications. Sixty-five mice were evenly divided into thirteen groups of five each: Twelve for treatments (M1 - M12), with the twelve *M. beaches* honey samples and one control group treated with pirfenidone (KitosCell®). Mice were housed in individual cages with unlimited access to food and water. They were intraperitoneally anesthetized with sodium pentobarbital (0.05 mg/g, Merck CAS 57-33-0). A circular incision was made on their shaved dorsum using a 5 mm diameter biopsy punch, marking day 0. The wounds received 100 µL of the designated honey treatment initially, followed by 50 µL treatments on days 1, 3, 6, 8, 10, 13, and 15. The control group received a 5 mg application of KitosCell®. Wound photographs were captured with a digital camera attached to a stereo microscope (Stemi DV4, ZeissR), and the wound area was measured using Image J software.

3.4. Evaluation of Anti-Inflammatory Activity

The methodology employed was based on the model reported by Garcia-Argaez et al. (21). Nine *M. beecheii* honey samples were tested. Sixty male CD1 mice, weighing 25 - 30 g, were divided into nine treatment groups (M1-M9) of five mice each, one negative control group (NC) which received a 2.5 μ g solution of 2-O-teradecanoyl-13-acetate (TPA, Sigma) in ethanol (J.T. Baker®), and two positive control groups treated with a solution of indomethacin (Sigma-Aldrich®, St. Louis, MO, USA, CAS 53-861) in acetone at concentrations of 1

mg/mL (Control-I 1) and 30 mg/mL (Control-I 2). The mice were housed in individual cages with unlimited access to food and water. They were anesthetized intraperitoneally with sodium pentobarbital (0.05 mg/g), and 5 μ L of TPA solution was topically applied to both the inner and outer sides of the right ear, while the left ear received only 10 µL of ethanol. Ten minutes later, 10 µL of honey was applied in the same manner to the right ear. The control mice were treated solely with the corresponding solvent. Four hours after application, the mice were euthanized by cervical dislocation, and a segment of ear tissue (5 mm in diameter) was excised from both the treated and untreated ears. The difference in weight between the two ear segments served as a measure of the edematous response. The percentage of inhibition relative to the negative control was calculated using the formula:

Inhibition
$$\% = \frac{C_0 - C_t}{C_0} \times 100$$
 (1)

Where C_0 and C_t represent the inflammatory response in the control group (TPA alone) and the treatment groups (TPA plus honey samples), respectively.

3.5. Statistical Analysis

The Kolmogorov-Smirnov test verified the normality of the collected data. A one-way analysis of variance (ANOVA) assessed differences between experimental groups, with Dunnett's test used for multiple mean comparisons against control groups. Results were presented as mean \pm standard deviation. The R (version 4.0.2) statistical software (https://rstudio.com/) was utilized for data analysis.

4. Results

4.1. Wound-Healing Activity

In this study, the progress of wound healing was monitored through measurements of wound area reduction, with wound edge contraction indicating a decrease in the original wound area.

No antigenic reactions to any of the honey treatments were observed in the wounds. From the first day, granulation tissue formation and continuous contraction of the wound edges were noted across all



Figure 1. Macroscopic analysis of skin wounds on days 0, 1, 3, 6, 8, 10, 13, and 15, featuring treatments with the control (KitosCell®) and M. beecheii honey samples (M1 - M12)

groups (Figure 1). A yellowish residue, likely from unabsorbed honey, was visible on the wound edges of those treated with honey. By day 6, all groups showed complete coverage of the wounds with granulation tissue and epithelialization, although the control group displayed a greater amount of epithelial tissue at the wound edges. By day 15, wounds in all groups had fully epithelialized.

For quantitative analysis, Figure 2 illustrates the wound area reduction over time for different honey



Figure 2. The behavior of wound area over time with treatments including the control (KitosCell®) and M. beaches honey samples (M1 - M12)

treatments compared with the healing agent KitosCell®.

Notably, the M11 group's curve showed a wound reduction percentage comparable to the control group,

marking the highest rate of wound area reduction in the healing process among all groups. The timeframe for an 80% wound size reduction (Figure 3) showed that the M2 group was the only one with a statistically significant difference (P < 0.05) compared to the control



Figure 3. Time to achieve 80% wound reduction. Bars represent the mean \pm SD. Different letters above the standard deviation bars denote a significant difference (P < 0.05) according to Dunnett's test

group, with wounds closing 80% in an average of 6.3 ± 0.9 days (%RSD 14.28).

4.2. Anti-Inflammatory Activity

The anti-inflammatory activity assessment revealed no statistical difference (P > 0.05) between the honey treatments and control-I 1 (indomethacin at 1 mg/mL). However, significant differences were noted (P < 0.05) in groups M3, M4, and M5 compared to Control-I 2 (indomethacin at 30 mg/mL), showing lower inhibition percentages (16.6 \pm 7.9, RSD 47.59%; 9.6 \pm 6.0, RSD 62.5%; 11.13 \pm 5.1, RSD 45.82%) (Figure 4).

5. Discussion

The medicinal properties of honey have been recognized for centuries (22), and while bee honey is produced worldwide, its therapeutic efficacy varies significantly. This variation depends on factors such as the bee species (entomological origin), the geographical location, the sources of nectar (botanical origin), and

other aspects like the timing of honey extraction, processing, and storage conditions (6).

In this research involving *M. beecheii* honey, the samples demonstrated efficacy comparable to that of wounds treated with KitosCell®, except for the M2 group, which exhibited a slower wound reduction rate than that of the healing agent. Similar outcomes were reported by Nisbet et al. (23) after assessing *A. mellifera* honey on rats and rabbits, noting no significant differences between the honey-treated groups and the control regarding wound diameter, granulation tissue formation, and epithelialization.

The study noted an early-phase wound-healing effect of honey comparable to the control drug. Haryanto et al. (20) reported analogous findings using *Apis dorsata* honey in mouse wounds, with hydrocolloid gel as a control. Mekkaoui et al. (24) observed significant healing of chemically induced wounds in mice treated with Moroccan honey compared to control groups (Vaseline and Madecassol®).



Figure 4. Percentage of inflammation inhibition, with bars showing the mean ± SD. Different letters above the standard deviation bars signify a significant difference (P < 0.05) as determined by Dunnett's test

The groups treated with honey displayed healthy granulation tissue formation, which promoted wound contraction, facilitated wound closure, and reduced scarring. This aligns with the conclusions of Jull et al. (7), who found that honey accelerates healing in burns and infected surgical wounds compared to conventional treatments.

Despite honey's anticoagulant properties contributing to the initial stage of wound healing (25), it acts as a barrier that maintains a moist wound environment, preventing eschar formation, reducing dermal necrosis, and encouraging epithelial cell migration to the wound surface (3), as observed during the first week (days 1 and 3) across all honey-treated groups. Other contributing factors might include honey's high acidity, which enhances blood oxygenation (5), and its low water activity, promoting oxygen and nutrient transport from deeper tissues to the wound area (22).

On day 1, the inflammatory phase emerged, likely due to honey's content of phenolic compounds (26, 27), glycopeptides (28), and low molecular weight proteins (29). These components stimulate monocytes and macrophages, enhancing the release of cytokines, chemokines, and enzymes that break down the extracellular matrix. This results in the activation of proinflammatory pathways markers, such as interleukins (IL-1, IL-10, and IL6), cyclooxygenases (COX-2), tumor necrosis factor (TNF- α), nuclear factors (NF- κ B and I κ B κ), transforming growth factor beta (TGF- β), the primary regulator of fibrosis lipoxygenase (LOX), nitric oxide (NO), nitric oxide synthase (iNOS), and prostaglandins (26).

Although inflammation is a critical biological response in the wound-healing process (14), it must be managed to prevent the condition from evolving into a chronic or excessive state (28). Therefore, the antiinflammatory activity of the honey was evaluated. No significant difference was observed in the inflammation inhibition percentage among any of the treated groups compared to control 1 (1 mg/mL of indomethacin); however, Groups M3, M4, and M5 showed a lower inhibition percentage compared to control 2 (30 mg/mL of indomethacin). The highest inflammation inhibition was seen in group M2 with 33.8 \pm 5.3%, which is considerably lower than the 54 \pm 5% reported by Borsato et al. (30) for mice topically treated with *M. marginata* honey. Researchers such as Hussein et al. (13) and Mohammed et al. (31) have reported the in vivo antiinflammatory activity of different types of honey, but these were administered orally.

The anti-inflammatory activity noted in the honeys studied here could be attributed to various mechanisms identified in previous research. For instance, Biluca et al. (32) demonstrated that meliponid honey's anti-inflammatory activity in vitro stemmed from the inhibition of TNF- α , IL-6, MCP-1, IL-12p70, INF- γ , and IL-10 secretion. Similarly, Ranneh et al. (33) reported a reduction in CRP, TNF- α , IL-1 β , IL-6, IL-8, MCP-1, NF- κ B, and p38 MAPK in blood levels in rat tissues with LPS-induced inflammation treated with honey from the genus Trigona.

This successfully study demonstrated the proangiogenic activity of honey during the proliferative phase by noting the development of granulation tissues in all groups from day 1 to day 13. Granulation tissues consist of fibroblasts growing in areas rich in capillaries that provide oxygen, spurred by mediators such as TNF- α and IL-1 β , which stimulate the release of various growth factors (34). It has also been established that honey's enzymatic regulation of hydrogen peroxide, combined with its antibacterial properties, facilitates calcium (Ca^{2+}) entry into cells (35), promoting the expression of early growth genes essential for wound healing. This process contributes to the modulation of adjacent non-inflammatory cells, such as fibroblasts (28), ultimately enhancing the rate of wound closure (36).

In the remodeling stage, all groups treated with honey in this study showed reduced scabbing and the development of thin scars. This outcome is attributed to the presence of amino acids in honey, such as arginine and glutamic acid, which serve as precursors to proline, a primary amino acid that enhances collagen synthesis. Additionally, the presence of iron, copper, and ascorbic acid in honey aids in the maturation of collagen fibers (36).

Figure 5 illustrates the key components in honey thatcontribute to the four phases of the wound-healingprocess:Hemostasis,inflammation,proliferation/epithelialization, and tissue remodeling.

Honey may contain over 200 compounds; its antiinflammatory and antibacterial effects, which are crucial for wound healing, are mainly due to its antioxidant properties. These properties are significantly influenced by the phenolic compounds' concentration and composition (21, 37), which vary based on factors like the bee species, the botanical sources of nectar, and the timing of honey collection. Variations in phenolic compound content in *M. beecheii* honey from Yucatan have been documented by Ruiz-Ruiz et al. (27), with 632.2 mg/kg GAE, and by Miranda et al. (38), with 1 317 mg/kg GAE post-harvest and 780 mg/kg GAE during the harvest season.

Although the basic composition of phenolic compounds in different honey varieties is somewhat consistent, differences in the phenolic profiles of honey from stingless bees have been reported by researchers such as Tomás-Barberán et al. (39), Sousa et al. (40), and Ranneh et al. (41). For instance, Ramon-Sierra et al. (42) identified three phenolic acids—chlorogenic, caffeic, and ellagic—and four flavonoids—catechin, myricetin, quercetin, and apigenin—in *M. beecheii* honey from the Yucatan Peninsula, Mexico. Conversely, Alvarez-Suarez et al. reported the presence of caffeic and coumaric acids, as well as apigenin, quercetin, luteolin, isorhamnetin, and kaempferol in the same type of honey from Cuba (43).

Honey has been shown to possess strong antioxidant activity (44, 45) and may serve as a natural antioxidant remedy, contributing to the prevention of various health issues (46). Stingless bee honey, undergoing natural fermentation during maturation, is a richer source of antioxidants compared to Apis honey, attributed to higher concentrations of aglycones and a greater oxygen radical absorbance capacity (ORAC) (47). Antioxidant activity in A. mellifera honey is reported with DPPH values ranging from 3.17 - 8.79 IC₅₀ mg/mL in Brazilian honey (48), 13.26 - 48.95 IC₅₀ mg/mL in Portuguese honey (49), and 5.24 - 17.51 IC₅₀ mg/mL in Malaysian honey (50). These values are comparable to those found in honeys from native bees like Melipona subnitida, with 10.6 - 12.9 IC₅₀ mg/mL (51), but are lower than those of *M. beecheii* honey from Yucatan, Mexico, which has $173.0 - 361.8 \text{ IC}_{50} \text{ mg/mL}(38)$.

This consideration allows us to view each honey as a unique product, explaining the variance in bioactive effects observed in some honey samples analyzed during this research. These differences are due to the concentration of compounds and their degree of



Hemostasis

- Protective effect: formation of an antimicrobial barrier.
- Moisture: prevention of pressure ulcer formation.
- · Osmotic effect: extraction of lymph to the wound surface.
- Acidity and phenols; improvement of tissue oxygenation.

Inflammation

- · Phenols: antioxidant effect, elimination of free radicals.
- · H,O, and peptides: regulation of proteases.
- H₂O₂: neutrophil activation and fibroblast differentiation.

Proliferation

- pH: promotion of epithelialization and wound closure.
- · H2O2: promotion of fibroblast proliferation.
- Moisture: rapid development of granulation tissue.

Remodeling

. H,O,: stimulation of neutrophils for the elimination of damaged tissue.

· Phenols: regulation of proteases and macrophages for the reduction of scar tissue.

Figure 5. Key components in honey that support the four stages of the wound-healing process.

isomerization or hydrolysis, which may affect their absorption. Lin et al. (52) observed that aglyconic compounds like quercetin have lower skin absorption compared to its polymethoxylated form (quercetin 3, 5, 7, 3', 4'-pentamethyl ether) in mice.

It's notable that the floral origin and the abundance of flowers throughout the year influence the phenolic content in honey. In this study, the honey samples were collected during the post-harvest period, coinciding with the Yucatan Peninsula's rainy season, which features minimal flowering. This scarcity makes it challenging for bees to forage, leading beekeepers to sometimes feed them with A. mellifera honey or sugar syrup. This supplementary feeding can reduce the phenolic compound content and other elements in the honey (53), possibly contributing to the lack of significant differences in the honey samples.

The scientific community has extensively documented the composition and therapeutic effects of honey (7), including its anti-inflammatory and healing properties (7, 20, 23, 30, 32). The composition and medicinal value of honey vary among bee species, influenced by their foraging behavior and the way they collect nectar and pollen from diverse plants and locations (54). This study evaluated honeys from the native stingless species *M. beecheii*, adding to the body of knowledge on honeys produced by traditional meliponiculturists in Yucatan. Nonetheless, research is still developing to scientifically validate the ancestral use of this honey. It's crucial to continue characterizing these honeys, assessing their medicinal properties, and correlating them with their physicochemical composition, bioactive compounds, and floral origin.

5.1. Conclusions

The present study found that the wound-healing and anti-inflammatory effects of honey produced by M. beecheii bees in vivo were comparable to those of the positive controls (pirfenidone and indomethacin). This suggests that it could serve as an alternative for topical wound treatment, either directly or as part of a formulation, without inducing the side effects associated with some current medications. However, the varying compositions of different honeys, which influence their therapeutic effectiveness, pose a

challenge to standardization. Further research is needed to explore the mechanisms of action in the inflammatory and wound-healing processes prompted by *M. beecheii* honey, including studies on growth factors, pro-inflammatory cytokines, tissue staining, oxidative enzymes, and the relationship with bioactive compounds in the honey. Continued verification of its use as a topical agent is warranted, as, despite the higher bioactivity related to its antioxidant properties reported in stingless bee honey, significant clinical evaluations in wound healing are yet to be realized.

Footnotes

Authors' Contribution: Study concept and design: D.B. and Y.M.; acquisition of data: J.A. and E.N. (animal model design for in vivo evaluations) and I.R. (sample analysis and in vivo tests); analysis and interpretation of data: I.R., J.A., and D.B.; drafting of the manuscript: I.R. and Y.M.; critical revision of the manuscript for important intellectual content: D.B. and Y.M.; statistical analysis: I.R. and D.B; administrative, technical, and material support: Y.M and D.B.; Study supervision: J.A. and D.B.

Conflict of Interests: No conflicts of interest.

Data Availability: The dataset presented in the study is available on request from the corresponding author during submission or after publication.

Ethical Approval: The experimental protocols were approved by the Institutional Animal Care Committee of the Faculty of Medicine of the UAEM (approval number: 005/2016), complying with NOM-062-ZOO-1999-Technical Specifications for the Care and Use of Laboratory Animals, as well as the guidelines for the care and use of experimental animals of the European Community Council Directive 86/609/EEC.

Funding/Support: This study was supported by a doctoral scholarship from the Consejo Nacional de Ciencia y Tecnología (CONACyT).

References

- Jackson WM, Nesti LJ, Tuan RS. Concise review: clinical translation of wound healing therapies based on mesenchymal stem cells. *Stem Cells Transl Med*. 2012;1(1):44-50. [PubMed ID: 23197639]. [PubMed Central ID: PMC3727688]. https://doi.org/10.5966/sctm.2011-0024.
- 2. Monika P, Chandraprabha MN, Rangarajan A, Waiker PV, Chidambara Murthy KN. Challenges in Healing Wound: Role of Complementary

 and Alternative Medicine. Front Nutr. 2022;8:791899. [PubMed ID:

 35127787].
 [PubMed Central ID: PMC8811258].

 https://doi.org/10.3389/fnut.2021.791899.

- Yupanqui Mieles J, Vyas C, Aslan E, Humphreys G, Diver C, Bartolo P. Honey: An Advanced Antimicrobial and Wound Healing Biomaterial for Tissue Engineering Applications. *Pharmaceutics*. 2022;14(8). [PubMed ID: 36015289]. [PubMed Central ID: PMC9414000]. https://doi.org/10.3390/pharmaceutics14081663.
- Kwakman PH, Zaat SA. Antibacterial components of honey. *IUBMB* Life. 2012;64(1):48-55. [PubMed ID: 22095907]. https://doi.org/10.1002/iub.578.
- Minden-Birkenmaier BA, Bowlin GL. Honey-Based Templates in Wound Healing and Tissue Engineering. *Bioengineering (Basel)*. 2018;5(2). [PubMed ID: 29903998]. [PubMed Central ID: PMC6027142]. https://doi.org/10.3390/bioengineering5020046.
- Nweze JA, Olovo CV, Nweze EI, Obi OJ, Chidebelu P. Therapeutic Properties of Honey. *Honey Analysis - New Advances and Challenges*. IntechOpen; 2019. https://doi.org/10.5772/intechopen.86416.
- Jull AB, Cullum N, Dumville JC, Westby MJ, Deshpande S, Walker N. Honey as a topical treatment for wounds. *Cochrane Database Syst Rev.* 2015;2015(3). CD005083. [PubMed ID: 25742878]. [PubMed Central ID: PMC9719456]. https://doi.org/10.1002/14651858.CD005083.pub4.
- 8. Stephen-Haynes J, Callaghan R. Properties of honey: Its mode of action and clinical outcomes. *Wounds UK*. 2011;7(1):50-7.
- Chen L, Deng H, Cui H, Fang J, Zuo Z, Deng J, et al. Inflammatory responses and inflammation-associated diseases in organs. *Oncotarget*. 2018;9(6):7204-18. [PubMed ID: 29467962]. [PubMed Central ID: PMC5805548]. https://doi.org/10.18632/oncotarget.23208.
- Landen NX, Li D, Stahle M. Transition from inflammation to proliferation: a critical step during wound healing. *Cell Mol Life Sci.* 2016;73(20):3861-85. [PubMed ID: 27180275]. [PubMed Central ID: PMC5021733]. https://doi.org/10.1007/s00018-016-2268-0.
- Cicala C, Morello S. Signaling Pathways in Inflammation and Its Resolution: New Insights and Therapeutic Challenges. *Int J Mol Sci.* 2023;24(13). [PubMed ID: 37446232]. [PubMed Central ID: PMC10341743]. https://doi.org/10.3390/ijms241311055.
- Krischak GD, Augat P, Claes L, Kinzl L, Beck A. The effects of nonsteroidal anti-inflammatory drug application on incisional wound healing in rats. *J Wound Care*. 2007;16(2):76-8. [PubMed ID: 17319622]. https://doi.org/10.12968/jowc.2007.16.2.27001.
- Hussein SZ, Mohd Yusoff K, Makpol S, Mohd Yusof YA. Gelam Honey Inhibits the Production of Proinflammatory, Mediators NO, PGE(2), TNF-alpha, and IL-6 in Carrageenan-Induced Acute Paw Edema in Rats. *Evid Based Complement Alternat Med.* 2012;2012:109636.
 [PubMed ID: 22919407]. [PubMed Central ID: PMC3418690]. https://doi.org/10.1155/2012/109636.
- Abd Jalil MA, Kasmuri AR, Hadi H. Stingless Bee Honey, the Natural Wound Healer: A Review. *Skin Pharmacol Physiol*. 2017;**30**(2):66-75. [PubMed ID: 28291965]. https://doi.org/10.1159/000458416.
- Shamsudin S, Selamat J, Sanny M, A R, Jambari NN, Khatib A. A Comparative Characterization of Physicochemical and Antioxidants Properties of Processed Heterotrigona itama Honey from Different Origins and Classification by Chemometrics Analysis. *Molecules*. 2019;24(21). [PubMed ID: 31671885]. [PubMed Central ID: PMC6864699]. https://doi.org/10.3390/molecules24213898.

- González-Acereto JA. [La importancia de la meliponicultura en México, con énfasis en la Península de Yucatán. Bioagrociencias]. *Bioagrociencias*. 2012;5(1):34-42. Spa.
- Pat FLA, Hernández BP, Pat FJM, Guizar VF, Ramos RR. Primera edición, editor. [Cría y manejo tradicional de la abeja Melipona beecheii (ko'olel kaab) en comunidades aledañas a la Reserva de la Biosfera Los Petenes, Campeche, México]. 52. México: El Colegio de la Frontera Sur; 2018. 227 p. Spa.
- Vit P, Vargas O, Lopez T, Maza F. Meliponini biodiversity and medicinal uses of pot-honey from El Oro province in Ecuador. *Emir J Food Agric*. 2015;27(6). https://doi.org/10.9755/ejfa.2015.04.079.
- Bratman E. Saving the Other Bees: The Resurgence of Stingless Beekeeping in the Zona Maya. *Conserv Soc.* 2020;18(4). https://doi.org/10.4103/cs.cs_20_66.
- Haryanto H, Urai T, Mukai K, Gontijo Filho PP, Suriadi S, Sugama J, et al. Effectiveness of indonesian honey on the acceleration of cutaneous wound healing: an experimental study in mice. *Wounds*. 2012;24(4):110-9. [PubMed ID: 25876247].
- Garcia-Argaez AN, Ramirez Apan TO, Parra Delgado H, Velazquez G, Martinez-Vazquez M. Anti-inflammatory activity of coumarins from Decatropis bicolor on TPA ear mice model. *Planta Med.* 2000;66(3):279-81. [PubMed ID: 10821059]. https://doi.org/10.1055/s-2000-14894.
- Tashkandi H. Honey in wound healing: An updated review. Open Life Sci. 2021;16(1):1091-100. [PubMed ID: 34708153]. [PubMed Central ID: PMC8496555]. https://doi.org/10.1515/biol-2021-0084.
- Nisbet HO, Nisbet C, Yarim M, Guler A, Ozak A. Effects of three types of honey on cutaneous wound healing. *Wounds*. 2010;**22**(11):275-83. [PubMed ID: 25901517].
- 24. Mekkaoui M, Assaggaf H, Qasem A, El-Shemi A, Abdallah EM, Bouidida EH, et al. Ethnopharmacological Survey and Comparative Study of the Healing Activity of Moroccan Thyme Honey and Its Mixture with Selected Essential Oils on Two Types of Wounds on Albino Rabbits. *Foods.* 2021;11(1). [PubMed ID: 35010154]. [PubMed Central ID: PMC8750595]. https://doi.org/10.3390/foods11010028.
- Ahmed A, Khan RA, Azim MK, Saeed SA, Mesaik MA, Ahmed S, et al. Effect of natural honey on human platelets and blood coagulation proteins. *Pak J Pharm Sci.* 2011;24(3):389-97. [PubMed ID: 21715274].
- Mm A, Sm H. Honey and propolis for wound healing. Int J Complement Altern Med. 2023;16(1):12-7. https://doi.org/10.15406/ijcam.2023.16.00626.
- Ruiz-Ruiz JC, Matus-Basto AJ, Acereto-Escoffié P, Segura-Campos MR. Antioxidant and anti-inflammatory activities of phenolic compounds isolated from Melipona beecheii honey. *Food Agric Immunol.* 2017;28(6):1424-37. https://doi.org/10.1080/09540105.2017.1347148.
- Schencke C, Vásquez B, Sandoval C, del Sol M. El Rol de la Miel en los Procesos Morfofisiológicos de Reparación de Heridas. *Int J Morphol.* 2016;34(1):385-95. https://doi.org/10.4067/s0717-95022016000100056.
- Ruiz-Ruiz JC, Acereto-Escoffié P, Segura-Campos MR. In vitro antioxidant and anti-inflammatory activities of Melipona beecheii honey protein fractions. J Food Meas Charact. 2018;12(4):2636-42. https://doi.org/10.1007/s11694-018-9881-6.
- 30. Borsato DM, Prudente AS, Doll-Boscardin PM, Borsato AV, Luz CF, Maia BH, et al. Topical anti-inflammatory activity of a monofloral honey of Mimosa scabrella provided by Melipona marginata during winter in

southern Brazil. *J Med Food*. 2014;**17**(7):817-25. [PubMed ID: 24650139]. https://doi.org/10.1089/jmf.2013.0024.

- Mohammed SAA, Talal Ali Hussein A, Abduljalil DSG, Marwan MAR. Anti-inflammatory activities of different types of Yemeni honey on rats. N Am Acad Res. 2020;3(11):170-89. https://doi.org/10.5281/zenodo.4279955.
- Biluca FC, da Silva B, Caon T, Mohr ETB, Vieira GN, Gonzaga LV, et al. Investigation of phenolic compounds, antioxidant and antiinflammatory activities in stingless bee honey (Meliponinae). *Food Res Int.* 2020;**129**:108756. [PubMed ID: 32036884]. https://doi.org/10.1016/j.foodres.2019.108756.
- Ranneh Y, Akim AM, Hamid HA, Khazaai H, Fadel A, Mahmoud AM. Stingless bee honey protects against lipopolysaccharide inducedchronic subclinical systemic inflammation and oxidative stress by modulating Nrf2, NF-kappaB and p38 MAPK. *Nutr Metab (Lond)*. 2019;**16**:15. [PubMed ID: 30858869]. [PubMed Central ID: PMC6391794]. https://doi.org/10.1186/s12986-019-0341-z.
- Rossiter K, Cooper AJ, Voegeli D, Lwaleed BA. Honey promotes angiogeneic activity in the rat aortic ring assay. J Wound Care. 2010;19(10):440. 442-6. [PubMed ID: 20948492]. https://doi.org/10.12968/jowc.2010.19.10.79091.
- Martinotti S, Laforenza U, Patrone M, Moccia F, Ranzato E. Honey-Mediated Wound Healing: H(2)O(2) Entry through AQP3 Determines Extracellular Ca(2+) Influx. *Int J Mol Sci.* 2019;20(3). [PubMed ID: 30754672]. [PubMed Central ID: PMC6387258]. https://doi.org/10.3390/ijms20030764.
- Sarkar S, Chaudhary A, Saha TK, Das AK, Chatterjee J. Modulation of collagen population under honey assisted wound healing in diabetic rat model. *Wound Medicine*. 2018;20:7-17. https://doi.org/10.1016/j.wndm.2017.12.001.
- Martinello M, Mutinelli F. Antioxidant Activity in Bee Products: A Review. Antioxidants (Basel). 2021;10(1). [PubMed ID: 33430511]. [PubMed Central ID: PMC7827872]. https://doi.org/10.3390/antiox10010071.
- Miranda IR, Ancona DB, Ordóñez YM. Influence of phenolic compounds and flavonoids on the colour and antioxidant activity of Melipona beecheii honey from deciduous forest of Yucatan, Mexico. *Emir J Food Agric.* 2023;12(35):1-10. https://doi.org/10.9755/ejfa.2023.3198.
- Tomás-Barberán FA, Truchado P, Ferreres F. Flavonoids in Stingless-Bee and Honey-Bee Honeys. *Pot-Honey*. New York, NY: Springer; 2013. p. 461-74. https://doi.org/10.1007/978-1-4614-4960-7_33.
- 40. Sousa JM, de Souza EL, Marques G, Meireles B, de Magalhães Cordeiro ÂT, Gullón B, et al. Polyphenolic profile and antioxidant and antibacterial activities of monofloral honeys produced by Meliponini in the Brazilian semiarid region. *Food Res Int.* 2016;**84**:61-8. https://doi.org/10.1016/j.foodres.2016.03.012.
- Ranneh Y, Ali F, Zarei M, Akim AM, Hamid HA, Khazaai H. Malaysian stingless bee and Tualang honeys: A comparative characterization of total antioxidant capacity and phenolic profile using liquid chromatography-mass spectrometry. *Lwt.* 2018;89:1-9. https://doi.org/10.1016/j.lwt.2017.10.020.
- 42. Ramon-Sierra J, Martinez-Guevara JL, Pool-Yam L, Magana-Ortiz D, Yam-Puc A, Ortiz-Vazquez E. Effects of phenolic and protein extracts from Melipona beecheii honey on pathogenic strains of Escherichia coli and Staphylococcus aureus. *Food Sci Biotechnol.* 2020;**29**(7):1013-

21. [PubMed ID: 32582463]. [PubMed Central ID: PMC7297938]. https://doi.org/10.1007/s10068-020-00744-4.

- Alvarez-Suarez JM, Giampieri F, Brenciani A, Mazzoni L, Gasparrini M, González-Paramás AM, et al. Apis mellifera vs Melipona beecheii Cuban polifloral honeys: A comparison based on their physicochemical parameters, chemical composition and biological properties. *Lwt.* 2018;87:272-9. https://doi.org/10.1016/j.lwt.2017.08.079.
- 44. Poulsen-Silva E, Gordillo-Fuenzalida F, Velasquez P, Llancalahuen FM, Carvajal R, Cabana-Brunod M, et al. Antimicrobial, Antioxidant, and Anti-Inflammatory Properties of Monofloral Honeys from Chile. *Antioxidants (Basel).* 2023;12(9). [PubMed ID: 37760088]. [PubMed Central ID: PMC10526086]. https://doi.org/10.3390/antiox12091785.
- Ahmed S, Othman NH. Review of the medicinal effects of tualang honey and a comparison with manuka honey. *Malays J Med Sci.* 2013;20(3):6-13. [PubMed ID: 23966819]. [PubMed Central ID: PMC3743976].
- Ahmed S, Sulaiman SA, Baig AA, Ibrahim M, Liaqat S, Fatima S, et al. Honey as a Potential Natural Antioxidant Medicine: An Insight into Its Molecular Mechanisms of Action. Oxid Med Cell Longev. 2018;2018:8367846. [PubMed ID: 29492183]. [PubMed Central ID: PMC5822819]. https://doi.org/10.1155/2018/8367846.
- Santos ACD, Biluca FC, Braghini F, Gonzaga LV, Costa ACO, Fett R. Phenolic composition and biological activities of stingless bee honey: An overview based on its aglycone and glycoside compounds. *Food Res Int.* 2021;147:110553. [PubMed ID: 34399530]. https://doi.org/10.1016/j.foodres.2021.110553.
- Pontis JA, Costa LAMAD, Silva SJRD, Flach A. Color, phenolic and flavonoid content, and antioxidant activity of honey from Roraima, Brazil. *Food Sci Technol.* 2014;**34**(1):69-73. https://doi.org/10.1590/s0101-20612014005000015.

- Ferreira ICFR, Aires E, Barreira JCM, Estevinho LM. Antioxidant activity of Portuguese honey samples: Different contributions of the entire honey and phenolic extract. *Food Chem.* 2009;**114**(4):1438-43. https://doi.org/10.1016/j.foodchem.2008.11.028.
- Khalil MI, Alam N, Moniruzzaman M, Sulaiman SA, Gan SH. Phenolic acid composition and antioxidant properties of Malaysian honeys. J Food Sci. 2011;76(6):C921-8. [PubMed ID: 22417491]. https://doi.org/10.1111/j.1750-3841.2011.02282.x.
- 51. Silva TMS, dos Santos FP, Evangelista-Rodrigues A, da Silva EMS, da Silva GS, de Novais JS, et al. Phenolic compounds, melissopalynological, physicochemical analysis and antioxidant activity of jandaíra (Melipona subnitida) honey. J Food Compos Anal. 2013;29(1):10-8. https://doi.org/10.1016/j.jfca.2012.08.010.
- 52. Lin CF, Leu YL, Al-Suwayeh SA, Ku MC, Hwang TL, Fang JY. Antiinflammatory activity and percutaneous absorption of quercetin and its polymethoxylated compound and glycosides: the relationships to chemical structures. *Eur J Pharm Sci.* 2012;**47**(5):857-64. [PubMed ID: 22609526]. https://doi.org/10.1016/j.ejps.2012.04.024.
- Becerril-Sanchez AL, Quintero-Salazar B, Dublan-Garcia O, Escalona-Buendia HB. Phenolic Compounds in Honey and Their Relationship with Antioxidant Activity, Botanical Origin, and Color. *Antioxidants* (*Basel*). 2021;10(11). [PubMed ID: 34829570]. [PubMed Central ID: PMC8614671]. https://doi.org/10.3390/antiox10111700.
- May-Canche I, Moguel-Ordonez Y, Valle-Mora J, Gonzalez-Cadenas R, Toledo-Nunez B, Arroyo-Rodriguez L, et al. Sensory and physicochemical analysis of honeys of nine stingless bee species of Mexico and Guatemala. *J Food Sci Technol.* 2022;**59**(12):4772-81. [PubMed ID: 36276552]. [PubMed Central ID: PMC9579246]. https://doi.org/10.1007/s13197-022-05561-7.