Published online: 2024 April 29.

**Research Article** 



# Enhancing Mycosporine-Like Amino Acids (MAAs) Extraction Yield from *Fischerella* sp. F5 using Experimental Design

Melika Saadatbakht 🔟 <sup>1</sup>, Tahereh Hosseinabadi 🔟 <sup>1,\*\*</sup>, Maryam Tabarzad 🔟 <sup>2,\*</sup>

<sup>1</sup> Department of Pharmacognosy, School of Pharmacy, Shahid Beheshti University of Medical Sciences, Tehran, Iran <sup>2</sup> Protein Technology Research Center, Shahid Beheshti University of Medical Sciences, Tehran, Iran

\* Corresponding Author: Protein Technology Research Center, Shahid Beheshti University of Medical Sciences, Tehran, Iran. Email: m\_tabarzad@sbmu.ac.ir \*\* Corresponding Author: Department of Pharmacognosy, School of Pharmacy, Shahid Beheshti University of Medical Sciences, Tehran, Iran. Email: t.hosseinabadi@sbmu.ac.ir

Received 2024 January 28; Revised 2024 March 17; Accepted 2024 March 28.

#### Abstract

**Background:** Mycosporine-like amino acids (MAAs) are ultraviolet-absorbing metabolites found in cyanobacteria, fungi, algae, and animals, known for their antioxidant and anti-inflammatory properties.

Objectives: This study aimed to optimize MAAs extraction from Fischerella sp. F5 using experimental design techniques.

**Methods:** The central composite design was employed to determine the optimal levels of five factors: Total extraction time, temperature, sonication time, methanol ratio, and solvent volume. Each run involved extracting 100 mg of dried biomass under the specified conditions. Mycosporine-like amino acids content was quantified using high performance liquid chromatography (HPLC), and the antioxidant activity of the extract was assessed using the DPPH radical scavenging test. The best model was

selected based on the highest R<sup>2</sup> value, and the optimized extraction conditions were determined.

**Results:** Mycosporine-like amino acids content in the final extracts ranged from 27.7% to 89%, with antioxidant activities ranging from 15% to 59.3%. Overall, all factors showed a positive correlation with MAAs content in the final extract. However, no significant model was established to correlate extraction conditions with MAAs purity ratio (MAA%) or extract antioxidant activity.

**Conclusions:** The bioactivity of partially purified MAAs may be influenced by factors such as associated impurities extracted alongside MAAs, which were not investigated in this study. Further research is necessary to identify the exact factors affecting MAAs extraction yield and purity, aiming to obtain partially purified MAAs extracts with optimal bioactivity.

Keywords: Antioxidant Activity, Fischerella sp., Mycosporine-Like Amino Acids, Central Composite Design

#### 1. Background

In recent years, much focus has been placed on developing efficient sunscreen products to prevent irradiation-induced skin damage. There are two types of sunscreens: Physical and chemical. Physical sunscreens, like zinc oxide, can cover a wide UV spectrum; however, they require frequent reapplication due to limitations. Chemical sunscreens work by absorbing light and releasing heat from the skin surface but may have side effects due to their chemical origin. Recently, manufacturers have shifted their focus from chemical agents to natural resources (1). Among the extensively researched natural sunscreens are mycosporine-like amino acids (MAAs) molecules. Mycosporine-like amino acids and gadusols are molecules produced by organisms to prevent sunlight damage, with microalgae and cyanobacteria being the main sources. Numerous researchers have evaluated these molecules for efficient sunscreens with antioxidant, repairing, and anti-aging properties (2).

Mycosporine-like amino acids are low molecular weight, water-soluble, nitrogen-rich molecules with maximum UV absorption. They make good sunscreens due to their high optical and thermal stability, strong UV radiation absorption, energy dispersion as heat, and excitation state in short wavelengths, preventing adverse photochemical reactions, including

Copyright © 2024, Saadatbakht et al. This open-access article is available under the Creative Commons Attribution 4.0 (CC BY 4.0) International License (https://creativecommons.org/licenses/by/4.0/), which allows for unrestricted use, distribution, and reproduction in any medium, provided that the original work is properly cited.

photoproduct formation. Mycosporine-like amino acids have been identified in cyanobacteria, microalgae, macroalgae (mainly in *Rhodophyta*), and marine animals. Their UV radiation absorption, antioxidant capacity, and physical and chemical properties enable their use in preventing and treating human diseases related to free radicals and UV radiation (3).

Various studies have reported the antioxidant effect of MAAs. For instance, a study in South Korea reported the antioxidant effect of MAAs extracted from *Chlamydomonas hedleyi* (4). In a review article, Kageyama and Waditee-Sirisattha documented the antioxidant effect of different MAAs investigated to date (5).

Culture conditions can significantly impact the yield of MAAs production in microalgae and cyanobacteria. Sharma investigated the effect of carbon content, salt, and pH on the production of UV-absorbing compounds by Spirulina. They found that using 0.4 M sodium chloride salt at pH 7 and the absence of carbon sources could lead to the production of compounds such as Phycocyanin and Allophycocyanin, while also increasing Phycoerythrin in *Spirulina plantesis* (6). Additionally, we reported that high salinity in Desmodesmus sp. culture induced the production of MAAs-related compounds (7). Furthermore, optimizing nitrogen and phosphate concentrations in the culture medium of Chlorella *vulgaris* could affect the type and yield of MAAs production (8). In our previous research, we identified two types of Fischerella sp. as sources for MAAs production. One of them produced at least two different types of MAAs, while another produced one main compound with characteristics similar to shinorine (9).

Moreover, factors influencing the extraction process, such as time, temperature, and type of solvent, can affect the extraction and purification yield. Studies by Karsten et al. demonstrated that pure distilled water was the most efficient solvent for extracting MMAs from *Peperomia crispa* and *Porphyra umbilicalis* (10). Similarly, Nishida's studies showed that pure distilled water yielded the best results for extracting MAAs from a red alga *Palmaria palmata* during a 6-hour extraction process (11). Additionally, Chaves-Pena et al. suggested that MAAs extraction with pure methanol from *Rhodophyta* could result in the detection of six types of MMAs (12).

#### 2. Objectives

The primary aim of this study was to optimize the extraction process of MAAs from Fischerella sp. F5, a strain previously investigated in our laboratory, and to evaluate its antioxidant effect. Various factors have been identified to influence the extraction yield of MAAs from cyanobacteria and other natural sources, including solvent polarity and volume, mechanical enhancers for cell disruption such as ultrasonic waves, thermal treatment, and extraction duration. Hence, we considered five quantitative factors-solvent type and volume, sonication time, extraction temperature, and duration-for the optimization of MAAs extraction from Fischerella sp. F5 using experimental design. Utilizing experimental design enables the simultaneous assessment of various factors on process outcomes, thereby reducing the number of experiments required to optimize processes. Therefore, the current study was designed employing experimental design methods.

#### 3. Methods

#### 3.1. Fischerella sp. Cultivation

Forty milliliters of *Fischerella* sp. F5 (accession No. OR228690) 25-day culture was cultivated in 200 mL of BG-11 culture medium (consisting of NaNO<sub>3</sub>, K<sub>2</sub>HPO<sub>4</sub>•H<sub>2</sub>O, MgSO<sub>4</sub>•7H<sub>2</sub>O, CaCl<sub>2</sub>•2H<sub>2</sub>O, citric acid, ferric ammonium citrate, EDTA, Na<sub>2</sub>CO<sub>3</sub>, H<sub>3</sub>BO<sub>3</sub>, MnCl<sub>2</sub>•4H<sub>2</sub>O, ZnSO<sub>4</sub>•7H<sub>2</sub>O, Na<sub>2</sub>MnO<sub>4</sub>•2H<sub>2</sub>O, CuSO<sub>4</sub>•5H<sub>2</sub>O, and Co (NO<sub>3</sub>)<sub>2</sub>•6H<sub>2</sub>O from Merck Chemicals, Germany). The cultures were harvested after 28 - 30 days at ambient temperature.

## *3.2. Mycosporine-Like Amino Acids Extraction and Partial Purification*

The biomass was separated by centrifugation (2-16KL, Sigma, Germany) at 9200 RPM for 10 minutes, followed by lyophilization. Subsequently, 10 mg of biomass was extracted for each run at varying concentrations of methanol (HPLC grade, Merck Chemical, Germany). The solvent type and volume, sonication time, extraction temperature, and duration were designed using DesignExpert® software. All sample mixtures were sonicated in an ultrasound bath (S 60 H, Elma, Germany). Extraction was carried out for 24, 48, and 72 hours according to the experimental design. The supernatant was separated by centrifugation at 12 500 RPM for 20 minutes, after which the samples were dried at room temperature. To dissolve the dried samples, 1 mL of deionized water (Millipore, Germany) was added, and each sample solution was transferred to a new microtube separately. For each sample, 0.5 mL of chloroform was added, followed by vigorous shaking and vortexing to remove non-polar content, including pigments and lipids. Samples were then passed through a 0.22 µm syringe filter (MS® MES Syringe Filter) before characterization high-performance by liquid chromatography (HPLC).

#### 3.3. Characterization of Mycosporine-Like Amino Acids

High Performance Liquid Chromatography was performed using a Shimadzu HPLC system equipped with a C18 HPLC column (5  $\mu$ m, 25 × 4.6 mm) and an isocratic mobile phase consisting of 12.5% methanol with 0.02% trifluoroacetic acid (TFA) aqueous solution plus 10% acetonitrile, with a flow rate of one mL.min<sup>-1</sup>. Using a Hamilton syringe, 100  $\mu$ L of each sample was injected into the injection loop. The absorbance spectrum was scanned in the wavelength range of 200 to 800 nm, and the chromatograms of samples were recorded at 254 and 330 nm. The best extract was further analyzed by direct mass spectrometry (MS) using an Agilent triple QQQ (USA) instrument.

### *3.4. Experimental Design to Optimize Mycosporine-Like Amino Acids Extraction*

A 1/2 factorial plan was designed using DesignExpert<sup>®</sup> 12 software to optimize five factors: Solvent type and volume, sonication time, extraction temperature, and duration (all quantitative). The features of the Experimental Design model are shown in Table 1. Applying the central composite design model, 32 runs in two blocks were designed to optimize these five factors. The total area under the curve (AUC) and its relative percentage, extracted from samples' HPLC chromatograms, were used for the optimization of the MAAs extraction process. For optimization, the central composite design (CCD) was utilized. The software applied an ANOVA test to verify the significance level of the models, considering a P-value less than 0.05 as significant.

#### 3.5. Determination of the Antioxidant Effect

The free radical scavenging of 2, 2-diphenyl-1picrylhydrazyl (DPPH) was used to determine the antioxidant activity. Briefly, 100  $\mu$ L of samples, positive controls (ascorbic acid 0.1%), and negative controls (methanol) were added to the 100  $\mu$ L DPPH reagent. For each sample, a blank was considered, which had methanol instead of DPPH reagents. Then, the absorption of each sample was measured at 517 nm after a 30-minute incubation. Ascorbic acid at a concentration of 1 mg/mL and a DPPH methanolic solution at a concentration of 0.1 mM were used as positive control and DPPH reagents, respectively. The antioxidant activity was calculated based on the DPPH scavenging activity using Equation 1.

DPPH scavenging effect 
$$\% = \left[A_0 - \frac{A_1}{A_0}\right] \times 100$$
 (1)

Where A0 and A1 represent control and sample absorptions, respectively.

#### 4. Results

#### *4.1. Characterization of Fischerella sp. F5 Extracts by High-performance Liquid Chromatography*

The HPLC chromatograms of samples extracted from *Fischerella* sp. F5 were analyzed to discover peaks showing similar features to MAAs. Most peaks with maximum absorbance between 330 - 340 nm exhibited a retention time of about 3 minutes. The sum of the AUCs of these peaks and their relative percentage compared to the sum of all peaks' AUCs were reported as MAA content and MAAs%, respectively, in Table 2. Runs 18 and 32 exhibited the highest value for MAA content, and run 32 had the highest value for MAA%. Accordingly, it seems that increasing the temperature for a short time and using a low percentage of methanol resulted in increased MAAs extraction efficiency. However, at room temperature, the extraction process with 55% methanol could result in promising extraction efficiency.

Some samples contained more than one peak with maximum absorption between 300 to 400 nm, as recorded for run 11 (Figure 1). As seen in the HPLC chromatogram, a peak with a retention time (Rt) of 2.905 and another at 8.8 min were recorded. This run had the highest value for total extraction time,

able 1. Characteristics of the Experimental Design Model					
Factor	Name	Minimum	Maximum		
А	Time (h)	24.00	72.00		
В	Temperature (°C)	4.00	50.00		
С	Sonication time (min)	30.00	120.00		
D	Methanol ratio (%)	10.00	100.00		
E	Solvent volume (mL)	1.00	3.00		

Table 2.	Data of tl	he Experiment	al Design an	d Extraction	Process										
Runs	Time	Temperature	Sonication Time	Methanol %	Solvent/Biomass	MAA Content	MAA%	Runs	Time	Temperature	Sonication Time	Methanol %	Solvent/Biomass	MAA Content	MAA%
1	24	4	120	100	3	141 733	54.998	17	24	4	30	10	3	240 635	67.058
2	24	4	120	10	1	160 578	44.867	18	24	50	120	10	3	4 612 592	67.887
3	48	27	75	55	2	226 182	59.142	19	24	50	30	10	1	302 597	54.977
4	24	50	120	100	1	231 576	62.844	20	48	27	75	55	2	226 182	59.142
5	72	4	120	10	3	190 063	52.898	21	48	27	120	55	2	306 473	41.585
6	24	4	30	100	1	151 182	63.272	22	48	27	75	55	2	226 182	59.142
7	24	50	30	100	3	195 858	64.796	23	48	27	75	55	2	226 182	59.142
8	72	4	30	100	3	230746	59.288	24	48	27	75	55	1	198 758	52.553
9	72	4	120	100	1	182 871	69.024	25	48	4	75	55	2	250 591	54.111
10	72	4	30	10	1	149 577	60.511	26	72	27	75	55	2	247 931	59.323
11	72	50	120	10	1	344 045	63.335	27	24	27	75	55	2	254 364	67.532
12	48	27	75	55	2	226 182	59.142	28	48	27	75	55	3	161 552	60.536
13	48	27	75	55	2	226 182	59.142	29	48	50	75	55	2	150 254	54.786
14	72	50	120	100	3	172 193	52.104	30	48	27	75	100	2	212 992	60.145
15	72	50	30	10	3	218 731	47.053	31	48	27	75	10	2	371 487	59.059
16	72	50	30	100	1	7 3745	27.73	32	48	27	30	55	2	119 1720	89.608

sonication time, and temperature at low concentrations of methanol, suggesting that increasing these three factors resulted in more efficient extraction of compounds with less polar characteristics. Previously, three peaks with retention times around 3.5 - 5 min had been observed in our study on this species of *Fischerella* sp.

For run 2, in which the extraction process was performed at high temperature and an extended time in the presence of 10% methanol, similar to run 11, two closely recorded peaks with maximum absorptions at 300 - 400 nm were observed, and the peak with Rt = 8.8 min was not recorded (Figure 2). This indicates that time and temperature also influenced the chemical type of extracted compounds.

It was observed that at higher temperatures with a decrease in sonication time (run 15), the peak of the less

polar compound with Rt = 8.8 min reappeared (Figure 3).

### *4.2. Modeling the Extraction Process and Optimized Model for Mycosporine-Like Amino Acids Extraction*

The data were analyzed using DesignExpert software, and a two-level five-factor method CCD was applied for optimization based on the interaction between different factors.

Moreover, the 2FI model was applied to extract the coefficient numbers of each factor influencing the MAA content of the extracted sample. The final formula was presented in the form of Equation 2, with the adjusted  $R^2$  equal to 0.77. This means that the model is roughly suitable for predicting the extraction yield by modulating these five factors.



Figure 1. High performance liquid chromatography chromatogram of the extract from run 11; A, chromatogram of the total extract; B, the UV spectrum of the two main peaks with maximum absorption at 300-400 nm (around 340 nm)



Figure 2. High performance liquid chromatography chromatogram of the extract from run 2. A, chromatogram of the total extract; B, the UV spectrum of the two main peaks with maximum absorption at 300 - 400 nm (around 340 nm)

$$\begin{split} MAA\ content &= -144955 + 29655.78\ (A) \\ &+ 8571.58\ (B) + 4294.25\ (C) + 11191.19\ (D) \\ &+ 383849\ (E) - 520.04\ (AB) - 233.79\ (AC) \\ &+ 251.92\ (AD) - 11154.66\ (AE) \\ &+ 281.78563\ (BC) - 288.05990\ (BD) \\ &+ 11109.90\ (BE) - 269931\ (CD) \\ &+ 249290.125\ (CE) \\ &- 262754.25(DE)\ (Adjusted\ R^2 = 0.7754) \end{split}$$

As seen in the proposed model for predicting MAA content from extraction factors, all single factors had a positive effect on the extraction yield; nevertheless, the interactions between these factors (combined factors) did not essentially have a positive effect on the extraction yield. Additionally, regarding the P-values reported for each component of the final model, the combined factors had non-significant values on the model designed for extracted MAA content, except for the interactions of temperature with time, sonication



Figure 3. High performance liquid chromatography chromatogram of the extract from run 15. A, chromatogram of the total extract; B, the UV spectrum of the two main peaks with maximum absorption at 300 - 400 nm (around 340 nm)

Table 3	. The Opt	imized Model for	for Mycosporine-Like /	Amino Acids Extractio	on				
Run	Time	Temperature	Sonication Time	Methanol Ratio	Solvent volume	MAA Content Predicted	%MAA Predicted	Desirability	
18	24	50	120.0	10.0	3.0	4301865.16	67.73	0.93	Selected





time, and methanol ratio. Finally, the optimized factors were predicted based on this model (Table 3).

The optimization for MAA% and antioxidant activity was also considered using CCD. The statistical data showed that none of these models could present a satisfactory adjusted R<sup>2</sup>, and the coefficients calculated for each factor did not have significant value (data not presented).

# *4.3. LC/MS Analysis of Extract Using the Optimized Method*

The optimized factors were considered for a new run of MAA extraction from 10 mg dried biomass of



Figure 5. The chemical structure and characteristics of mycosporine-glycine-valine (A), and porphyra; (B), drawn by ChemBioDraw Ultra

*Fischerella* sp. F5, and then, the extracted sample was analyzed by HPLC once more. The extracted sample of the optimized model was further analyzed by LC/MS. The main peak with  $\lambda$ max = 335 nm has m/z of 344 (Figure 4), which showed the most similarity with the features corresponding to mycosporine-glycine-valine (Figure 5A) or porphyra (Figure 5B). The peak of 887, which was seen in the MS spectrum, may be related to MAA derivatives, such as glycosylated MAAs.

### *4.4. Antioxidant Effect of Selected Mycosporine-Like Amino Acids Extracts*

The results of the antioxidant investigation of the selected extracts were reported in Table 4.

Accordingly, it seems that the samples which were extracted with 100% methanol have a higher antioxidant

effect. The highest value of antioxidant activity was calculated for runs 31 and 32, in which the highest amount of MAAs was extracted. However, it gives the impression that the antioxidant activity was not directly related to the amount of MAAs content, as seen in the sample of run 18, which showed 21% antioxidant activity compared to the 54.9% of run 32, despite both having the highest amount of MAAs content. Mycosporine-like amino acids extracts% in the extract of run 18 was about 67% compared to 89% for run 32; therefore, the antioxidant activity might be negatively influenced by impurities rather than MAAs, which were present in the extract. According to the experimental design conditions applied for MAA extractions, it might be concluded that a low concentration of methanol, either in low or high temperature of the extraction process,

Runs	Antioxidant (%)	Runs	Antioxidant (%)
1	$59.38 \pm 2.8$	17	$21.58\pm8.5$
2	$32.84\pm9.7$	18	$21.98\pm0.57$
3	$25.40\pm1.4$	19	$27.81 \pm 1.99$
4	$47.92 \pm 1.99$	20	$25.60 \pm 1.4$
5	$39.88 \pm 7.1$	21	$31.03 \pm 8.25$
6	$35.45 \pm 5.97$	22	$25.80 \pm 1.4$
7	$39.88 \pm 1.4$	23	$25.10\pm1.4$
3	$41.29\pm0.57$	24	$20.78\pm3.4$
)	$31.63 \pm 3.41$	25	$14.95\pm1.4$
0	$47.92 \pm 1.42$	26	$32.44 \pm 18.2$
1	46.71±1.99	27	$39.48\pm19.1$
2	$26.10\pm1.4$	28	$36.66 \pm 2.6$
3	$25.50\pm1.4$	29	34.85±15.3
4	$35.45 \pm 5.4$	30	$42.49\pm0.1$
5	$28.82\pm5.7$	31	$54.15 \pm 8.5$
6	$34.25 \pm 6.5$	32	$54.96 \pm 9.7$

resulted in similar antioxidant activities, as seen in samples of run 17 and 18, respectively. However, statistical analysis could not find any significant relationship between the amount or percentage of MAAs and antioxidant activity. It might be concluded that a change in extraction conditions might change the type of chemical compound in the final extract.

#### 5. Discussion

One of the important metabolites extracted from microalgae is MAAs. Mycosporine-like amino acids consist of approximately 20 relatively similar watersoluble compounds, and various methods for their extraction and identification have been reported (13). According to previously published work, *Fischerella* sp. F5 was found to produce UV absorbent compounds. The MAA compounds discovered had a retention time of about 3.5 to 5 minutes with UV absorption at 330 - 340 nm (9). Additionally, the effect of culture optimization on MAA production showed that increasing nitrate and phosphate ion concentrations could enhance the yield of MAAs production (14). We used the experimental design method to optimize five influential factors on extraction efficiency. To minimize the number of tests required, the central composite design was applied. This approach has previously been used for optimizing MAAs

production (8) and extracting R-phycoerythrin-enriched extracts from *Sarcopeltis skottsbergii* (15).

Metabolites extracted from cyanobacteria and microalgae have been widely studied and have shown promising applicability in the fields of agriculture, energy supply, pharmaceuticals, nutrition, and cosmetic-sanitary industries. Cyanobacterial metabolites exhibit diverse and valuable biological activities, including antimicrobial, anti-inflammatory, and UV-absorbing properties (16). The main compound detected in the optimized extraction product showed a maximum absorbance at around 335 nm, with a mass-tocharge ratio of 344. These features are highly similar to the characteristics of Mycosporine-glycine-valine. This MAA was previously reported from Euphausia superba under photosynthetically active radiation (PAR) with additional ultraviolet radiation (UVR) (17), Phaeocystis antarctica (18), and Artemia urmiana under PAR and low salinity conditions (19). Recently, an in silico study claimed that mycosporine-glycine-valine has the potential to effectively inhibit angiotensin-converting enzyme (ACE2), and consequently, it could be explored as a therapeutic agent for coronavirus disease 19 (COVID-19) (20). Moreover, a peak of higher molecular weight was also observed in the MS spectrum; it may be a glycosylated MAA. Glycosylated MAAs have been previously reported in *Nostoc commune* with m/z of

1051 and 721, which exhibited 27% antioxidant activity (21).

According to the results, it seems that this type of experimental design could not precisely lead to the best condition with the maximum amount of MAA (as an index for extraction) and MAA% (as an index for purification). However, this approach has previously been applied to optimize the extraction process of a biliprotein from a red macroalgae *Sarcopeltis skottsbergii* (15), as well as to obtain extracts with high antioxidant activity from a red seaweed *Gracilaria mammillaris* (22).

The effects of solid-liquid ratio, extraction time, and temperature on the yield of MAA extraction from four red macroalgae, including *Bangia fusco-purpurea, Gelidium amansii, Gracilaria confervoides,* and *Gracilaria* sp., were evaluated. It was found that increasing these factors could lead to an increase in MAA extraction yields, reaching a maximum level at specific values of these factors (23). Our study also revealed that elevating the extraction time and temperature and increasing the solvent/biomass ratio could enhance the yield of MAA extraction.

#### 5.1. Conclusions

According to our findings, the optimal conditions predicted for the maximum yield of MAA extraction from *Fischerella* sp. F5 indicate that using a 3-volume ratio (related to biomass) of 10% methanol, 120 minutes of sonication time, and maintaining the temperature of  $50^{\circ}$ C for 24 hours could result in the best MAA extraction. Verification of the best condition showed that this process could yield MAA content similar to the best run of experiments, with an MAA% of  $60 \pm 2.2$ %, falling within the medium range.

#### Footnotes

**Authors' Contribution:** M. Tabarzad and T. Hosseinabadi contributed to the study conception and design. Material preparation, data collection, and analysis were performed by M. Saadatbakht. The first draft of the manuscript was written by M. Saadatbakht and M. Tabarzad, and all authors commented on previous versions of the manuscript. All authors read and approved the final manuscript.

**Conflict of Interests Statement:** All authors declare that they have no conflicts of interest.

**Data Availability:** Additional data will be available upon request from the corresponding author.

EthicalApproval:Code:IR.SBMU.PHARMACY.REC.1401.241.

**Funding/Support:** This study was financially supported by the research deputy of the School of Pharmacy, Shahid Beheshti University of Medical Sciences, Tehran, Iran (Grant# 43004376).

#### References

- 1. Priyadarshani I, Rath B. Commercial and industrial applications of micro algae-A review. *J Algal Biomass Util*. 2012;**3**(4):89-100.
- Singh A, Cizkova M, Bisova K, Vitova M. Exploring Mycosporine-Like Amino Acids (MAAs) as Safe and Natural Protective Agents against UV-Induced Skin Damage. *Antioxidants (Basel)*. 2021;**10**(5). [PubMed ID: 33925517]. [PubMed Central ID: PMC8145676]. https://doi.org/10.3390/antiox10050683.
- Figueroa FL. Mycosporine-Like Amino Acids from Marine Resource. Mar Drugs. 2021;19(1). [PubMed ID: 33406728]. [PubMed Central ID: PMC7824388]. https://doi.org/10.3390/md19010018.
- Suh SS, Hwang J, Park M, Seo HH, Kim HS, Lee JH, et al. Antiinflammation activities of mycosporine-like amino acids (MAAs) in response to UV radiation suggest potential anti-skin aging activity. *Mar Drugs.* 2014;**12**(10):5174-87. [PubMed ID: 25317535]. [PubMed Central ID: PMC4210892]. https://doi.org/10.3390/md12105174.
- Kageyama H, Waditee-Sirisattha R. Antioxidative, Anti-Inflammatory, and Anti-Aging Properties of Mycosporine-Like Amino Acids: Molecular and Cellular Mechanisms in the Protection of Skin-Aging. *Mar Drugs.* 2019;**17**(4). [PubMed ID: 31013795]. [PubMed Central ID: PMC6521297]. https://doi.org/10.3390/md17040222.
- Sharma G. Effect of Carbon Content, Salinity and pH on Spirulina platensis for Phycocyanin, Allophycocyanin and Phycoerythrin Accumulation. J Microbial Biochemical Tec. 2014;6(4). https://doi.org/10.4172/1948-5948.1000144.
- Gharib R, Tabarzad M, Hosseinabadi T. Effect of High Salinity on Mycosporine-Like Amino Acid Production in Desmodesmus sp.: MAA production in Desmodesmus sp. *Trends Pept Protein Sci.* 2020;5:1-6 (e2).
- Hosseinabadi T, Gharib R, Salehian S, Tabarzad M. A study on the effect of nitrate and phosphate concentrations on the production of mycosporine-like amino acids by Chlorella vulgaris. *Iran J Biotechnol.* 2022;20(3). e3194.
- Tabarzad M, Baktash S, Atabaki V, Hosseinabadi T. Primary Assessment of Mycosporine-Like Amino Acids Production by Two Species of Fischerella sp. *Iran J Pharm Res.* 2021;20(4):405-14. [PubMed ID: 35194455]. [PubMed Central ID: PMC8842606]. https://doi.org/10.22037/ijpr.2021.115100.15194.
- 10. Karsten U, Escoubeyrou K, Charles F. The effect of re-dissolution solvents and HPLC columns on the analysis of mycosporine-like amino acids in the eulittoral macroalgae Prasiola crispa and

Porphyra umbilicalis. *Helg Mar Res.* 2009;**63**(3):231-8. https://doi.org/10.1007/s10152-009-0152-0.

- Nishida Y, Kumagai Y, Michiba S, Yasui H, Kishimura H. Efficient Extraction and Antioxidant Capacity of Mycosporine-Like Amino Acids from Red Alga Dulse Palmaria palmata in Japan. *Mar Drugs.* 2020;18(10). [PubMed ID: 33008002]. [PubMed Central ID: PMC7599624]. https://doi.org/10.3390/md18100502.
- Chaves-Pena P, de la Coba F, Figueroa FL, Korbee N. Quantitative and Qualitative HPLC Analysis of Mycosporine-Like Amino Acids Extracted in Distilled Water for Cosmetical Uses in Four Rhodophyta. *Mar Drugs.* 2019;**18**(1). [PubMed ID: 31905630]. [PubMed Central ID: PMC7024326]. https://doi.org/10.3390/md18010027.
- Carreto JI, Carignan MO. Mycosporine-like amino acids: relevant secondary metabolites. Chemical and ecological aspects. *Mar Drugs.* 2011;9(3):387-446. [PubMed ID: 21556168]. [PubMed Central ID: PMC3083659]. https://doi.org/10.3390/md9030387.
- Salehian S, Saadatbakht M, Tabarzad M, Hosseinabadi T. Culture Optimization to Produce High Yields of Mycosporine-Like Amino Acids by Fischerella sp. F5. *Mol Biotechnol.* 2023. [PubMed ID: 37597118]. https://doi.org/10.1007/s12033-023-00854-4.
- Castro-Varela P, Celis-Pla PS, Figueroa FL, Rubilar M. Highly Efficient Water-Based Extraction of Biliprotein R-Phycoerythrin From Marine the Red-Macroalga Sarcopeltis skottsbergii by Ultrasound and High-Pressure Homogenization Methods. *Front Mar Sci.* 2022;9. https://doi.org/10.3389/fmars.2022.877177.
- Vijayakumar S, Menakha M. Pharmaceutical applications of cyanobacteria—A review. J Acute Med. 2015;5(1):15-23. https://doi.org/10.1016/j.jacme.2015.02.004.
- 17. Newman SJ, Dunlap WC, Nicol S, Ritz D. Antarctic krill (Euphausia superba) acquire a UV-absorbing mycosporine-like amino acid from

dietary algae. *J Exp Mar Biol Ecol.* 2000;**255**(1):93-110. [PubMed ID: 11090854]. https://doi.org/10.1016/s0022-0981(00)00293-8.

- Moisan TA, Mitchell BG. UV absorption by mycosporine-like amino acids in Phaeocystis antarctica Karsten induced by photosynthetically available radiation. *Marine Biol.* 2001;138(1):217-27. https://doi.org/10.1007/s002270000424.
- Khosravi S, Khodabandeh S, Agh N, Bakhtiarian M. Effects of salinity and ultraviolet radiation on the bioaccumulation of mycosporinelike amino acids in Artemia from Lake Urmia (Iran). *Photochem Photobiol.* 2013;89(2):400-5. [PubMed ID: 22998644]. https://doi.org/10.1111/j.1751-1097.2012.01245.x.
- Sahu N, Mishra S, Kesheri M, Kanchan S, Sinha RP. Identification of Cyanobacteria-Based Natural Inhibitors Against SARS-CoV-2 Druggable Target ACE2 Using Molecular Docking Study, ADME and Toxicity Analysis. *Indian J Clin Biochem*. 2023;**38**(3):361-73. [PubMed ID: 35812791]. [PubMed Central ID: PMC9255548]. https://doi.org/10.1007/s12291-022-01056-6.
- Matsui K, Nazifi E, Kunita S, Wada N, Matsugo S, Sakamoto T. Novel glycosylated mycosporine-like amino acids with radical scavenging activity from the cyanobacterium Nostoc commune. *J Photochem Photobiol B.* 2011;105(1):81-9. [PubMed ID: 21813286]. https://doi.org/10.1016/j.jphotobiol.2011.07.003.
- 22. Ospina M, Castro-Vargas HI, Parada-Alfonso F. Antioxidant capacity of Colombian seaweeds: 1. Extracts obtained from Gracilaria mammillaris by means of supercritical fluid extraction. *The Journal of Supercritical Fluids*. 2017;**128**:314-22. https://doi.org/10.1016/j.supflu.2017.02.023.
- Sun Y, Han X, Hu Z, Cheng T, Tang Q, Wang H, et al. Extraction, Isolation and Characterization of Mycosporine-like Amino Acids from Four Species of Red Macroalgae. *Mar Drugs*. 2021;**19**(11). [PubMed ID: 34822486]. [PubMed Central ID: PMC8625514]. https://doi.org/10.3390/md19110615.