



The Kinetics of 3-Chymotrypsin-Like Cysteine Protease (3CLpro) Inhibition by *Otostegia persica* and *Otostegia aucheri* Extracts and Fractions: An In Vitro Model for Coronavirus Enzyme Inhibition

Soheila Moein ^{1,*}, Zahra Zebarjad ¹, Mahmoodreza Moein ^{1,2}, Zahra Sabahi ¹

¹ Medicinal Plants Processing Research Center, Shiraz University of Medical Sciences, Shiraz, Iran

² Department of Pharmacognosy, School of Pharmacy, Shiraz University of Medical Sciences, Shiraz, Iran

*Corresponding Author: Medicinal Plants Processing Research Center, Shiraz University of Medical Sciences, Shiraz, Iran. Email: smoein231@gmail.com

Received: 7 May, 2025; Revised: 7 July, 2025; Accepted: 7 July, 2025

Abstract

Background: As of 2025, COVID-19 has caused over 700 million cases globally (WHO, 2025). The COVID-19 epidemic has created a severe global health crisis due to the broad loss of human lives and economic impact. The 3-Chymotrypsin-like cysteine protease (3CLpro, which cleaves viral polyproteins to produce functional proteins essential for replication), performing critical functions in the duplication of severe acute respiratory syndrome (SARS)-CoV, is necessary for its life cycle. Certainly, 3CLpro (as a model for coronavirus protease) is a previously approved drug target for SARS-CoV and Middle East respiratory syndrome (MERS)-CoV. This emphasizes the significant role of 3-chymotrypsin-like cysteine protease in the construction of powerful medicine against COVID-19. Nowadays, finding a novel drug derived from a plant with better treatment is extremely desirable due to the lack of specific treatments for SARS-CoV, MERS-CoV, and SARS-CoV-2.

Objectives: The present study aimed to evaluate the efficacy of *Otostegia persica* and *O. aucheri* extracts and their fractions in inhibiting COVID-19 marker enzymes, which are called proteases.

Methods: Ethanol extracts of these plants were fractionated using petroleum ether (petr), chloroform (chlo), ethyl acetate (eta), and n-butanol (buta) solutions, respectively. For the measurement of shrimp-derived cysteine protease as a 3CLpro model retardation, 200 μ L of the enzyme (at 25°C), 400 μ L of a substrate, and plant extracts and fractions were used. After determining the percentage of inhibition and IC_{50} , K_m and V_{max} were also calculated.

Results: The results showed that the eta fraction of *O. aucheri* (with $IC_{50} = 0.002 \pm 0.0001$ mg/mL) had the greatest inhibitory effect on protease activity compared to iodoacetamide (with $IC_{50} = 0.01 \pm 0.01$ mg/mL, $P = 0.04$) as the standard. Also, the chlo fraction of *O. aucheri* (with an $IC_{50} = 0.09 \pm 0.007$ mg/mL) and the buta fraction of *O. aucheri* (with an $IC_{50} = 0.07 \pm 0.002$ mg/mL) possessed significant inhibitory effects in protease retardation. *Otostegia persica* fractions had no significant inhibition on protease activity ($P > 0.05$). The mode of protease inhibition of all samples was uncompetitive, except for the petr fraction of *O. aucheri*, which was mixed inhibition.

Conclusions: The eta, chlo, and buta fractions of *O. aucheri* had remarkable effects in the inhibition of protease. In vitro results may require in vivo validation.

Keywords: Ethanol Extracts, *Otostegia persica*, *Otostegia aucheri*, Protease Inhibition, SARS-CoV, 3CLpro, In Vitro

1. Background

In December 2019, pneumonia related to the 2019 novel severe acute respiratory syndrome (SARS)-CoV-2 (2019-nCoV) was reported in Wuhan, China (1). Human SARS-CoV is one of the major pathogens of respiratory

infection (2). The two most pathogenic viruses, SARS-CoV and Middle East respiratory syndrome (MERS)-CoV, induce severe respiratory syndrome in humans, and four human coronaviruses, including HCoV-OC43, HCoV-229E, HCoV-NL63, and HCoV-HKU1, cause mild upper respiratory disorders (1). The SARS-CoV depends on the

Copyright © 2025, Moein et al. This open-access article is available under the Creative Commons Attribution 4.0 (CC BY 4.0) International License (<https://creativecommons.org/licenses/by/4.0/>), which allows for unrestricted use, distribution, and reproduction in any medium, provided that the original work is properly cited.

How to Cite: Moein S, Zebarjad Z, Moein M, Sabahi Z. The Kinetics of 3-Chymotrypsin-Like Cysteine Protease (3CLpro) Inhibition by *Otostegia persica* and *Otostegia aucheri* Extracts and Fractions: An In Vitro Model for Coronavirus Enzyme Inhibition. Jundishapur J Nat Pharm Prod. 2025; 20 (3): e162409. <https://doi.org/10.5812/jjnpp-162409>.

host cell for replication and enters the host cell with protease activity (3). Therefore, natural compounds with antiprotease activity can neutralize the effect of SARS-CoV (4). The *Otostegia* genus possesses 31 species and is a member of the Lamiaceae family, which grows in the Asian and Mediterranean regions. Lamiaceae species are used traditionally (5) for the treatment of many chronic diseases (6); they possess a diterpenoid component (otostegindiol), and many usages of this compound are observed (7). Moreover, the species of the genus *Otostegia* are widely used by traditional Persian medicine practitioners for the treatment of various disorders, and its components have been shown to possess antispasmodic, antidepressant, antiulcer, anxiolytic, and sedative properties (7). One of the *Otostegia* species is *Otostegia persica*, which is traditionally consumed in some parts of Iran as an antidiabetic plant. *Otostegia persica* is used to treat diabetes in both modern and traditional medicine (8). Thus, its reported potential has been examined by some researchers. Also, diterpenoids such as otostegindiol are compounds that have been separated from the aerial parts of *O. persica*. Ayatollahi et al. (9, 10) demonstrated that a dichloromethane extract of aerial parts of *O. persica* could be a significant source of various terpenoids.

Triterpenes are commonly biosynthesized following the isoprene rule of Ruzicka. Triterpenes have been reported to have a broad range of therapeutic uses, including antiviral activities (11). Moreover, some of the significant constituents of *O. persica* are flavonoids and phenolic compounds, including morin, quercetin, kaempferol, and isovitexin (C-glucoflavone), as well as trans-cinnamic acid (12). Another species of *Otostegia* is *O. aucheri*, which exhibited anti-hyperglycemic properties (8). It is reported that *O. aucheri* possesses 6,8-di-C-glycosylapigenin and friedelin (8). Recent studies have also exhibited the antiviral activity of apigenin against several viruses (13). In addition, Khandelwal et al. have reported that apigenin possesses antiviral effects against the buffalopox virus (14). Another investigation reported that apigenin analogs bind to COVID-19's main protease substrates. The interactions between the analogs of apigenin and SARS-CoV-2 Mpro inhibitors were detected as electronic bonds, hydrogen bonds, and hydrophobic interactions. Several studies show that apigenin 7-glucoside-4'-p-coumarate is the best candidate for SARS-CoV-2 Mpro inhibition. Apigenin (4',5,7-trihydroxyflavone), a flavone glycoside, is found in a variety of vegetables and fruits (15). Antiprotease activities of some plants have been reported (16).

2. Objectives

This study evaluated the kinetic parameters (Km, Vmax) of shrimp protease as a 3CLpro model inhibition and determined the mode of inhibition of *O. persica* and *O. aucheri* extracts and different fractions.

3. Methods

3.1. Materials

One material, Potassium sodium tartrate (Cat. No. 217255), was obtained from Sigma Aldrich, USA. Trichloroacetic acid (10% w/v, Cat. No. 76-03-9) was purchased from Merck Chemicals (Darmstadt, Germany). All chemicals used, such as casein (1% w/v, Cat. No. C7078), tris (Cat. No. 77-86-1), and di-potassium hydrogen phosphate-3 hydrate (Cat. No. 16788-57-1), were of analytical grade ($\geq 99\%$ purity). Casein was dissolved in a pH 7.4 phosphate buffer.

3.2. Specimen Collection

Otostegia persica and *O. aucheri* were gathered from around Bandar Abbas, Hormozgan province, Iran, by Mr. Asadipour in March 2020. These plants were identified by the botanists at Shiraz University of Medical Sciences and were then extracted. The voucher number was saved in the Medicinal Plants Processing Research Center, School of Pharmacy, Shiraz University of Medical Sciences, Shiraz, Iran (MPRCM 94-86, and MPPRC-96-9 for *O. persica* and *O. aucheri*, respectively).

3.3. Extraction and Fractionation of the Studied Plants

First, the leaves of the two plants (3000 g per species) were air-dried at 25°C, chopped, and macerated in 80% v/v ethanol for 72 hours at room temperature. Then, the extracts were concentrated using a rotary evaporator (Eyela model N-1100, 40°C, 100 mbar). Subsequently, dried extracts were suspended in a 70:30 methanol:water solution and then filtered through Whatman No. 1 filter paper. After extraction, these extracts were subjected to fractionation by various solvents in order of increasing polarity; in other words, fractionation was performed with petroleum ether (petr), chloroform (chlo), ethyl acetate (eta), and n-butanol (buta), respectively (9). Ethanol extracts, petr fractions, chlo fractions, eta fractions, and buta fractions yielded 166.5, 80.7, 19.9, 15.1, and 1.1 g from 3000 g of powdered plant, respectively. The source of the enzyme was shrimp-*Penaeus vannamei* protease as a model for 3CLpro. Professor Ahmad Homaie (Hormozgan University, Bandar Abbas, Iran) donated the protease.

3.4. Determination of Protease Inhibition

Two hundred microliters of phosphate buffer (50 mM, pH = 7.4) was blended with 200 μ L of cysteine-like protease from shrimp. Then, 400 μ L of casein (1%, Cat. No. 9000-71-9), as a substrate, was added, and the mixture was incubated for 10 minutes at 25°C. Afterward, 800 μ L of trichloroacetic acid (TCA, 10%) was added as a stopper of the reaction. After 21 minutes, the mixture was centrifuged (Peco FD-T6-4000, Iran) at 12000 rpm for 5 minutes to ensure complete precipitation, and a UV-visible spectrophotometer (PG instrument T90, England) was used to determine the absorbance of the supernatant (17) at 280 nm \pm 2 nm.

Protease inhibitory activity was determined as follows: Inhibition (%) = $100 - [(OD \text{ sample} - OD \text{ blank}) / (OD \text{ control} - OD \text{ blank}) \times 100]$. The above formula was adapted from Kunitz, 1947 (18). A blank solution was prepared to omit the background absorbance and contained plant extract without the enzyme. In the control solution, the plant extract was replaced with ethanol. The control contained the enzyme without the plant extract. The absorbances of 0.1 to 550 micromol/mL of tyrosine were detected at 280 nm by spectrophotometer, and the standard curve was plotted (Figure 1).

3.5. Kinetics of Protease Inhibition

The inhibition modes of the protease were evaluated based on the method delineated by Rouzbehan et al. (9). For this evaluation, constant amounts of protease were blended with increasing concentrations of its substrates (casein concentrations from 0.1 to 2.0 mg/mL) at 25°C for 10 minutes in a pH 7.4 phosphate buffer, and afterward mixed with and without the IC₅₀ concentrations of the samples. Then, the reactions were stopped, and absorption was detected using the Michaelis-Menten curve. After that, the absorbance was converted into the activity of the enzyme using the standard curve of tyrosine (Figure 1); then, different concentrations of the sample were plotted against the concerned activity. All assessments were repeated three times. The Michaelis-Menten equation is: $V = V_{max} [S] / (K_m + [S])$.

3.6. Statistical Analysis

The results are shown as the mean \pm SD of the biological triplicates. One-way Analysis of Variance (ANOVA) with repeated measures and post-hoc Tukey ($\alpha = 0.05$) were employed to detect the variations among the means. A P-value of less than 0.05 was regarded as statistically significant. After that, kinetic parameters including V_{max} and K_m were determined using GraphPad Prism 7.0. For this determination, we

principally converted the amount of reading absorption into enzyme activity using the product (tyrosine) standard curve. Next, the values of K_m and V_{max} were determined employing GraphPad Prism 7.0. Statistical analysis was done with GraphPad InStat 3.0 software.

4. Results

4.1. The Yield of the Extraction

In both *Otostegia* species, 3000 g of dried leaves yielded 166 g of dried extract. The yields were 5.55% for ethanol extracts of *O. aucheri* and *O. persica*. Additionally, 3000 g of petr, chlo, eta, and n-buta fractions gave 80.7, 19.9, 15.14, and 1.1 g of lyophilized post-fractionation, respectively. The yields ranged from 1.1 g (n-buta) to 80.7 g (petr) across replicates. Yield percentages of petr, chlo, eta, and buta fractions were 2.69, 0.66, 0.50, and 0.06, respectively.

4.2. The Effects of the Extracts and Fractions of *Otostegia persica* and *Otostegia aucheri* in Inhibition of Protease

Samples tested ranged from 0.015 to 0.5 mg/mL. The results of the present research showed that at 0.125 mg/mL, a buta fraction of *O. persica* had 31.2 \pm 4.03% inhibition on protease activity, and at 0.25 mg/mL, a petr fraction of *O. persica* had 37.05 \pm 4.8% inhibition on protease activity. Other fractions of *O. persica* had no effects on the inhibition of protease (Figure 2A). Additionally, the IC₅₀ of *O. persica* extract was 0.46 \pm 0.005 mg/mL (Table 1). However, the eta fraction of *O. aucheri* had the most inhibitory effect on protease activity in comparison to iodoacetamide (IC₅₀ 0.002 \pm 0.0001 mg/mL, P = 0.04, Table 1). Besides, buta and chlo fractions of *O. aucheri* also had significant inhibitory effects on protease inhibition (Figure 2B). Figure 2C showed the percentage of protease inhibition of iodoacetamide as a standard. The saturation diagrams (Michaelis-Menten curve) of protease inhibition are shown in Figure 3A - G. All experiments were performed three times.

4.3. The Kinetics of Protease Inhibition of Samples

The IC₅₀ concentrations of the samples were 0.002 - 0.46 mg/mL based on Table 1. Table 1 shows the K_m and V_{max} of the samples against protease. Compared to the control without inhibitor, the V_{max} reduction was significant for all the samples except the petr fraction (P < 0.05). The K_m was decreased for all the samples except for the petr fraction of *O. aucheri*, in which K_m = 0.03 \pm 0.003, remained constant (Table 1, P < 0.05). The types of




Figure 1. The standard curve of tyrosine




Figure 2. A, percentage of protease inhibition of *Otostegia persica* extract (sample sizes, n = 3 per concentration); B, percentage of protease inhibition of *O. aucheri* extract and its fractions. Ot. Ethano, ethanol extract of *O. aucheri*; Ot. Petroll, petroleum ether (petr) fraction of *O. aucheri*; Ot. Chlor, chloroform (chlo) fraction of *O. aucheri*; Ot. Ethyl, ethyl acetate (eta) fraction of *O. aucheri*; Ot. Buthan, butanol (buta) fraction of *O. aucheri* (sample sizes, n = 3 per concentration); C, percentage of protease inhibition of iodoacetamide as a standard (sample sizes, n = 3 per concentration).

protease inhibition by all the samples and by iodoacetamide as a positive standard were uncompetitive (Table 1), except for the petr fraction of *O. aucheri*, which exhibited mixed inhibition (Table 1).

5. Discussion

Coronaviruses can cause infections in the respiratory tract, such as SARS and MERS, in different animals and humans (17, 19, 20). The consumption of natural compounds, including antioxidant agents to control the immune system, can be useful for afflicted patients. Throughout human history, many epidemic disorders have appeared. Ibn Sina's Canon of Medicine (21) and other physicians proposed solutions to these epidemics that were effective in controlling these diseases. Herbal

recommendations (use of plants, extracts, and compounds) for the treatment of COVID-19 symptoms are according to traditional Persian medicine (21). As a result, it is recommended that these patients use natural compounds such as flavonoids and tannins (22) as anti-inflammatory and antioxidant factors to regulate their immune systems. Therefore, the utilization of natural compounds, including flavonoids and tannins (22) as anti-inflammatory and antioxidant factors, to regulate the immune system, is suggested for these patients (23-26). The herbal components, especially terpenoids, possess protease inhibition activities, so these may be potentially effective due to low IC_{50} values for the management of viral epidemics. These components can be efficiently utilized as antiprotease drugs for SARS-CoV-2 in the future (27).

Table 1. Percentage of Maximum Inhibition of Protease, IC₅₀, Kinetic Parameters, and the Kind of Protease Inhibition by the Samples^a

Samples	Maximum of Protease Inhibition % at 0.015 mg/mL of the Sample	IC ₅₀ mg/mL	Vmax ± SD (μmol/min) ^b	Km ± SD (mmol) ^c	Kind of Inhibition
<i>Otostegia persica</i> extract	88.95 ± 1.3	0.46 ± 0.005	1.03 ± 0.07	0.0125 ± 0.003	Uncompetitive
<i>O. aucheri</i> extract	88.6 ± 1.9	ND	ND	ND	ND
Petr fraction of <i>O. aucheri</i>	79.3 ± 1.6	0.325 ± 0.007	1.2 ± 0.005	0.03 ± 0.003	Mixed ^d
Chlo fraction of <i>O. aucheri</i>	77.15 ± 1.4	0.09 ± 0.007	1.12 ± 0.007	0.019 ± 0.0007	Uncompetitive
Eta fraction of <i>O. aucheri</i>	61.5 ± 0.6	0.002 ± 0.0001	1.19 ± 0.2	0.025 ± 0.004	Uncompetitive
Buta fraction of <i>O. aucheri</i>	77.95 ± 0.5	0.07 ± 0.002	1.15 ± 0.15	0.025 ± 0.003	Uncompetitive
Iodoacetamide (standard)	At 0.03 mg/mL of iodoacetamide possessed 98.2 ± 2.2% inhibition	0.01 ± 0.002	1.13 ± 0.002	0.024 ± 0.003	Uncompetitive
Without inhibition			1.2 ± 0.003	0.031 ± 0.002	

Abbreviations: ND, not determined; petr, petroleum ether; chlo, chloroform; eta, ethyl acetate; buta, butanol.

^a Values are expressed as mean ± SD.

^b Vmax values of *Otostegia persica* extract and petr, chlo, and eta fractions of *O. aucheri* were significantly different in comparison control (reaction without inhibition; $P < 0.05$). Vmax values of buta fraction and iodoacetamide were not significantly different in comparison control ($P > 0.05$).

^c Km values of all of the samples and iodoacetamide were not significantly different in comparison control ($P > 0.05$).

^d Mixed inhibition such as petr fraction, altered both Km and Vmax.




Figure 3. A, Michaelis-Menten curve of ethanol extract of *Otostegia persica*; B, Michaelis-Menten curve of the control; C, Michaelis-Menten curve of the petroleum ether (petr) fraction of *O. aucheri*; D, Michaelis-Menten curve of the butanol (buta) fraction of *O. aucheri*; E, Michaelis-Menten curve of the chloroform (chlo) fraction of *O. aucheri*; F, Michaelis-Menten curve of the ethyl acetate (eta) fraction of *O. aucheri*; G, Michaelis-Menten curve of iodoacetamide as a standard.

Otostegia aucheri Boiss is endemic to Pakistan and Iran. This genus includes eight species that are commonly scattered in the Mediterranean and southwestern Asia (28, 29). Phenolic compounds are recognized in *Otostegia* species. These compounds are well-known for their anti-inflammatory and antibacterial activities (30). Apigenin in *O. aucheri* may contribute to the inhibition of the enzyme (15). The

results of this study showed that the IC₅₀ of *O. persica* extract was 0.46 ± 0.005 mg/mL (Table 1). On the other hand, different fractions of *O. persica* had no significant inhibitory effects on the inhibition of protease. Also, the IC₅₀ of buta and chlo fractions of *O. aucheri* were 0.07 ± 0.002 mg/mL and 0.09 mg/mL, respectively (Table 1), and the IC₅₀ of iodoacetamide as the standard was 0.01 ±

0.002 mg/mL (Table 1). The eta fraction of *O. aucheri* has the greatest inhibition of protease activity ($IC_{50} = 0.002 \pm 0.0001$), which is more than iodoacetamide ($IC_{50} = 0.01 \pm 0.002$ mg/mL) as a standard (Table 1, $P = 0.04$). The IC_{50} values of petr, chlo, and buta fractions of *O. aucheri* were significantly different in comparison to iodoacetamide ($P = 0.037$). Also, in other research, it is reported that the eta fraction of *O. persica* (30) had the most α -amylase inhibitory activity, which was similar to acarbose used as a standard (30).

For the first time (Table 1), the mode of protease inhibition of *O. persica*, *O. aucheri* extracts, and their different fractions were evaluated, and the results showed that all of the samples, except for the petr fraction of *O. aucheri*, inhibited protease through an uncompetitive mechanism. Also, iodoacetamide, as a positive standard, inhibited protease through an uncompetitive mechanism in which the inhibitor binds to the enzyme-substrate complex, reducing both V_{max} and K_m (Table 1). The petr fraction of *O. aucheri* inhibited protease through a mixed mechanism (Table 1). One advantage of uncompetitive and mixed inhibitors is that they would be effective at lower concentrations of substrate (9); in other words, their inhibitory power is not affected by the concentration of the substrate (8). The comparison of K_m and V_{max} in the presence and absence of inhibitors showed the mode of retardation. If the value of V_{max} is fixed and that of K_m increases, the inhibition is competitive, and if the value of V_{max} is reduced and that of K_m is constant, the inhibition is noncompetitive. If the values of V_{max} and K_m are reduced, the inhibition is uncompetitive; otherwise, the inhibition is mixed (8). It was reported that the methanol extracts of *Salvia macilenta* and *Salvia officinalis* revealed uncompetitive inhibition on α -glucosidase as a marker enzyme of diabetes (31). In another research, anti-protease activities from different parts of *Salvia aculeatissimum* fruits were examined, and Dixon plots and Lineweaver-Burk double reciprocal plots revealed a competitive mode of inhibition (32).

5.1. Conclusions

The genus *Otostegia* is a significant source of biologically active natural compounds and active secondary metabolites. The crude extracts and fractions of different plants belonging to the Lamiaceae family contain good anti-inflammatory and antimicrobial factors (7). Different fractions of *O. aucheri* can significantly inhibit shrimp protease as a 3CLpro model. The eta fraction of *O. aucheri*, with an $IC_{50} = 0.002$ mg/mL compared to iodoacetamide at 0.01 mg/mL,

showed the highest protease inhibition, although different fractions of *O. persica* did not significantly inhibit protease activity. This difference may be attributed to the flavonoids and terpenoids in the *O. persica* and *O. aucheri* extracts. Molecular docking studies could identify active compounds, and HPLC-MS could identify active flavonoids. In addition, the purification of active compounds and clinical trials are warranted to validate therapeutic potential, pending in vivo pharmacokinetic studies.

5.2. Limitations

This study has five limitations. First, the fractionation of ethanol extracts of the studied plants was only performed with petr, chlo, buta, and eta. Future studies should compare different methods of extraction, such as the Soxhlet method, which is a common method for the extraction of lipids, sterols, and fatty acids (33), and fractionation. Second, the experiments were only performed on crude extracts and different fractions of *O. persica* and *O. aucheri*, not on purified compounds of these two plants. Future studies should purify the active compounds of these two plants and evaluate protease inhibition and the mode of inhibition. Also, differences in harvest time or location may affect flavonoid and terpenoid content. Third, in vitro results may not fully reflect in vivo efficacy due to potential metabolism in vivo. Thus, animal or clinical studies are needed to confirm protease inhibition efficacy. Fourth, shrimp protease may not fully mimic 3CLpro kinetics and may differ in the active site from 3CLpro. Fifth, there is a limitation on assay specificity as casein may not mimic viral polyproteins.

Footnotes

Authors' Contribution: Design the study: S. Moein and M. M.; Preparation of the sample: M. M. and Z. Z.; Performing the experiments: Z. Z.; Analysis and interpretation of results: S. Moein; Drafting the manuscript: S. Moein and Z. S.; Statistical analysis: S. Moein.

Conflict of Interests Statement: The authors declare no conflict of interest.

Data Availability: Raw data are available upon request via email to S. Moein et al. (email: SMoein231@gmail.com).

Ethical Approval: This study was approved by the Ethics Committee of Shiraz University of Medical Sciences (IR.SUMS.REC.1400.624).

Funding/Support: The current study was supported by Shiraz University of Medical Sciences (Grant No 24332 for natural product research).

References

- Chen N, Zhou M, Dong X, Qu J, Gong F, Han Y, et al. Epidemiological and clinical characteristics of 99 cases of 2019 novel coronavirus pneumonia in Wuhan, China: a descriptive study. *Lancet*. 2020;**395**(10223):507-13. [PubMed ID: 32007143]. [PubMed Central ID: PMC7135076]. [https://doi.org/10.1016/S0140-6736\(20\)30211-7](https://doi.org/10.1016/S0140-6736(20)30211-7).
- Haagmans BL, Al Dhahiry SH, Reusken CB, Raj VS, Galiano M, Myers R, et al. Middle East respiratory syndrome coronavirus in dromedary camels: an outbreak investigation. *Lancet Infect Dis*. 2014;**14**(2):140-5. [PubMed ID: 24355866]. [PubMed Central ID: PMC7106553]. [https://doi.org/10.1016/S1473-3099\(13\)70690-X](https://doi.org/10.1016/S1473-3099(13)70690-X).
- Corman VM, Muth D, Niemeyer D, Drosten C. Hosts and Sources of Endemic Human Coronaviruses. *Adv Virus Res*. 2018;**100**:163-88. [PubMed ID: 29551135]. [PubMed Central ID: PMC7112090]. <https://doi.org/10.1016/bs.aivir.2018.01.001>.
- Dinda B, Dinda M, Dinda S, Chakraborty M. Some natural compounds and their analogues having potent anti-SARS-CoV-2 and anti-proteases activities as lead molecules in drug discovery for COVID-19. *Eur J Med Chem Rep*. 2022;**6**:100079. [PubMed ID: 36060987]. [PubMed Central ID: PMC9420082]. <https://doi.org/10.1016/j.ejmcr.2022.100079>.
- Hussain AI, Anwar F, Iqbal T, Bhatti IA. Antioxidant attributes of four Lamiaceae essential oils. *Pak. J. Bot*. 2011;**43**(2):1315-21.
- Rashid R, Murtaza G, Khan AK, Mir S. Antioxidant and hypoglycemic effect of *Otostegia aucheri* methanolic extract in streptozotocin-induced diabetic male long-Evans rats. *Acta Pol Pharm*. 2014;**71**(4):631-5. [PubMed ID: 25272889].
- Harley RM, Reynolds T. Advances in labiate science. (No Title). 1992.
- Khan S, Syed F. Bioactive constituents from genus *Otostegia*. *SARJ of Physical Sci*. 2013;**1**(1160):15-25.
- Rouzbehan S, Moein S, Homaei A, Moein MR. Kinetics of alpha-glucosidase inhibition by different fractions of three species of Labiatae extracts: a new diabetes treatment model. *Pharm Biol*. 2017;**55**(1):1483-8. [PubMed ID: 28367665]. [PubMed Central ID: PMC7011978]. <https://doi.org/10.1080/13880209.2017.1306569>.
- Ayatollahi SAM, Kobarfard F, Asgarpanah J, Ahmed Z. Chemical constituents from *Otostegia persica*. *Journal of the Chemical Society of Pakistan*. 2007;**29**(1):61-3.
- Darshani P, Sen Sarma S, Srivastava AK, Baishya R, Kumar D. Anti-viral triterpenes: a review. *Phytochem Rev*. 2022;**21**(6):1761-842. [PubMed ID: 25050304]. [PubMed Central ID: PMC7011978]. <https://doi.org/10.1080/13880209.2017.1306569>.
- Shariffifar F, Yasa N, Shafiei A. Antioxidant activity of *Otostegia persica* (Labiatae) and its constituents. *Iran J Pharmaceut Res*. 2003.
- Villa-Rodriguez JA, Kerimi A, Abranko L, Tumova S, Ford L, Blackburn RS, et al. Acute metabolic actions of the major polyphenols in chamomile: an in vitro mechanistic study on their potential to attenuate postprandial hyperglycaemia. *Sci Rep*. 2018;**8**(1):5471. [PubMed ID: 29615674]. [PubMed Central ID: PMC5882934]. <https://doi.org/10.1038/s41598-018-23736-1>.
- Khandelwal N, Chander Y, Kumar R, Riyesh T, Dedar RK, Kumar M, et al. Antiviral activity of Apigenin against buffalopox: Novel mechanistic insights and drug-resistance considerations. *Antiviral Res*. 2020;**181**:104870. [PubMed ID: 32707051]. <https://doi.org/10.1016/j.antiviral.2020.104870>.
- Farhat A, Ben Hlima H, Khemakhem B, Ben Halima Y, Michaud P, Abdelkafi S, et al. Apigenin analogues as SARS-CoV-2 main protease inhibitors: In-silico screening approach. *Bioengineered*. 2022;**13**(2):3350-61. [PubMed ID: 35048792]. [PubMed Central ID: PMC8974217]. <https://doi.org/10.1080/21655979.2022.2027181>.
- Bachar SC, Mazumder K, Bachar R, Aktar A, Al Mahtab M. A Review of Medicinal Plants with Antiviral Activity Available in Bangladesh and Mechanistic Insight Into Their Bioactive Metabolites on SARS-CoV-2, HIV and HBV. *Front Pharmacol*. 2021;**12**:732891. [PubMed ID: 34819855]. [PubMed Central ID: PMC8606584]. <https://doi.org/10.3389/fphar.2021.732891>.
- Mirza MAHDI, Nik ZB. [Extraction and identification of chemical components of the essential oil of *Salvia compressa* Vent]. *Research Institute of Forests and Rangelands*. 2006. FA.
- Kunitz M. Isolation of a Crystalline Protein Compound of Trypsin and of Soybean Trypsin-Inhibitor. *J Gen Physiol*. 1947;**30**(4):311-20. [PubMed ID: 19873497]. [PubMed Central ID: PMC2142839]. <https://doi.org/10.1085/jgp.30.4.311>.
- Laparra JM, Haros CM. Plant seed protease inhibitors differentially affect innate immunity in a tumor microenvironment to control hepatocarcinoma. *Food Funct*. 2019;**10**(7):4210-9. [PubMed ID: 31257391]. <https://doi.org/10.1039/c9fo00795d>.
- Zumla A, Alagaili AN, Cotten M, Azhar EI. Infectious diseases epidemic threats and mass gatherings: refocusing global attention on the continuing spread of the Middle East Respiratory syndrome coronavirus (MERS-CoV). *BMC Med*. 2016;**14**(1):132. [PubMed ID: 27604081]. [PubMed Central ID: PMC5015245]. <https://doi.org/10.1186/s12916-016-0686-3>.
- Mohammadi Kenari H, Yousefsani BS, Eghbalian F, Ghobadi A, Jamshidi A, Mahroozade S. Herbal recommendations for treatment of COVID-19 symptoms according to Persian medicine. *Journal of Medicinal Plants*. 2021;**20**(77):1-14. <https://doi.org/10.52547/jmp.20.77.1>.
- Nile SH, Keum YS, Nile AS, Jalde SS, Patel RV. Antioxidant, anti-inflammatory, and enzyme inhibitory activity of natural plant flavonoids and their synthesized derivatives. *J Biochem Mol Toxicol*. 2018;**32**(1). [PubMed ID: 28972678]. <https://doi.org/10.1002/jbt.22002>.
- Park M, Cho H, Jung H, Lee H, Hwang KT. Antioxidant and Anti-Inflammatory Activities of Tannin Fraction of the Extract from Black Raspberry Seeds Compared to Grape Seeds. *Journal of Food Biochemistry*. 2014;**38**(3):259-70. <https://doi.org/10.1111/jfbc.12044>.
- Wang SC, Chou IW, Hung MC. Natural tannins as anti-SARS-CoV-2 compounds. *Int J Biol Sci*. 2022;**18**(12):4669-76. [PubMed ID: 35874955]. [PubMed Central ID: PMC9305278]. <https://doi.org/10.7150/ijbs.74676>.
- Russo M, Moccia S, Spagnuolo C, Tedesco I, Russo GL. Roles of flavonoids against coronavirus infection. *Chem Biol Interact*. 2020;**328**:109211. [PubMed ID: 32735799]. [PubMed Central ID: PMC7385538]. <https://doi.org/10.1016/j.cbi.2020.109211>.
- Ullah S, Munir B, Al-Sehemi AG, Muhammad S, Haq IU, Aziz A, et al. Identification of phytochemical inhibitors of SARS-CoV-2 protease 3CL(pro) from selected medicinal plants as per molecular docking, bond energies and amino acid binding energies. *Saudi J Biol Sci*. 2022;**29**(6):103274. [PubMed ID: 35345871]. [PubMed Central ID: PMC8944115]. <https://doi.org/10.1016/j.sjbs.2022.03.024>.
- Scheen A, Albert VA. Nomenclatural and taxonomic changes within the Leucas clade (Lamiaceae; Lamiaceae). *Systematics and Geography of Plants*. 2007:229-38.
- Scheen A, Albert VA. Molecular Phylogenetics of the Leucas Group (Lamiaceae; Lamiaceae). *Systematic Botany*. 2009;**34**(1):173-81. <https://doi.org/10.1600/036364409787602366>.
- Sadeghi Z, Akaberi M, Valizadeh J. *Otostegia persica* (Lamiaceae): A review on its ethnopharmacology, phytochemistry, and pharmacology. *Avicenna J Phytomed*. 2014;**4**(2):79-88. [PubMed ID: 25050304]. [PubMed Central ID: PMC4103708].
- Moein S, Pimoradloo E, Moein M, Vessal M. Evaluation of Antioxidant Potentials and alpha-Amylase Inhibition of Different Fractions of

- Labiatae Plants Extracts: As a Model of Antidiabetic Compounds Properties. *Biomed Res Int.* 2017;**2017**:7319504. [PubMed ID: 29082253]. [PubMed Central ID: PMC5610887]. <https://doi.org/10.1155/2017/7319504>.
31. Javid H, Moein S, Moein MR. Evaluating the Inhibitory Effects of Dichloromethane and Methanol Extracts of *Salvia macilenta* and *Salvia officinalis* on the Diabetes Marker Enzyme, Alpha-Glucosidase: An Approach for the Treatment of Diabetes. *Jundishapur Journal of Natural Pharmaceutical Products.* 2021;**16**(3). <https://doi.org/10.5812/jjnpp.109694>.
 32. Meenu Krishnan VG, Murugan K. Purification, characterization and kinetics of protease inhibitor from fruits of *Solanum aculeatissimum* Jacq. *Food Science and Human Wellness.* 2015;**4**(3):97-107. <https://doi.org/10.1016/j.fshw.2015.06.003>.
 33. Yu X, Tu X, Tao L, Daddam J, Li S, Hu F. Royal Jelly Fatty Acids: Chemical Composition, Extraction, Biological Activity, and Prospect. *Journal of Functional Foods.* 2023;**111**. <https://doi.org/10.1016/j.jff.2023.105868>.