Published online 2022 July 13.

Research Article

The Interactive Effect of Home-Based Aerobic Training and Turmeric Supplementation on the Quality of Life in Type 2 Diabetic Women

Mahsa Ahmadi Darmian 💿¹, Rastegar Hoseini 💿^{1,*}, Ehsan Amiri 💿¹ and Sanam Golshani 💿²

¹Department of Exercise Physiology, Faculty of Sport Sciences, Razi University, Kermanshah, Iran
²Kermanshah University of Medical Sciences, Army Hospital No. 520, Kermanshah, Iran

Corresponding author: Department of Exercise Physiology, Faculty of Sport Sciences, Razi University, Kermanshah, Iran. Email: r.hoseini@razi.ac.ir

Received 2021 July 02; Revised 2022 April 23; Accepted 2022 May 07.

Abstract

Background: Type 2 diabetes is the most common health problem worldwide. Exercise and dietary interventions can help individuals with type 2 diabetes to improve their condition.

Objectives: The purpose of this study was to determine the effects of aerobic training (AT) and turmeric supplement (TS) on the quality of life (QoL) of middle-aged hyperlipidemic women with type 2 diabetes.

Methods: This study investigated 42 women with type 2 diabetes aged 45 to 60 years, who were volunteered and divided into four groups, including control + placebo (C; n = 11), AT + placebo (AT; n = 10), TS (n = 11), and AT + TS (n = 11). Aerobic training program was conducted for eight weeks (3 sessions per week for 20 to 40 minutes with 60 - 75% of maximum heart rate). The TS and AT + TS consumed 2100 mg per day of turmeric for eight weeks. The 26-item WHO QoL questionnaire (WHO QoL-BREF) was used to assess QoL. Finally, the data were statistically analyzed at a significant level of P < 0.05 using one-way analysis of variance (ANOVA), Tukey's post hoc, and an independent *t*-test with SPSS.

Results: AT + TS (P = 0.001), AT (P = 0.023), and TS (P = 0.039) significantly improved the average QoL after eight weeks. However, the improvement in AT + TS was more significant for the QoL of type 2 diabetic patients with hyperlipidemia compared to other groups. **Conclusions:** According to the results, exercise and turmeric supplementation can be used to improve the QoL of middle-aged women with type 2 hyperlipidemia during COVID-19 quarantine.

Keywords: Exercise, Curcumin, Quality of Life, Coronavirus Disease

1. Background

Hyperlipidemic diabetes (HD) is one of the most common chronic diseases in the present century (1). Hyperlipidemic diabetes is characterized by an improper response to insulin, disruption of fat, carbohydrate, and protein metabolism, and eventually hyperglycemia (2). According to the latest data on diabetes incidence, there are 230 million cases of diabetes worldwide and 3 million cases in Iran (3, 4). Hyperlipidemic diabetes affects various aspects of individuals' life causing physical and psychological problems by increasing the risk of comorbidities (5). The most significant predictors of HD are sedentary lifestyles, unhealthy diets, and reduced physical activity (6). A general concept called Quality of Life (QoL), which is impaired in patients with HD, can shed light on the overall health status of a person (7-9). The primary goal of HD management is achieving higher

levels of QoL and preventing the onset of comorbidities (10). As a non-pharmacological solution, a regular exercise program has been known as a critical component of HD improvement to reduce the complications of diabetes and improve the mental status and QoL (11, 12). Exercise benefits QoL by increasing the average scores of psychosocial physical domains in patients with HD (13), among which aerobic training (AT) plays an essential role in increasing insulin-induced glucose uptake, strengthening skeletal muscles, reducing depression, and thus enhancing QoL (14, 15).

As a result of COVID-19 onset, individuals with chronic diseases such as diabetes are now at higher risk of developing viral infections (16, 17). Quarantine recommendations focusing on staying home impacted social behaviors and physical activity levels, leading to inactivity and reduced QoL (18). Aerobic training is an appropriate approach to reduce inactivity and increase

Copyright © 2022, Journal of Kermanshah University of Medical Sciences. This is an open-access article distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (http://creativecommons.org/licenses/by-nc/4.0/) which permits copy and redistribute the material just in noncommercial usages, provided the original work is properly cited.

QoL in COVID-19 susceptible populations (19).

Herbal supplement, such as turmeric supplementation (TS), is another strategy to control and prevent HD (20-22), which has been considered due to its lower side effects, availability, and cost-effectiveness (23). Due to the presence of curcumin, TS is known to have the most therapeutic effects on glycemic control and to reduce the complications of HD via antioxidant properties, which might also increase the QoL (24-26). Maithili Karpaga Selvi et al. reported that TS enhances blood glucose, oxidative stress, and inflammation in patients with T2DM (27). However, Adab et al. observed the improvement of bodyweight following TS in patients with HD and reported no significant effect on total antioxidant capacity, hs-CRP, and glycemic status (28). Although numerous studies have examined the effects of TS on the physiological status of diabetics, no study has examined the effects of TS on the quality of life of HD patients.

2. Objectives

Due to the limited results in the literature investigating the simultaneous effect of AT + TS, the purpose of the present study was to evaluate the interactive effect of AT + TS on the QoL during COVID-19 quarantine in type 2 diabetic women with hyperlipidemia.

3. Methods

3.1. Trial Design and Participants

This quasi-experimental (single-blinded) study was conducted with three experimental groups and one control group. Forty-four women with type 2 diabetes were selected (The Diabetes Center of Taleghani Hospital, Kermanshah, Iran) by purposive and convenient sampling. In addition, G. POWER 3.1 was used at a significance level of 0.05 with a statistical power = 80 % and an effect size = 95%. Then, the subjects were randomly divided into four groups, including control + placebo (C; n = 11), AT + placebo (AT; n = 11), TS (n = 11), and AT + TS (n = 11). Two subjects refused to continue the study (one in AT and one in C; Figure 1).

The inclusion criteria included women diagnosed with type 2 diabetes, aged 40 to 65 years, and classified as overweight (body mass index (BMI), 25 to 29.99 kg/m²). The exclusion criteria included having regular exercise, a history of other diseases, smoking, and COVID-19 infection during the intervention.

The Ethics Committee of Kermanshah Razi University approved this trial (IR.RAZI.REC.1400.008). This study was also registered in the Iranian Clinical Trial Registration Center (IRCT20201129049524N1). All participants completed and signed a written informed consent to participate or leave the study voluntarily. The medical history questionnaire revealed no evidence of coronary artery disease, renal failure, or thyroid disease among the subjects. The information about research conduction, possible risks, the online home-based exercise sessions, and controlling the exercise intensity during the training sessions was explained to the subjects. In addition, the informed consent and physical activity readiness questionnaire were completed. Then, the initial variables, including height, weight, and body fat percentage, were measured.

3.2. Intervention

3.2.1. Aerobic Training

Aerobic training was conducted at home under the online supervision of exercise physiologists three times per week for eight weeks. The AT started with 20 min at 60% of HR_{max} , and increased to 40 min at 75% of HR_{max} according to the American Diabetes Association (ADA) (29) (Table 1). Warming up and cooling down time were each 10 minutes.

The HR_{max} formula [$HR_{max} = 220$ - age] determined the target heart rate (30). The participants learned the pulse palpation method to count their pulse or heart rate in an instructional session. To assure achieving the desired exercise intensity, the rating of perceived exertion (RPE) (6 - 20 scale) was considered during the exercise session (31) (Table 1).

3.2.2. Diet and Turmeric Supplementation

The participants filled out two questionnaires to attend the training program. First, a questionnaire assessed demographic data, physical activity, and health status. Then, a detailed semi-quantitative food record questionnaire adapted to the Iranian population was used before and after the intervention that was composed of a limited checklist of foods and beverages (everyday food items, serving sizes, meals) with a separate frequency response section and portion size for subjects. Participants were requested to consume the same macronutrient composition one day before the post-test. The Food Processor Nutritionist Four (FPN4) software determined the food intake via multiplying the reported frequency of each food by the macronutrient amount (protein, fat, and carbohydrates) in a serving size. The diet consisted of 15% protein, 30% fat, and 55% carbohydrates. Table 2 shows the carbohydrate, protein, and fat consumption. The reliability and validity of this questionnaire are also evaluated.



Figure 1. Flow chart of the study population

Table 1. Aerobic Training Protocol Week Variables 1 2 3 4 5 6 7 8 Intensity (HR_{Max},%) 60 - 65 60 - 65 60 - 65 65 - 70 65 - 70 65 - 70 70 - 75 70 - 75 Time (min) 40 40 20 25 30 30 35 35 Borg scale 10 10 11 11 12 12 13 13

J Kermanshah Univ Med Sci. 2022; 26(1):e115917.

Table 2. Mean ± Standard Deviation Food Intake Between Groups										
Groups -	Carbohydrate (g)		Protein (g)		Fat (g)					
	Pre	Post	Pre	Post	Pre	Post				
AT + TS	387.8 ± 123.22	360.4 ± 138.13	112.3 ± 18.01	116.6 ± 22.04	95.15 ± 7.01	92.7± 8.02				
AT	360.7±133.41	347.7± 98.42	107.2 ± 23.08	112.62 ± 16.02	98.18 ± 6.03	96.14 ± 5.04				
TS	383.3± 88.29	365.6 ± 117.18	115.35 ± 24.06	118.63 ± 13.03	100.16 ± 8.07	99.11± 4.03				
С	357.8±123.31	366.6 ± 89.23	110.3 ± 27.11	111.5 ± 11.07	88.14 ± 11.02	100.11± 8.01				

Abbreviations: C, the control group; TS, turmeric supplement group; AT, aerobic training group; AT + TS, aerobic training + turmeric supplement.

According to previous research, consuming 1 - 3 g of turmeric per day is safe. Therefore, 2100 mg per day (2.1 g per day) was determined based to human studies (32). In the TS and AT + TS groups, 700 mg of turmeric powder was consumed as part of each meal for eight weeks. The AT and C received three capsules containing 700 mg of cornstarch flour, which were identical in shape, color, and packaging.

3.2.3. Anthropometric Measurements and Body Composition

Three days before the intervention, anthropometric parameters and body composition were measured. A stadiometer (DETECTO, Model 3PHTROD-WM, USA) was used to measure height to the nearest 0.5 cm. A non-elastic tape measure was utilized to determine waist circumference to the nearest 0.5 cm. In addition, the BMI and weight were determined using bioelectric impedance analysis by the INBODY test (Zeus 9.9 PLUS; Jawon Medical Co., Ltd., Kungsang Bukdo, South Korea) in the morning after at least 12 hours of fasting. Subjects were asked to refrain from taking diuretic drugs and diuretics 48 hours before the measurements.

3.3. Quality of Life

The SF-12 was used to evaluate QoL. It has two summary components of the mental component summary (MCS) and physical component summary (PCS), 12 questions, and eight scales, including mental health (MH), role limitations due to emotional problems (RE), general health perception (GHP), vitality (VT), social functioning (SF), physical functioning (PF), role limitations due to physical health (PH), and body pain (BP). The final score of the SF-12 ranged from 0 to 100, with a higher score indicating a better QoL. The SF-12 was adapted and validated to Persian conditions by Rohani et al. (33).

3.4. Data Analysis

The data were analyzed at a significant P < 0.05 using the SPSS (Version 21; SPSS Inc., Chicago, IL, USA) Software. To evaluate the normality of distribution, the Shapiro-Wilk's test was used. The *t*-test and ANOVA were

used to compare the mean variables within and between groups. In addition, Tukey's post hoc test was used in the case of significant differences.

4. Results

The descriptive characteristics of the participants are presented in Table 3. As shown in Table 3, the mean \pm SD of BMI and body fat percentage of all participants were in the overweight category. Furthermore, all subjects in the present study had the indicators of type 2 diabetes and hyperlipidemia.

The correlated *t*-test shows that AT + TS, AT, and TS decreased the QoL in the post-test compared to the pre-test, which was not statistically significant. In addition, the C group showed decreased QoL (Figure 2). The results of one-way ANOVA showed no significant difference in the mean QoL in the pre-test between groups, while a significant difference was observed in the post-test. Tukey's post hoc test results showed a significant difference in the QoL between the AT + TS, AT, and TS with C. Moreover, there was a significant difference between AT + TS with TS but not with AT. Besides, a significant difference was observed between the AT and TS.

5. Discussion

Most of the evidence about the health benefits of physical activity is based on epidemiological studies, and there are very few interventional studies. However, our knowledge of the interactive effects of AT + TS on the QoL in women with HD is limited.

The present study indicated that eight weeks of AT + TS, AT, and TS significantly increased QoL. However, the interactive effect of AT + TS was higher than the separate effect. Our results are in line with Cai et al. (34), Chiang et al. (35), and Delevatti et al. (36), reporting improved QoL in patients with HD following separate AT or TS interventions. The positive effect of AT on improving the QoL in patients with HD might

Variables	AT + TS	AT	TS	С	P-Value ^a
Age (y)	43.02 ± 3.04	42.13 ± 2.39	44.33 ± 1.23	44.22 ± 3.07	0.336
Height	160.12 ± 4.11	163.13 ± 3.03	159.24 ± 2.79	161.13 ± 5.08	0.393
Body weight (kg)	73.12 ± 2.91	75.13 ± 2.07	74.13 ± 2.68	75.09 ± 3.20	0.061
Body mass index (kg/m ²)	28.15 ± 1.02	28.12 ± 1.23	29.30 ± 0.30	29.02 ± 1.04	0.223
Body fat percent (%)	30.14 ± 1.26	32.09 ± 2.01	29.23 ± 1.27	31.16 ± 2.52	0.206
WHR	0.90 ± 0.03	0.92 ± 0.04	0.91± 0.05	0.92 ± 0.02	0.229

Abbreviations: C, the control group; TS, turmeric supplement group; AT, aerobic training group; AT + TS, aerobic training + turmeric supplement. ^a P values is calculated using one-way analysis of variance test followed by post hoc Tukey's test.





be probably due to the improved biochemical structure of muscle, increased oxygen consumption, and reduced insulin resistance (37). According to other studies, AT reduces the required proinsulin mRNA to synthesize insulin and decreases glucokinase levels (29, 30). This, in turn, decreases blood insulin levels and phosphorylated insulin receptors (38). Furthermore, AT activates the central nervous system and increases the secretion of endorphins, norepinephrine, serotonin synthesis, and endorphins, which reduces complications and improves QoL in patients with HD (39).

In addition, TS has been shown to have antioxidant, anti-inflammatory, antibacterial, antiviral, and anti-cancer characteristics with a potential role in treating many chronic diseases such as diabetes (40). Possible mechanisms by which turmeric improves the QoL in patients with HD include increased acyl-CoA cholesterol acyltransferase (ACAT) and carnitine palmitoyltransferase 1 (CPT1) (41, 42), decreased IL-6, HbA1c, and TNF- α (43, 44) inhibited MCP-1, and suppressed NF κ B signaling pathway (45). Thus, AT + TS might ultimately improve QoL in HD patients through the mechanisms outlined above.

5.1. Conclusions

Eight weeks of separate and interactive AT and TS improved the QoL in women with HD. However, the mechanisms involved in these changes have not yet been fully elucidated. According to results, AT + TS might be an appropriate non-pharmacological treatment to improve the QoL in patients with HD, given the positive effects of AT + TS. The main limitations of this study included the small sample size and not controlling the food intake within the eight weeks. More accurate results could be achieved by more control.

Footnotes

Authors' Contribution: R. H. designed the study. M. A. D. experimented. R. H. analyzed the data and wrote the manuscript. E. A. and S. G. involved in the interpretation of data, reviewed and edited the manuscript. All authors read and approved the final manuscript.

Clinical Trial Registration Code: This study was registered in the Iranian Clinical Trial Registration Center under the code IRCT20201129049524N1.

Conflict of Interests: The authors do not have any competing interests.

Data Reproducibility: The data presented in this study are openly available in one of the repositories or will be available on request from the corresponding author by this journal representative at any time during submission or after publication. Otherwise, all consequences of possible withdrawal or future retraction will be with the corresponding author.

Ethical Approval: This study was approved by the ethics committee of Razi University of Kermanshah (IR.RAZI.REC.1400.013), Iran.

Funding/Support: The author(s) received no specific funding for this work.

Informed Consent: All participants completed and signed a written informed consent to participate or leave the study voluntarily.

References

- Liu Y, Wang Y, Ni Y, Cheung CKY, Lam KSL, Wang Y, et al. Gut Microbiome Fermentation Determines the Efficacy of Exercise for Diabetes Prevention. *Cell Metab.* 2020;**31**(1):77–91 e5. [PubMed ID: 31786155]. https://doi.org/10.1016/j.cmet.2019.11.001.
- American Diabetes Association. Diagnosis and classification of diabetes mellitus. *Diabetes Care*. 2005;**28 Suppl 1**:S37-42. [PubMed ID: 15618111]. https://doi.org/10.2337/diacare.28.suppl_1.s37.
- Singh AK, Gupta R, Ghosh A, Misra A. Diabetes in COVID-19: Prevalence, pathophysiology, prognosis and practical considerations. *Diabetes Metab Syndr*. 2020;14(4):303–10. [PubMed ID: 32298981]. [PubMed Central ID: PMC7195120]. https://doi.org/10.1016/j.dsx.2020.04.004.

- Esteghamati A, Larijani B, Aghajani MH, Ghaemi F, Kermanchi J, Shahrami A, et al. Diabetes in Iran: Prospective Analysis from First Nationwide Diabetes Report of National Program for Prevention and Control of Diabetes (NPPCD-2016). *Sci Rep.* 2017;7(1):13461. [PubMed ID: 29044139]. [PubMed Central ID: PMC5647418]. https://doi.org/10.1038/s41598-017-13379-z.
- Graham JE, Stoebner-May DG, Ostir GV, Al Snih S, Peek MK, Markides K, et al. Health related quality of life in older Mexican Americans with diabetes: a cross-sectional study. *Health Qual Life Outcomes*. 2007;5:39. [PubMed ID: 17626634]. [PubMed Central ID: PMC1947953]. https://doi.org/10.1186/1477-7525-5-39.
- 6. Riyahi Malayeri S, Abdolhay S, Behdari R, Hoseini M. [The combined effect of resveratrol supplement and endurance training on IL-10 and TNF- α in type 2 diabetic rats]. *Razi Journal of Medical Sciences*. 2019;**25**(12):140–9. Persain.
- Anderson BJ, Laffel LM, Domenger C, Danne T, Phillip M, Mazza C, et al. Factors Associated With Diabetes-Specific Health-Related Quality of Life in Youth With Type 1 Diabetes: The Global TEENs Study. *Diabetes Care*. 2017;40(8):1002–9. [PubMed ID: 28546221]. [PubMed Central ID: PMC5864137]. https://doi.org/10.2337/dc16-1990.
- Goh SG, Rusli BN, Khalid BA. Diabetes quality of life perception in a multiethnic population. *Qual Life Res.* 2015;24(7):1677-86. [PubMed ID: 25492728]. [PubMed Central ID: PMC4483180]. https://doi.org/10.1007/s11136-014-0885-3.
- Speight J, Holmes-Truscott E, Hendrieckx C, Skovlund S, Cooke D. Assessing the impact of diabetes on quality of life: what have the past 25 years taught us? *Diabet Med.* 2020;**37**(3):483–92. [PubMed ID: 31797443]. https://doi.org/10.1111/dme.14196.
- Chalmers J, Cooper ME. UKPDS and the legacy effect. N Engl J Med. 2008;359(15):1618–20. [PubMed ID: 18843126]. https://doi.org/10.1056/NEJMe0807625.
- 11. Bakhshalipour V, Azizi B, Khodaparast Sareshkeh S. [The effect of a 6-months moderate-intensity walking exercise program on mental health, quality of life, and sleep subjective quality in non-active elderly people with type 2 diabetes]. RBNE-Revista Brasileira de Nutrição Esportiva. 2018;12(75):905–14. Portuguese.
- Morgan AJ, Parker AG, Alvarez-Jimenez M, Jorm AF. Exercise and Mental Health: An Exercise and Sports Science Australia Commissioned Review. J Exerc Physiol Online. 2013;16(4).
- Heydari M, Elhani F, Kazemnejad A, Moezzi F. [The effect of empowerment model on quality of life of diabetic adolescents]. *Iran J Pediatr.* 2007;**17**(1 (supplement)). Persian.
- Bridle C, Spanjers K, Patel S, Atherton NM, Lamb SE. Effect of exercise on depression severity in older people: systematic review and meta-analysis of randomised controlled trials. *Br J Psychiatry*. 2012;**201**(3):180–5. [PubMed ID: 22945926]. https://doi.org/10.1192/bjp.bp.111.095174.
- Gilani SRM, Feizabad AK. The effects of aerobic exercise training on mental health and self-esteem of type 2 diabetes mellitus patients. *Health Psychol Res.* 2019;7(1):6576. [PubMed ID: 30997432]. [PubMed Central ID: PMC6441819]. https://doi.org/10.4081/hpr.2019.6576.
- Cure E, Cumhur Cure M. Angiotensin-converting enzyme inhibitors and angiotensin receptor blockers may be harmful in patients with diabetes during COVID-19 pandemic. *Diabetes Metab Syndr.* 2020;14(4):349–50. [PubMed ID: 32311651]. [PubMed Central ID: PMC7159862]. https://doi.org/10.1016/j.dsx.2020.04.019.
- Sohrabi C, Alsafi Z, O'Neill N, Khan M, Kerwan A, Al-Jabir A, et al. World Health Organization declares global emergency: A review of the 2019 novel coronavirus (COVID-19). *Int J Surg.* 2020;**76**:71–6. [PubMed ID: 32112977]. [PubMed Central ID: PMC7105032]. https://doi.org/10.1016/j.ijsu.2020.02.034.
- Mora-Gonzalez J, Esteban-Cornejo I, Cadenas-Sanchez C, Migueles JH, Rodriguez-Ayllon M, Molina-Garcia P, et al. Fitness, physical activity, working memory, and neuroelectric activity in children with overweight/obesity. *Scand J Med Sci Sports*. 2019;**29**(9):1352–63. [PubMed ID: 31058358]. https://doi.org/10.1111/sms.13456.

- Chekroud SR, Gueorguieva R, Zheutlin AB, Paulus M, Krumholz HM, Krystal JH, et al. Association between physical exercise and mental health in 1-2 million individuals in the USA between 2011 and 2015: a cross-sectional study. *The Lancet Psychiatry*. 2018;5(9):739–46. https://doi.org/10.1016/s2215-0366(18)30227-x.
- Arumugam G, Manjula P, Paari N. A review: Anti diabetic medicinal plants used for diabetes mellitus. J Acute Dis. 2013;2(3):196–200. https://doi.org/10.1016/s2221-6189(13)60126-2.
- Bindu J, Narendhirakannan RT. Role of medicinal plants in the management of diabetes mellitus: a review. 3 *Biotech*. 2019;9(1):4. [PubMed ID: 30555770]. [PubMed Central ID: PMC6291410]. https://doi.org/10.1007/s13205-018-1528-0.
- Elekofehinti OO. Saponins: Anti-diabetic principles from medicinal plants - A review. *Pathophysiology*. 2015;22(2):95–103. [PubMed ID: 25753168]. https://doi.org/10.1016/j.pathophys.2015.02.001.
- 23. Tarighat esfanjani A, Namazi N, Bahrami A. [Effect of Hydro-alcoholic Nettle Extract on Lipid Profiles and Blood Pressure in Type 2 Diabetes Patients]. *Iran J Endocrinol Metab.* 2012;**13**(5):449–58. Persian.
- Srinivasan K. Plant foods in the management of diabetes mellitus: spices as beneficial antidiabetic food adjuncts. *Int J Food Sci Nutr.* 2005;56(6):399–414. [PubMed ID: 16361181]. https://doi.org/10.1080/09637480500512872.
- Zahid Ashraf M, Hussain ME, Fahim M. Antiatherosclerotic effects of dietary supplementations of garlic and turmeric: Restoration of endothelial function in rats. *Life Sci.* 2005;77(8):837–57. [PubMed ID: 15964306]. https://doi.org/10.1016/j.lfs.2004.11.039.
- De Man FH, Cabezas MC, Van Barlingen HH, Erkelens DW, de Bruin TW. Triglyceride-rich lipoproteins in non-insulin-dependent diabetes mellitus: post-prandial metabolism and relation to premature atherosclerosis. *Eur J Clin Invest.* 1996;**26**(2):89-108. [PubMed ID: 8904518]. https://doi.org/10.1046/j.1365-2362.1996.114256.x.
- Maithili Karpaga Selvi N, Sridhar MG, Swaminathan RP, Sripradha R. Efficacy of Turmeric as Adjuvant Therapy in Type
 Diabetic Patients. Indian J Clin Biochem. 2015;30(2):180–6. [PubMed ID: 25883426]. [PubMed Central ID: PMC4393385]. https://doi.org/10.1007/s12291-014-0436-2.
- Adab Z, Eghtesadi S, Vafa MR, Heydari I, Shojaii A, Haqqani H, et al. Effect of turmeric on glycemic status, lipid profile, hs-CRP, and total antioxidant capacity in hyperlipidemic type 2 diabetes mellitus patients. *Phytother Res.* 2019;33(4):1173–81. [PubMed ID: 30859660]. https://doi.org/10.1002/ptr.6312.
- Aylin K, Arzu D, Sabri S, Handan TE, Ridvan A. The effect of combined resistance and home-based walking exercise in type 2 diabetes patients. Int J Diabetes Dev Ctries. 2009;29(4):159–65. [PubMed ID: 20336198]. [PubMed Central ID: PMC2839130]. https://doi.org/10.4103/0973-3930.57347.
- Branco BHM, de Oliveira Mendes F, Ladeia GF, Bertolini SMMG, Badilla PV, Andreato LV. Maximum heart rate predicted by formulas versus values obtained in graded exercise tests in Brazilian jiu-jitsu athletes. Sport Sci Health. 2019;16(1):39–45. https://doi.org/10.1007/s11332-019-00570-0.
- 31. Rahimi E. [Physical activity and type 2 diabetes: A narrative review]. *Journal of Physical Activity and Hormones*. 2019;**2**(4):51–62. Persian.
- 32. Khajehdehi P, Zanjaninejad B, Aflaki E, Nazarinia M, Azad F, Malekmakan L, et al. Oral supplementation of turmeric decreases proteinuria, hematuria, and systolic blood pressure in patients suffering from relapsing or refractory lupus nephritis: a randomized and placebo-controlled study. J Ren Nutr. 2012;22(1):50–7. [PubMed ID: 21742514]. https://doi.org/10.1053/j.jrn.2011.03.002.
- 33. Rohani C, Abedi HA, Langius A. The Iranian SF-12 Health Survey Version

2 (SF-12v2): Factorial and Convergent Validity, Internal Consistency and Test-Retest in a Healthy Sample. *Iranian Rehabilitation Journal*. 2010;**8**(2):4-14. eng.

- 34. Cai H, Li G, Zhang P, Xu D, Chen L. Effect of exercise on the quality of life in type 2 diabetes mellitus: a systematic review. Qual Life Res. 2017;26(3):515-30. [PubMed ID: 27990609]. https://doi.org/10.1007/s11136-016-1481-5.
- 35. Chiang S, Chiou C, Lin C, Ho C, Chen L, Chang C, et al. Effects of a 12-week exercise training on insulin sensitivity, quality of life, and depression status in patients with type 2 diabetes. J Med Sci. 2017;37(6):227. https://doi.org/10.4103/jmedsci.jmedsci_68_17.
- 36. Delevatti RS, Schuch FB, Kanitz AC, Alberton CL, Marson EC, Lisboa SC, et al. Quality of life and sleep quality are similarly improved after aquatic or dry-land aerobic training in patients with type 2 diabetes: A randomized clinical trial. *J Sci Med Sport*. 2018;21(5):483–8. [PubMed ID: 28935128]. https://doi.org/10.1016/j.jsams.2017.08.024.
- Tadibi V. [Effect of eight weeks aerobic training and drug intervention on quality of life in women with type 2 diabetes]. J Gorgan Uni Med Sci. 2012;14(2):30–6. Persian.
- Colberg SR, Sigal RJ, Fernhall B, Regensteiner JG, Blissmer BJ, Rubin RR, et al. Exercise and type 2 diabetes: the American College of Sports Medicine and the American Diabetes Association: joint position statement. *Diabetes Care*. 2010;**33**(12):e147–67. [PubMed ID: 21115758]. [PubMed Central ID: PMC2992225]. https://doi.org/10.2337/dc10-9990.
- Eckert K. Impact of physical activity and bodyweight on health-related quality of life in people with type 2 diabetes. *Diabetes Metab Syndr Obes.* 2012;5:303–11. [PubMed ID: 22952412]. [PubMed Central ID: PMC3430085]. https://doi.org/10.2147/DMSO.S34835.
- Perrone D, Ardito F, Giannatempo G, Dioguardi M, Troiano G, Lo Russo L, et al. Biological and therapeutic activities, and anticancer properties of curcumin. *Exp Ther Med.* 2015;**10**(5):1615–23. [PubMed ID: 26640527]. [PubMed Central ID: PMC4665301]. https://doi.org/10.3892/etm.2015.2749.
- Seo KI, Choi MS, Jung UJ, Kim HJ, Yeo J, Jeon SM, et al. Effect of curcumin supplementation on blood glucose, plasma insulin, and glucose homeostasis related enzyme activities in diabetic db/db mice. *Mol Nutr Food Res*. 2008;**52**(9):995–1004. [PubMed ID: 18398869]. https://doi.org/10.1002/mnfr.200700184.
- Kaur G, C M. Amelioration of obesity, glucose intolerance, and oxidative stress in high-fat diet and low-dose streptozotocin-induced diabetic rats by combination consisting of "curcumin with piperine and quercetin". *ISRN Pharmacol.* 2012;2012:957283. [PubMed ID: 22474599]. [PubMed Central ID: PMC3317057]. https://doi.org/10.5402/2012/957283.
- El-Moselhy MA, Taye A, Sharkawi SS, El-Sisi SF, Ahmed AF. The antihyperglycemic effect of curcumin in high fat diet fed rats. Role of TNF-alpha and free fatty acids. *Food Chem Toxicol*. 2011;49(5):1129–40. [PubMed ID: 21310207]. https://doi.org/10.1016/j.fct.2011.02.004.
- 44. Na LX, Zhang YL, Li Y, Liu LY, Li R, Kong T, et al. Curcumin improves insulin resistance in skeletal muscle of rats. *Nutr Metab Cardiovasc Dis.* 2011;21(7):526–33. [PubMed ID: 20227862]. https://doi.org/10.1016/j.numecd.2009.11.009.
- 45. Jain SK, Rains J, Croad J, Larson B, Jones K. Curcumin supplementation lowers TNF-alpha, IL-6, IL-8, and MCP-1 secretion in high glucose-treated cultured monocytes and blood levels of TNF-alpha, IL-6, MCP-1, glucose, and glycosylated hemoglobin in diabetic rats. *Antioxid Redox Signal*. 2009;**11**(2):241–9. [PubMed ID: 18976114]. [PubMed Central ID: PMC2933148]. https://doi.org/10.1089/ars.2008.2140.