



Factors Affecting Sleep Problems Among Young People During COVID-19 Pandemic: A Report from Western Iran

Habibolah Khazaie ¹, Masoumeh Rostampour¹, Leeba Rezaie¹, Saeid Komasi ¹ and Ali Zakiei ^{1,*}

¹Sleep Disorders Research Center, Kermanshah University of Medical Sciences, Kermanshah, Iran

*Corresponding author: Sleep Disorders Research Center, Kermanshah University of Medical Sciences, Kermanshah, Iran. Email: zakiei.1@gmail.com

Received 2023 September 20; Revised 2023 November 27; Accepted 2023 November 28.

Abstract

Background: Sleep disorders are a public health problem, and sleep studies among ethnic groups are few. Therefore, this study aimed to evaluate sleep quality and sleep problems among Iranian people.

Methods: This cross-sectional study was conducted on 4748 young people aged 18 to 35 (average: 27.58 years) in western Iran during the COVID-19 pandemic. In addition, 60.4% of the participants were men. The data were collected using the Pittsburgh Sleep Quality Inventory (PSQI), the Insomnia Severity Index (ISI), and the Demographic Information Questionnaire and analyzed with SPSS software version 22.

Results: About 44.4% of the participants had unfavorable sleep quality and had sleep problems. Besides, the mean score of sleep problems (measured with PSQI) reported by the participants was 5.68 ± 2.67 . The logistic regression analysis showed that the participant's age, education, marital status, occupation, drug abuse, alcohol consumption, use of mobile phones and the internet, and a history of previous sleep problems accounted for sleep problems during the COVID-19 pandemic.

Conclusions: Based on the results, there was a high prevalence of sleep problems among Iranian young people during the COVID-19 pandemic.

Keywords: COVID-19, Insomnia, Sleep Quality, Sleep Problems, Western Iran

1. Background

Sleep problems are a significant public health issue that can increase the mortality rate among people and impose a substantial financial burden on society from an economic perspective (1). Sleep problems are prevalent among adolescents and young adults (2, 3), including a wide range, including insomnia, reduced total sleep time, increased sleep onset latency, nighttime awakenings (4), reduced sleep quality, delayed sleep onset, sleep efficiency, sleep cycle disturbances, sleep fragmentation, sleep medications, and daytime sleepiness (5). Nightmares can also be considered as part of sleep problems (6). Sleep problems are associated with mental disorders such as depression and post-traumatic stress disorder (PTSD) (7) as risk factors for self-harming and suicidal behaviors (8, 9) and contribute to the development of physical problems and diseases such as heart disease and diabetes (10, 11). Furthermore, sleep problems make a person susceptible to poor metabolic health and weight gain (12), affecting systemic inflammation (13).

Studies have indicated that sleep problems are complex issues, which should be understood (14). Furthermore, sleep problems may lead to potentially life-threatening symptoms, which are the consequences of this and primary triggers of other diseases (15). Thus, more comprehensive studies are needed to explore these problems. In addition, natural or social crises always have consequences that affect social, family, and psychological life (16), including the COVID-19 pandemic, which has adverse social and economic consequences and health threats. Therefore, these conditions may have affected people's sleep health. Studies have shown that sleep problems increased during the COVID-19 pandemic, and these problems have had consequences for people (17-19). A study showed that sleep problems play a role in the prevalence and persistence of PTSD and depression in university students during the COVID-19 pandemic (20). A longitudinal study also showed that sleep problems had a significant relationship with the increase in suicide during the COVID-19 pandemic (21).

Many studies have addressed risk factors for sleep problems (22-25). The results of one study indicated that alcohol consumption, inadequate physical activity, and current smoking are risk factors for the occurrence of sleep problems (26). However, these studies seem insufficient, especially in the young population. A review of previous research indicates that studies on the prevalence of sleep problems in Iran have been previously conducted (27-30). However, some of these studies are from more than a decade ago (27, 29), during which significant social changes have occurred. Issues such as the COVID-19 pandemic create the need to gather new information. A study on the adolescent population (28) highlighted the need to investigate the prevalence of sleep problems in the young population. Some other studies (29, 30) have only examined the severity of insomnia, but we have explored various dimensions of sleep problems in this study.

In recent decades, Iran has faced rapid social, economic, and demographic changes. Currently, the urban population in this country is over 70%, and young people constitute a significant part of the country's population (31). Effective planning to provide mental health services to the community requires basic information, epidemiological studies, and an understanding of the magnitude of the problem. Hence, studying the epidemiology of mental disorders can raise the awareness of health, educational, and medical officials and workers about the importance and scale of mental disorders and contribute to providing essential mental health services to citizens in the country (32) as a case for sleep problems. Thus, identifying the severity of health problems for public health planning is one of the vital requirements of the public health system. Considering the importance of the young population in the development of society and the role of sleep quality in young people's health, this study aims to investigate the quality of sleep and sleep problems among Iranian young people.

2. Methods

2.1. Study Design

This study was conducted using a descriptive cross-sectional design.

2.2. Study Population and Sample

The research population consisted of all young people aged 18 to 35 living in Kermanshah, Sanandaj, Ilam, and Hamadan (western Iran) in 2020. A total of 5000 questionnaires were distributed in these cities among people who met the inclusion criteria, of whom 4748 persons with an average age of 27.58 years were selected as

the participants in the research sample. About 60.4% of the participants were men.

The inclusion criteria were: (1) Willingness to participate in the study; (2) being in the age range of 18 to 35 years; (3) living at least five years in the western cities of Iran; and (4) having no chronic physical diseases such as diabetes, cardiovascular diseases, respiratory diseases, and thyroid problems. Besides, the exclusion criteria included having (1) non-Iranian nationality, (2) an incurable disease, (3) chronic mental disorders, especially personality disorders, and (4) an addiction to all kinds of drugs.

2.3. Procedure

A notice containing general information about the research objectives and the research procedure was posted on online social networks and channels allowed in Iran, such as WhatsApp and Instagram, from October 2020 to March 2021 to invite people to participate in this study. A total of 5452 persons who volunteered to participate in the study were identified. The questionnaires and informed consent forms were sent to 5000 persons who met the criteria for entering the study. Finally, 4748 questionnaires were returned, and the data from these questionnaires were used for analysis.

2.4. Instruments

2.4.1. Pittsburgh Sleep Quality Inventory

This 19-item instrument was developed by Buysse et al. (1989). Pittsburgh Sleep Quality Inventory (PSQI) is a standard questionnaire with 18 items scored from 0 to 21. A higher score indicates a poorer quality of sleep. A score higher than 5 indicates poor sleep quality. The reliability of the PSQI was assessed using Cronbach's alpha, and its value was 0.83 (33). A meta-analysis study showed that the inventory is a suitable tool for screening sleep problems (14). In the current study, Cronbach's alpha of this inventory was 0.91.

2.4.2. Insomnia Severity Index

This instrument was used to measure the severity of insomnia with seven items that assess the severity of sleep onset, sleep maintenance, early morning awakening problems, sleep dissatisfaction, interference of sleep difficulties with daytime functioning, and distress caused by sleep difficulties. Each item is scored from 0 to 4 depending on the severity of insomnia, and a respondent's total score equals the sum of the scores given to the seven items (34). In the current study, Cronbach's alpha for Insomnia Severity Index (ISI) was 0.92.

2.4.3. Nightmare Questionnaire

The intensity of nightmares was measured using the question: "Have you had any scary dreams and nightmares in the last six months?" The participants were asked to choose one option: Never (1), rarely (2), some nights (3), often (4), most of the nights (5), and almost every night (6). This assessment had been used in a similar study (8).

2.5. Data Analysis

The collected data were analyzed using SPSS software version 22. The mean and standard deviation were used to describe the data, and the frequencies were compared using the chi-square test. Pearson's correlation test was used to check the relationship between the variables after establishing the statistical assumptions. Moreover, a one-way analysis of variance (ANOVA) was used to compare the group's means. Logistic regression analysis was also used to predict sleep problems.

3. Results

The participants were 4748, of whom 2866 persons (60.4%) were men and 1882 persons (39.6%) were women. The participants' ages ranged from 18 to 35 years, with a mean age of 27.58 ± 5.27 . Table 1 shows the participants' demographic characteristics

3.1. Sleep Problems Reported by the Participants

Table 2 shows the descriptive statistics for the sleep quality subscales (sleeping problems in the last month) and the severity of insomnia by gender.

Table 2 indicates that the total mean score for sleep problems among the Pittsburgh Sleep Quality Inventory participants was 5.68 ± 2.67 , showing no significant difference between the male and female participants. However, this value confirmed the poor sleep quality among the participants. Furthermore, there were significant differences between the male and female participants regarding the sleep quality subscales, except for daytime dysfunction. However, insomnia was more severe in the female participants than the male participants. Table 3 shows the prevalence of sleep problems reported by the participants.

As shown, 44.4% of the participants reported poor sleep quality and suffered from sleep problems. However, there was no significant difference between the male and female participants regarding the prevalence of sleep problems. Table 3 also presents that 4.1% of the participants had moderate insomnia, and 0.9% reported severe insomnia. Moderate and severe insomnia was reported by 5.8% and 1.2% of the female participants, and

Table 1. The Participants' Demographic Characteristics

Variables	No. (%)
Sex	
Male	1882 (39.6)
Female	2866 (60.4)
Age (y)	
18 - 35	1740 (36.6)
25 - 30	1178 (24.8)
30 - 35	1830 (38.5)
Marital status	
Single	1420 (29.9)
Married	2934 (61.8)
Others	394 (8.3)
Education	
Under diploma	812 (17.1)
Diploma	1854 (39)
Bachelor's degree	1528 (32.2)
Higher	554 (11.7)
Occupation	
Unemployed	842 (17.7)
Student	1324 (27.9)
Employed	2582 (54.4)
Smoking	
Yes	770 (16.2)
No	3978 (83.8)
Alcohol use	
Yes	936 (19.7)
No	3812 (80.3)

the corresponding values were 2.9% and 0.7% for the male participants, indicating significant differences between the male and female participants with insomnia being more severe in the women.

As shown in Table 3, in the last six months, 29% of the participants have nightmares on some nights, 10.2% have a lot of nightmares, 3.1% have nightmares most nights, and 3.3% have nightmares every night. The data also confirm a significant difference between the male and female participants in the intensity of nightmares and scary dreams in the last six months, with a higher frequency of nightmares among men. Table 4 presents the prevalence of sleep complaints throughout life.

As shown in Table 4, 44% of the participants reported sleep problems such as insomnia, nightmares, hypersomnia, apnea, sleepwalking, etc. during their

Table 2. The Descriptive Statistics for Sleep Quality Components and Insomnia Severity

Domains	Total	Female	Male	F	P-Value
Subjective sleep quality	1.07 ± 0.77	1.01 ± 0.77	1.11 ± 0.78	19.96	0.001
Sleep latency	0.89 ± 0.78	1.03 ± 0.89	0.79 ± 0.70	107.64	0.001
Sleep duration	0.38 ± 0.78	0.43 ± 0.87	0.36 ± 0.72	9.91	0.002
Sleep efficiency	0.25 ± 0.74	0.40 ± 0.90	0.15 ± 0.58	135.56	0.001
Sleep disturbances	1.22 ± 0.66	1.18 ± 0.68	1.25 ± 0.64	13.32	0.001
Sleeping medication	0.82 ± 0.82	0.63 ± 0.80	0.95 ± 0.81	183.44	0.001
Daytime dysfunction	1.04 ± 0.20	1.03 ± 0.18	1.04 ± 0.21	3.01	0.08
Sleep quality (total)	5.26 ± 2.67	5.71 ± 2.95	5.66 ± 2.46	0.47	0.49
Insomnia Severity Index	6.19 ± 4.33	6.75 ± 5.82	5.82 ± 3.95	980.31	0.001

Table 3. Sleep Problems, the Severity of Insomnia, and Scary Dreams and Nightmares Reported by the Participants

Variables	Total	Female	Male	P-Value
Sleep Quality				0.40
0 - 5 (good sleeper)	2638 (55.6)	1060 (56.3)	1578 (55.1)	
6 - 21 (poor sleeper)	2110 (44.4)	822 (43.7)	1288 (44.9)	
Insomnia severity				0.001
0 - 7 (normal)	3344 (70.4)	1226 (65.1)	2118 (73.9)	
8 - 14 (subthreshold)	1168 (24.6)	524 (27.8)	644 (22.5)	
15 - 21 (moderate severity)	194 (4.1)	110 (5.8)	84 (2.9)	
22 - 28 (severe)	42 (0.9)	22 (1.2)	20 (0.7)	
Nightmare				0.001
Has not experienced	928 (19.5)	512 (27.2)	416 (14.5)	
Rarely happened	1654 (34.8)	596 (31.7)	1058 (36.9)	
Experienced some nights	1378 (29)	504 (26.8)	874 (30.5)	
Experienced a lot	482 (10.2)	162 (8.6)	320 (11.2)	
Happened most nights	148 (3.1)	40 (2.1)	108 (3.8)	
Experienced every night	158 (3.3)	68 (3.6)	90 (3.1)	

Table 4. The Prevalence of Sleep Complaints (Insomnia, Nightmares, Hypersomnia, Apnea, Sleepwalking, etc.) Throughout Life

Group	Total	Female	Male	P-Value
With sleep problem	2090 (44)	864 (45.9)	1226 (42.8)	0.02
Without sleep problem	2658 (56)	1018 (54.1)	1640 (57.2)	

lifetime. Moreover, 45.9% of the female and 42.8% of the male participants reported sleep problems, indicating a significant difference between the men and women.

3.2. Prediction of Sleep Problems

Multilevel logistic regression was used to predict sleep problems in the participants, as shown in Table 5.

The logistic regression analysis showed that the participant's age, education, marital status, occupation,

drug abuse, alcohol consumption, use of mobile phones and the internet, and a history of previous sleep problems played a role in predicting sleep problems during the COVID-19 pandemic ($P < 0.05$). The data indicated that sleep problems were more prevalent in younger people, people with lower education, single and unemployed people, people who consumed alcohol, and those who had more substance abuse. Furthermore, the results indicated those with a previous history of sleep problems were

Table 5. Multilevel Logistic Regression Results for Predicting Sleep Problems

Variables	No. (%)	B	OR (95% [CI])	P-Value
Sex				
Female	822 (43.7)	0.12	1.13 (0.99 - 1.30)	0.064
Male	1288 (44.9)	1	1	-
Age				
18 - 25	834 (47.9)	-0.22	0.80 (0.69 - 0.94)	0.005
25 - 30	480 (40.7)	0.12	1.12 (0.96 - 1.31)	0.146
30 - 35	796 (43.5)	1	1	-
Education				
Compulsory school	460 (56.7)	-0.28	0.75 (0.59 - 0.96)	0.024
High school	798 (43)	-0.09	0.92 (0.74 - 1.13)	0.431
Bachelor	640 (41.9)	-0.09	0.92 (0.74 - 1.13)	0.416
Masters and PhD	212 (38.3)	1	1	-
Marital status				
Never married	628 (44.2)	0.36	1.43 (1.11 - 1.84)	0.005
Married	1266 (43.1)	0.14	1.15 (0.911 - 1.45)	0.242
Previously married	216 (54.8)	1	1	-
Occupation				
Unemployed	488 (58)	-0.37	0.69 (0.57 - 0.83)	0.001
Student	562 (42.4)	-0.02	0.98 (0.86 - 1.14)	0.790
Employed	1060 (41.1)	1	1	-
Addiction				
Yes	422 (69.4)	-0.70	0.50 (0.36 - 0.68)	0.001
No	1562 (40.4)	1	1	-
Smoking				
Yes	458 (59.5)	-0.19	0.83 (0.69 - 1)	0.053
No	1652 (41.5)	1	1	-
Alcohol use				
Yes	536 (57.3)	-0.25	0.78 (0.66 - 0.92)	0.003
No	1574 (41.3)	1	1	-
Cell phone				
Never	66 (45.8)	0.54	1.72 (1.18 - 2.49)	0.004
Some times	352 (37.4)	0.55	1.73 (1.45 - 2.06)	0.001
Most nights	890 (41.6)	0.35	1.41 (1.22 - 1.64)	0.001
Every night	802 (52.6)	1	1	-
Sleep problem (in past)				
Yes	1132 (54.2)	-0.54	0.58 (0.52 - 0.66)	0.001
No	978 (36.8)	1	1	-

Abbreviations: B, unstandardized coefficient; OR, odds ratio.

probably more involved in sleep problems.

4. Discussion

Young people experienced poor sleep quality during the COVID-19 pandemic. Specifically, 44.4% of the

participants reported unfavorable sleep quality and encountered sleep-related problems. However, there was no significant difference in the prevalence of sleep problems between women and men. Sleep disorders are common in the general adult population and are associated with various adverse effects (35). One study categorized sleep problems in adults into three groups: Difficulty falling asleep, behavioral and movement disorders during sleep, and excessive sleepiness during the day (35). Sleep problems are widespread, affecting 9 - 20% of adults in the United States and 7% in Europe (36). These results show a significant difference in the prevalence of sleep problems in the present study, which was estimated to be higher than 44%.

A study conducted on young people aged 17 to 24 years (37) showed that sleep problems are highly prevalent in this population. The prevalence of sleep problems measured by the PSQI was higher than 60%. The results also indicated that sleep and wake times were delayed on weekends, and participants frequently used prescribed, over-the-counter, and psychoactive drugs to alter sleep/wakefulness (37). The prevalence of sleep problems reported in this study (37) was significantly higher than the values estimated in the present study. Another study reported the frequency of sleep problems was about 7% in people aged 19 to 33 years (38). Another study reported the prevalence of sleep problems to be 22.3% (39). These conflicting findings can be attributed to sociocultural and economic issues, differences in the research sample, and time. However, studies conducted during the COVID-19 pandemic also reported different rates of sleep problems. For example, a study showed that the prevalence of sleep problems was about 30% among the young population during the COVID-19 pandemic (40), which was lower than the prevalence rate reported in the present study.

The present study showed that 4.1% of the participants had moderate insomnia, and 0.9% had severe insomnia, which was 34.9% in another study (41). A further study showed that almost 41% of the young population suffered from severe insomnia during the COVID-19 pandemic (42).

The data in this study revealed that 29% of the participants had nightmares some nights in the last six months, 10.2% had a lot of nightmares in the last six months, 3.1% had nightmares most nights, and 3.3% of them had nightmares every night. The data also confirmed a significant difference between the male and female participants in the intensity of nightmares and scary dreams in the last six months, with over 45% of the participants having nightmares in the previous six months. A study (43) showed that 45.2% of the participants reported nightmares at least once in the last

month, and 7.9% had nightmares at least once a week. However, contrary to the present study, girls reported more nightmares than boys (43). A few studies have addressed the prevalence of nightmares in young people, so it is difficult to draw general conclusions.

The age below 25 years, lower education levels, being single, unemployment, drug abuse, alcohol consumption, excessive use of mobile phones and the internet, and having a history of previous sleep problems accounted for sleep problems among young people during the COVID-19 pandemic. However, another study addressing the factors related to sleep problems among young people during the COVID-19 pandemic showed that factors such as age over 24 years, higher education, and negative attitudes towards COVID-19 could predict sleep problems (44). Another study revealed that exercise, alcohol, and caffeine consumption and stability of sleep schedule are not significant predictors of sleep quality in young people (37). These results were not consistent with the findings of the present study. However, one study showed that alcohol consumption is related to sleep problems (45), and another study reported that smoking is related to sleep problems, such as maintaining sleep, breathing problems during sleep, and difficulty in sleep onset (46), as was confirmed in the present study.

The results of the present study concerning the role of using mobile phones and surfing the internet (before sleep) in sleep problems are contradictory because the prevalence of sleep problems in people who have never used a mobile phone an hour before sleep was higher than those who used mobile phones "some nights" and "most nights." However, sleep problems were less prevalent in these people than those who used mobile phones every night. Therefore, drawing solid conclusions about the role of using mobile phones in sleep problems is not possible. Nevertheless, contrary to previous studies, those who used mobile phones every night reported more sleep problems. A review study showed that screen time has a negative relationship with sleep outcomes (mainly a shorter sleep time and sleep latency) in 90% of studies (3). Another study found that dependence on mobile phones is related to poor subjective sleep quality and latency (47). These results have been confirmed in another study (48). Thus, further research is needed to explore this issue.

4.1. Conclusions

Based on the results, there was a high prevalence of sleep problems among Iranian young people during the COVID-19 pandemic, but there was no significant difference between men and women regarding the prevalence of sleep problems. The data in this study showed that age below 25 years, lower education levels,

being single, unemployment, drug abuse, alcohol consumption, excessive use of mobile phones and the internet, and having a history of previous sleep problems can account for sleep problems among young people during the COVID-19 pandemic.

4.2. Suggestions

(1) This study was conducted using a cross-sectional design, and the data were collected through self-report instruments, which could affect behaviors such as sleep. Thus, future studies need to use other instruments.

(2) The present study was conducted on people living in urban areas in western Iran. Thus, we need to be cautious in generalizing the results to other groups and populations. Furthermore, these variables should be addressed in other societies as well.

(3) The participants in this study were selected from people living in urban areas. Thus, it is also necessary to conduct similar studies in rural populations.

(4) Following the present study's findings, effective planning is needed to improve sleep quality and reduce sleep problems among young people.

Acknowledgments

The authors would like to thank those who contributed to this research project, especially the Deputy Office of Research and Technology staff of Kermanshah University of Medical Sciences and the Sleep Disorders Research Center.

Footnotes

Authors' Contribution: Ali Zakiei and Habibollah Khazaie designed the study. Leeba Rezaie and Saeid Komasi collected the data. Masoumeh Rostampour and Ali Zakiei processed the data and wrote the manuscript. Ali Zakiei and Habibollah Khazaie helped with statistical analysis and revised the manuscript. All authors reviewed the paper and approved the final version to be published.

Conflict of Interests: The researchers declared that they had no conflict of interest in this study.

Data Availability: Data used and analyzed in this study are available on request.

Ethical Approval: To comply with the protocol for research ethics, the participants were assured their data would not be made available to individuals or public or private organizations, and the information they provided would be used only in a research study. The participants were also told that their participation was voluntary and they could leave the study if they

wished. The Medical Research and Ethical Committee of Kermanshah University of Medical Sciences (Kermanshah, Iran; registration No. [IR.KUMS.REC.1399.125](#)) approved the study, which was performed under the ethical principles laid down in the seventh and current edition (2013) of the Declaration of Helsinki.

Funding/Support: This research project was financed by the Deputy Office of Research and Technology of Kermanshah University of Medical Sciences (grant number: 990085).

Informed Consent: The selected people participated in the survey after declaring their consent and receiving the assurance that their information would remain confidential. Informed consent was obtained from all individual participants included in the study.

References

1. Kelley GA, Kelley KS. Exercise and sleep: a systematic review of previous meta-analyses. *J Evid Based Med*. 2017;**10**(1):26–36. [PubMed ID: [28276627](#)]. [PubMed Central ID: [PMC5527334](#)]. <https://doi.org/10.1111/jebm.12236>.
2. Gradisar M, Gardner G, Dohnt H. Recent worldwide sleep patterns and problems during adolescence: a review and meta-analysis of age, region, and sleep. *Sleep Med*. 2011;**12**(2):110–8. [PubMed ID: [21257344](#)]. <https://doi.org/10.1016/j.sleep.2010.11.008>.
3. Hale L, Guan S. Screen time and sleep among school-aged children and adolescents: a systematic literature review. *Sleep Med Rev*. 2015;**21**:50–8. [PubMed ID: [25193149](#)]. [PubMed Central ID: [PMC4437561](#)]. <https://doi.org/10.1016/j.smrv.2014.07.007>.
4. Schreck KA, Richdale AL. Sleep problems, behavior, and psychopathology in autism: inter-relationships across the lifespan. *Curr Opin Psychol*. 2020;**34**:105–11. [PubMed ID: [31918238](#)]. <https://doi.org/10.1016/j.copsyc.2019.12.003>.
5. Stewart CA, Auger RR, Enders FT, Felmlee-Devine D, Smith GE. The effects of poor sleep quality on cognitive function of patients with cirrhosis. *J Clin Sleep Med*. 2014;**10**(1):21–6. [PubMed ID: [24426816](#)]. [PubMed Central ID: [PMC3869064](#)]. <https://doi.org/10.5664/jcsm.3350>.
6. Stefani A, Hogl B. Nightmare Disorder and Isolated Sleep Paralysis. *Neurotherapeutics*. 2021;**18**(1):100–6. [PubMed ID: [33230689](#)]. [PubMed Central ID: [PMC8116464](#)]. <https://doi.org/10.1007/s13311-020-00966-8>.
7. Geng F, Liang Y, Li Y, Fang Y, Pham TS, Liu X, et al. Bidirectional associations between insomnia, posttraumatic stress disorder, and depressive symptoms among adolescent earthquake survivors: a longitudinal multiwave cohort study. *Sleep*. 2019;**42**(11). [PubMed ID: [31328781](#)]. <https://doi.org/10.1093/sleep/zsz162>.
8. Khazaie H, Khazaie S, Zakiei A, Dursteler KM, Bruhl AB, Brand S, et al. When Non-Suicidal Self-Injury Predicts Non-Suicidal Self-Injury and Poor Sleep-Results from a Larger Cross-Sectional and Quasi-Longitudinal Study. *Int J Environ Res Public Health*. 2021;**18**(24). [PubMed ID: [34948620](#)]. [PubMed Central ID: [PMC8701371](#)]. <https://doi.org/10.3390/ijerph182413011>.
9. Pigeon WR, Pinquart M, Conner K. Meta-analysis of sleep disturbance and suicidal thoughts and behaviors. *J Clin Psychiatry*. 2012;**73**(9):e1160–7. [PubMed ID: [23059158](#)]. <https://doi.org/10.4088/JCP.11r07586>.
10. Javaheri S, Redline S. Insomnia and Risk of Cardiovascular Disease. *Chest*. 2017;**152**(2):435–44. [PubMed ID: [28153671](#)]. [PubMed Central ID: [PMC5577359](#)]. <https://doi.org/10.1016/j.chest.2017.01.026>.

11. Espie CA, Emsley R, Kyle SD, Gordon C, Drake CL, Siriwardena AN, et al. Effect of Digital Cognitive Behavioral Therapy for Insomnia on Health, Psychological Well-being, and Sleep-Related Quality of Life: A Randomized Clinical Trial. *JAMA Psychiatry*. 2019;**76**(1):21-30. [PubMed ID: 30264137]. [PubMed Central ID: PMC6583463]. <https://doi.org/10.1001/jamapsychiatry.2018.2745>.
12. Chaput JP, McHill AW, Cox RC, Broussard JL, Dutil C, da Costa BGG, et al. The role of insufficient sleep and circadian misalignment in obesity. *Nat Rev Endocrinol*. 2023;**19**(2):82-97. [PubMed ID: 36280789]. [PubMed Central ID: PMC9590398]. <https://doi.org/10.1038/s41574-022-00747-7>.
13. Irwin MR, Olmstead R, Carroll JE. Sleep Disturbance, Sleep Duration, and Inflammation: A Systematic Review and Meta-Analysis of Cohort Studies and Experimental Sleep Deprivation. *Biol Psychiatry*. 2016;**80**(1):40-52. [PubMed ID: 26140821]. [PubMed Central ID: PMC4666828]. <https://doi.org/10.1016/j.biopsych.2015.05.014>.
14. Mollaveya T, Thuraiajah P, Burton K, Mollaveya S, Shapiro CM, Colantonio A. The Pittsburgh sleep quality index as a screening tool for sleep dysfunction in clinical and non-clinical samples: A systematic review and meta-analysis. *Sleep Med Rev*. 2016;**25**:52-73. [PubMed ID: 26163057]. <https://doi.org/10.1016/j.smrv.2015.01.009>.
15. Fabbri M, Beracci A, Martoni M, Meneo D, Tonetti L, Natale V. Measuring Subjective Sleep Quality: A Review. *Int J Environ Res Public Health*. 2021;**18**(3). [PubMed ID: 33530453]. [PubMed Central ID: PMC7908437]. <https://doi.org/10.3390/ijerph18031082>.
16. Zakiei A, Korani D, Sahraei Z, Rostampour M, Khazaie H. Predicting sleep quality and insomnia severity using the components of the acceptance and commitment therapy (ACT) model: A new perspective. *J Context Behav Sci*. 2022;**26**:227-33. <https://doi.org/10.1016/j.jcbs.2022.10.006>.
17. Luciano F, Cenacchi V, Vegro V, Pavei G. COVID-19 lockdown: Physical activity, sedentary behaviour and sleep in Italian medicine students. *Eur J Sport Sci*. 2021;**21**(10):1459-68. [PubMed ID: 33108970]. <https://doi.org/10.1080/17461391.2020.1842910>.
18. Xiao H, Zhang Y, Kong D, Li S, Yang N. The Effects of Social Support on Sleep Quality of Medical Staff Treating Patients with Coronavirus Disease 2019 (COVID-19) in January and February 2020 in China. *Med Sci Monit*. 2020;**26**:e923549. <https://doi.org/10.12659/msm.923549>.
19. Eleftheriou A, Rokou A, Arvaniti A, Nena E, Steiropoulos P. Sleep Quality and Mental Health of Medical Students in Greece During the COVID-19 Pandemic. *Front Public Health*. 2021;**9**:775374. [PubMed ID: 34869189]. [PubMed Central ID: PMC8639533]. <https://doi.org/10.3389/fpubh.2021.775374>.
20. Wang D, Zhao J, Zhai S, Ye H, Bu L, Fan F. Does sleep disturbance predicts posttraumatic stress disorder and depression among college students during COVID-19 lockdown? A longitudinal survey. *Front Public Health*. 2022;**10**:986934. [PubMed ID: 36176529]. [PubMed Central ID: PMC9514232]. <https://doi.org/10.3389/fpubh.2022.986934>.
21. Wang D, Ross B, Zhou X, Meng D, Zhu Z, Zhao J, et al. Sleep disturbance predicts suicidal ideation during COVID-19 pandemic: A two-wave longitudinal survey. *J Psychiatr Res*. 2021;**143**:350-6. [PubMed ID: 34571319]. [PubMed Central ID: PMC8460177]. <https://doi.org/10.1016/j.jpsychires.2021.09.038>.
22. Fang H, Tu S, Sheng J, Shao A. Depression in sleep disturbance: A review on a bidirectional relationship, mechanisms and treatment. *J Cell Mol Med*. 2019;**23**(4):2324-32. [PubMed ID: 30734486]. [PubMed Central ID: PMC6433686]. <https://doi.org/10.1111/jcmm.14170>.
23. Alonso-Alvarez ML, Cordero-Guevara JA, Teran-Santos J, Gonzalez-Martinez M, Jurado-Luque MJ, Corral-Penafiel J, et al. Obstructive sleep apnea in obese community-dwelling children: the NANOS study. *Sleep*. 2014;**37**(5):943-9. [PubMed ID: 24790273]. [PubMed Central ID: PMC3985101]. <https://doi.org/10.5665/sleep.3666>.
24. de Zambotti M, Goldstone A, Colrain IM, Baker FC. Insomnia disorder in adolescence: Diagnosis, impact, and treatment. *Sleep Med Rev*. 2018;**39**:12-24. [PubMed ID: 28974427]. [PubMed Central ID: PMC5931364]. <https://doi.org/10.1016/j.smrv.2017.06.009>.
25. Lin CY, Su SB, Hu YT, Peng CJ, Chen KH, Chen KT. Prevalence and risk factors associated with poor sleep among firefighters in Taiwan: An observational study. *Medicine (Baltimore)*. 2022;**101**(44):e31570. [PubMed ID: 36343071]. [PubMed Central ID: PMC9646591]. <https://doi.org/10.1097/MD.00000000000031570>.
26. Bedford LE, Tang EHM, Dong W, Wong CKH, Tse ETY, Ng APP, et al. Who reports insufficient and disturbed sleep? Results from a representative population-based health survey in Hong Kong. *BMJ Open*. 2022;**12**(9):e058169. [PubMed ID: 36115682]. [PubMed Central ID: PMC9486237]. <https://doi.org/10.1136/bmjopen-2021-058169>.
27. Malakouti SK, Foroughan M, Nojomi M, Ghalebandi MF, Zandi T. Sleep patterns, sleep disturbances and sleepiness in retired Iranian elders. *Int J Geriatr Psychiatry*. 2009;**24**(11):1201-8. [PubMed ID: 19343736]. <https://doi.org/10.1002/gps.2246>.
28. Shamsadini A, Amizadeh M, Motamed S, Rajabi F. Epidemiology of Sleep Disorders Among Adolescents in Kerman, Iran, in 2019. *Soc Work Public Health*. 2022;**37**(8):775-80. [PubMed ID: 35695466]. <https://doi.org/10.1080/19371918.2022.2089310>.
29. Yazdi Z, Sadeghniai-Haghighi K, Zohal MA, Elmizadeh K. Validity and reliability of the Iranian version of the insomnia severity index. *Malays J Med Sci*. 2012;**19**(4):31-6. [PubMed ID: 23613647]. [PubMed Central ID: PMC3629678].
30. Khojasteh-Kaffash S, Fanoodi A, Mozayani F, Dastjerdi R, Riahi SM. Health Status of Iranian Healthcare Workers During the First Peak of COVID-19 Epidemic in a High-Incidence Area (South Khorasan Province, Iran). *Curr Res Med Sci*. 2022;**6**(2):11-22.
31. Statistical Centre of Iran. *Iran Statistical Yearbook*. Tehran, Iran: Statistical Centre of Iran; 2015, [cited 2023].
32. Khazaie H, Hamzeh B, Najafi F, Chehri A, Rahimi-Movaghar A, Amin-Esmaili M, et al. Prevalence of Psychiatric Disorders and Associated Factors among the Youth in Ravansar, Iran. *Arch Iran Med*. 2019;**22**(8):435-42. [PubMed ID: 31679346].
33. Davoodi I, Neisi A, Zakiei A, Khazaei Arshadi. [The Roles of Some Cognitive and Emotional Factors in Predicting Insomnia]. *J Maz Univ Med Sci*. 2017;**27**(150):107-21. Persian.
34. Bastien CH, Vallieres A, Morin CM. Validation of the Insomnia Severity Index as an outcome measure for insomnia research. *Sleep Med*. 2001;**2**(4):297-307. [PubMed ID: 11438246]. [https://doi.org/10.1016/s1389-9457\(00\)00065-4](https://doi.org/10.1016/s1389-9457(00)00065-4).
35. Holder S, Narula NS. Common Sleep Disorders in Adults: Diagnosis and Management. *Am Fam Physician*. 2022;**105**(4):397-405. [PubMed ID: 35426627].
36. Shinjyo N, Waddell G, Green J. Valerian Root in Treating Sleep Problems and Associated Disorders-A Systematic Review and Meta-Analysis. *J Evid Based Integr Med*. 2020;**25**:2515690X20967323. [PubMed ID: 33086877]. [PubMed Central ID: PMC7585905]. <https://doi.org/10.1177/2515690X20967323>.
37. Lund HG, Reider BD, Whiting AB, Prichard JR. Sleep patterns and predictors of disturbed sleep in a large population of college students. *J Adolesc Health*. 2010;**46**(2):124-32. [PubMed ID: 20113918]. <https://doi.org/10.1016/j.jadohealth.2009.06.016>.
38. Veldi M, Aluoja A, Vasar V. Sleep quality and more common sleep-related problems in medical students. *Sleep Med*. 2005;**6**(3):269-75. [PubMed ID: 15854858]. <https://doi.org/10.1016/j.sleep.2004.12.003>.
39. Angelone AM, Mattei A, Sbarbati M, Di Orio F. Prevalence and correlates for self-reported sleep problems among nursing students. *J Prev Med Hyg*. 2011;**52**(4):201-8. [PubMed ID: 22442926].
40. Shrestha D, Adhikari SP, Rawal N, Budhathoki P, Pokharel S, Adhikari Y, et al. Sleep quality among undergraduate students of a medical college in Nepal during COVID-19 pandemic: an online survey. *F1000Res*. 2021;**10**:505. [PubMed ID: 34249344]. [PubMed Central ID: PMC8261762]. <https://doi.org/10.12688/f1000research.53904.2>.
41. Alrashed FA, Sattar K, Ahmad T, Akram A, Karim SI, Alsubiheeh AM. Prevalence of insomnia and related psychological factors with coping strategies among medical students in clinical years

- during the COVID-19 pandemic. *Saudi J Biol Sci.* 2021;**28**(11):6508-14. [PubMed ID: 34764766]. [PubMed Central ID: PMC8568835]. <https://doi.org/10.1016/j.sjbs.2021.07.022>.
42. AlHadi AN, Alhuwaydi AM. Insomnia Prevalence and Associated Factors Among University Students in Saudi Arabia During the COVID-19 Pandemic and Lockdown: A Large-Scale Survey. *Nat Sci Sleep.* 2022;**14**:1651-63. [PubMed ID: 36164410]. [PubMed Central ID: PMC9507980]. <https://doi.org/10.2147/NSS.S380972>.
43. Liu X, Liu ZZ, Liu BP, Jia CX. Nightmare frequency and psychopathological problems in a large sample of Chinese adolescents. *Soc Psychiatry Psychiatr Epidemiol.* 2022;**57**(4):805-16. [PubMed ID: 35064282]. <https://doi.org/10.1007/s00127-022-02224-y>.
44. Yang Y, Zhu JF, Yang SY, Lin HJ, Chen Y, Zhao Q, et al. Prevalence and associated factors of poor sleep quality among Chinese returning workers during the COVID-19 pandemic. *Sleep Med.* 2020;**73**:47-52. [PubMed ID: 32771927]. [PubMed Central ID: PMC7358171]. <https://doi.org/10.1016/j.sleep.2020.06.034>.
45. Huang R, Ho SY, Lo WS, Lai HK, Lam TH. Alcohol consumption and sleep problems in Hong Kong adolescents. *Sleep Med.* 2013;**14**(9):877-82. [PubMed ID: 23777753]. <https://doi.org/10.1016/j.sleep.2013.03.022>.
46. Mak KK, Ho SY, Thomas GN, Lo WS, Cheuk DK, Lai YK, et al. Smoking and sleep disorders in Chinese adolescents. *Sleep Med.* 2010;**11**(3):268-73. [PubMed ID: 20176504]. <https://doi.org/10.1016/j.sleep.2009.07.017>.
47. Ibrahim NK, Baharoon BS, Banjar WF, Jar AA, Ashor RM, Aman AA, et al. Mobile Phone Addiction and Its Relationship to Sleep Quality and Academic Achievement of Medical Students at King Abdulaziz University, Jeddah, Saudi Arabia. *J Res Health Sci.* 2018;**18**(3). e00420. [PubMed ID: 30270211]. [PubMed Central ID: PMC6941644].
48. Amra B, Shahsavari A, Shayan-Moghadam R, Mirheli O, Moradi-Khaniabadi B, Bazukar M, et al. The association of sleep and late-night cell phone use among adolescents. *J Pediatr (Rio J).* 2017;**93**(6):560-7. [PubMed ID: 28257717]. <https://doi.org/10.1016/j.jped.2016.12.004>.