Published online: 2024 July 1.

**Research Article** 



# Assessment of Occupational Hazards and Health Status (Respiratory and Non-respiratory) Among Sawmillers in Akure Metropolis, Nigeria

Sunday Olakunle Olarewaju (1)<sup>1</sup>, Adewale Allen Sokan-Adeaga (1)<sup>2,\*</sup>, Akinlolu Oluwafemi Fansanmi (1)<sup>3</sup>, Olufunke Julianah Ogidan (1)<sup>4</sup>, Micheal Ayodeji Sokan-Adeaga (1)<sup>5</sup>, Joy Stephen Amusan (1)<sup>6</sup>

<sup>1</sup> Department of Community Medicine, Faculty of Clinical Sciences, College of Health Sciences, Osun State University, Osogbo, Nigeria

<sup>2</sup> Department of Environmental Health Science, Faculty of Basic Medical Sciences, Ajayi Crowther University, Oyo, Nigeria

<sup>3</sup> The Aurum Institute, Parktown, Johanesburg, South Africa

<sup>4</sup> School of Postgraduate Studies, Ladoke Akintola University of Technology (LAUTECH), Ogbomosho, Nigeria

<sup>5</sup> Department of Community Health and Primary Health Care, Faculty of Clinical Sciences, College of Medicine, University of Lagos, Lagos, Nigeria
 <sup>6</sup> Department of Public Health, Faculty of Basic Medial Sciences, College of Health Sciences, Osun State University, Osogbo, Nigeria

\* Corresponding author: Department of Environmental Health Science, Faculty of Basic Medical Sciences, Ajayi Crowther University, Oyo, Nigeria. Email: sokanadeaga.adewaleallen@yahoo.com

Received 2023 December 23; Accepted 2024 May 14.

#### Abstract

**Background:** Sawmilling exposed workers to diverse occupational mishaps and subsequent health complications. Investigating workplace risk and health status can help mitigate morbidness and death among this group.

**Objectives:** This study aimed to evaluate the occupational hazards and health status of sawmill workers in Akure, Nigeria. **Methods:** This cross-sectional descriptive survey was conducted on 304 sawmill workers in Akure metropolis selected via multistage sampling methodology. Data were collected using a pre-tested, standardized questionnaire. In addition, the body mass index (BMI), blood pressure (mmHg), and respiratory function parameters (spirometry) were measured. Descriptive and inferential statistics (chi-square) were used to evaluate and summarize the data, with a significance level of P < 0.05.

**Results:** The respondents' average age was  $42.0 \pm 2.8$  years. The common reported workplace hazards included noise 273 (89.8%), manual lifting of objects 221 (72.7%), heat 192 (63.2%), and wood dust 192 (63.2%), while common reported injuries were bruises 292 (96.1%), hearing impairment 281 (92.4%), electric shock 250 (82.2%), and fainting attack 232 (76.3%). The common prevalent ailments suffered by respondents in the last month were cough 158 (52.0%), phlegm production 149 (49.0%), chest pain 63 (20.7%), skin problems 100 (32.9%), and eyes irritation 111 (36.5%). The anthropometric measurement showed that most 272 (89.5%) sawmill workers are overweight and obese. Spirometry measurements revealed that one-third of the respondents had a forced expiratory ratio (FER) measurement below 70.00% and an abnormal respiratory rate. The bivariate analysis showed a significant correlation between the respondents' self-reported prior workplace injury, knowledge of workplace dangers, and sociodemographic characteristics (job category and educational level).

**Conclusions:** The results showed that the respondents suffered from impaired respiratory function related to workplace hazard exposure and the non-usage of protective devices during operational activities. In addition, the majority are at risk of developing diverse cardiovascular and respiratory diseases due to workplace exposures.

Keywords: Occupational Hazards, Sawmill Workers, Respiratory Function Defects, Health Problems, Injuries, Safety Practices, Nigeria

#### 1. Background

Sawmilling is one of the most traditional occupations in Nigeria (1). The primary wood products produced, used, and marketed in Nigeria are plywood, sawn wood, newsprint, particle board, and paper materials (2). A variety of human actions in sawmilling

processes accentuates worker exposure to higher levels of risk (3). Noise pollution and unfavorable weather are physical threats, while mechanical hazards like being struck by or entangled in machinery and chemical hazards such as breathing in wood dust and chlorophenols are chemical threats (4). Cancer and respiratory problems are two negative effects associated

Copyright © 2024, Journal of Kermanshah University of Medical Sciences. This open-access article is available under the Creative Commons Attribution-NonCommercial 4.0 (CC BY-NC 4.0) International License (https://creativecommons.org/licenses/by-nc/4.0/), which allows for the copying and redistribution of the material only for noncommercial purposes, provided that the original work is properly cited. with wood dust exposure. Wood dust's health effects include dermatitis, eye irritation, nose and throat irritation, and pulmonary system harm (5-7). Besides being stifling and restricted, the ambient air that sawmill workers breathed during work is full of particulates, making breathing difficult (8). According to several authors (9-11), exposure to wood dust from various wood species causes respiratory problems. In Ile-Ife, Nigeria, workers exposed to wood dust had a higher prevalence of respiratory symptoms (chest pain, sputum production), conjunctivitis, hearing loss, and skin irritation than the general population. Ige and Onadeko (12) discovered that employees exposed to wood dust have higher respiratory symptoms, eye and nose irritations, and skin inflammation rates than the control group in a study conducted in Oyo State, Nigeria.

Verifiable reports have shown that sawmillers in third-world countries are still highly susceptible to occupational dangers, even if some professions have created cutting-edge facilities to lessen workers' vulnerability to taxing activities (13). The Akure metropolis's sawmill industry consists of several modestly sized, privately owned sawmills without occupational health and safety regulations. There is no published data or documented information on the safety procedures and occupational risks/injuries among these workers.

#### 2. Objectives

This study aims to assess employees' awareness of safety protocols and possible workplace risks and injuries to pinpoint the health issues (both respiratory and non-respiratory) that these workers experience.

#### 3. Methods

#### 3.1. Population

The cross-sectional study was conducted on saw millers in Akure metropolis, Akure Nigeria, who were from different marketing points: Ajikowo (85), Agagu Road (75), Ondo Road (135), Ondo Road area (125), Igbatoro Road (115), Oda Road (120), Cultural Center road (95), and Oba Ile road (50). Among the 800 saw millers, 304 participants were selected using multistage sampling. The exclusion criteria were passers-by and buyers of wood.

#### 3.2. Ethical Considerations

The Health Research and Ethics Committee of the College of Health Sciences, Osun State University, Osogbo, Nigeria, accepted the protocol and granted ethical clearance for the study, which the Declaration of Helsinki conducted. In addition, all willing participants verbally agreed to participate in the study after being informed about the research. Those who gave their consent were interviewed after signing or thumbprinting in the case of illiterate participants. The respondents' confidentiality and privacy were strictly respected.

#### 3.3. Research Instrument

The data were collected using a pre-tested with a Cronbach alpha score of 0.89 semi-structured questionnaires, divided into four segments viz: Section A – respondents' socio-demographic characteristics. Section B – occupational history and exposure among respondents. Section C – respondents' knowledge of safety/protective devices and health hazards/injuries in the sawmilling industry. Section D deals with respondents' occupational health problems in the last month.

#### 3.4. Anthropometric Measurements

The anthropometric measurements were taken using a Seca electronic bathroom weighing scale measuring weight in kilograms (kg) and a customized stadiometer measuring height. The weighing scale was standardized using weights every morning before measurements were taken, and the stadiometer was calibrated using a GPM anthropometer. The body mass index (BMI) was computed as body weight in kilograms divided by height in meters squared (kg/m<sup>2</sup>). An inelastic dressmaker's tape was used to measure waist and hip circumferences.

#### 3.5. Blood Pressure and Pulse Rate Measurement

The blood pressure (BP) was measured in millimeters of mercury (mmHg) and reported to the nearest 2 mmHg using a validated automated blood pressure monitor (Omron M6 Comfort; Omron Corporation). This equipment automatically displays the participants' pulse rate (PR) during the BP measurement process.

#### 3.6. Respiratory Status Assessment

A standard field spirometer (SpirobankG MIR Model number: SNA23-048.06769), calibrated from 0.10 to 9.99 L and measured to the nearest 0.01 L, was employed to calculate the forced vital capacity (FVC) and forced

expiratory volume (FEV<sub>1</sub>) in the first second. The forced expiratory ratio (FER), a calculated ratio expressed in %, was derived by dividing each participant's FEV<sub>1</sub> by their FVC to determine the possible presence of a restrictive or obstructive lung disease. The instrument was precalibrated and programmed to measure FEV<sub>1</sub> and FVC before usage.

#### 3.7. Data Management and Analysis

The surveys were carefully reviewed for accuracy following data collection. Version 23 of IBM statistical product and service solutions was used to analyze the collected data manually. Frequency tables, percentages, charts, and descriptive statistics (mean and standard deviation) were used to summarize the analyzed data. Inferential statistics were utilized for the bivariate analysis of the gathered data, namely chi-square ( $\chi^2$ ). One-tailed P < 0.05 was used as the threshold for statistical significance for all inferential analyses.

#### 4. Results

Following administration, completion, retrieval, and analysis of all 304 surveys, a 100% response rate was obtained.

#### 4.1. Socio-demographic Characteristics of Respondents

The average age of the respondents was  $42.0 \pm 2.8$  years, with the bulk being male -262 (86.2%) and falling within the 40 - 59 age range. The majority 275 (90.5%) of the respondents were married, and 159 (52.3%) were Christians. Of the interviewees, 62 (20.4%) had postsecondary education, and 117 (45.1%) had only completed secondary school. Nearly 300 (98.7%) respondents were from Akure South Local Government Area of Ondo State, Nigeria. Table 1.

Table 1. Respondents' Socio-demographic Cl	haracteristics
Variables	No. (%)
Gender	
Male	262 (86.2)
Female	42 (13.8)
Total	304 (100.0)
Age group (y)	
20 - 39	113 (37.2)
40 - 59	181 (59.5)
60 and above	10 (3.3)
Total	304 (100.0)
Marital status	
Single	20 (6.6)

Variables	No. (%)			
Married	275 (90.5)			
Separated/divorced	7 (2.3)			
Widowed	2 (0.7)			
Total	304 (100.0)			
Religion				
Christianity	159 (52.3)			
Islam	145 (47.7)			
Total	304 (100.0)			
Local govt area				
Afijio LG Oyo State	1(0.3)			
Akure South	300 (98.7)			
Akoko South West	1(0.3)			
Ado Ekiti	2 (0.7)			
Total	304 (100.0)			
Level of education				
Primary and below	105 (34.5)			
Secondary education	137 (45.1)			
Tertiary education	62 (20.4)			
Total	304 (100.0)			

#### 4.2. Occupational History and Exposure of Respondents

In Table 2, barely more than half, 159 (52.3%) sawmillers worked more than 9 hours a day, and the majority, 296 (97.4%), worked more than five days a week. Most 237 (78%) of the respondents are involved in operation activities. The majority, 194 (63.8%) of the study participants received official tutelage for the job, and 296 (78.3%) were not engaged in any other time of job. About 175 (57.6%) of the respondents were currently smoking, and 159 (52.3%) were passive smokers. The following proportions of the respondents were exposed to various sources of smoke, including wood fire smoke – 99 (32.6%), coal fire smoke – 56 (18.4%), sawdust smoke – 286 (94.1%), and kerosene stove smoke – 294 (96.7%).

W	N - (0/)
Variables	No.(%)
Working hours per day	
Less than 9 hours	145 (47.7)
Above 9 hours	159 (52.3)
Total	304 (100.0)
Days worked per week	
Less than five days	8 (2.6)
Above five days	296 (97.4)
Total	304 (100.0)
Category of Job	
Operation aspect	237 (78.0)
Administrative aspect	67 (22.0)
Total	304 (100.0)
Were you formally trained for this job you are do	ing?
Yes	194 (63.8)

Variables	No. (%)
No	110 (32)
Total	304 (100.0)
Have you ever done any type of job before?	
Yes	66 (21.7)
No	238 (78.3)
Total	304 (100.0)
Are you presently engaged in any other (part-time) job apart from this one you are doing?	
Yes	8 (2.6)
No	296 (97.4)
Total	304 (100.0)
Are you currently smoking?	
Yes	175 (57.6)
No	129 (42.4)

Total	304 (100.0)
Are you a past smoker?	
Vec	28 (12 5)

Yes	38 (12.5)			
No	266 (87.5)			
Total	304 (100.0)			
Are you a passive smoker?				
Yes	159 (52.3)			
No	145 (47.7)			
Total	304 (100.0)			
Are you exposed to wood dust smoke at work?				

Yes	99 (32.6)
No	205 (67.4
Total	304 (100.0)
Are you exposed to coal fire smoke?	
Yes	56 (18.4)
No	248 (81.6
Total	304 (100.0)
Are you exposed to sawdust smoke?	
Yes	286 (94.1
No	18 (5.9)
Total	304 (100.0)
Are you exposed to kerosene stove smoke?	

#### Yes 294 (96.7) No 10 (3.3) 304 Total (100.0)

### 4.3. Knowledge of Safety Practices and Occupational Hazards/Injuries Among Respondents

As shown in Table 3, 100 (32.9%) of the respondents have attended training on safety before, and the

majority claimed that their training source is from the Ministry of Health. The majority, 221 (72.7%), can identify personal protective equipment (PPE), but only 104 (34.5%) use it. Reasons given by respondents for nonusage of PPE include non-availability 48 (24.1%), not necessary 57 (28.6%), inconvenient 5 (2.5%), forgetfulness 6 (3.1%), and other reason 83 (41.7%). Some of the common workplace hazards reported by respondents were noise 273 (89.8%), animal/insect bite 264 (86.8%), manual lifting of objects 221 (72.7%), heat 192 (63.2%), and wood dust and smoke 192 (63.2%). The respondents reported the following as some of the common injuries in the workplace: Bruises 292 (96.1%), lacerations 281 (92.4%), hearing impairment 281 (92.4%), fracture 262 (86.2%), electric shock 250 (82.2%), fainting attack 232 (76.3%), and burns 229 (75.3%). Some of the common respiratory and non-respiratory problems experienced in the workplace reported by the respondents were cough 285 (93.8%), phlegm production 288 (94.7%), chest pain 280 (92.1%), dyspnoea 249 (81.9%), skin problem 268 (88.2%), and red eye 271 (89.1%). According to Figure 1, 172 (57.0%) respondents have good erudition of workplace safety procedures and health risks/injuries, while 132 (44.0%) do not know enough.

Variables	No. (%)
Have you attended any training on safety before?	
Yes	100 (32.9)
No	204 (67.1)
Total	304 (100.0)
If yes, above. Type of training and organizer.	
Preventing health hazards by the Ministry of Health.	65 (65.0)
Effect of industrial gadgets by health agencies.	14 (14.0)
Training on the importance of safety by medical personnel	21 (21.0)
Total	100 (100.0)
How many such training have you attended?	
2	16 (5.3)
5	36 (11.8)
4	40 (13.2)
3	4 (1.3)
6	4 (1.3)
Total	100 (100.0)
I can identify PPE when I see one.	
Yes	221 (72.7)
No	83 (27.3)
Total	304 (100.0)
Usage of PPEs	
Yes	105 (34.5)
No	199 (65.5)
Total	304 (100.0)
If no, state the reason for not using safety gadgets.	
Not available	48 (24.1)

Variables	No. (%)
Not necessary	57 (28.6)
Not convenient	5 (2.5)
Forgetfulness	6 (3.1)
Other reason	83 (41.7)
Total	199 (100.0)
Common hazards in the workplace	
Noise	273 (89.8)
Heat	192 (63.2)
Chemical	121 (39.8)
Wood dust and smoke	192 (63.2)
Fire	128 (42.1)
Machine	174 (57.2)
Electricity	150 (49.3)
Manual lifting of objects	221 (72.7)
Animal/insect bite	264 (86.8)
Common injuries in the workplace	
Bruises	292 (96.1)
Sprain	280 (92.1)
Laceration	281 (92.4)
Fracture	262 (86.2)
Fainting attack	232 (76.3)
Burns	229 (75.3)
Electric shock	250 (82.2)
Hearing impairment	281 (92.4)
Common respiratory and non-respiratory problems in the workplace	
Cough	285 (93.8)
Phlegm production	288 (94.7)
Wheeze	284 (93.4)
Chest pain	280 (92.1)
Chest tightness	252 (82.9)
Shortness of breath	250 (82.2)
Dyspnoea	249 (81.9)
Sneezing/running nose	279 (91.8)
Skin problem	268 (88.2)
Eyes problem/red eye	271 (89.1)

Abbreviation: PPE, personal protective equipment.



Knowledge level of respondents

Figure 1. Knowledge of respondents on safety/protective devices and occupational hazards/injuries in sawmilling industry

## 4.4. Occupational Health Problems Suffered by Respondents in the Last One Month

Respondents reported having suffered from some of the following common respiratory problems in the last month: Cough 158 (52.0%), phlegm production 149 (49.0%), wheezing 114 (37.5%), chest pain 63 (20.7%), and running nose 154 (50.7%) while non-respiratory problems encountered in the last one month were skin problem 100 (32.9%) and eyes problem 111 (36.5%). Other various occupational health challenges experienced by the respondents in the last month were sprain 18 (5.9%), bruises 108 (35.5%), lacerations 4 (1.3%), fractures 14 (4.6%), heat exhaustion 4 (1.3%), electric shock 3 (1.0%), and fainting attack 8 (2.6%) (Table 4). Of 304 respondents, 130 (43%) had previously suffered one type of illness or the other at work, while 174 (57%) had not (Figure 2).

Variables	No. (%)
Respiratory problems	
Cough	158 (52.0)
Phlegm production	149 (49.0)
Wheeze	114 (37.5)
Chest pain	63 (20.7)
Chest tightness	46 (15.1)
Shortness of breath	47 (15.5)
Dyspnoea	55 (18.1)
Sneezing/running nose	154 (50.7)
Non-respiratory problems	
Skin problem	100 (32.9)
Eyes problem/red eye	111 (36.5)
Suffered from sprain	
Yes	18 (5.9)
No	286 (94.1)
Total	304 (100.0)
Suffered from bruises	
Yes	108 (35.5)
No	196 (64.5)
Total	304 (100.0)
Suffered from laceration	
Yes	4 (1.3)
No	300 (98.7)
Total	304 (100.0)
Suffered from fracture	
Yes	14 (4.6)
No	290 (95.4)
Total	304 (100.0)
Suffered from heat exhaustion	
Yes	4 (1.3)
No	300 (98.7)

Variables	No. (%)
Total	304 (100.0)
Suffered from electric shock	
Yes	3 (1.0)
No	301 (99.0)
Total	304 (100.0)
Suffered from fainting attack	
Yes	8 (2.6)
No	296 (97.4)
Total	304 (100.0)



Figure 2. Summarized score on injury suffered by respondents

## 4.5. Anthropometric Measurement and Vital Signs of Respondents

Table 5 depicts the respondents' BMI, blood pressure, pulse rate, and respiratory status measurements. A larger proportion of the respondents are obese 141 (46.4%) and overweight 131 (43.1%) from the BMI readings. The majority, 294 (96.7%) of the respondents have normal blood pressure, while a few, 10 (3.3%), are hypertensive. Likewise, most 281 (92.4%) respondents have normal pulse rate. Barely more than one-third, 119 (39.1%) of the respondents have a FER (%) lower than 70%, while 111 (36.5%) have respiratory rates outside the normal range (12-18 breaths per minute).

Table	5.	Parameter	Measurements	of	Body	Mass	Index,	Blood	Pressure	and
Respir	ato	ry Function	Test							

Variables	<b>No.</b> (%)
Using BMI	
Underweight	1(0.3)
Normal	31 (10.2)
Overweight	131 (43.1)
Obesity	141 (46.4)
Total	304 (100.0)
Blood Pressure	
Hypertension	10 (3.3)
Normal blood pressure	294 (96.7)
Total	304 (100.0)
Pulse rate (beats per minute)	

Variables	No. (%)
Normal rate	281 (92.4)
Abnormal rate	23 (7.6)
Total	304 (100.0)
FER (%)	
Greater than 70.00%	185 (60.9)
Lesser than 70.00%	119 (39.1)
Respiratory rate	
12 - 18 breaths per minute	193 (63.5)
Outside the range (12 - 18 breaths per minute)	111 (36.5)

Abbreviation: BMI, body mass index.

4.6. Relationship Between Respondents' Sociodemographic Characteristics and Their Summed Knowledge of Safety Procedures and Occupational Hazards and Accidents

Table 6 indicates the association between respondents' socio-demographic variables and summarized knowledge of safety practices and workplace hazards/injuries. Knowledge of hazards and safety procedures was statistically correlated with sociodemographic factors, including job category, gender, and educational attainment.

 
 Table 6. Association Between Demographic Status and Summarized Knowledge of Safety Practices and Workplace Hazard/Injuries

Variables	Good Knowledge <sup>a</sup>	Poor Knowledge <sup>a</sup>	Df	$\chi^2$	P- Value
Age group (y)			2	0.34	0.840
20 - 39	66 (58.4)	47 (41.6)			
40 - 59	100 (55.2)	81 (44.8)			
≥60	6(60.0)	4 (40.0)			
Job category			1	3.72	0.040
Administrative aspect	31 (46.3)	36 (53.7)			
Operation aspect	141 (59.5)	96 (40.5)			
Gender			1	8.64	0.003
Male	157 (59.9)	105 (40.1)			
Female	15 (35.7)	27 (64.3)			
Religion			1	2.6	0.107
Christianity	83 (52.2)	76 (47.8)			
Islam	89 (61.4)	56 (38.6)			
Marital status			3	4.20	0.240
Single	12 (60.0)	8 (40.0)			
Married	152 (55.3)	123 (44.7)			
Separated/divorced	6 (85.7)	1 (14.3)			
Widowed	2 (100.0)	0(0.0)			
Level of education			4	32.14	0.0001
Primary and below	43 (41.0)	62 (59.0)			

Variables	Good Knowledge <sup>a</sup>	Poor Knowledge a	Df	$\chi^2$	P- Value
education	85 (62.0)	52 (38.0)			
Tertiary education	44 (71.0)	18 (29.0)			
<sup>a</sup> Values are express	sed as No. (%).				

#### 4.7. Association Between Summarized Knowledge and Summarized Injury Occurrence Among Respondents

There is no association between summarized knowledge and summarized injury occurrence among respondents. However, according to Table 7, the percentage of those who had previously suffered injuries among those with good knowledge is reduced compared to those with poor knowledge.

Variables	Previously Suffer Injury <sup>a</sup>	Non-suffers of Injury <sup>a</sup>	Df	$\chi^2$	P- Value
Good knowledge	95 (55.2)	77 (44.8)		0.65	0.42
Poor knowledge	79 (59.8)	53 (40.2)	1		

#### 5. Discussion

This study assessed the awareness of safety procedures, recognition of workplace hazards, and understanding of potential injuries among Akure sawmill workers. Most respondents, young adult males with a mean age of  $42.0 \pm 2.8$  years, were predominantly engaged in operational activities, reflecting the physically demanding nature of the sawmilling occupation. Although the average ages of the study participants were higher than in similar studies in Kwara state and Opa Ile-Ife (14, 15), many had only completed secondary school, often through internships, which aligns with the occupation's unskilled nature (14, 16). The drawback of apprenticeship is that learners are constrained to the knowledge and methods of the mentors.

Regarding occupational history and exposure, participants worked longer hours than recommended safety standards, highlighting economic pressures and poverty in the nation (17). Sensitivity to workplace hazards such as smoke, sawdust, wood dust, and coal fire smoke was confirmed. Workplace exposure to wood dust and other similar compounds is a severe occupational risk that requires regular inspection and set guidelines (18). Although the majority (72.7%) of the participants in this study were knowledgeable about PPE, their use of PPE did not correspond to their knowledge, which was consistent with previous authors' observations (8, 19).

Major job dangers identified included noise (89.8%), heat (63.2%), wood dust and smoke (63.2%), machines (57.2%), physical lifting (72.7%), and animal or bug bites (86.8%) (4). Noise, recognized as a risk, was consistent with previous studies, and hearing loss emerged as a prevalent occupational ailment due to long-term exposure (3, 4). This result was in contrast to Osagbemi et al. (8), showing a lower percentage indicating nose as a risk. The respondents listed sprains, fractures, bruises, fatigue, fractures, and fainting attacks as common industrial injuries due to the respondents' constant manual lifting and handling of heavy goods. The results of earlier studies that addressed ergonomic risks have supported this claim (2-4). Only slightly more than half of the participants believed that machines were dangerous in their jobs, but this percentage was lower than in earlier research (13, 17). This knowledge gap could be attributed to sawmill workers learning more about the machines over time. Most participants agreed with Adeove et al. (4) that animal and insect bites are a prevalent biological danger.

Some hazards like chemical exposure (39.8%), fire (42.1%), and electricity (49.3%) were less acknowledged. The report by Adeoye et al. (4) is consistent with the present respondents' minimal impression of chemical hazards in sawmills. Numerous studies have confirmed that being exposed to chemicals coming from exposed wood species surfaces can have dangerous effects (2, 9). Respondents identified electricity as a risk that could result in electric shock due to carelessly exposed and dangerously exposed cut-outs, wires, and fittings (4, 8) and burns brought on by fire dangers as a result of carelessly and indiscriminately burning a pile of wood dust in the open (4, 6). The overall categorization of respondents' knowledge of safety procedures and awareness of workplace dangers and injuries is shown in Figure 1. This study showed that over half (57%) of participants were strongly aware of safety procedures, workplace dangers, and injuries, which was greater than that reported by earlier authors (14, 15).

The frequency of certain illnesses and injuries among sawmill workers over the previous month was also evaluated. The respondents' top three most common respiratory illnesses were coughing, sneezing, running nose, and phlegm production. This result was consistent with what Fatusi and Erhabor (20) reported, who identified cough as a typical respiratory symptom among sawmill workers. Similar to Ige and Onadeko (12). The prevalence of chest pain as a common symptom was very low compared to similar studies (20, 21).

Anthropometric measurements and vital signs, including a high prevalence of overweight issues (90%) and respiratory problems in one-third of the respondents, revealed potential health concerns (22). Approximately two-fifths had a decreased FER, indicating potential obstructive airway issues linked to wood dust exposure (23).

Additionally, nine out of ten responders were either obese or had excess body weight. The hypertension and irregular pulse rate identified in a small number of employees may be caused by these disorders. Obesity has been identified as a major risk factor for the development of stroke, hypertension, asthma, coronary artery disease, and several cardiovascular disorders, according to various studies (24-26). Most study participants came from a low socioeconomic background and had little or no formal education. Numerous authors have noted that due to subpar sanitation, a breakdown in public health, and unfavorable environmental and working conditions, people from the lower strata of society are more susceptible to a variety of health issues. However, this finding aligned with prior descriptive studies demonstrating a substantial correlation between sociodemographic characteristics and self-reported occupational injuries (27, 28).

#### 5.1. Conclusions

This survey indicated that respondents possess good knowledge of safety procedures and workplace risks. Despite awareness, many exhibit poor compliance with protective gear, leading to self-reported work-related injuries and health issues. Predominantly respiratory problems were reported in the past month, with some non-respiratory issues. Job type and education level significantly influenced safety knowledge and selfreported injuries. Sawmill workers in Akure experienced decreased respiratory function due to wood dust exposure and inadequate protective equipment. The majority of people dealing with obesity are at risk of various health issues.

Several recommendations emerge from the research to enhance the well-being and productivity of sawmill workers in Akure and nationwide. First, fostering collaboration between the National Union of Civil Engineering, Construction, Furniture, and Wood Workers (NUCECFWW) and government bodies like the Ministries of Labour & Productivity and Health is essential. This partnership should facilitate regular training sessions to educate sawmill workers on workplace hazards, proper use of PPE, and measures to prevent occupational injuries. Second, technological upgrades are crucial, including advanced equipment like fabric filters, exhaust ventilation devices, and noise abatement devices. Establishing training and innovation centers for technical skills is equally important. Third, promoting toolbox discussions between management and employees can prevent avoidable workplace injuries. Implementing safety procedures such as risk assessments and job safety analyses is imperative. Fourth, government agencies like the Environmental Protection Agency should routinely monitor atmospheric aerosols around sawmill sites to gauge occupational exposure levels. Finally, effective occupational health programs necessitate informed management and employees aware of wood dust's health and safety implications.

#### Footnotes

Authors' Contribution: Study concept and design, and administrative, technical, and material support: Sunday Olakunle Olarewaiu and Olufunke Julianah Ogidan: acquisition of data: Olufunke Julianah Ogidan; analysis and interpretation of data: Olufunke Julianah Ogidan and Adewale Allen Sokan-Adeaga; drafting of the manuscript: Adewale Allen Sokan-Adeaga; critical revision of the manuscript for important intellectual content: Sunday Olakunle Olarewaju, Adewale Allen Sokan-Adeaga, Akinlolu Oluwafemi Fansanmi, Olufunke Julianah Ogidan, Micheal Ayodeji Sokan-Adeaga, and Joy Stephen Amusan; statistical analysis: Sunday Olakunle Olarewaju, Olufunke Julianah Ogidan, and Adewale Allen Sokan-Adeaga; study supervision: Sunday Olakunle Olarewaju. All authors read and approved the final version of the manuscript.

**Conflict of Interests Statement:** The authors declare that they have no competing interest.

**Data Availability:** The dataset presented in the study is available on request from the corresponding author during submission or after publication. The data are not publicly available due to confidentiality.

**Funding/Support:** This research was personally funded. No funding was received from any external source or institution.

**Informed Consent:** Informed consent was obtained from all individual who participated in this study.

#### References

- Olorunnisola AO; International Union of Forestry Research Organizations Division 5; Chinese Forest Products Association; United States Forest Service. A Characterization of the Small -Scale Sawmilling Industry in Nigeria. 5.04.08-sawing, milling and machining; 2007 D5 IUFRO All Division 5 Conference: Forest Products and Environment : a Productive Symbiosis: Proceedings: October 29th -November 2nd, 2007, Grand Hotel, Taipei, Taiwan. Beijing: Chinese Forest Products Association; 2007. 273 p.
- Bello SR, Mijinyawa Y. Assessment of injuries in small scale sawmill industry of south western Nigeria. Agric Eng Int CIGR J. 2010;12(1):151-7.
- 3. Aruofor RO. Review and improvement of data related to wood-products in Nigeria. 2000. Available from: https://www.fao.org/4/X6762E/X6762E00.htm.
- 4. Adeoye OA, Adeomi AA, Abodunrin OL, Olugbenga-Bello AI, Abdulsalam ST. Awareness of occupational hazards and health problems among sawmill workers in Osun state, Nigeria. *Int J Res Rev.* 2015;**2**(1):1-14.
- Rongo LMB, Besselink A, Barten F, Msamanga GI, Dolmans WMV, Heederik D. Lung function among low-and high-exposure workers in small-scale wood industries in Tanzania. *East Afr J Public Health*. 2005;2(1):17-20.
- 6. US Department of Labor Occupational Safety & Health Administration. *Wood Dust.* 2008. Available from: https://www.osha.gov/wood-dust.
- Shamssain MH. Pulmonary function and symptoms in workers exposed to wood dust. *Thorax*. 1992;47(2):84-7. [PubMed ID: 1549828].
   [PubMed Central ID: PMC463576]. https://doi.org/10.1136/thx.47.2.84.
- Osagbemi GK, La-Kadri RT, Aderibigbe SA. Awareness of Occupational Hazards, Health Problems and Safety Measures among Sawmill Workers in North Central Nigeria. *TAF Prev Med Bull*. 2010;9(4):325-8.
- 9. Bean TL, Butcher TW, Lawrence T. *Wood dust exposure hazard*. 2006. Available from: https://ohioline.osu.edu/.
- Meo SA. Lung function in Pakistani wood workers. *Int J Environ Health Res.* 2006;16(3):193-203. [PubMed ID: 16611564]. https://doi.org/10.1080/09603120600641375.
- Pontier JP, Popin E, Kopferschmitt-Kubler MC, Bessot JC, Pauli G. [Asthma from tropical abachi wood]. *Rev Pneumol Clin.* 2002;**58**(5 Pt 1):282-5. French. [PubMed ID: 12486378].
- Ige OM, Onadeko OB. Respiratory symptoms and ventilatory function of the sawmillers in Ibadan, Nigeria. *Afr J Med Med Sci.* 2000;**29**(2):101-4. [PubMed ID: 11379437].
- International Finance Corporation. Environmental, health, and safety guidelines for sawmilling and manufactured wood products. 2007. Available from: http://documents.worldbank.org/curated/en/378521489661822778/En vironmental-health-and-safety-guidelines-for-sawmilling-andmanufactured-wood-products.
- 14. Agbana BE, Joshua AO, Daikwo MA, Metiboba LO. Knowledge of occupational hazards among sawmill workers in Kwara state,

Nigeria. Niger Postgrad Med J. 2016;**23**(1):25-32. [PubMed ID: 27098946]. https://doi.org/10.4103/1117-1936.180176.

- Faremi F, Ogunfowokan A, Mbada C, Olatubi M, Ogungbemi A. Occupational hazard awareness and safety practices among Nigerian sawmill workers. Int J Med Sci Public Health. 2014;3(10):1244. https://doi.org/10.5455/ijmsph.2014.260620142.
- Sabitu K, Iliyasu Z, Dauda MM. Awareness of occupational hazards and utilization of safety measures among welders in Kaduna metropolis, northern Nigeria. *Ann Afr Med.* 2009;8(1):46-51. [PubMed ID: 19763007]. https://doi.org/10.4103/1596-3519.55764.
- Olarewaju SO, Sokan-Adeaga AA, Awoaafe FB, Akinyele CA, Omobuwa O. Assessment of Occupational Injuries and Safety Practices among Automobile Repair Artisans in Apo and Gudu Area of Abuja, Nigeria. J Epidemiol Soc Nigeria. 2021;4(2):71-86. https://doi.org/10.5281/zenodo.6632724.
- Mahmood N, Karadaky K, Hussain S, Ali A, Mohammad G, Mahmood O. Respiratory function among sawmill workers in different areas of Sulaimani city. *Int J Med Sci Public Health*. 2016;5(2):351. https://doi.org/10.5455/ijmsph.2016.25082015128.
- 19. Agu AP, Umeokonkwo CD, Nnabu RC, Odusanya OO. Health Problems among Sawmill Workers in Abakaliki and Workplace Risk Assessment. J Community Med Prim Health Care. 2016;**28**(2):1-10.
- 20. Fatusi A, Erhabor G. Occupational health status of sawmill workers in Nigeria. J R Soc Health. 1996;**116**(4):232-6. [PubMed ID: 8783853]. https://doi.org/10.1177/146642409611600408.
- 21. Ugheoke A, Wahab K, Erhabor G. Prevalence of respiratory symptoms among sawmill workers in Benin city, Nigeria. *Int J Trop Med.* 2009;**4**(1):1-3.
- 22. Allen SAA, Ree AG, Ayodeji SAM, Deborah SAE, Ejike OM. Secondary inorganic aerosols: impacts on the global climate system and human health. *Biodivers Int J.* 2019;**3**(6):249-59. https://doi.org/10.15406/bij.2019.03.00152.
- Pellegrino R, Viegi G, Brusasco V, Crapo RO, Burgos F, Casaburi R, et al. Interpretative strategies for lung function tests. *Eur Respir J.* 2005;26(5):948-68. [PubMed ID: 16264058]. https://doi.org/10.1183/09031936.05.00035205.
- 24. Centers for Disease Control and Prevention. *Overweight and obesity*. 2011. Available from: www.cdc.gov/obesity.
- Poirier P, Giles TD, Bray GA, Hong Y, Stern JS, Pi-Sunyer FX, et al. Obesity and cardiovascular disease: pathophysiology, evaluation, and effect of weight loss. *Arterioscler Thromb Vasc Biol*. 2006;26(5):968-76. [PubMed ID: 16627822]. https://doi.org/10.1161/01.ATV.0000216787.85457.f3.
- Din-Dzietham R, Liu Y, Bielo MV, Shamsa F. High blood pressure trends in children and adolescents in national surveys, 1963 to 2002. *Circulation*. 2007;116(13):1488-96. [PubMed ID: 17846287]. https://doi.org/10.1161/CIRCULATIONAHA.106.683243.
- 27. Thangaraj S, Shireen N. Occupational health hazards among automobile mechanics working in an urban area of Bangalore -a cross sectional study. *Int J Med Sci Public Health*. 2017;**6**(1):18. https://doi.org/10.5455/ijmsph.2016.31052016540.
- Johnson OE, Bassey EA. Work Habits and Health Problems of Automobile Technicians at Mechanic Village, Uyo, Nigeria. *Glob Adv Res J Med Med Sci.* 2016;5(5):136-42.