



Factors Associated with Cesarean Section in Preterm Births at a Tertiary Hospital in Rasht, Iran

Forozan Milani ¹, Seyedeh Hajar Sharami ^{1,*}, Seyedeh Maryam Attari ², Zahra Rafiee Sorouri ¹, Sara Farzadi ¹, Soodabeh Kazemi ¹

¹ Department of Obstetrics & Gynecology, Reproductive Health Research Center, Al-zahra Hospital, School of Medicine, Guilan University of Medical Sciences, Rasht, Iran

² Reproductive Health Research Center, Guilan University of Medical Sciences, Rasht, Iran

*Corresponding author: Department of Obstetrics & Gynecology, Reproductive Health Research Center, Al-zahra Hospital, School of Medicine, Guilan University of Medical Sciences, Rasht, Iran. Email: sharami@gums.ac.ir

Received 2024 January 7; Revised 2024 May 8; Accepted 2024 May 14.

Abstract

Background: Cesarean section (C-section) is increasing globally. Many studies have investigated factors associated with C-sections, but predictors of C-sections in preterm birth are not well documented.

Objectives: This study aimed to evaluate Factors associated with C-sections in preterm births at a Tertiary hospital in Rasht, Iran.

Methods: A retrospective cross-sectional study was conducted on all neonates born between 2017 and 2022. The data were collected from the medical records. Descriptive statistics, chi-square, and Fisher's exact test were performed. Multiple logistic regression was applied to identify factors associated with C-section.

Results: The sample community included 1145 preterm neonates. Factors associated with C-section in preterm births included the mother's age ($P < 0.001$, OR = 1.06), gestational age ($P < 0.001$, OR = 1.24), neonate's weight ($P = 0.042$, OR = 0.62), infertility history, ($P = 0.043$, OR = 1.58), abortion history ($P = 0.006$, OR = 1.88), and iatrogenic preterm birth ($P < 0.001$, OR = 4.99).

Conclusions: Clinicians and policymakers should give special attention to these factors for C-sections in preterm births, especially iatrogenic preterm birth.

Keywords: Preterm Birth, Cesarean Section, Mode of Childbirth

1. Background

Preterm birth is defined as birth before the 37th week of pregnancy (1), which is classified into three subgroups, extremely preterm (< 28 weeks), very preterm (28 - 32 weeks), and moderate preterm (32 - 37 weeks) (2). Preterm birth accounts for 3.1% of the global burden of diseases (3). Preterm birth in Iran has been estimated to be a total of 10% (4).

Cesarean section (C-section) is increasing worldwide, making it a global concern (5, 6). This trend in Iran is also rising, especially in private hospitals (7). The World Health Organization does not consider C-section to be more than 10% beneficial for the mother and baby (8). Meanwhile, the C-section rate in Iran is estimated at 48% (9). The literature generally states that vaginal delivery (VD) is a potential risk for neonatal mortality for

preterm infants, and C-section serves as a preventive strategy for its reduction (10). However, there is controversy regarding the effect of the mode of childbirth on the death of preterm infants (11). Although C-section may reduce neonatal death, they can increase maternal morbidity and mortality (12, 13). A study in Brazil showed that C-section is associated with an increased risk of infant mortality unless there is a clear indication for the procedure (14).

Many factors predict C-section in preterm birth. A large cross-sectional study showed that the mother's age, socioeconomic status, body mass index, parity, type of pregnancy, preeclampsia, infant height, and baby's head circumference are associated with C-section in Tehran, Iran. Sometimes, C-section is the preferred procedure of mothers and doctors to save the baby (15,

16). A study in preterm neonates showed that women who delivered vaginally were younger, had lower rates of hypertensive disorders and diabetes mellitus, and had higher rates of drug abuse as compared to those who delivered by C-section. Neonates who were born by VD had higher birth weights and a lower rate of SGA than those who were born by C-section. However, VD neonates had higher Apgar scores, as compared with C-section, the neonatal composite outcome was similar between the two modes of delivery (17). The factors associated with C-section (15, 18, 19) have been extensively addressed, but factors associated with C-section in preterm birth are not well documented.

2. Objectives

Accordingly, this paper evaluates the factors associated with C-section in preterm births at a Tertiary Hospital in Rasht, the center of the province of Gilan, Iran.

3. Methods

3.1. Study Design and Setting

This retrospective medical record review was conducted on neonatal subjects born between 2017 and 2022 at the Al-Zahra Maternity Hospital, Rasht, Iran, to determine the factors associated with C-section among preterm births.

Rasht is a populous city in the north of Iran equipped with one public referral maternity hospital. This hospital has two wards for neonatal hospitalization (Premature Neonate Unit and Neonatal Intensive Care Unit). Most pre-term births are performed because Al-Zahra is a tertiary hospital. According to Rezaeian et al. (20) which reported the prevalence of preterm delivery to be 7.1% and with an error of 5% using ratio estimation, the minimum sample size was determined to be 1126 people.

3.2. Sampling, Data Collection Procedure, and Tool

All medical records of neonates admitted between May 2017 and March 2022 were included. Two midwives and a trained intern used a structured checklist to collect data from the medical records of 1161 preterm infants. The checklist was adapted after reviewing various literature on similar studies. In this study, the characteristics of the mother's age, gestational age,

pregnancy and childbirth history, C-section cause, and neonate complications were investigated.

3.3. Ethics Considerations

Ethics approval was obtained from the Ethics Committee of Guilan University of Medical Sciences, Rasht, Iran (IR.GUMS.REC.1399.302). All reviewed medical records had a signed informed consent form for the use of patient medical data completed by the neonates' parents.

3.4. Data Analysis

The gathered data were analyzed in SPSS software version 25. The chi-square and Fisher's exact tests were performed to analyze the relationships between nominal scale variables. Factors with ≤ 0.2 significant level in the bivariate logistic regression analysis were considered in the multivariable logistic regression analysis. The multivariable analyses were conducted using multiple logistic regression models at a 95% confidence interval (CI) and $P < 0.05$ statistical significance.

4. Results

The mean age of mothers was 30.90 ± 6.24 years old. Sixteen of the reviewed records were excluded from the study (five revisits and eleven out-of-hospital admissions). Finally, 1145 medical documents were imported into the study. Most of the preterm births (77%) were C-sections of which, 44.4% were repeated. The rest were performed primarily due to multiple pregnancies and fetal distress. Table 1 presents the C-section causes among mothers who had their first C-section.

Table 1. Cesarean Section (C-section) Causes Among First C-section Mothers

C-section Cause	No. (%)
Multiple pregnancies	154 (33.8)
Fetal distress	145 (31.9)
Breech presentation	67 (14.7)
Preeclampsia	34 (7.5)
Fetal growth restriction	23 (5.1)
Others	21 (4.6)
Placenta abruption	11 (2.4)

Table 2 represents the maternal and neonatal characteristics of neonates based on the mode of childbirth. The primary analysis showed a significant

difference between the mother's age ($P < 0.001$), gestational age ($P < 0.001$), abortion history ($P = 0.007$), hypertension ($P < 0.001$), preeclampsia ($P = 0.003$), fetal growth restriction ($P = 0.001$), fetal distress ($P < 0.001$), meconium ($P = 0.013$), and the preterm birth cause ($P < 0.001$) with the mode of childbirth (Table 1).

Table 2. Maternal and Neonatal Characteristics of Neonates Based on the Mode of Childbirth

Variables	Vaginal Delivery (N = 263)	C-section (N = 882)	P-Value ^a
Age			< 0.001
< 35	184 (24.4)	570 (75.6)	
≥ 35	47 (14.6)	275 (85.4)	
Infertility			0.054
Yes	36 (17.1)	174 (82.9)	
No	215 (23.3)	708 (76.5)	
History of abortion			0.007
Yes	39 (15.8)	208 (84.2)	
No	212 (23.9)	674 (76.1)	
History of death child			> 0.999
Yes	17 (21.5)	62 (78.5)	
No	234 (22.2)	820 (77.8)	
Hypertension			< 0.001
Yes	41 (13.8)	257 (86.2)	
No	210 (25.1)	625 (74.9)	
Preeclampsia			0.003
Yes	12 (11.1)	96 (88.9)	
No	239 (23.3)	786 (76.7)	
Diabetes			0.085
Yes	63 (18.8)	272 (81.2)	
No	188 (23.6)	610 (76.4)	
Gestational age (w)			< 0.001
≤ 28	41 (49.4)	42 (50.6)	
29 - 32	50 (25.5)	146 (74.5)	
≥ 33	159 (19.4)	659 (80.6)	
Sex			0.885
Female	113 (22.3)	394 (77.7)	
Male	136 (22.7)	462 (77.3)	
Birth weight (gr)			0.118
< 2500	156 (23.1)	518 (76.9)	
≥ 2500	71 (18.9)	304 (81.1)	
Congenital malformation			0.428
Yes	10 (27)	27 (73)	
No	241 (22)	855 (78)	
Fetal Growth restriction			0.001
Yes	18 (12.1)	131 (87.9)	
No	233 (23.7)	751 (76.3)	
Fetal distress			< 0.001
Yes	0 (0)	132 (100)	
No	251 (25.1)	750 (74.9)	
Meconium			0.013

Variables	Vaginal Delivery (N = 263)	C-section (N = 882)	P-Value ^a
Yes	5 (8.8)	52 (91.2)	
No	246 (29.9)	830 (77.1)	
Preterm birth cause			< 0.001
Spontaneous	232 (27.4)	616 (72.6)	
Iatrogenic	13 (6.1)	201 (93.9)	

^a Chi-square or Fisher's exact test.

^b Values are expressed as No. (%).

The variables with $P < 0.2$ were introduced to the logistic regression model. The analysis identified the following as factors associated with C-section: The mother's age ($P < 0.001$), gestational age ($P < 0.001$), neonate's weight ($P = 0.042$), infertility history ($P = 0.042$), abortion history ($P = 0.006$), and preterm birth cause ($P < 0.001$) (Table 3).

Table 3. Multivariate Logistic Regression Analysis of Factors Associated with Cesarean Section

Variables	β	SD	Wald	P-Value	OR (95% CI)
Mother's age	0.063	0.015	18.186	< 0.001	1.06 (1.034, 1.09)
Gestational age	0.216	0.036	35.794	< 0.001	1.24 (1.15, 1.33)
Birth weight	-0.464	0.229	4.125	0.042	0.62 (0.40, 0.98)
Hypertension	0.196	0.204	0.015	0.430	1.21 (0.74, 1.97)
Preeclampsia	0.026	0.467	0.003	0.955	1.02 (0.41, 2.56)
Diabetes	0.025	0.204	0.015	0.902	1.02 (0.68, 1.52)
Infertility history	0.461	0.228	4.083	0.043	1.58 (1.01, 2.48)
Abortion history	0.636	0.232	7.481	0.006	1.88 (1.19, 2.97)
Fetal growth restriction	-0.162	0.313	0.267	0.605	0.85 (0.46, 1.57)
Fetal malformation	-0.435	0.427	1.039	0.308	0.64 (0.28, 1.49)
Preterm birth reason	1.609	0.380	17.957	< 0.001	4.99 (2.37, 10.53)

5. Discussion

Given the high rate of C-sections in the study setting in recent years and public preference for C-section to vaginal delivery due to attitudinal and operational barriers (21), the C-section rate was high among preterm births. The most common reason for the C-section procedure was repeated C-sections. Mother's age, gestational age, birth weight, infertility and abortion history, and preterm birth were found to be the factors associated with C-section in preterm births.

5.1. Neonatal Factors

The results of this study indicated that the number of C-section operations increases as the gestational age increases with each week, which can be attributed to the

the risk of C-section for the mother more easily. In line with this finding, a study on term infants also found that older gestational age is a risk factor associated with an increased risk of C-section (22).

Birth weight is another neonatal factor that predicts C-section. Preterm neonates $\geq 2,500$ g were 1.5 times more likely to deliver through C-section than neonates $< 2,500$ g. This finding is in line with the study of Alhassan in Ghana, which indicated that the odds of birth by C-section decreased among pregnant women with low birth weight (23).

5.2. Maternal Factors

The current study found that mothers ≥ 35 years old were more likely to have C-section compared to mothers < 35 years old. This finding broadly supports the work of other studies in this area. Manyeh et al. found that mothers aged 30 - 34 years were more likely to have C-section compared to those < 20 years (22). Another study showed that maternal age ≥ 35 years is one of the factors associated with increased odds of C-section (24).

In addition, a history of infertility and abortion were predictors of C-sections. Another large study in southwest Iran also showed that previous abortion and previous infertility were the risk factors for C-section (25). Sinnott et al. also concluded that a prior history of miscarriage is a predictor for C-section (26). On the other hand, some studies have shown that C-section is one of the risk factors for infertility and abortion (27, 28). This cause-and-effect relationship cannot be definitively determined due to the study's cross-sectional nature. However, abortion and infertility are predictors of C-sections after splitting repeated C-sections. Maybe women with a history of abortion and infertility and their doctors are more concerned about saving the baby and are more inclined to choose C-section.

5.3. Iatrogenic Preterm Birth

Although hypertension and preeclampsia were found to have a stronger association with C-sections in the primary analysis, no association was identified between these conditions and C-sections in the regression analysis results. The iatrogenic preterm births were about five times more likely to take the C-section procedure than spontaneous preterm births. By

the present results, Dehaene et al. found higher C-section rates in iatrogenic preterm births (29). Considering the role of iatrogenic preterm birth in C-sections, it seems that a possible reduction in iatrogenic births could decrease the C-section rate. In addition, iatrogenic preterm births had a higher rate of neonatal death at discharge compared to spontaneous preterm births. In general, some neonatal complications are associated with etiologic types of preterm birth. Thus, proper management of different etiological subgroups of preterm births can prevent adverse neonatal outcomes.

This study has several strengths, including a large sample of preterm births in a referral tertiary hospital. However, several limitations should be noted. First, some key factors related to C-sections (the mother's level of education and insurance status) were not explored since the data were mainly gathered from only the medical records of neonates. Second, a definite opinion about the causal relationships can be made due to the cross-sectional design. Third, missing data may have affected the results due to the incompleteness of some of the neonates' medical records.

5.4. Conclusions

The incidence of C-sections is significant among preterm births in the Alzahra tertiary care hospital. Mother's age, gestational age, birth weight, infertility and abortion history, and iatrogenic preterm birth were the factors associated with C-section in preterm births. Physicians must pay attention to these factors in managing preterm births. In addition, preterm birth management based on their etiology can prevent adverse neonatal outcomes.

Footnotes

Authors' Contribution: F.M., S.KA, and SH.Sh contributed to conception and design; F.M., Z.RS, and S.F. contributed to data acquisition; SM.A performed the statistical analysis; SM.A and S.KA drafted the manuscript. F.M. and SH.Sh critically revised the manuscript. All authors read and approved the manuscript and agree to be accountable for all aspects of the work, ensuring integrity and accuracy.

Conflict of Interests Statement: The authors declared that they have no competing interests.

Data Availability: The dataset presented in the study is available on request from the corresponding author during submission or after publication.

Ethical Approval: Ethics approval was obtained from the Ethics Committee of Guilan University of Medical Sciences, Rasht, Iran (IR.GUMS.REC.1399.302).

Funding/Support: The study was financially supported by the Research Deputy of Guilan University of Medical Sciences.

Informed Consent: All reviewed medical records had an informed consent form for the use of patient medical data completed by the neonates' parents.

References

- Manuck TA, Rice MM, Bailit JL, Grobman WA, Reddy UM, Wapner RJ, et al. Preterm neonatal morbidity and mortality by gestational age: a contemporary cohort. *Am J Obstet Gynecol*. 2016;**215**(1):103 e1-103 e14. [PubMed ID: 26772790]. [PubMed Central ID: PMC4921282]. <https://doi.org/10.1016/j.ajog.2016.01.004>.
- Tucker J, McGuire W. Epidemiology of preterm birth. *BMJ*. 2004;**329**(7467):675-8. [PubMed ID: 15374920]. [PubMed Central ID: PMC517653]. <https://doi.org/10.1136/bmj.329.7467.675>.
- Murray CJ, Vos T, Lozano R, Naghavi M, Flaxman AD, Michaud C, et al. Disability-adjusted life years (DALYs) for 291 diseases and injuries in 21 regions, 1990-2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet*. 2012;**380**(9859):2197-223. [PubMed ID: 23245608]. [https://doi.org/10.1016/S0140-6736\(12\)61689-4](https://doi.org/10.1016/S0140-6736(12)61689-4).
- Sharifi N, Khazaeian S, Pakzad R, Fathnezhad Kazemi A, Chehreh H. Investigating the Prevalence of Preterm Birth in Iranian Population: A Systematic Review and Meta-Analysis. *J Caring Sci*. 2017;**6**(4):371-80. [PubMed ID: 29302576]. [PubMed Central ID: PMC5747595]. <https://doi.org/10.15171/jcs.2017.035>.
- Le Ray C, Blondel B, Prunet C, Khireddine I, Deneux-Tharoux C, Goffinet F. Stabilising the caesarean rate: which target population? *BJOG*. 2015;**122**(5):690-9. [PubMed ID: 25412695]. <https://doi.org/10.1111/1471-0528.13199>.
- Assari Arani A, Atashbar T, Antoun J, Bossert T. Iran's Health Reform Plan: Measuring Changes in Equity Indices. *Iran J Public Health*. 2018;**47**(3):390-6. [PubMed ID: 29845027]. [PubMed Central ID: PMC5971176].
- Pourshirazi M, Heidarzadeh M, Taheri M, Esmaily H, Babaey F, Talkhi N, et al. Caesarean delivery in Iran: a population-based analysis using the Robson classification system. *BMC Pregnancy Childbirth*. 2022;**22**(1):185. [PubMed ID: 35260106]. [PubMed Central ID: PMC8903666]. <https://doi.org/10.1186/s12884-022-04517-1>.
- World Health Organization Human Reproduction Programme A. WHO Statement on caesarean section rates. *Reprod Health Matters*. 2015;**23**(45):149-50. [PubMed ID: 26278843]. <https://doi.org/10.1016/j.rhm.2015.07.007>.
- Rafiei M, Saei Ghare M, Akbari M, Kiani F, Sayehmiri F, Sayehmiri K, et al. Prevalence, causes, and complications of caesarean delivery in Iran: A systematic review and meta-analysis. *Int J Reprod Biomed*. 2018;**16**(4):221-34. [PubMed ID: 29942930]. [PubMed Central ID: PMC6004597].
- Iyanda AE, Osayomi T. Association Between Childbirth Modes and Neonatal and Maternal Deaths Using a Negative Binomial Model. *Women's Reproductive Health*. 2019;**6**(2):128-40. <https://doi.org/10.1080/23293691.2019.1601905>.
- Daemi A, Ravaghi H, Jafari M. Risk factors of neonatal mortality in Iran: a systematic review. *Med J Islam Repub Iran*. 2019;**33**:87. [PubMed ID: 31696081]. [PubMed Central ID: PMC6825378]. <https://doi.org/10.34171/mjiri.33.87>.
- Moudi Z, Arabnezhad L, Ansari H, Tabatabaei SM. Severe maternal morbidity among women with a history of cesarean section at a tertiary referral teaching hospital in the southeast of Iran. *Public Health*. 2019;**175**:101-7. [PubMed ID: 31470235]. <https://doi.org/10.1016/j.puhe.2019.07.016>.
- Changizi N, Rezaeizadeh G, Janani L, Shariat M, Habibelahi A. In Depth Analysis of the Leading Causes of Maternal Mortality Due to Cesarean Section in Iran. *J Family Reprod Health*. 2017;**11**(1):1-6. [PubMed ID: 29114262]. [PubMed Central ID: PMC5664984].
- Paixao ES, Bottomley C, Pescarini JM, Wong KLM, Cardim LL, Ribeiro Silva RC, et al. Associations between cesarean delivery and child mortality: A national record linkage longitudinal study of 17.8 million births in Brazil. *PLoS Med*. 2021;**18**(10). e1003791. [PubMed ID: 34637451]. [PubMed Central ID: PMC8509988]. <https://doi.org/10.1371/journal.pmed.1003791>.
- Amini P, Mohammadi M, Omani-Samani R, Almasi-Hashiani A, Maroufizadeh S. Factors Associated with Cesarean Section in Tehran, Iran using Multilevel Logistic Regression Model. *Osong Public Health Res Perspect*. 2018;**9**(2):86-92. [PubMed ID: 29740533]. [PubMed Central ID: PMC5935148]. <https://doi.org/10.24171/j.phrp.2018.9.2.08>.
- Ochieng Arunda M, Agardh A, Asamoah BO. Cesarean delivery and associated socioeconomic factors and neonatal survival outcome in Kenya and Tanzania: analysis of national survey data. *Glob Health Action*. 2020;**13**(1):1748403. [PubMed ID: 32345146]. [PubMed Central ID: PMC7241493]. <https://doi.org/10.1080/16549716.2020.1748403>.
- Gluck O, Tairy D, Bar J, Barda G. The impact of mode of delivery on neonatal outcome in preterm births. *J Matern Fetal Neonatal Med*. 2021;**34**(8):1183-9. [PubMed ID: 31154888]. <https://doi.org/10.1080/14767058.2019.1627319>.
- Maroufizadeh S, Bagheri-Lankarani N, Almasi-Hashiani A, Amini P, Esmaeilzadeh A, Navid B, et al. Prevalence of Cesarean Section and its Related Factors among Primiparas in Tehran Province, Iran, in 2015. *J Isfahan Med Sch*. 2017;**35**(423):303-9. Persian.
- Sabetghadam S, Keramat A, Goli S, Malary M, Rezaie Chamani S. Assessment of Medicalization of Pregnancy and Childbirth in Low-risk Pregnancies: A Cross-sectional Study. *Int J Community Based Nurs Midwifery*. 2022;**10**(1):64-73. [PubMed ID: 35005042]. [PubMed Central ID: PMC8724730]. <https://doi.org/10.30476/IJCBNM.2021.90292.1686>.
- Rezaeian M, Goujani R, Sheikh Fathollahi M, Vaziri Nejad R, Manshori A, Razi S. A Comparative Study on Prevalence of Preterm Birth and Low Birth Weight in Iranians and Afghans Races in Rafsanjan NIK-Nafs Hospital in 2011-2012. *J Rafsanjan Univ Med Sci*. 2014;**13**(1):67-82. Persian.
- Firoozi M, Tara F, Ahanchian MR, Latifnejad Roudsari R. Clinician's and women's perceptions of individual barriers to vaginal birth after cesarean in Iran: A qualitative inquiry. *Caspian J Intern Med*. 2020;**11**(3):259-66. [PubMed ID: 32874432]. [PubMed Central ID: PMC67442460]. <https://doi.org/10.22088/cjim.11.3.259>.
- Manyeh AK, Amu A, Akpakli DE, Williams J, Gyapong M. Socioeconomic and demographic factors associated with caesarean section delivery in Southern Ghana: evidence from INDEPTH

- Network member site. *BMC Pregnancy Childbirth*. 2018;**18**(1):405. [PubMed ID: 30326869]. [PubMed Central ID: PMC6191905]. <https://doi.org/10.1186/s12884-018-2039-z>.
23. Alhassan AR. Prevalence and socioeconomic predictive factors of cesarean section delivery in Ghana. *Menoufia Med J*. 2022;**35**:190-5. https://doi.org/10.4103/mmj.mmj_444_20.
24. Adewuyi EO, Auta A, Khanal V, Tapshak SJ, Zhao Y. Cesarean delivery in Nigeria: prevalence and associated factors-a population-based cross-sectional study. *BMJ Open*. 2019;**9**(6). e027273. [PubMed ID: 31213450]. [PubMed Central ID: PMC6596937]. <https://doi.org/10.1136/bmjopen-2018-027273>.
25. Rajabi A, Maharlouei N, Rezaianzadeh A, Rajaeefard A, Gholami A. Risk factors for C-section delivery and population attributable risk for C-section risk factors in Southwest of Iran: a prospective cohort study. *Med J Islam Repub Iran*. 2015;**29**:294. [PubMed ID: 26913257].
26. Sinnott SJ, Brick A, Layte R, Cunningham N, Turner MJ. National Variation in Caesarean Section Rates: A Cross Sectional Study in Ireland. *PLoS One*. 2016;**11**(6). e0156172. [PubMed ID: 27280848]. [PubMed Central ID: PMC4900579]. <https://doi.org/10.1371/journal.pone.0156172>.
27. Sharif SN, Azizi Kutenae M, Darsareh F, Roozbeh N. Prevalence and Risk Factors of Infertility in a Southern Port City of Iran. *Hormozgan Med J*. 2020;**24**(2). e99412. <https://doi.org/10.5812/hmj.99412>.
28. Diao J, Gao G, Zhang Y, Wang X, Zhang Y, Han Y, et al. Caesarean section defects may affect pregnancy outcomes after in vitro fertilization-embryo transfer: a retrospective study. *BMC Pregnancy Childbirth*. 2021;**21**(1):487. [PubMed ID: 34229640]. [PubMed Central ID: PMC8261987]. <https://doi.org/10.1186/s12884-021-03955-7>.
29. Dehaene I, Scheire E, Steen J, De Coen K, Decruyenaere J, Smets K, et al. Obstetrical characteristics and neonatal outcome according to aetiology of preterm birth: a cohort study. *Arch Gynecol Obstet*. 2020;**302**(4):861-71. [PubMed ID: 32621250]. <https://doi.org/10.1007/s00404-020-05673-5>.