Published Online: 2025 June 23

Research Article



Molecular Docking of Silymarin as a Ligand for Aromatase Enzyme and HIV-1 Reverse Transcriptase Using Molegro Virtual Docker

Ahmad Hafezi 🔟 ¹, Zahra Khamar 🔟 ^{1,*}

¹ Department of Biology, Faculty of Science, Ferdowsi University of Mashhad, Mashhad, Iran

Corresponding Author: Department of Biology, Faculty of Science, Ferdowsi University of Mashhad, Mashhad, Iran. Email: zahrakhamr@um.ac.ir

Received: 27 April, 2025; Revised: 27 May, 2025; Accepted: 7 June, 2025

Abstract

Background: Molegro Virtual Docker (MVD) software is specifically designed for executing the molecular docking process. It has gained recognition among docking software for its user-friendliness, affordability, and quick result evaluation. The docking procedure can utilize multiple algorithms, and different molecules can be treated as ligands and receptors. Selin marine is one compound that, owing to its appropriate structure, can significantly function as a ligand and inhibitor for key enzymes involved in pathogenesis.

Objectives: The primary objective of this research is to perform in silico analyses to explore the potential of silibinin, silidianin, and silychristin molecules as ligands for interaction with the active sites of the Aromatase enzyme and human immunodeficiency virus-1 (HIV-1) reverse transcriptase enzymes.

Methods: In this research, the ligand structures were initially obtained from the PubChem database, while the enzyme structures were sourced from the PDB database. Subsequently, the docking process was conducted using the MVD software. Any structural errors in the molecules were rectified utilizing the 'if missing' approach. The cavities associated with the active sites of each enzyme were identified based on the positioning of the primary ligand. To ensure the docking process was executed with maximum precision, the MolDock Score (GRID) and the MolDock simplex evolution (SE) algorithm were employed in the settings, with ten runs conducted for each molecule.

Results: The findings from the docking process revealed that the molecules silibinin, silidianin, and silychristin can occupy the active site of the aromatase enzyme, exhibiting free energies of -141.972, -127.805, and -148.425 kcal/mol, respectively. Their ability to form steric interactions and hydrogen bonds suggests that they could serve as effective ligands for interaction with this active site. Furthermore, these ligands exhibit significant potential for establishing hydrogen and ester bonds within the active site of HIV-1 reverse transcriptase, with calculated free energies of -140.701, -138.956, and -135.365 kcal/mol for silibinin, silidianin, and silychristin, respectively.

Conclusions: Silymarin demonstrated the ability to interact with the active site of the aromatase enzyme and HIV-1 reverse transcriptase, potentially competing with their primary ligands for binding at this location, thereby inhibiting the activity of both enzymes in affected cells.

Keywords: Silymarin, Molecular Docking, Aromatase Enzyme, HIV-1 Reverse Transcriptase

1. Background

Flavonoids are a family of benzo gamma-pyrones, of which more than 4,000 different flavonoids have been identified to date. These compounds are abundant in both the animal and plant kingdoms, especially in photosynthetic cells. They have been used for centuries for their numerous therapeutic properties. Compounds such as quercetin, taxifolin, and silymarin have been used as active ingredients, both alone and as components of complex chemical preparations (1-3). This compound is extracted from the seeds and fruits of Silybum marianum (Compositae) due to its well-defined therapeutic properties and is actually a mixture of three structural components: Silibinin, silidianin, and silychristin (Figure 1) (3-5). Silymarin is a flavonolignan

Copyright @ 2025, Journal of Microbiota. This open-access article is available under the Creative Commons Attribution-NonCommercial 4.0 (CC BY-NC 4.0) International License (https://creativecommons.org/licenses/by-nc/4.0/), which allows for the copying and redistribution of the material only for noncommercial purposes, provided that the original work is properly cited.

How to Cite: Hafezi A, Khamar Z. Molecular Docking of Silymarin as a Ligand for Aromatase Enzyme and HIV-1 Reverse Transcriptase Using Molegro Virtual Docker. J Microbiota. 2025; 2 (2): e162254. https://doi.org/10.5812/jmb-162254.

that has recently been identified as a hepatoprotective agent (3, 6, 7). Silymarin is used to treat many liver disorders (1, 3, 8). Flavonolignans in Silybum marianum (silymarin) protect liver cells by preventing cytotoxins from entering the cells (2, 8, 9). This compound protects liver cells from damage caused by ischemia, radiation, iron overload, and viral hepatitis (3). Silymarin has also shown antihyperglycemic, anti-inflammatory, cardioprotective, and antioxidant effects (3, 6, 9), as well as cytoprotective, hypoglycemic, hypocholesterolemia, anticarcinogenic effects, gastro-protective, and/or antiulcer properties (5, 10).

The mechanism of action of silymarin involves inhibiting the absorption of toxins such as phalloidin or α -amanitin, preventing them from binding to the cell surface and inhibiting membrane transport systems. In addition, by interacting with the lipid components of the cell membrane, it can affect their chemical and physical properties (3). Carcinogenesis is a process involving steps that are activated by changing the expression of transcription factors and proteins involved in proliferation, cell cycle regulation, differentiation, apoptosis, angiogenesis, invasion, and metastasis (1, 6). Accordingly, factors that can disrupt this cycle are effective anticancer agents (5). Silymarin is able to disrupt this balance by interfering with the expression of cell cycle regulators and proteins involved in apoptosis. Also, the anti-inflammatory and antimetastatic activity of silymarin is due to the modulation of specific proteins (1, 6, 7).

1.1. The Effect of Silymarin on the Aromatase Enzyme

Aromatase is an enzyme that catalyzes the final step in estrogen biosynthesis. The regulation of aromatase activity and levels of estrogens determines the levels of estrogens, which subsequently have endocrine, paracrine, and autocrine effects on tissues (11). Estrogen is a steroid hormone that maintains female characteristics in the body. In addition to the reproductive system, estrogen plays an important role in the cardiovascular system and musculoskeletal system (12), regulating reproduction, metabolism, behavior, and brain function (11, 12). In the brain, cell survival and neuronal activity are influenced by estrogens and, as a result, aromatase (11). However, estrogen production itself affects female diseases. Therefore, its inhibition is effective in treating diseases (12). The aromatase reaction is the final step in the estrogen biosynthetic pathway, and inhibition of this final step inactivates the biosynthesis of other steroid classes (12). The aromatase enzyme plays a key role in the

pathogenesis of polycystic ovary syndrome (PCOS) (13), breast cancer (12), and endometriosis (14). For example, aromatase converts precursor steroids to estradiol and estrogen and promotes the proliferation of endometrial tissue. Therefore, aromatase contributes to disease progression by stimulating an inflammatory response (15). As a result, inhibition of the aromatase enzyme may provide an effective way to treat and control endometrial tissue proliferation by reducing estrogen production (15). Silymarin, which is extracted directly from the dried seeds of S. marianum and has antiinflammatory (4, 13), antioxidant (6, 16), and anti-fibrotic properties, acts as the main ingredient (13). Silvmarin is effective in treating endometrial lesions with antioxidant and anti-inflammatory effects (15) and plays its antioxidant role by eliminating free radicals as well as increasing the levels of glutathione peroxidase and superoxide dismutase (SOD) (6, 10, 16).

1.2. The Effect of Silymarin on HIV

Human immunodeficiency virus-1 (HIV-1) has infected nearly a million people worldwide. Antiretroviral therapy (ART) has now prolonged the lives of people with chronic infection (17). Herbal use is common among HIV patients, often used as selfmedication without medical supervision (18). Silymarin extract contains silibinin, which is a mixture of the flavonolignans silvbin A (SA) and silvbin B (SB). Silibinin has a wide range of antioxidant, immunomodulatory, antiproliferative, antifibrotic, and antiviral activities in tissues and organs. Silymarin and silibinin both inhibit HCV infection in cell culture by variably blocking viral entry, viral fusion, viral RNA and protein synthesis, HCV N5SB RNA-dependent RNA polymerase activity, and virus transmission (17). Although herbal supplements are often harmless, they can actually modulate various cytochrome P450 enzymes (such as CYP3A4) and drug transporters [such as P-glycoprotein (P-gp)], providing evidence for herbal-drug interactions. The most wellknown herbal remedy is St. John's wort (Hypericum perforatum), which has drug interactions with and is a potent inducer of CYP3A4 and P-gp activity (18). Laboratory studies have shown that silymarin may interact with many drugs, including HIV protease inhibitors. However, clinical studies with milk thistle have shown conflicting results regarding its ability to modulate these proteins in vivo, and evidence of interactions between milk thistle and antiretroviral drugs from clinical trials has prevented its current use in treatment regimens (18).

2. Objectives



Figure 1. The three structural components of silymarin (3)

The main objective of this research is to utilize the Molegro Virtual Docker (MVD) software to explore the potential of silibinin, silidianin, and silychristin molecules as ligands within the active site of the Aromatase enzyme and HIV-1 reverse transcriptase and examine the interactions formed by each ligand with other amino acids present in these sites.

3. Methods

Initially, the ligands silibinin (ID = 31553), silidianin (ID = 11982272), and silvchristin (ID = 441764) were obtained in the form of structure-data files (SDFs) from https://pubchem.ncbi.nlm.nih.gov/. the website Subsequently, the website http://www.rcsb.org/ was employed to search for the HIV-1 reverse transcriptase and Aromatase enzymes, identified by IDs 1JLQ and 3EQM, respectively. Afterwards, to conduct the molecular docking analysis, version 6 of MVD was installed on a Windows operating system. To perform the docking process, the downloaded proteins and ligands were first entered into the software. Water molecules were eliminated, polar hydrogen atoms were introduced, and energy minimization was conducted. Afterwards, the option for molecule preparation was accessed in the preparation tab. In the Assign all Below section, the setting was modified to "if missing," which allows the software to correct any structural errors in the molecule that may interfere with the docking process. Following that, in the same section, the option for detecting cavities was selected, leading to the choice of the molecular surface in the expanded Van Der Waals mode. The maximum number of cavities was established at 100 to pinpoint the most extensive areas available for the molecular-level docking process, and the cavities associated with the main ligand of each molecule were selected. For the docking procedure, the binding sites of the principal ligands were chosen: The 76.288 cavity for the HIV enzyme and the 304.64 cavity for the aromatase enzyme. Subsequently, the Docking wizard part was chosen from the Docking section of the software, and the imported ligands were selected to configure the parameters associated with the docking procedure. The score was calculated using the MolDock Score (GRID) mode, with the origin in the binding site section configured to the designated cavities. The algorithm employed for this procedure was MolDock simplex evolution (SE), and ten runs were conducted for each ligand. Ultimately, the software was instructed to save the poses file in Mol2 format. After the docking procedure was completed, the outcomes were reviewed (19, 20).

4. Results

The active site of the aromatase enzyme was identified through its native ligand. This cavity was measured by MVD, which perfectly accommodates the active site (Figure 2). After the docking process, the results show that silibinin possesses considerable potential to bind effectively within the aromatase active site by hydrogen and steric interactions (Figure 3), with



Figure 2. Active site of the aromatase enzyme according to the original ligand. The green area is a cavity estimated by Molegro virtual docker (MVD) (Vol = 304.64), and the stick structure is the original ligand.



Figure 3. A, Silibinin hydrogen bonds with cysteine 437, proline 429, valine 370, glutamine 367 and serine 314; B, steric interaction bonds with the other residues in the active site of aromatase.

a total energy measurement of -141.972 kcal/mol (Table 1).

The binding free energy of silychristin at the active site of the aromatase enzyme was computed using software, yielding a value of -148.425 kcal/mol (Table 1). This result suggests that silychristin exhibits a strong affinity for the active site, allowing it to compete effectively with the primary ligand for binding. The high

Table 1. MolDock Score and Rerank Score of Each Ligand with Aromatase Enzyme			
Descriptors	MolDock Score	Rerank Score	
Silibinin	-141.972	-79.431	
Siladianin	-127.805	-105.274	
Silychristin	-148.425	-113.008	



Figure 4. Steric interaction bonds of slychristin with Leu 152, Pro 368 and 429, Val 370, Met 303, 311, and 364

binding affinity is attributed to the establishment of hydrogen bonds with the amino acids cysteine 437, valine 369, serine 314, and methionine 303, along with the potential formation of steric interaction bonds with other amino acids (Figure 4), which may further enhance the binding strength.



Figure 5. Steric interaction bonds of Slidianin with: Gly 436, Arg 115, Ile 133 and Cys 437

During the molecular docking analysis of the silidianin ligand within the active site of the aromatase enzyme, it was found that this compound exhibits a notable capacity to bind to the active site as a ligand, achieving a binding energy of -127.805 kcal/mol in this site (Table 1). Figures 5 and 6 reveal the amino acids that play a crucial role in forming hydrogen and ester bonds.

During the docking procedure with the HIV-1 reverse transcriptase molecule, the software identified 41 potential cavities, among which the 76.288 cavity, recognized as the active site for the primary ligand of this enzyme, was chosen (Figure 7). All three molecules demonstrated considerable binding potential at this site. The findings indicated that the silibinin molecule could occupy this site with a free energy of -140.701 kcal/mol, establishing hydrogen bonds with the amino acids Ile 180, Tyr 318, and Arg 172 in the enzyme's A chain. Additionally, the silidianin molecule exhibited promising results, achieving a binding energy of -138.956 kcal/mol (Figure 8). Lastly, silychristin also showed effective binding by forming hydrogen bonds with Tyr 318 in the A chain and Glu 138 in the B chain, along with other ester bonds in this site (Figure 9), resulting in a free energy of -135.365 kcal/mol (Table 2).

5. Discussion



Figure 6. Four amino acids: Cysteine 437, arginine 115, proline 429, and tyrosine 310 have interacted with the Slidianin molecule through the formation of hydrogen bonds.

The MVD is a molecular docking software utilized in numerous studies owing to its exceptional accuracy, affordability, and rapid processing capabilities (21). This software can automatically detect potential ligand binding sites, which can be modified within the software settings (20). Furthermore, users can select between two algorithms, the MolDock optimizer and the MolDock SE, based on their specific requirements. The primary distinction between these algorithms lies in the capacity to recombine poses, a feature that is exclusive to the SE mode. Users have the option to utilize if-missing statement to rectify potential the inaccuracies in their ligand or receptor structures. This approach was employed in the current study, despite the absence of any errors in the ligand and enzyme structures examined.

Recognizing the significance of this software, Kusumaningrum and colleagues utilized it to conduct molecular docking of NO11 (a modified naphthoquinone) as a potential inhibitor of Polo-like kinase 1 (Plk1), which is associated with various cancers. Their findings indicated that NO11 did not establish any hydrogen bonds with Plk1. Nevertheless, the software calculated that the MolDock score for compound NO11 is -134.73 kcal/mol, attributed to the steric interaction bonds formed with Phe 133, Asp 194, Glu 101, Lys 82, Cys 133, and Glu 140 of Plk1 (20).

In a separate research study, Naeem and associates examined chlorogenic acid as a ligand targeting aldose reductase, a crucial enzyme in the polyol pathway that aids in the transformation of glucose into sorbitol with the involvement of NADPH, which is linked to



Figure 7. From 41 cavities, the 76.288 cavity in the active site was selected for molecular docking (yellow = original ligands; green = caviteies).



Figure 8. A, Steric interaction bonds; and B, hydrogen bonds of Siladianin in the active site of human immunodeficiency virus-1 (HIV-1) reverse transcriptase

complications arising from diabetes (19, 22). The findings indicated that chlorogenic acid is capable of establishing hydrogen bonds with the amino acids Tyr 48, His 110, and Trp 111 located in the enzyme's active site. The average MolDock score for chlorogenic acid was -119.34 kcal/mol, while the MolDock Re-rank score was estimated at -114.92 kcal/mol (19).

5.1. Conclusions

This research utilized three molecules — silibinin, silidianin, and silychristin — derived from the seeds and fruits of silymarin (3-5), as effective ligands to explore their potential binding effects on HIV-1 reverse transcriptase and the Aromatase enzyme. Numerous studies have demonstrated that silymarin offers protection to liver cells from ischemic damage and radiation exposure (3). Furthermore, it has exhibited antioxidant properties in various investigations (3, 6). Consequently, this study employed these compounds in a docking process, revealing their potential as suitable ligands for binding to the active sites of both HIV-1 reverse transcriptase and the Aromatase enzyme, thereby destroying their enzymatic activities. The inhibition of the Aromatase enzyme, which is crucial in the context of cancer cell proliferation, could serve as a foundation for developing drugs based on silymarin compounds. Additionally, these molecules may facilitate the creation of potent medications aimed at inactivating the HIV-1 reverse transcriptase in infected cells, thereby preventing viral replication and



Figure 9. Steric interaction bonds of Silychristin in the active site of human immunodeficiency virus-1 (HIV-1) reverse transcriptase

Table 2. MolDock Score and Rerank Score of Each Ligand Human Immunodeficiency Virus-1 Reverse Transcriptase Molecule			
Descriptors	MolDock Score	Rerank Score	
Silibinin	-140.701	-92.856	
Siladianin	-138.956	-16.734	
Silychristin	-135.365	-67.577	

addressing the global HIV epidemic, which affects approximately one million individuals. While the findings of this study suggest that these molecules can bind to the active site of the enzymes, additional in vivo research is necessary to verify the accurate binding of these ligands.

Footnotes

Authors' Contribution: Analysis and interpretation of data: A. H.; Study concept and design: Z. Kh.

Conflict of Interests Statement: The authors declared no conflict of interest.

Data Availability: The dataset presented in the study is available on request from the corresponding author during submission or after publication. The data are not publicly available due to being published first in this journal.

Funding/Support: The present study received no funding/support.

References

- 1. Ramasamy K, Agarwal R. Multitargeted therapy of cancer by silymarin. Cancer Lett. 2008;269(2):352-62. [PubMed ID: 18472213].

 [PubMed Central ID: PMC2612997].

 https://doi.org/10.1016/j.canlet.2008.03.053.
- Saller R, Melzer J, Reichling J, Brignoli R, Meier R. An updated systematic review of the pharmacology of silymarin. *Forsch Komplementmed*. 2007;14(2):70-80. [PubMed ID: 17464157]. https://doi.org/10.1159/000100581.
- 3. Fraschini F, Demartini G, Esposti D. Pharmacology of Silymarin. *Clinical Drug Investigation*. 2002;**22**(1):51-65. https://doi.org/10.2165/00044011-200222010-00007.
- Al-Moziel MS, Kadhum HJ, Ghanim WK. Effects of Tadalafil and/or Silymarin On Gonadal Function in Adults Male Albino Rats. *BasJVetRes*. 2018;17(2):147-60.
- 5. Khalil EA. Hormonal Profile And Histopathological Study On The Influence Of Silymarin On Both Female And Male Albino Rats. *The Egyptian J Hospital Med.* 2003;**13**(1):112-22. https://doi.org/10.21608/ejhm.2003.18236.
- Surai PF. Silymarin as a Natural Antioxidant: An Overview of the Current Evidence and Perspectives. *Antioxidants (Basel)*. 2015;4(1):204-47. [PubMed ID: 26785346]. [PubMed Central ID: PMC4665566]. https://doi.org/10.3390/antiox4010204.
- Eghbali A, Sadeghian M, Ghasemi A, Afzal RR, Eghbali A, Ghaffari K. Effect of oral silymarin on liver function in pediatric acute lymphoblastic leukemia in the maintenance phase: a double-blind randomized clinical trial. *Front Pharmacol.* 2024;**15**:1295816. [PubMed ID: 38283627]. [PubMed Central ID: PMC10811082]. https://doi.org/10.3389/fphar.2024.1295816.
- Karimi G, Vahabzadeh M, Lari P, Rashedinia M, Moshiri M. "Silymarin", a promising pharmacological agent for treatment of diseases. *Iran J Basic Med Sci.* 2011;14(4):308. [PubMed ID: 23492971]. [PubMed Central ID: PMC3586829].
- Akheratdoost V, Panahi N, Safi S, Mojab F, Akbari G. Protective effects of silymarin-loaded chitosan nanoparticles in the diet-induced hyperlipidemia rat model. *Iran J Basic Med Sci.* 2024;27(6):725-32. [PubMed ID: 38645495]. [PubMed Central ID: PMC11024415]. https://doi.org/10.22038/IJBMS.2024.74490.16179.

- Abedi H, Jahromi HK, Hashemi SMA, Jashni HK, Jahromi ZK, Pourahmadi M. The effect of silymarin on spermatogenesis process in rats. *Int J Med Res Health Sci.* 2016;5(6):146-50.
- Boon WC, Chow JD, Simpson ER. The multiple roles of estrogens and the enzyme aromatase. *Prog Brain Res.* 2010;**181**:209-32. [PubMed ID: 20478440]. https://doi.org/10.1016/S0079-6123(08)81012-6.
- 12. Sayyad NB, Sabale PM, Umare MD, Bajaj KK. Aromatase Inhibitors: Development and Current Perspectives. *Indian J Pharmaceutical Edu* Res. 2022;**56**(2):311-20. https://doi.org/10.5530/ijper.56.2.51.
- Negm S, Aboraya A. Therapeutic Effects of Milk thistle Seeds (Silybum marianum) and Red Ginseng roots on Polycystic Ovary Syndrome Induced by Letrozole in Female Rats. *EKB J Manage System*. 2023;**2023**(75):199-234. https://doi.org/10.21608/mbse.2023.188764.1298.
- Mirzaei N, Jahanian Sadatmahalleh S, Rouholamin S, Nasiri M. A randomized trial assessing the efficacy of Silymarin on endometrioma-related manifestations. *Sci Rep.* 2022;**12**(1):17549. [PubMed ID: 36266431]. [PubMed Central ID: PMC9584967]. https://doi.org/10.1038/s41598-022-22073-8.
- Jouhari S, Mohammadzadeh A, Soltanghoraee H, Mohammadi Z, Khazali S, Mirzadegan E, et al. Effects of silymarin, cabergoline and letrozole on rat model of endometriosis. *Taiwan J Obstet Gynecol.* 2018;57(6):830-5. [PubMed ID: 30545536]. https://doi.org/10.1016/j.tjog.2018.10.011.
- Koksal E, Gulcin I, Beyza S, Sarikaya O, Bursal E. In vitro antioxidant activity of silymarin. J Enzyme Inhib Med Chem. 2009;24(2):395-405. [PubMed ID: 18830883]. https://doi.org/10.1080/14756360802188081.
- McClure J, Lovelace ES, Elahi S, Maurice NJ, Wagoner J, Dragavon J, et al. Silibinin inhibits HIV-1 infection by reducing cellular activation and proliferation. *PLoS One*. 2012;7(7). e41832. [PubMed ID: 22848626]. [PubMed Central ID: PMC3404953]. https://doi.org/10.1371/journal.pone.0041832.
- Molto J, Valle M, Miranda C, Cedeno S, Negredo E, Clotet B. Effect of milk thistle on the pharmacokinetics of darunavir-ritonavir in HIVinfected patients. *Antimicrob Agents Chemother*. 2012;**56**(6):2837-41. [PubMed ID: 22430963]. [PubMed Central ID: PMC3370734]. https://doi.org/10.1128/AAC.00025-12.
- Naeem S, Hylands P, Barlow D. Docking studies of chlorogenic acid against aldose redutcase by using molgro virtual docker software. J Applied Pharmaceut Sci. 2013;3(1):13-20.
- 20. Kusumaningrum S, Budianto E, Kosela S, Sumaryono W, Juniarti F. The molecular docking of 1, 4-naphthoquinone derivatives as inhibitors of Polo-like kinase 1 using Molegro Virtual Docker. J Applied Pharmaceut Sci. 2014;4(11):47-53.
- Thomsen R, Christensen MH. MolDock: a new technique for highaccuracy molecular docking. *J Med Chem.* 2006;49(11):3315-21. [PubMed ID: 16722650]. https://doi.org/10.1021/jm051197e.
- Yabe-Nishimura C. Aldose reductase in glucose toxicity: a potential target for the prevention of diabetic complications. *Pharmacol Rev.* 1998;**50**(1):21-33. [PubMed ID: 9549756].