



The Relationship Between Retained Primitive Reflexes and Motor Proficiency of Elementary School Children

Soraya Goodarzi ¹, Hasan Khalaji ², Jalil Moradi ^{2,*}

¹ Arak University, Arak, Iran

² Department of Motor Behavior and Sport Psychology, Faculty of Sport Sciences, Arak University, Arak, Iran

*Corresponding Author: Department of Motor Behavior and Sport Psychology, Faculty of Sport Sciences, Arak University, Arak, Iran. Email: j-moradi@araku.ac.ir

Received: 25 November, 2024; Revised: 25 April, 2025; Accepted: 27 April, 2025

Abstract

Background: The asymmetrical tonic neck reflex (ATNR), symmetrical tonic neck reflex (STNR), and tonic labyrinthine reflex (TLR) are essential primitive reflexes for infant motor development, and their lack of integration can significantly impact a child's motor skills and overall development.

Objectives: The present study aimed to investigate the relationship between retained primitive reflexes and motor proficiency in elementary school children.

Methods: This study involved 110 primary school children aged 6 to 9 years (57 girls and 53 boys) who were randomly selected. Motor proficiency was assessed using the Bruininks-Oseretsky test, and retained reflexes were measured with the profile test of primary reflexes. Spearman's rank correlation coefficient was used to examine relationships between variables, while independent samples *t*-test and Mann-Whitney U test were employed to compare mean values between girls and boys, with significance set at $P \leq 0.05$.

Results: The findings indicated a significant negative correlation between retained ATNR and fine motor skills, as well as between retained STNR and fine motor skills. Additionally, no significant difference was found in motor proficiency between girls and boys.

Conclusions: The results suggest that reducing retained primitive reflexes may improve children's motor proficiency in both gross and fine motor tasks. Given the negative relationship between retained primitive reflexes and motor skills, early identification of these reflexes could enable timely intervention.

Keywords: Early Intervention, Motor Skills, Primitive Reflexes, Tonic Labyrinthine Reflex, Gross Motor Skills, Fine Motor Skills

1. Background

It is well known that primitive reflexes, which play a crucial role in infant survival and development, emerge by the 25th week of gestation. These reflexes – including spinal and brainstem reflexes – develop during the fetal period and typically integrate fully by 12 months of age (1). Primitive reflexes facilitate essential functions such as navigating the birth canal, initiating breathing, avoiding harmful stimuli, urination, crawling, grasping, head control, feeding (sucking/swallowing), and kicking (2). A critical question arises: Why are infants born with these reflexes? For skilled movement development,

primitive reflexes must fade or integrate, allowing voluntary motor control to replace them (3).

The timing of primitive reflex appearance and disappearance serves as a key marker of typical neurodevelopment, with deviations indicating potential atypical development (4). Persistence beyond the expected age suggests central nervous system immaturity and is associated with developmental delays (5, 6). This study focuses on retained primary reflexes – asymmetrical tonic neck reflex (ATNR), symmetrical tonic neck reflex (STNR), and tonic labyrinthine reflex (TLR) – in children aged 6 - 9 years (7). Prior research reports high prevalence rates of retained reflexes in this population (8, 9), with ATNR linked to gait asymmetry

(10) and even healthy school-aged children often exhibiting at least one active primitive reflex, which may impair motor coordination and academic performance (11). Persistent reflexes during preschool and school years can disrupt neuromotor maturation, hindering overall motor skill development (12).

The ATNR is an involuntary response to head rotation: When the head turns, the limbs on the same side extend while the opposite limbs flex. Retention beyond 6 months may impair movement, balance, and hand-eye coordination (13). The TLR, present at birth, should integrate by 2 - 4 months; persistence compromises postural stability and vertical balance (14). The STNR is triggered by head flexion/extension: Flexion bends the upper limbs and extends the lower limbs, while extension reverses this pattern. Failure to inhibit STNR disrupts crawling coordination (5), sitting posture, and hand-eye control (15).

A key concern is the relationship between retained primitive reflexes and motor proficiency. Motor proficiency – reflected in the quality of gross and fine motor skills – serves as a benchmark for children's motor competence (16, 17). Gross motor skills (e.g., walking, jumping) involve large muscle groups, whereas fine motor skills (e.g., writing, playing the piano) require precise small-muscle coordination (18).

Proper primitive reflex integration is foundational for later motor skill acquisition. Retained reflexes may impede milestones like rolling, crawling, and walking (3). Understanding their impact on sensorimotor function enables accurate diagnosis and enhances treatment efficacy (19). Moreover, early motor competence establishes the groundwork for lifelong physical activity (20).

Relevant studies highlight that reflex integration techniques improve motor outcomes in children with spastic cerebral palsy (13). In typically developing preschoolers, higher reflex activity correlates with lower motor efficiency, underscoring the need for early screening to mitigate academic and social challenges (14). Targeted intervention may reduce developmental delays and improve cognitive-motor outcomes (14). Poorly integrated reflexes are also associated with task-specific difficulties, such as reading analog clocks (21). However, systematic reviews note limited evidence linking primitive reflexes to motor and learning deficits in preschoolers, calling for further research (22).

2. Objectives

Limited research has investigated the persistence of primitive reflexes and their association with motor difficulties in typically developing children, as most

studies have prioritized academic and social outcomes over motor skill assessment (1). Existing literature primarily examines the prevalence and clinical implications of primitive reflexes in children with neurodevelopmental disorders (13, 23, 24). Elucidating the link between retained primitive reflexes and motor proficiency could identify early markers of developmental delay and guide the design of targeted screening and intervention programs for motor deficits in educational and clinical settings (12). Consequently, this study aimed to explore this relationship in neurotypical children.

3. Methods

3.1. Subjects

This descriptive correlational study examined 110 typically developing children (57 girls, 53 boys) aged 6 - 9 years, randomly selected from Arak public schools. The sample size was determined via G*Power (v1.3), requiring 112 participants ($\alpha = 0.05$, power = 0.95, $R = 0.3$). Inclusion criteria comprised: (1) No special education/sports training beyond school activities, (2) normal medical history without developmental delays, prematurity (< 37 weeks), cerebral palsy, ADHD, or Developmental Coordination Disorder (DCD) (verified through records). From an initial pool of 130 children, 20 were excluded for ADHD (5 boys and 5 girls)/DCD (7 boys and 3 girls), yielding the final sample.

3.2. Apparatus and Task

3.2.1. Primitive Reflexes

Evaluation methods of Blythe were used to evaluate reflexes and diagnose children's motor neurodevelopment. To assess the ATNR, the child stands with arms extended forward at shoulder level and wrists relaxed. With the child's eyes closed, the examiner slowly rotates the child's head to each side so the chin is parallel with the shoulder, holding each position for 5 - 10 seconds before returning to the center. This process is repeated four times on each side (25). To assess the STNR, the child is placed in a four-point kneeling position, and the examiner gives the instructions and asks the child to slowly bend the head; after a 5-second pause, the head is slowly moved upward. All steps are repeated six times (25). To assess the TLR, the child is placed in a standing position with legs next to each other and hands next to the body. Then the child is asked to close the eyes and extend the head; then after 10 seconds, slowly flexes the head. All steps are repeated 4 times. The response is

scored from 0 (non-retained) to 4 (completely retained) after each reflection (25).

Goddard-Blythe's reflex assessment methods are widely recognized in the field of neurological research for their systematic approach to the evaluation of primitive reflexes. These methods are based on decades of research on the role of primitive reflexes in the development of children and their impact on motor, cognitive, and behavioral results. This instrument has been used in many studies and has acceptable validity and reliability (12, 14).

3.2.2. Motor Proficiency

The motor skills of children were assessed using the Bruininks-Oseretsky Test of Motor Proficiency Second Edition (16). The four composites along with the subtests are: (1) Fine manual control, including (A) fine motor precision and (B) fine motor integration; (2) Manual coordination, including (A) manual dexterity and (B) upper limb coordination; (3) Body coordination, including (A) bilateral coordination and (b) balance; (4) Strength and agility, including (A) speed and (B) strength. The short form of BOT-2 contains 14 items, which include five fine motor skills and nine gross motor skills (26-28). According to the instructions of the Bruininks-Oseretsky motor skills manual, how to perform each subtest was explained to each student.

3.2.3. Developmental Coordination Disorder Questionnaire for Parents

The parent-completed DCD Questionnaire has 15 items assessing motor control, general coordination, and fine movements on a 5-point scale. It is used to identify and exclude children prone to DCD from the sample. In the end, according to the points obtained, children are classified as either prone to DCD or without DCD based on their scores (29).

3.2.4. Conners' Parent Rating Scale

To identify children with ADHD, the Conners' Parent Rating Scale (CPRS) was used. This scale consists of 48 items, which evaluate the Hyperactivity Index according to a 4-point Likert scale (30).

3.3. Procedure

Before starting the research, the parents attended a briefing session, and the purpose of the research was explained to them. The parents completed the consent form, the CPRS, and the DCD Questionnaire. Afterwards, the reflex evaluation test and then the BOT-2 test were administered. The testing environment was secure, safe,

and completely silent, and each individual child was tested for 30 - 40 minutes.

3.4. Data Analysis

The data were analyzed using SPSS 26, with a significance level set at $P \leq 0.05$. The variable of motor proficiency and gross motor tasks of boys had a normal distribution, while other variables, including fine motor tasks, gross motor tasks of girls, and all three reflexes (ATNR, STNR, and TLR), had a non-normal distribution. As a result, non-parametric tests such as Spearman's correlation and the Mann-Whitney U test were used, alongside the independent samples *t*-test, to analyze relationships and mean differences between boys and girls.

4. Results

In Table 1, we discussed the demographic and descriptive characteristics of male and female students.

4.1. Prevalence of Reflex Activity

The prevalence of primitive reflexes was analyzed based on response scores (range: 0 - 4) among male and female students. Across all participants, ATNR showed the highest reflex activity (65.4%), while STNR exhibited the lowest (63.6%). Mann-Whitney U tests revealed statistically significant gender differences in STNR prevalence ($U = 1099$, $P = 0.004$), with boys demonstrating higher mean retained primitive reflex scores than girls. Gender-specific analysis showed ATNR as the most prevalent reflex in both boys (62.3%) and girls (68.4%), whereas STNR showed the lowest prevalence (boys: 49.1%; girls: 24.6%), indicating particularly low retention among female students.

4.2. Motor Proficiency

An independent samples *t*-test revealed no significant gender differences in overall motor proficiency [$t(108) = -1.02$, $P = 0.308$]. The Mann-Whitney U test was utilized to examine the mean difference of fine and gross motor skills as well as the subscales of the BOT-2 test between boys and girls. While no significant differences emerged for the fine ($U = 1498$, $P = 0.943$) or gross motor ($U = 1199$, $P = 0.062$) scores, there was a significant difference in the subscales of Dribbling a Ball ($U = 993$, $P = 0.002$), Tapping Feet and Fingers ($U = 1206$, $P = 0.033$), and Knee Push-ups ($U = 1078$, $P = 0.008$).

4.3. Correlation between Retained Primitive Reflexes and Motor Proficiency by Gender

Table 1. Demographic and Descriptive Characteristics of Participants ^a

Variables	Boys (N = 53)	Girls (N = 57)
Age (y)	7.87 ± 0.94	84 ± 0.90.7
Weight (kg)	28.00 ± 6.23	39 ± 7.57.27
Height (m)	128.42 ± 9.29	05 ± 7.53.127
Gross motor skill	29.23 ± 8.22	49 ± 6.34.26
Fine motor skill	23.25 ± 4.45	33 ± 4.47.23
Motor proficiency	51.75 ± 10.90	82 ± 9.35.49
ATNR	1.30 ± 1.21	40 ± 1.19.1
STNR	0.89 ± 1.17	33 ± 0.69.0
TLR	0.89 ± 1.08	88 ± 1.15.0

Abbreviations: ATNR, asymmetrical tonic neck reflex; STNR, symmetrical tonic neck reflex; TLR, tonic labyrinthine reflex.

^a Values are expressed as mean ± SD.

Table 2 shows the relationship between retained primitive reflexes and the subscales of BOT-2. The results indicate that in both girls and boys, there are significant negative correlations between the persistence of primitive reflexes and motor proficiency. In particular, reducing retained reflexes is associated with higher motor proficiency scores, affecting both fine and gross motor skills.

5. Discussion

Retained primitive reflexes can disrupt the development of voluntary motor control, leading to issues with movement patterns, coordination, and timing in children (12, 14). This study aimed to investigate the relationship between retained primitive reflexes and motor proficiency among elementary school children and found that boys had higher levels of the STNR reflex than girls. Overall, there was a significant negative relationship between the persistence of these reflexes and both fine and gross motor skills, indicating that retained reflexes are associated with lower motor proficiency in children.

The key finding of this study is the significant negative relationship between the persistence of reflexes and motor skills. This result is consistent with the findings of Pecuch et al. (12) and Callcott (31). In this regard, Callcott (8) found that 65% had moderate to high levels of persistent ATNR reflexes, and this retention was linked with lower motor and academic skills. If primitive reflexes are not properly integrated as the child grows, a range of functions – including cognitive, academic, social, and motor abilities – can be negatively affected (12). Richards et al. (32) also concluded that children with the persistence of primitive reflexes are

likely to have lower typical word and letter legibility scores.

The findings show that ATNR is the most commonly retained primitive reflex among students (65.4%), while STNR is the least common (63.6%), with previous research supporting ATNR's high prevalence and STNR's lower prevalence in children (14). A high rate of retained reflexes suggests delayed integration, highlighting the importance of assessment by professionals and the need for reflex integration therapy. Such therapy, using targeted exercises, can improve motor skills, coordination, and balance, ultimately supporting children's broader development (12).

Previous studies have pointed out differences between boy and girl participants and acknowledged that retained primitive reflexes are higher in boys than in girls (19, 31). Our findings also partially confirm these results, although only the STNR reflex was statistically significant; boys had a higher level of motor proficiency than girls. Therefore, while it has been shown that the presence of some reflexes is associated with a decrease in motor skills subtests in boys, still, their performance is better than in girls.

Early motor control theories posited that persistent primitive reflexes may contribute to developmental delays, with empirical evidence linking uninhibited reflexes to learning challenges and underscoring the critical need for early intervention (33). Our findings align with this literature, demonstrating that children with retained primary reflexes exhibit weaker motor proficiency. This supports the dual importance of reflex integration programs and targeted motor skills training, both of which have demonstrated efficacy in improving motor coordination and emotional regulation (12, 14). Notably, our results suggest that heightened primitive reflex activity may impair sensory-

Table 2. Correlation Between Retained Primitive Reflexes and the Subscales of BOT-2 of Elementary School Children by Gender

Variables	Boys (N = 53)			Girls (N = 57)			Total Students (N = 110)		
	ATNR	STNR	TLR	ATNR	STNR	TLR	ATNR	STNR	TLR
Fine motor skills									
Drawing, lines through paths-crooked	-0.204	-0.191	-0.103	-0.115	-0.184	-0.183	-0.160	-0.153	-0.141
Folding Paper	-0.384 ^a	-0.152	-0.052	-0.226	-0.149	-0.179	-0.293 ^a	-0.183	-0.120
Copying a square	-0.210	0.059	-0.162	-0.230	-0.057	-0.296 ^b	-0.224 ^a	0.044	-0.226 ^b
Copying a star	-0.254	-0.435 ^a	-0.121	-0.160	-0.067	-0.122	-0.199 ^b	-0.244	-0.118
Transferring pennies	-0.282 ^b	-0.307	0.044	0.095	-0.104	-0.101	-0.115	-0.202 ^b	-0.035
Total fine motor	-0.388 ^b	-0.343 ^b	-0.141	-0.192	-0.150	-0.177	-0.295 ^a	-0.242	-0.162
Gross motor skills									
Dropping and catching a ball-both hands	-0.188	-0.271 ^b	-0.078	-0.121	-0.378 ^a	-0.443 ^a	-0.168	-0.244 ^b	-0.267 ^b
Dribbling a ball-alternating hand	-0.297 ^b	-0.546 ^a	-0.369 ^a	-0.072	-0.147	-0.136	-0.199 ^b	-0.253 ^a	-0.226 ^b
Jumping in place-same sides synchronized	-0.192	-0.136	-0.009	-0.198	-0.329 ^b	-0.374 ^a	-0.197 ^b	-0.195 ^b	-0.199 ^b
Tapping feet and fingers-same sides synchronize	-0.319 ^b	-0.205	-0.045	-0.206	-0.196	-0.317 ^b	-0.251 ^b	-0.257 ^a	-0.176
Walking forward on a line	0.138	-0.019	0.001	-0.069	-0.468 ^a	-0.288 ^b	0.054	-0.193	-0.122
Standing on one leg on a balance beam-eyes open	-0.125	-0.086	-0.197	-0.025	-0.381 ^a	-0.354 ^a	-0.081	-0.224	-0.268 ^a
One-legged stationary hop	-0.010	-0.164	-0.083	-0.054	-0.251	-0.073	-0.032	0.209	-0.080
Knee push-ups	-0.202	-0.176	-0.150	-0.255	-0.113	-0.285 ^b	-0.237 ^b	-0.073	-0.213 ^b
Sit-ups	-0.344 ^a	-0.171	-0.087	-0.254	-0.011	-0.053	-0.296 ^a	-0.055	-0.019
Total gross motor	-0.380 ^a	-0.410 ^a	-0.221	-0.258	-0.341 ^a	-0.315 ^b	-0.321 ^a	-0.302 ^a	-0.266 ^a
Motor proficiency	-0.415 ^a	-0.421 ^a	-0.232	-0.283 ^b	-0.311 ^b	-0.301 ^b	-0.359 ^b	-0.311 ^a	-0.268 ^a

Abbreviations: ATNR, asymmetrical tonic neck reflex; STNR, symmetrical tonic neck reflex; TLR, tonic labyrinthine reflex.

^a Correlation is significant at the 0.01 level (2-tailed).

^b Correlation is significant at the 0.05 level (2 tailed).

motor development, potentially affecting multiple domains including motor competence, social functioning, and academic achievement. These findings emphasize the necessity of comprehensive diagnostic approaches that identify underlying causes rather than solely addressing symptomatic behaviors, thereby enabling more effective intervention strategies (19).

Given that primitive reflexes constitute the neurodevelopmental foundation for subsequent motor milestones (34), their assessment represents a valuable screening tool for identifying at-risk children. Several limitations warrant consideration. First, our sample (N = 110) was drawn from a single school, potentially limiting generalizability. Second, the study focused exclusively on three primary reflexes (ATNR, STNR, and TLR); future investigations should incorporate a broader array of primitive reflexes. We recommend subsequent research with larger, more diverse samples to enhance external validity. Furthermore, in light of our findings and previous research demonstrating an inverse relationship between primitive reflex retention and motor proficiency, future studies should investigate the

efficacy of combined reflex integration protocols and motor skill interventions. Such investigations could establish evidence-based practices for addressing these neurodevelopmental concerns.

Footnotes

Authors' Contribution: S. G., H. K., and J. M. designed the study. S. G. and H. K. collected and analyzed data. J. M. and S. G. wrote the manuscript. All authors read and approved the final manuscript.

Conflict of Interests Statement: The authors declare no conflict of interest.

Data Availability: The dataset presented in the study is available on request from the corresponding author during submission or after publication.

Ethical Approval: The ethical number is [IR.ARAKU.REC.1401.047](#), which was approved by the Ethics Committee of the Arak University, Arak, Iran. All

the methods included in this study are in accordance with the declaration of Helsinki.

Funding/Support: The present study received no funding/support.

Informed Consent: All the parents have completed the informed consent form before the study.

References

1. Zafeiriou DI. Primitive reflexes and postural reactions in the neurodevelopmental examination. *Pediatr Neurol.* 2004;**31**(1):1-8. [PubMed ID: 15246484]. <https://doi.org/10.1016/j.pediatrneurol.2004.01.012>.
2. McDonald KG. *Integrating primitive reflexes through play and exercise: An interactive guide to the asymmetrical tonic neck reflex (ATNR)*. New York, US: Polaris Therapy; 2020.
3. Haywood K, Getchell N. *Life span motor development*. Illinois, USA: Human kinetics; 2024.
4. Futagi Y, Toribe Y, Suzuki Y. The grasp reflex and moro reflex in infants: hierarchy of primitive reflex responses. *Int J Pediatr.* 2012;**2012**:191562. [PubMed ID: 22778756]. [PubMed Central ID: PMC3384944]. <https://doi.org/10.1155/2012/191562>.
5. Goddard Blythe S, Duncombe R, Preedy P, Gorely T. Neuromotor readiness for school: the primitive reflex status of young children at the start and end of their first year at school in the United Kingdom. *Education* 3-13. 2021;**50**(5):654-67. <https://doi.org/10.1080/03004279.2021.1895276>.
6. Schott JM, Rossor MN. The grasp and other primitive reflexes. *J Neurol Neurosurg Psychiatry.* 2003;**74**(5):558-60. [PubMed ID: 12700289]. [PubMed Central ID: PMC1738455]. <https://doi.org/10.1136/jnnp.74.5.558>.
7. Blythe SG, Hyland D. Screening for Neurological Dysfunction in the Specific Learning Difficulty Child. *British Journal of Occupational Therapy.* 1998;**61**(10):459-64. <https://doi.org/10.1177/030802269806101008>.
8. Callcott D. *The effect of a reflex replication program on retained primary reflexes, motor coordination, vocabulary, visual motor ability and rapid naming in preprimary aged children [dissertation]*. Edith Cowan University; 2008.
9. Gieysztor E, Sadowska L, Choińska A. The degree of primitive reflexes integration as a diagnostic tool to assess the neurological maturity of healthy preschool and early school age children. *Nursing and Public Health.* 2017;**7**(1):5-11. <https://doi.org/10.17219/pzp/69471>.
10. Gieysztor E, Pecuch A, Kowal M, Borowicz W, Paprocka-Borowicz M. Pelvic Symmetry Is Influenced by Asymmetrical Tonic Neck Reflex during Young Children's Gait. *Int J Environ Res Public Health.* 2020;**17**(13). [PubMed ID: 32630679]. [PubMed Central ID: PMC7370024]. <https://doi.org/10.3390/ijerph17134759>.
11. Feldhacker DR, Cosgrove R, Feiten B, Schmidt K, Stewart M. The Correlation between Retained Primitive Reflexes and Scholastic Performance among Early Elementary Students. *Journal of Occupational Therapy, Schools, & Early Intervention.* 2021;**15**(3):288-301. <https://doi.org/10.1080/19411243.2021.1959482>.
12. Pecuch A, Gieysztor E, Wolanska E, Telenga M, Paprocka-Borowicz M. Primitive Reflex Activity in Relation to Motor Skills in Healthy Preschool Children. *Brain Sci.* 2021;**11**(8). [PubMed ID: 34439585]. [PubMed Central ID: PMC8394673]. <https://doi.org/10.3390/brainsci11080967>.
13. Wagh SC, Malawade MR, Vardharajulu G. Effect of Specific Reflex Integration Approach on Primitive Reflexes in Spastic Cerebral Palsy Children. *Int. J. Health Sci. Res.* 2019;**9**:87-93.
14. Gieysztor EZ, Choinska AM, Paprocka-Borowicz M. Persistence of primitive reflexes and associated motor problems in healthy preschool children. *Arch Med Sci.* 2018;**14**(1):167-73. [PubMed ID: 29379547]. [PubMed Central ID: PMC5778413]. <https://doi.org/10.5114/aoms.2016.60503>.
15. O'Dell NE, Cook P. *Stopping ADHD*. Penguin; 2004.
16. Bruininks RH, Bruininks BD. Bruininks-Oseretsky test of motor proficiency. Circle Pines, MN: American Guidance Service; 1978.
17. Venetsanou F, Kambas A. Motor Proficiency in Young Children. *Sage Open.* 2016;**6**(1). <https://doi.org/10.1177/2158244015626226>.
18. Wang L, Wang L. Relationships between Motor Skills and Academic Achievement in School-Aged Children and Adolescents: A Systematic Review. *Children (Basel).* 2024;**11**(3). [PubMed ID: 38539371]. [PubMed Central ID: PMC10969452]. <https://doi.org/10.3390/children11030336>.
19. Pecuch A, Gieysztor E, Telenga M, Wolanska E, Kowal M, Paprocka-Borowicz M. Primitive Reflex Activity in Relation to the Sensory Profile in Healthy Preschool Children. *Int J Environ Res Public Health.* 2020;**17**(21). [PubMed ID: 33172138]. [PubMed Central ID: PMC7664452]. <https://doi.org/10.3390/ijerph17218210>.
20. Loprinzi PD, Davis RE, Fu YC. Early motor skill competence as a mediator of child and adult physical activity. *Prev Med Rep.* 2015;**2**:833-8. [PubMed ID: 26844157]. [PubMed Central ID: PMC4721422]. <https://doi.org/10.1016/j.pmedr.2015.09.015>.
21. Kalembe A, Lorent M, Blythe SG, Gieysztor E. The Correlation between Residual Primitive Reflexes and Clock Reading Difficulties in School-Aged Children-A Pilot Study. *Int J Environ Res Public Health.* 2023;**20**(3). [PubMed ID: 36767689]. [PubMed Central ID: PMC9915247]. <https://doi.org/10.3390/ijerph20032322>.
22. McWhirter K, Steel A, Adams J. The association between learning disorders, motor function, and primitive reflexes in pre-school children: A systematic review. *J Child Health Care.* 2024;**28**(2):402-28. [PubMed ID: 35830652]. <https://doi.org/10.1177/1367493522114187>.
23. S DF, G DA, Bilbilaj DS. Measuring Primitive Reflexes in Children with Learning Disorders. *European Journal of Multidisciplinary Studies.* 2017;**5**(1). <https://doi.org/10.26417/ejms.v5i1.p285-298>.
24. Konicarova J, Bob P, Raboch J. Persisting primitive reflexes in medication-naïve girls with attention-deficit and hyperactivity disorder. *Neuropsychiatr Dis Treat.* 2013;**9**:1457-61. [PubMed ID: 24092983]. [PubMed Central ID: PMC3788695]. <https://doi.org/10.2147/NDT.S49343>.
25. Blythe SG. *Assessing neuromotor readiness for learning: The INPP developmental screening test and school intervention programme*. Hoboken, NJ: John Wiley & Sons; 2024.
26. Carmosino K, Grzeszczak A, McMurray K, Olivo A, Slutz B, Zoll B, et al. Test items in the complete and short forms of the BOT-2 that contribute substantially to motor performance assessments in typically developing children 6-10 years of age. *Journal of Student Physical Therapy Research.* 2014;**7**(2).
27. Goddard S, Blythe SG. *The well balanced child: Movement and early learning*. Victoria, Australia: Hawthorn; 2005.
28. Radanovic D, Dordevic D, Stankovic M, Pekas D, Bogataj S, Trajkovic N. Test of Motor Proficiency Second Edition (BOT-2) Short Form: A Systematic Review of Studies Conducted in Healthy Children. *Children (Basel).* 2021;**8**(9). [PubMed ID: 34572219]. [PubMed Central ID: PMC8471722]. <https://doi.org/10.3390/children8090787>.
29. Wilson BN, Crawford SG, Green D, Roberts G, Aylott A, Kaplan BJ. Psychometric properties of the revised Developmental Coordination Disorder Questionnaire. *Phys Occup Ther Pediatr.* 2009;**29**(2):182-202. [PubMed ID: 19401931]. <https://doi.org/10.1080/01942630902784761>.

30. Pendergast LL, Vandiver BJ, Schaefer BA, Cole PM, Murray-Kolb LM, Christian P. Factor Structure of Scores from the Conners' Rating Scales-Revised Among Nepali Children. *Int J Sch Educ Psychol*. 2014;**24**:261-70. [PubMed ID: 25574454]. [PubMed Central ID: PMC4283941]. <https://doi.org/10.1080/21683603.2013.878678>.
31. Callcott D. Retained primary reflexes in preprimary-aged Indigenous children: The effect on movement ability and school readiness. *Australasian Journal of Early Childhood*. 2012;**37**(2):132-40. <https://doi.org/10.1177/183693911203700218>.
32. Richards L, Avery R, Gray S, Price R. Relationship of Retained Primitive Reflexes and Handwriting Difficulty in Elementary-Age Children. *The American Journal of Occupational Therapy*. 2022;**76**(Supplement_1):7610505010p1. <https://doi.org/10.5014/ajot.2022.76SI-RP10>.
33. Melillo R, Leisman G, Machado C, Machado-Ferrer Y, Chinchilla-Acosta M, Kamgang S, et al. Retained Primitive Reflexes and Potential for Intervention in Autistic Spectrum Disorders. *Front Neurol*. 2022;**13**:922322. [PubMed ID: 35873782]. [PubMed Central ID: PMC9301367]. <https://doi.org/10.3389/fneur.2022.922322>.
34. Payne VG, Isaacs LD. *Human motor development: A lifespan approach*. Oxfordshire, UK: Routledge; 2024.