




Predicting Missed Nursing Care Based on Moral Sensitivity and Care-Related Factors

Elmira Fouladi¹, Mohammad Ali Mohammadi ², Naser Mozaffari ^{3,*}

¹ School of Nursing and Midwifery, Ardabil University of Medical Sciences, Ardabil, Iran

² Department of Medical-Surgical Nursing, School of Nursing and Midwifery, Ardabil University, Ardabil, Iran

³ Department of Nursing, School of Nursing and Midwifery, Ardabil University of Medical Sciences, Ardabil, Iran

*Corresponding author: Department of Nursing, School of Nursing and Midwifery, Ardabil University of Medical Sciences, Ardabil, Iran. Email: naser.m51@gmail.com

Received 2023 December 6; Revised 2024 April 9; Accepted 2024 April 21.

Abstract

Background: Care is one of the essential metaparadigms of nursing. The personal and professional character of the nurse, along with attention to human, financial, and communication factors, can provide an essential part of standard care. Moral sensitivity, as a part of professional ethics, offers responsible and accurate care.

Objectives: This study was conducted with the aim of predicting missed nursing care based on moral sensitivity and care-related factors in 2021.

Methods: In this cross-sectional-analytical study, 345 nurses working in medical education centers in Ardabil were selected by stratified random sampling from internal, surgical, special care, emergency, and pediatric departments. Data were collected using the modified Lutzen Moral Sensitivity Questionnaire, Kalish's Missed Nursing Care, and Blackman's Missed Care Related factors. The collected data were analyzed using SPSS 19 with descriptive statistics (frequency, percentage, mean, and standard deviation) and inferential statistics (ANOVA, regression, and Pearson).

Results: The results of the present study showed that nurses had an average level of moral sensitivity (70.75 ± 10.83). The mean and standard deviation of missed nursing care were (33.93 ± 10.35), which was less than the median of the questionnaire. The components of moral sensitivity (0.32) and related factors (0.08) explain missed care. There was a significant relationship between age, work experience, overtime, and second job with missed care ($P < 0.001$).

Conclusions: Identifying influential factors in reducing missed care is crucial in care, education, and management. It can be said that the dimensions of moral sensitivity and related factors such as human, financial, and communication resources are predictors of missed care. Increasing workload and having a second job increase forgetfulness in caregiving. It is suggested to utilize the experiences of nurses with expertise in care, education, and management, in addition to enhancing moral sensitivity, to ensure comprehensive care while reducing overtime and avoiding second jobs.

Keywords: Moral Status, Ethics, Moral Sensitivity, Missed Nursing Care, Factors

1. Background

In today's world, the significance of an ethical approach in medical professions is widely recognized and deemed essential (1). The nursing profession encompasses health promotion activities, disease prevention, and care for patients of all ages (2). Ethics and clinical practice are not separate (3). Professional ethics are an essential component for individuals aspiring to become professionals (4). It is crucial to pay attention to professional behavior and ethics, including

moral sensitivity (5). Moral sensitivity is one of the essential dimensions of professional ethics, providing the basis for decision-making and clinical practice (6). Therefore, moral sensitivity and a sense of responsibility are significant for nurses providing care (7). Nurses should be accountable for their actions according to their moral obligations (8).

Nursing care is widely recognized as an indispensable component of healthcare services, requiring a comprehensive and holistic approach in all

aspects, with no room for neglect (9). Nevertheless, nurses encounter situations where they inadvertently or intentionally fail to fulfill certain responsibilities due to factors such as organizational, personal, and psychological influences. This failure to provide necessary care can be referred to as missed nursing care (MNC), an unfortunate omission or oversight of care required by the patient (10). The missed care model was presented by Kalisch et al. in 2009 (11). According to the mentioned model, factors such as human resources, financial resources, and communication can lead to failure in care (12). Missed care leads to consequences such as increasing the length of hospitalization, reducing patient satisfaction and safety, and lowering the hospital's reputation from the patients' perspective (13). The ability to predict facing these issues and recognize ethical problems and the appropriate decision-making solution are essential factors in providing standard care (7).

2. Objectives

This study aimed to predict missed nursing care based on moral sensitivity and care-related factors in 2021.

3. Methods

3.1. Study Design

This research is a cross-sectional study aimed at predicting missed nursing care based on moral sensitivity and care-related factors. The study was conducted from July 11, 2021, to October 30, 2021. The study aimed to predict missed nursing care based on moral sensitivity and care-related factors.

3.2. Study Settings and Participants

The study was conducted with a population of working nurses from the medical education centers affiliated with Ardabil University of Medical Sciences. To determine the sample size, Cochran's formula (as shown in Equation 1) was utilized, with ($d = 0.05$), ($P = 0.5$), ($z = 1.96$), and ($N = 1368$). A sample size of 345 nurses was calculated, accounting for a 15% estimated attrition rate.

$$n = \frac{\frac{z^2 - pq}{d^2}}{1 + \frac{1}{n} \left[\frac{z^2 - pq}{d^2} - 1 \right]}$$

The ratio of nurses in various hospitals was determined as follows: Imam Khomeini (154), Fatemi (65), Alavi (48), Bu Ali (53), and Imam Reza (25). Random samples were then taken from each of the five departments: Surgery (77), internal medicine (65), emergency (94), special care (72), and pediatric (37). Questionnaires were distributed to nurses who met the inclusion criteria, which included having a bachelor's degree or higher, at least six months of work experience in their relevant department, and a willingness to participate in the study. Exclusion criteria consisted of incomplete questionnaire responses. The questionnaires were distributed randomly across different shifts and classes. A total of 345 participants were enrolled in the study, providing a diverse and representative sample of nurses from the five departments.

3.3. Data Collection and Measurement

Data collection tools used in this study included a demographic information questionnaire, Lutzen's modified moral sensitivity questionnaire (MSQ), Kalish's missed care questionnaire, and a questionnaire assessing factors related to Blackman's missed care. The demographic questionnaire gathered information on participants' age, sex, marital status, work experience, income, education level, workplace sector, and shift work status, including whether they had a second job. The MSQ, developed by Lutzen, is a 30-item questionnaire that measures nurses' moral sensitivity using a five-point Likert scale. The questionnaire was designed to assess the level of moral sensitivity among nurses in various clinical settings (14, 15). A new 25-item version of the Moral Sensitivity Questionnaire (MSQ) was introduced in a 2021 study, which assesses moral sensitivity using a five-point Likert scale. The questionnaire demonstrated strong reliability in the Iranian version, with a Cronbach's alpha coefficient of 0.89. The MSQ categorizes respondents' moral sensitivity into three levels based on their scores: Low (0 - 50), medium (51 - 75), and high (76 - 100) (16). The missed nursing care questionnaire (MNC), initially developed by Kalisch in 2006, underwent further

psychometric evaluation by the same author in 2009 (11). It has been widely utilized for patient movement, turning, evaluation, training, planning, discharge, and drug prescription (17). This tool assesses the missing elements of nursing care and the reasons for their omission. It consists of 24 items that ask respondents to indicate how frequently they missed providing care during their last working shift using a four-point Likert scale with scores ranging from 1 to 4. The questionnaire scores range from 24 to 96, with higher scores indicating a greater likelihood of missed care. The validity and reliability of the questionnaire were evaluated using Cronbach's alpha coefficient, with estimated values of 0.99 and 0.91, respectively (18).

Blackman et al. designed a questionnaire of factors related to MNC in Australia in 2014 (12). The 17-question questionnaire consists of three sub-scales: Human resources, material resources, and communication. Upon validation in the context of Iran's nursing system, three additional items were incorporated, resulting in a total of 20 items. Responses are measured on a four-point Likert scale, with scores ranging from 20 to 80. Questions pertaining to human resources (1-4, 7, 12-14), financial resources (6, 9, 10, 16), and communication (5, 7, 12-14) were identified. The validity of the questionnaire in Iran was confirmed by Khajooee et al., showing a reliability coefficient of 0.98 and internal consistency with a Cronbach's alpha of 0.98 (18).

3.4. Data Analysis

Descriptive statistics (frequency, percentage, mean, and standard deviation) and inferential statistics (ANOVA, regression, and Pearson correlation) were conducted using SPSS 19 software. The significance level for data analysis was set at 0.05. No outliers or missing data were identified in the study.

3.5. Ethical Considerations

This article presents the findings of a research project conducted under the code of ethics I.R.ARUMS.REC.1400.101. The study adhered to all ethical principles related to research, including obtaining informed consent from participants, ensuring anonymity, respecting autonomy, avoiding harm, and protecting the rights of the institution and researchers.

4. Results

The results showed that 83.2% of the participating nurses (287 samples) were female, 80% were married (276 samples), and 69.9% (241 samples) had official employment status. Moral sensitivity had a positive significant relationship with age and work experience and a negative significant relationship with overtime per month (Table 1).

Table 1. Relationship Between Moral Sensitivity and Demographic Characteristics ^a

Variables	Mean ± Standard Deviation	Sig.
Age	32.19 ± 3.98	P = 0.001 ^a ; R = 0.105
Work Experience (y)	6.23 ± 2.67	P = 0.001 ^a ; R = 0.184
Overtime per Month	73.99 ± 21.95	P > 0.001 ^a R = - 0.313

^aPearson Correlation.

Missed nursing care had a significant relationship with education, department of activity, and having a second job (Table 2).

Table 2. Relationship Between MNC and Demographic Characteristics

Variables	Number (%)	P-Value
Education		> 0.001 ^a
Bachelor	308 (89.3)	
Masters	37 (10.7)	
Department of activity		> 0.001 ^a
Internal	65 (18.8)	
Surgery	77 (22.3)	
Intensive care	72 (20.9)	
Emergency	94 (27.2)	
Children	37 (10.7)	
Second job		> 0.001 ^a
Yes	18 (5.2)	
No	327 (94.8)	

^aIndependent-sample t-test.

In investigating the moral sensitivity of nurses, the mean score of moral sensitivity was found to be 70.75 ± 10.83 (Table 3).

Findings related to MNC showed that the total mean and standard deviation of MNC were 33.93 ± 10.35. The highest average MNC was related to "supervising the preparation of food for a patient who can eat by himself," with an average of 1.97, and "doing oral care," with an average of 1.92, respectively.

Table 3. Mean and Standard Deviation of Dimensions and Total Moral Sensitivity

Moral Sensitivity	Mean ± SD	Mean ± SD of Item to Item
Respect for the client's independence	9.12 ± 1.90	0.63 ± 3.04
Knowledge of how to communicate with the patient	3.40 ± 14.72	0.68 ± 2.94
Professional knowledge	1.63 ± 4.46	0.81 ± 2.23
Experiencing moral problems and conflicts	1.70 ± 8.84	0.56 ± 2.94
Applying ethical concepts in ethical decisions	2.73 ± 14.19	0.54 ± 2.83
Honesty and benevolence	3.85 ± 19.41	0.55 ± 2.77
Total moral sensitivity score	10.83 ± 70.75	0.43 ± 2.83

Table 4. Correlation Between Nurses' Moral Sensitivity and MNC

Order	Variables	1	2	3	4	5	6	7	8
1	Total moral sensitivity	1							
2	Respect for independence	0.535	1						
3	Knowledge of how to communicate with the patient	0.675 ^a	0.095	1					
4	Professional knowledge	0.623 ^a	0.330	0.111	1				
5	Experience moral conflicts	0.678 ^a	0.474	0.309	0.484	1			
6	Applying ethical concepts	0.793 ^a	0.355	0.550	0.344	0.460	1		
7	Honesty and benevolence	0.823 ^a	0.324	0.393	0.608	0.425	0.508	1	
8	The whole care is missed	-0.253 ^a	-0.345 ^a	-0.034	-0.119 ^a	-0.280 ^a	-0.031	-0.330 ^a	1

^a Correlation is significant at the 0.01 level (2-tailed).

In the prediction of MNC based on the components of nurses' moral sensitivity, the results showed that the features of moral sensitivity of nurses explain 0.4% of MNC. The path coefficient of independent to dependent components is equal to 1.574. This value represents the Pearson correlation between the features and the variable (Table 4).

There is a significant inverse relationship between moral sensitivity and missed care. Professional knowledge, experience of moral dilemmas and conflicts, respect for client independence, honesty, and benevolence predict missed care. Significant relationships were found between age, amount of overtime, second job, work experience, department, and education with missed care (Table 5).

This study revealed that nurses exhibited a moderate level of moral sensitivity, consistent with findings from other studies conducted in Iran and elsewhere (19, 20). Mousavi et al. indicated medium to high levels of moral sensitivity among students and nurses at Aja University (21), whereas Filipova suggested that nurses lacked sufficient moral sensitivity for decision-making (22).

These discrepancies in reported results suggest that various factors may influence nurses' moral sensitivity, including workplace conditions, overtime, work experience, departmental dynamics, and the extent of nursing care (23). Research indicates that higher levels of moral sensitivity among nurses correlate with improved quality of nursing care (24). The average level of moral sensitivity among nurses reflects their relative emphasis on adhering to ethical principles in patient care. Therefore, it is imperative for nurses to enhance their understanding of nursing ethics. Among the dimensions of moral sensitivity, the highest score was attributed to "honesty and benevolence," while the lowest score was associated with "professional knowledge." This finding aligns with the results reported by Abdou, wherein the dimension of honesty and benevolence received the highest score among moral sensitivity dimensions, consistent with the current study (5). In the study by Comrie, the highest score was related to the field of "applying moral concepts," and the lowest score was related to the field of "experiencing moral problems and conflicts" (19). The

Table 5. Analysis of Regression Coefficients of Missed Care Based on Moral Sensitivity and Factors Related to Missed Care

Predictor Variables	Non-standard Coefficient		Standard Coefficient Beta	P-Value	Statistical Values
S.E	β				
Respect for the client's independence	-0.010	0.022	-0.119	0.027	R = 1.574; R ² = 0.4
Knowledge of how to communicate with the patient	-0.018	0.011	-0.034	0.529	R ² Adj = 0.379
Professional knowledge	-0.008	0.044	-0.280	< 0.001	F = 56.01; P < 0.001
Experiencing moral problems and conflicts	-0.008	0.057	-0.345	< 0.001	
Applying ethical concepts in ethical decisions	-0.014	0.008	-0.031	0.571	
Honesty and benevolence	-0.123	0.019	-0.330	< 0.001	
Total moral sensitivity	0.265	0.055	-0.253	< 0.001	
Human resources	3.42	19.65	0.22	0.001	
Funds	3.39	24.32	0.15	0.001	
Connections	3.37	25.45	0.13	0.01	

professional knowledge dimension of moral sensitivity refers to cases where decisions are made without the patient's participation. The results of the present study also showed that nurses obtained the lowest score in the professional knowledge dimension. Not paying attention to the patient's autonomy and the idea that the treatment team must help the patient's decision-making are still neglected. Nurses participate less in their patients' treatment and care decisions. Additionally, the findings related to the determination of MNCs showed that the occurrence of MNCs in this research also exists like other research conducted worldwide. The highest mean associated with MNC was related to "supervising the preparation of food for a patient who can eat by himself" and "doing oral care." Also, the lowest mean was related to the statement "General evaluation of the patient in each work shift" and "Evaluation and care of the peripheral and central venous routes of the patient." The results of Cho et al.'s research in Seoul regarding the effect of increasing nursing staff on unperformed nursing care are consistent with our study. They showed that monitoring meals and movement, oral care, bathing, and skincare preparation are part of the MNC (25). The study by Kalisch in America showed that "changing the patient's condition," "cooperation and supervision of going to the toilet in the first 15 minutes of the request," and "participating in interdisciplinary patient care conferences" are the most MNCs (17). We can attribute the differences in management and educational systems, as well as the variance in care styles across the studied departments, to explain the disparities observed

in the types of MNC. Regarding the prediction of MNCs based on the components of moral sensitivity and related factors, the results indicated that nurses' moral sensitivity components account for 0.4% of MNCs. Essentially, as nurses' moral sensitivity increases, MNC decreases, and vice versa. Additionally, components such as professional knowledge, experience of moral dilemmas, respect for client autonomy, honesty, and benevolence showed significant correlations with missed care. However, components like awareness of communication with the patient and application of ethical concepts in decision-making did not exhibit significant correlations. The highest correlation was noted with experiencing moral dilemmas and conflicts, while the lowest was observed with respect for client autonomy. Although nurses' clinical behaviors partly reflect their moral sensitivities, the research underscores the presence of other contributing factors to missed care, warranting further investigation. Policymakers and researchers in the healthcare system must take heed of these factors. Among the demographic variables, a statistically significant relationship was found between overtime and second jobs with moral sensitivity, indicating that nurses with increased overtime and workload experienced more missed care and received lower scores in moral sensitivity. Dehghani et al. also identified the frequency of work shifts and fatigue resulting from overtime as significant barriers to nurses' adherence to professional ethical standards (26). Extended weekly working hours lead to fatigue and dissatisfaction among nurses, resulting in reduced sensitivity to ethical issues. High

workload can foster indifference and diminish nurses' performance. Additionally, a statistically significant correlation was observed between nurses' work experience and missed care, indicating a decrease in missed care with increasing work experience. As nurses age and accumulate more experience, they develop greater clinical skills and moral sensitivity, resulting in enhanced accuracy in care provision. Lutzen et al. demonstrated that nurses' moral sensitivity increases with greater work experience (27), a finding consistent with the results of this study. However, studies by Abbaszadeh et al. and Izadi et al. did not reveal a statistically significant relationship between moral sensitivity and work history (20, 28). In justifying these differences, it can be stated that moral sensitivity tends to increase with accumulating work experience when it is accompanied by ongoing learning, experience growth, and knowledge development. This trend arises because if work experience doesn't coincide with continuous education, there's a possibility that monotony and routine tasks might decrease nurses' sensitivity and their ability to handle new ethical challenges, thus weakening their decision-making skills (20). Considering the influence of organizational and environmental factors, disparities in the research settings between these two studies could contribute to these differences. Additionally, due to the limitation in sample size, caution is warranted when extrapolating the results to other populations.

5. Discussion

The average level of moral sensitivity among nurses reflects their awareness and attentiveness to this aspect of professional ethics. However, the results suggest that this level of moral sensitivity may not suffice for delivering comprehensive patient care and could lead to instances of care being missed or overlooked.

Based on the findings, it can be concluded that moral sensitivity, including respect for client autonomy, professional knowledge, experience with moral dilemmas, honesty, and benevolence, along with related factors such as human resources, finances, and communication, serve as predictors of missed nursing care (MNC). This implies that the higher the level of sensitivity nurses exhibit in clinical care, coupled with favorable human, financial, and communication

resources, the fewer instances of missed care are likely to occur.

Ultimately, to mitigate missed care, fostering moral sensitivity among nurses should be considered a priority within the nursing system.

Acknowledgements

We are grateful to all the officials of Ardabil University of Medical Sciences and the participants who helped in this research.

Footnotes

Authors' Contribution: Concept and design of the study: N. Mozaffari, E. Fouladi, data collection: E. Fouladi, Data analysis and interpretation: Fouladi, Mozaffari, Mohammadi, Preparation of the manuscript: E. Fouladi, N. Mozaffari, Statistical analysis and consultant: Mohammad Ali Mohammadi, Critical revision of the manuscript for important intellectual content: Naser Mozaffari, Academic supervision: Naser Mozaffari

Conflict of Interests Statement: The article's authors declare that there was no conflict of interest in writing this article. Also, this study has not received financial resources from governmental or non-governmental institutions/universities.

Data Availability: It has not been declared by the authors.

Ethical Approval: This article results from a research project with the code of ethics [I.R.ARUMS.REC.1400.101](#), in which all the ethical principles related to research, including consent to participate in the study, maintaining anonymity, autonomy, non-harming, and protecting the rights of the institution and researchers, are included

Funding/Support: No financial aid was received from anywhere or anyone to conduct this research

References

1. Behmaneshpour F, Irandegani F, Miandoab NY. Relationship between the nurses' observance of professional ethics and quality of nursing care from the patients' point of view. *Drug Invention Today*. 2019;12(12).

2. Kane RA. Long-term care and a good quality of life: bringing them closer together. *Gerontologist*. 2001;**41**(3):293-304. [PubMed ID: 11405425]. <https://doi.org/10.1093/geront/41.3.293>.
3. Suhonen R, Stolt M, Virtanen H, Leino-Kilpi H. Organizational ethics: a literature review. *Nurs Ethics*. 2011;**18**(3):285-303. [PubMed ID: 21558106]. <https://doi.org/10.1177/0969733011401123>.
4. Roshanzadeh M, Kashani E, Tajabadi A. Knowledge and practice of nurses regarding the standards of professional nursing Ethics. *Quarterly J Med Ethics*. 2022.
5. Abdou HA, Baddar F, Alkorashy H. The relationship between work environment and moral sensitivity among the nursing faculty assistants. *World Applied Sciences Journal*. 2010;**11**(11):1375-87.
6. Hajibabae F, Salisu WJ, Akhlaghi E, Farahani MA, Dehi MMN, Haghani S. The relationship between moral sensitivity and caring behavior among nurses in Iran during COVID-19 pandemic. *BMC Nurs*. 2022;**21**(1):58. [PubMed ID: 35277159]. [PubMed Central ID: PMC8913325]. <https://doi.org/10.1186/s12912-022-00834-0>.
7. Schluter J, Winch S, Holzhauser K, Henderson A. Nurses' moral sensitivity and hospital ethical climate: a literature review. *Nurs Ethics*. 2008;**15**(3):304-21. [PubMed ID: 18388166]. <https://doi.org/10.1177/0969733007088357>.
8. Akter B, Chaowalit A, Nasae T, RN MS, Yai H. Moral behaviors of nurse supervisors expected and perceived by nurses in Bangladesh. *Medl Sci Nurs*. 2014.
9. Borhani F, Keshtgar M, Abbaszadeh A. Moral self-concept and moral sensitivity in Iranian nurses. *Journal of medical ethics and history of medicine*. 2015;**8**.
10. Scott PA, Harvey C, Felzmann H, Suhonen R, Habermann M, Halvorsen K, et al. Resource allocation and rationing in nursing care: A discussion paper. *Nurs Ethics*. 2019;**26**(5):1528-39. [PubMed ID: 29607703]. [PubMed Central ID: PMC6681425]. <https://doi.org/10.1177/0969733018759831>.
11. Kalisch BJ, Landstrom G, Williams RA. Missed nursing care: errors of omission. *Nurs Outlook*. 2009;**57**(1):3-9. [PubMed ID: 19150261]. <https://doi.org/10.1016/j.outlook.2008.05.007>.
12. Blackman I, Henderson J, Willis E, Hamilton P, Toffoli L, Verrall C, et al. Factors influencing why nursing care is missed. *J Clin Nurs*. 2015;**24**(1-2):47-56. [PubMed ID: 25265893]. <https://doi.org/10.1111/jocn.12688>.
13. John ME, Mgbekem MA, Nsemio AD, Maxwell GI. Missed nursing care, patient outcomes and care outcomes in selected hospitals in Southern Nigeria. *Journal of Nursing & Healthcare*. 2016;**1**(2):1-5.
14. Lützn K, Nordin C, Brolin G. Conceptualization and instrumentation of nurses' moral sensitivity in psychiatric practice. *Int J Methods Psychiatric Res*. 1994. <https://doi.org/10.1037/t60329-000>.
15. Hassanpoor M, Hosseini M, Fallahi Khoshknab M, Abbaszadeh A. Evaluation of the impact of teaching nursing ethics on nurses' decision making in Kerman social welfare hospitals in 1389. *Iranian Journal of Medical Ethics and History of Medicine*. 2011;**4**(5):58-64.
16. Afrasiabifar A, Mosavi A, Dehbanizadeh A, Khaki S. Nurses' caring behaviour and its correlation with moral sensitivity. *J Res Nurs*. 2021;**26**(3):252-61. [PubMed ID: 35251248]. [PubMed Central ID: PMC8895006]. <https://doi.org/10.1177/1744987120980154>.
17. Kalisch BJ, McLaughlin M, Dabney BW. Patient perceptions of missed nursing care. *Jt Comm J Qual Patient Saf*. 2012;**38**(4):161-7. [PubMed ID: 22533128]. [https://doi.org/10.1016/s1553-7250\(12\)38021-5](https://doi.org/10.1016/s1553-7250(12)38021-5).
18. Khajooee R, Bagherian B, Dehghan M, Azizzadeh FM. [Missed nursing care and its related factors from the points of view of nurses affiliated to Kerman University of Medical Sciences in 2017]. *Hayat*. 2019. Persian.
19. Comrie RW. An analysis of undergraduate and graduate student nurses' moral sensitivity. *Nurs Ethics*. 2012;**19**(1):116-27. [PubMed ID: 22183964]. <https://doi.org/10.1177/0969733011411399>.
20. Abbaszadeh A, Borhani F, Moazen NL. The comparison of the level of moral sensitivity in nursing students and nursing staffs in Kerman in 1389. *Medical Ethics Journal*. 2010.
21. Mousavi S, Mohsenpour M, Borhani F, Ebadi M. [Ethical Sensitivity of nurses and nursing students working in Aja University of Medical Sciences]. *Medical Ethics Quarterly*. 2015. Persian.
22. Filipova AA. Licensed nurses' perceptions of ethical climates in skilled nursing facilities. *Nurs Ethics*. 2009;**16**(5):574-88. [PubMed ID: 19671644]. <https://doi.org/10.1177/0969733009106650>.
23. Taghavi MR, Zibaei A, Jafarzadeh Hesari M, Razazan P, Jafarimoghadam A. Evaluation of the Moral Courage and Moral Sensitivity and Factors Affecting them in the Nurses of Selected Hospitals in Bojnourd. *Journal of North Khorasan University of Medical Sciences*. 2022;**13**(4):35-41. <https://doi.org/10.52547/nkums.13.4.35>.
24. Bilgic S. Does the compassion level of nursing students affect their ethical sensitivity? *Nurse Educ Today*. 2022;**109**:105228. [PubMed ID: 34864553]. <https://doi.org/10.1016/j.nedt.2021.105228>.
25. Cho SH, Kim YS, Yeon KN, You SJ, Lee ID. Effects of increasing nurse staffing on missed nursing care. *Int Nurs Rev*. 2015;**62**(2):267-74. [PubMed ID: 25762430]. <https://doi.org/10.1111/inr.12173>.
26. Dehghani A, Dastpak M, Gharib A. [Barriers to respect professional ethics standards in clinical care; viewpoints of nurses]. *Iran J Med Edu*. 2013;**13**(5):421-30. Persian.
27. Lutzen K, Blom T, Ewalds-Kvist B, Winch S. Moral stress, moral climate and moral sensitivity among psychiatric professionals. *Nurs Ethics*. 2010;**17**(2):213-24. [PubMed ID: 20185445]. <https://doi.org/10.1177/0969733009351951>.
28. Izadi A, Imani H, Khademi Z, FariAsadi Noughabi F, Hajizadeh N, Naghzadeh F. [Moral sensitivity of critical care nurses in clinical decision making and its correlation with their caring behavior in teaching hospitals of Bandar Abbas in 2012]. *Iran J Med Ethics History Medicine*. 2013;**6**(2):43-56. Persian.