Published online: 2024 May 1.

**Research Article** 



# Mental Health Literacy and Attitude Toward Seeking Professional Psychological Help Among Iranian Older Adults: A Cross-Sectional Study

Kosar Fazlifar 🝺 1, Abdolhakim Jorbonyan 🔟 2,3, Mina Rafati Lemraski 4, Shahab Papi 🔟 2,3,\*

<sup>1</sup> Department of Psychiatric Nursing, School of Nursing and Midwifery, Mazandaran University of Medical Sciences, Sari, Iran

<sup>2</sup> Department of Geriatric Health, School of Health, Mazandaran University of Medical Sciences, Sari, Iran

<sup>3</sup> Dental Research Center, Mazandaran University of Medical Sciences, Sari, Iran

<sup>4</sup> MSc student in geriatric nursing, department of geriatric nursing, school of nursing and midwifery, Mazandaran university of medical science, Sari, Iran

Corresponding author: Department of Geriatric Health, School of Health, Mazandaran University of Medical Sciences, Sari, Iran. Email: shahabpapi@yahoo.com

Received 2024 March 13; Revised 2024 April 13; Accepted 2024 April 14.

## Abstract

**Background:** Considering the rising prevalence of mental health issues among older adults and potential barriers to treatment seeking, investigating mental health literacy and attitudes toward professional psychological services in this demographic is crucial for enhancing access to care and fostering psychological well-being.

**Objectives:** This study aimed to explore the association between mental health literacy (MHL) and attitudes toward seeking professional psychological help (ATSPPH) among older adults attending health centers in Amol, Mazandaran, Iran, in 2023.

**Methods:** A cross-sectional study was conducted involving 128 older adults attending health centers in Amol, Mazandaran, Iran, in 2023. Participants were selected using cluster random sampling and completed research instruments including O'Connor and Casey's Mental Health Literacy Scale (MHLS) and the short form of Attitude Toward Seeking Professional Psychological Help (ATSPPH-S). Data were analyzed using descriptive statistics, independent t-tests, one-way analysis of variance, and Pearson's correlation coefficient in SPSS 24 software.

**Results:** The Pearson correlation coefficient revealed a positive and significant relationship between older adults' level of MHL and their ATSPPH (r = 0.28,  $P^* 0.05$ ). Tukey's post hoc test indicated a significant association between older adults' MHL and their educational level. Additionally, independent t-test results demonstrated a significant correlation between ATSPPH and gender as well as the history of visiting a psychologist or psychiatrist ( $P^* 0.05$ ).

**Conclusions:** The study findings suggest that higher levels of MHL among older adults are associated with an increased willingness to seek professional psychological help. Therefore, initiatives aimed at enhancing MHL through education and promotion may play a vital role in encouraging older adults to seek appropriate support.

Keywords: Mental Health, Help-Seeking Behavior, Aging, Health Literacy

# 1. Background

Mental disorders manifest as clinically significant impairments in cognition, affective regulation, or behavior, causing distress (1). These conditions notably affect older individuals, as aging brings about various physical and psychological changes (2). The World Health Organization (WHO) estimates that 15% of 60year-olds experience mental illnesses, primarily depression and dementia (3). Additionally, drug misuse, anxiety, and mood disorders are prevalent among older individuals (4). With the aging population expanding, experts anticipate a rise in mental health issues, necessitating the development of mental health systems and diagnostic approaches for this demographic (5). Given the higher healthcare costs associated with older age, early detection and treatment of mental illnesses are paramount (6), underscoring the importance of mental health literacy (MHL), which encompasses knowledge of mental illness prevention, symptoms, and treatments (7).

Copyright © 2024, Fazlifar et al. This open-access article is available under the Creative Commons Attribution 4.0 (CC BY 4.0) International License (https://creativecommons.org/licenses/by/4.0/), which allows for unrestricted use, distribution, and reproduction in any medium, provided that the original work is properly cited.

In 1997, Jorm defined MHL as the knowledge and beliefs about mental disorders that improve disease detection, management, and prevention (8). This concept encompasses awareness of specific disorders or mental distress, understanding of risk factors, knowledge about self-help interventions and available professional help, an attitude conducive to recognition, and appropriate help-seeking behavior, along with knowledge of mental health information (9). MHL plays a crucial role in reducing mental illnesses and associated social costs, as well as enhancing overall quality of life (10). Lower levels of MHL can detrimentally affect the quality of life of older individuals, who tend to exhibit poorer MHL and less knowledge about mental disorder diagnoses compared to younger adults, as observed by Bragg et al. Despite the growing population of older adults, developing countries have limited data on MHL (11). Despite several studies on mental health among older individuals in Iran, MHL remains inadequately researched.

One aspect influencing the understanding of mental illnesses and mental health literacy (MHL) among older adults is their cultural perspectives (12). Older individuals often fail to recognize their mental health issues due to the stigma surrounding such disorders, which hinders them from seeking necessary care and support (3). Seeking help entails reaching out to both formal and informal sources for knowledge, counseling, information, therapy, and general support in response to challenges or uncomfortable situations (13). The wellbeing of older individuals hinges on their willingness to seek help when needed, and the ability to manage their numerous symptoms with available services is essential within their communities (14). Despite facing various health challenges requiring heightened care, older individuals often opt not to seek assistance (15). According to Robb et al., younger individuals consult mental health specialists twice as often as older adults and possess greater health knowledge (16). The lack of help-seeking behavior and delayed diagnosis of mental issues in older individuals contribute to increased healthcare and societal burdens (5). Internal barriers such as stigma, fear of psychotherapy, and the perception that these symptoms are expected in old age hinder older adults from seeking help. External barriers include cost, insurance coverage, age discrimination, and transportation issues (17). Given that older individuals with mental disorders are at greater risk of physical ailments, their failure to seek early treatment poses a public health concern. Additionally, low MHL in older persons may impede the interpretation of health information and delay the utilization of mental health services (18). Yu et al. suggested that raising mental

health awareness could increase help-seeking behavior (19). However, despite the prevalence of mental disorders among older adults and their insufficient help-seeking behavior, it has not yet been evaluated through a literature review whether MHL, as a determinant of mental health behaviors, can influence older individuals' attitudes toward seeking psychological help. Thus, this study aims to investigate the association between MHL and attitudes toward seeking professional psychological help (ATSPPH) in older adults.

## 2. Objectives

This study seeks to determine the relationship between mental health literacy (MHL) and attitudes toward seeking professional psychological help (ATSPPH) among older adults referred to health centers in Amol, Iran, in 2023.

## 3. Methods

This cross-sectional study was conducted on individuals aged 60 and above who were referred to health centers in Amol City, Mazandaran, Iran, in 2023 (from July to November). The sample size for the study was determined based on the research by Almansef, which reported a correlation coefficient of 0.26 between mental health literacy (MHL) and help-seeking behavior (20). Using the correlation study algorithm, 113 samples were calculated with 95% confidence and 90% power. Considering a possible decrease in participants, the final sample size was set at 150. Cluster random sampling was employed, selecting 20 out of the 47 health centers in Amol City, and eligible elderly individuals attending these centers were included in the study. Exclusion criteria included failure to complete the questionnaire, while inclusion criteria comprised individuals aged 60 and older, absence of cognitive impairment, and provision of informed consent.

The questionnaires used in this study included sociodemographic characteristics, the Mental Health Literacy Scale (MHLS), and the short form of the Attitudes Toward Seeking Professional Psychological Help Scale (ATSPPH-S). Considering the varying literacy levels among elderly participants, the researcher administered all questionnaire items, and responses were recorded from the elderly individuals.

1. Socio-demographic characteristics encompassed variables such as age, gender, location, marital status, educational level, employment status, economic status, history of illness or physical disability (based on individual medical records), history of mental disorders in oneself or family (based on individual medical records), history of referral to a psychologist or psychiatrist, and experience of stressful events such as car accidents, falls, or acquiring a new disease in the last six months.

2. O'Connor and Casey developed the Mental Health Literacy Scale (MHLS) in 2015. This Likert scale questionnaire consists of 35 items rated on a scale of 1 to 5, with scores ranging from 35 to 160, where higher scores indicate higher levels of MHLS (21). In this study, the modified MHLS by O'Connor and Casey was utilized for the Iranian population. This version comprises 29 items categorized into six domains: Information seeking (4 items), self-help (2 items), recognition of disorders (8 items), understanding of risk factors and causes (2 items), attitudes conducive to recognition and help-seeking (10 items), and seeking assistance from professionals (3 items). Items 1 to 15 are rated on a 4point Likert scale, while items 16 to 29 are rated on a 5point scale, resulting in scores ranging from 29 to 130. MacDonald's omega and Cronbach's alpha coefficients for all MHLS domains were 0.79 and 0.78, respectively, indicating its reliability across different age groups in Iran (22). In the present study, the reliability of this instrument was also assessed, yielding a Cronbach's alpha coefficient of 0.74.

3. The Attitudes Toward Seeking Professional Psychological Help Scale (ATSPPH-S) is a 10-item scale developed by Fisher and Farnia in 1995 as a shorter version of the original 29-item scale. It assesses individuals' attitudes toward seeking professional mental health services using a 4-point Likert scale (0 =disagree, 1 = somewhat disagree, 2 = somewhat agree, 3 = agree). Scores range from 0 to 30, with higher scores indicating a more positive attitude toward seeking psychological help (23). Sharifi et al. evaluated the psychometric properties of this scale among Iranian adults in 2018, demonstrating its validity through confirmatory factor analysis and good reliability with a Cronbach's alpha coefficient of 0.84 (17). In the present study, the reliability of the ATSPPH-S was also assessed, yielding a Cronbach's alpha coefficient of 0.81.

This study used mean and standard deviation to calculate descriptive statistics. Also, considering the normal distribution of the data, which were in the range of (-2 -2) using skewness and kurtosis tests and Kolmogorov-Smirnov normality test results (P > 0.05), parametric. Parametric tests, including independent *t*-tests and ANOVA, were employed to compare the means of mental health literacy and attitudes toward seeking professional psychological help based on demographic variables, considering the normal distribution of the

data. The Pearson correlation coefficient was used to examine the relationship between mental health literacy and attitudes toward seeking professional psychological help. All analyses were conducted using SPSS 24, with a significance level set at 0.05.

Ethical considerations in this research were addressed by obtaining permission from the university ethics committee (Code: IR.MAZUMS.REC.1402.261), securing informed consent from the participants, and ensuring the anonymity of the participants. Participants were provided with an informed consent form and informed of their right to withdraw from the study at any time.

## 4. Results

The study included 128 older adults, with a mean age of 66.36 (standard deviation = 6.51), ranging from 60 to 86 years. Among the participants, 55.5% were male, and the majority (85.2%) were married, with 60.9% residing in urban areas. In terms of education, 56.3% had undergraduate degrees, and 34.4% were self-employed, while 62% reported an average economic status (Table 1).

The mean MHL score for older adults was 76.63 (standard deviation = 8.88). The mean scores for each MHL subscale were as follows: 8.11 (3.01) for knowledge of where to seek information, 23.28 (3.84) for ability to recognize information, 5.12 (1.15) for knowledge of risk factors and causes, 21.92 (6.19) for attitudes promoting recognition or appropriate help-seeking behavior, 9.06 (1.07) for knowledge of available professional help, and 5.39 (0.93) for knowledge of self-treatment. Additionally, the mean score for the Attitudes Toward Seeking Professional Psychological Help Scale (ATSPPH-S) among older adults was 16.51 (standard deviation = 4.29) (Table 2).

The Pearson correlation coefficient test revealed a statistically significant relationship (r = 0.28, P < 0.05) between MHL and ATSPPH-S in older adults. Further analysis showed that among the MHL subscales, only the ability to recognize mental disorders (r = 0.23, P < 0.05) and knowledge of where to seek information (r = 0.30, P < 0.05) had a positive and significant relationship with ATSPPH-S in older adults (Table 3).

Demographic factors significantly influenced MHL and ATSPPH-S in older adults. Education level had a considerable impact on average MHL, with individuals holding bachelor's degrees exhibiting significantly higher MHL scores than those with diplomas (P < 0.05). Moreover, women had significantly higher mean ATSPPH-S scores than men (P < 0.05), and participants who had visited a psychiatrist or psychologist had

Variables and Categories	Values <sup>a</sup>
Age, y	Min = 60; Max = 86
Gender	
Mean	66.3 ± 6.51; 71 (55.5
Female	57 (44.5)
farital status	
Married	100 (82.2)
Single	7 (5.5)
Widowed/divorced	12 (9.4)
ducational level	
Under diploma	72 (56.3)
Diploma	30 (23.4)
Associate degree	7 (5.5)
Bachelor's degree	14 (10.9)
Master's degree and higher	5 (3.9)
ocation	
Rural	50 (39.1)
Urban	78 (60.9)
conomic status	
Weak	8(6.3)
Medium	86 (67.2)
Good	34 (26.6)
ob status	
Free	44 (34.4)
Unemployed	43 (33.6)
Retired	41 (32.0)
listory of illness or physical disability	
Yes	46 (35.9)
No	82 (64.1)
listory of visiting a psychiatrist/psychologist	
Yes	26 (20.3)
No	102 (79.7)
amily history of mental disorder	
Yes	18 (14.1)
No	110 (85.9)
listory of stressful events in the last six months	
Yes	46 (35.9)
No	82 (64.1)

significantly higher mean ATSPPH-S scores compared to those who had not (P < 0.05).

No significant differences were found in MHL and ATSPPH-S means based on age, employment status, economic position, marital status, location, history of mental disorders, or stressful events (P > 0.05). The results of significant factors are presented in Table 4.

# 5. Discussion

We examined the relationship between MHL and ATSPPH in older individuals, also assessing MHL and ATSPPH levels within this demographic. The average MHL among older adults in this study was 76.63 (8.88), similar to the study average. Upon reviewing the literature, we found no studies measuring MHL among elderly Iranian adults (11). However, studies by Bragg et al. and Ding et al. observed lower MHL averages in elderly populations (24), consistent with our findings. Conversely, studies by Jafari et al. and Noroozi et al.

Fable 2. Status of MHL and ATSPPH-S in the participant (N = 128)			
Scale/Subscale	Mean <sup>a</sup>	Min	Max
MHL	76.63 (8.88)	53	105
knowledge of where to seek information	11.8 (3.01)	4	20
Ability to recognize information	23.28 (3.84)	8	31
Knowledge of risk factors and causes	5.12 (1.15)	2	8
Attitudes that promote recognition or appropriate help-seeking behavior	21.92 (6.19)	10	46
Knowledge of professional help available	9.06 (1.07)	6	12
Knowledge of self-treatment	5.39 (0.93)	4	8
ATSPPH-S	16.51 (4.29)	4	30

<sup>a</sup> Values are expressed as mean (SD).

<b>able 3.</b> The Results of Pearson's Correlation Analysis of the MHL Scale/Subscale with ATSPPH-S (N = 128)					
Scale/Subscale	ATSPPH-S	ATSPPH-S			
	Correlation Coefficient (r)	P-Value			
MHL	0.28	0.001 <sup>a</sup>			
The ability to find information	0.30	0.001 <sup>a</sup>			
Ability to recognize information	0.23	0.007 <sup>a</sup>			
Knowledge of risk factors and causes	0.08	0.31			
Help-seeking or recognition-promoting attitudes	0.09	0.3			
Professional help knowledge	0.05	0.53			
Self-care knowledge	0.04	0.62			
<sup>a</sup> P < 0.05.					

reported higher MHL averages (10, 25), contradicting our results. Our findings suggest that older individuals generally have lower MHL levels compared to the broader population, particularly younger individuals.

In our study, the mean ATSPPH score among older adults was 16.51 (4.29). A study by Sharifi et al. among Iranian older adults reported a similar average ATSPPH score (17). However, a study by Roh et al. among older American adults found a higher average ATSPPH score, indicating potential cultural and social differences between countries influencing older adults' attitudes toward seeking professional psychological help (26).

We found a positive correlation between MHL and ATSPPH among older individuals, indicating that those with higher MHL were more inclined to seek professional psychological help. While there is limited research on this relationship among older populations, studies by Almansef, Gorczynski and Sims-Schouten., Kantaş Yılmaz and ÜNKÜR have identified this link among broader populations and student groups, suggesting that increasing MHL and understanding of mental health symptoms among older individuals could improve their attitudes toward mental health care and increase service utilization (20, 27, 28).

Education level was found to be associated with MHL in this study, indicating that older individuals with higher education had higher MHL levels. This aligns with the findings of Ding et al., who observed a similar link between education and MHL in older populations (24). However, this contradicts the findings of Piper et al. in a study of older Australians (18). Such discrepancies may stem from differences in societal, economic, and cultural contexts. Noroozi et al. also identified this association in Iran's population (25), supporting the positive relationship between education and MHL in Iranian society.

The study revealed a significant association between gender and ATSPPH, with older women showing a greater inclination to seek professional psychological care. This finding is consistent with studies by Roh et al., Mackenzie et al., and Kessler et al. among older populations (26, 29, 30). Kantaş Yılmaz and ÜNKÜR similarly confirmed this link among student populations (28). To date, no study has reported a higher propensity for seeking psychological help among men.

ables	Mean (SD)	t	P-Value	DF	Levene's Test	Levene's Test P-Value
der		2.12	0.03 <sup>a</sup>	126	0.75	0.09
Woman	17.40 (4.16)					
Man	15.80 (4.29)					
tory of visiting a psychiatrist/psychologist		2.22	0.03 <sup>a</sup>	126	0.40	0.69
Yes	18.30 (4.72)					
No	16.05 (4.07)					
cational level	-	12.29	≥ 0.001	4295	≤ 0.05	2.51
cational level	-	12.29	≥0.001	4295	≤0.05	

According to these studies, women are more inclined to seek help as they tend to be more comfortable expressing emotions and discussing mental health issues openly compared to men.

The study also found a significant relationship between a history of visiting a psychologist or psychiatrist and participants' ATSPPH. Older adults with such a history demonstrated a greater willingness to seek professional psychological help than those without. This finding is consistent with Kessler et al.'s study among older German adults (30). It suggests that a positive past experience with mental health professionals leads to better understanding of the counseling and treatment process, reducing fears and misconceptions about seeking help and fostering greater comfort in expressing problems.

This study represents the first investigation into MHL and ATSPPH among Iranian older adults, highlighting its strength. However, a limitation is that the results may not be generalizable to all populations due to the cultural dependency of MHL and ATSPPH. Therefore, it is recommended that further research be conducted in diverse geographical locations to enhance understanding in this field.

## 5.1. Conclusions

The results of this study demonstrated a positive and significant relationship between older adults' MHL levels and their ATSPPH. Furthermore, a significant association was found between older adults' MHL and their educational levels, as well as between their ATSPPH and their gender and history of visiting a psychologist or psychiatrist. These findings have several implications. Firstly, interventions to enhance MHL and promote positive ATSPPH among older adults are crucial. Many senior citizens have low literacy levels and have not attended university. Consequently, the education sector requires support in addressing these issues. Communication barriers and limited acculturation can hinder low-literacy individuals' participation in health education and promotion activities. To effectively promote MHL and positive ATSPPH among older adults, educational interventions must be tailored to their age, literacy levels, and cultural backgrounds.

Moreover, the observed relationship between MHL and educational attainment highlights the importance of lifelong learning opportunities for older adults. Providing accessible and relevant educational programs can help improve their MHL and, subsequently, their willingness to seek professional psychological help when needed.

Additionally, the association between ATSPPH and gender underscores the need for gender-sensitive approaches in mental health promotion efforts targeting older adults. Addressing gender-specific barriers, stigma, and sociocultural norms may be necessary to encourage help-seeking behavior among both men and women in this age group. Implications also extend to the healthcare system, emphasizing the necessity for age-friendly and culturally competent mental health services. Ensuring accessibility, affordability, and understanding of the needs and preferences of older adults can enhance their engagement with professional psychological support when required.

## Acknowledgements

The authors wish to express their sincere gratitude to all the elderly individuals who participated in this study for their time and cooperation. We are also thankful to the staff of the health centers in Amol county for their assistance in facilitating this research project. Without their support, this study would not have been possible.

## Footnotes

**Authors' Contribution:** Shahab Papi and Abdolhakim Jourbanian designed the study and analyzed the data. Kosar Fazlifar gathered and interpreted the data. Shahab Papi and Kosar Fazlifar wrote the entire manuscript. All authors confirmed the final edited version of the manuscript.

**Conflict of Interests Statement:** The authors have no conflicts of interest to disclose regarding funding or research support, employment, personal financial interests, stocks or shares in companies, consultation fees, patents, personal or professional relations with organizations and individuals (family, spouse, children, etc.), or unpaid membership in government or non-governmental organizations. Two of the authors (Dr. Papi and Dr. Jorbanian) of this manuscript are reviewers for this journal; however, they were not involved in the review process for this manuscript.

**Data Availability:** The dataset presented in the study is available at the request of the corresponding author during submission or after publication. The data are not publicly available due to privacy or ethics.

**Ethical Approval:** All ethical considerations in this research, including obtaining permission from the university ethics committee (Code: IR.MAZUMS.REC.1402.261), securing informed consent from the participants to participate in the study, and ensuring the anonymity of the participants in the research, were addressed. The participants completed an informed consent form and were told they could withdraw from the study at any time.

**Funding/Support:** Mazandaran University of Medical Sciences supported this research study (project number: 18274).

**Informed Consent:** Written consent was also obtained from older adults before data collection.

#### References

- 1. World Health Organization. *Mental Disorder*. Geneva: World Health Organization; 2022, [cited 8 June 2022]. Available from: https://www.who.int/news-room/fact-sheets/detail/mental-disorders.
- 2. World Health Organization. *Aging and health*. Geneva: World Health Organization; 2023, [cited 1 October 2022]. Available from: https://www.who.int/news-room/fact-sheets/detail/ageing-and-health.
- 3. World Health Organization. *Mental health of older adults*. Geneva: World Health Organization; 2023, [cited 20 October 2023]. Available from: https://www.who.int/news-room/fact-sheets/detail/mentalhealth-of-older-adults.

- Andreas S, Schulz H, Volkert J, Dehoust M, Sehner S, Suling A, et al. Prevalence of mental disorders in elderly people: the European MentDis\_ICF65+ study. *Br J Psychiatry*. 2017;210(2):125-31. [PubMed ID: 27609811]. https://doi.org/10.1192/bjp.bp.115.180463.
- 5. Petrova NN, Khvostikova DA. [Prevalence, structure, and risk factors for mental disorders in older adults.]. *Adv Gerontol.* 2021;**34**(1):152-9. [PubMed ID: 33993676].
- Polacsek M, Boardman GH, McCann TV. Help-seeking experiences of older adults with a diagnosis of moderate depression. *Int J Ment Health Nurs.* 2019;28(1):278-87. [PubMed ID: 30120874]. [PubMed Central ID: PMC8351636]. https://doi.org/10.1111/inm.12531.
- Moreira WC, Sousa AR, Cardoso R, Queiroz AM, Oliveira MAF, Sequeira C. COVID-19 in Brazil: Are there any differences in Mental Health Literacy between young and aged men? *Rev Lat Am Enfermagem.* 2022;**30**. e3603. [PubMed ID: 35858006]. [PubMed Central ID: PMC9285197]. https://doi.org/10.1590/1518-8345.5651.3603.
- Jorm AF, Korten AE, Jacomb PA, Christensen H, Rodgers B, Pollitt P. "Mental health literacy": a survey of the public's ability to recognise mental disorders and their beliefs about the effectiveness of treatment. *Med J Aust.* 1997;166(4):182-6. [PubMed ID: 9066546]. https://doi.org/10.5694/j.1326-5377.1997.tb140071.x.
- Jorm AF. Mental health literacy. Public knowledge and beliefs about mental disorders. *Br J Psychiatry*. 2000;**177**:396-401. [PubMed ID: 11059991]. https://doi.org/10.1192/bjp.177.5.396.
- Jafari A, Nejatian M, Momeniyan V, Barsalani FR, Tehrani H. Mental health literacy and quality of life in Iran: a cross-sectional study. *BMC Psychiatry*. 2021;21(1):499. [PubMed ID: 34641793]. [PubMed Central ID: PMC8507341]. https://doi.org/10.1186/s12888-021-03507-5.
- Bragg TL, Segal DL, Coolidge FL. Mental Health Literacy and Attitudes about Mental Disorders among Younger and Older Adults: A Preliminary Study. *Open Journal of Geriatrics*. 2018;1(2):1-6. https://doi.org/10.22259/2639-359x.0102001.
- Lee HY, Lytle K, Yang PN, Lum T. Mental health literacy in Hmong and Cambodian elderly refugees: a barrier to understanding, recognizing, and responding to depression. *Int J Aging Hum Dev.* 2010;**71**(4):323-44. [PubMed ID: 21261138]. https://doi.org/10.2190/AG.71.4.d.
- van den Toren SJ, van Grieken A, Lugtenberg M, Boelens M, Raat H. Adolescents' Views on Seeking Help for Emotional and Behavioral Problems: A Focus Group Study. *Int J Environ Res Public Health*. 2019;**17**(1). [PubMed ID: 31892118]. [PubMed Central ID: PMC6981849]. https://doi.org/10.3390/ijerph17010191.
- Ohta R, Ryu Y, Sano C. Older People's Help-Seeking Behaviors in Rural Contexts: A Systematic Review. *Int J Environ Res Public Health.* 2022;19(6). [PubMed ID: 35328920]. [PubMed Central ID: PMC8951636]. https://doi.org/10.3390/ijerph19063233.
- Teo K, Churchill R, Riadi I, Kervin L, Wister AV, Cosco TD. Help-Seeking Behaviors Among Older Adults: A Scoping Review. J Appl Gerontol. 2022;41(5):1500-10. [PubMed ID: 35156428]. [PubMed Central ID: PMC9024019]. https://doi.org/10.1177/07334648211067710.
- Robb C, Haley WE, Becker MA, Polivka LA, Chwa HJ. Attitudes towards mental health care in younger and older adults: similarities and differences. *Aging Ment Health*. 2003;7(2):142-52. [PubMed ID: 12745392]. https://doi.org/10.1080/1360786031000072321.
- Sharifi M, Abolfathi Momtaz Y, Alizadeh T, Zanjari N, Mohammadi Safa N. Psychometric properties of the attitudes toward receiving professional psychological services scale in a sample of Iranian older adults. *Shenakht Journal of Psychology and Psychiatry*. 2019;6(5):83-95. https://doi.org/10.29252/shenakht.6.5.83.
- Piper SE, Bailey PE, Lam LT, Kneebone ,I. Predictors of mental health literacy in older people. *Arch Gerontol Geriatr*. 2018;**79**:52-6. [PubMed ID: 30107312]. https://doi.org/10.1016/j.archger.2018.07.010.

- Yu Y, Liu ZW, Hu M, Liu XG, Liu HM, Yang JP, et al. Assessment of mental health literacy using a multifaceted measure among a Chinese rural population. *BMJ Open*. 2015;5(10). e009054. [PubMed ID: 26438139]. [PubMed Central ID: PMC4606438]. https://doi.org/10.1136/bmjopen-2015-009054.
- 20. Almanasef M. Mental Health Literacy and Help-Seeking Behaviours Among Undergraduate Pharmacy Students in Abha, Saudi Arabia. *Risk Manag Healthc Policy*. 2021;**14**:1281-6. [PubMed ID: 33790673]. [PubMed Central ID: PMC8006949]. https://doi.org/10.2147/RMHP.S289211.
- O'Connor M, Casey L. The Mental Health Literacy Scale (MHLS): A new scale-based measure of mental health literacy. *Psychiatry Res.* 2015;**229**(1-2):511-6. [PubMed ID: 26228163]. https://doi.org/10.1016/j.psychres.2015.05.064.
- Nejatian M, Tehrani H, Momeniyan V, Jafari A. A modified version of the mental health literacy scale (MHLS) in Iranian people. *BMC Psychiatry*. 2021;**21**(1):53. [PubMed ID: 33485306]. [PubMed Central ID: PMC7824912]. https://doi.org/10.1186/s12888-021-03050-3.
- Fischer EH, Farina A. Attitudes toward seeking professional psychological help: A shortened form and considerations for research. *J College Student Develop.* 1995;**36**(4):368-73. https://doi.org/10.1037/t05375-000.
- Ding KR, Wang SB, Xu WQ, Lin LH, Liao DD, Chen HB, et al. Low mental health literacy and its association with depression, anxiety and poor sleep quality in Chinese elderly. *Asia Pac Psychiatry*. 2022;14(4). e12520. [PubMed ID: 36210054]. https://doi.org/10.1111/appy.12520.

- Noroozi A, Khademolhosseini F, Lari H, Tahmasebi R. The Mediator Role of Mental Health Literacy in the Relationship Between Demographic Variables and Health-Promoting Behaviours. *Iran J Psychiatry Behav Sci.* 2018;12(2). https://doi.org/10.5812/ijpbs.12603.
- 26. Roh S, Burnette CE, Lee KH, Lee YS, Martin JI, Lawler MJ. Predicting Help-Seeking Attitudes Toward Mental Health Services Among American Indian Older Adults: Is Andersen's Behavioral Model a Good Fit? *J Appl Gerontol.* 2017;**36**(1):94-115. [PubMed ID: 25416511]. https://doi.org/10.1177/0733464814558875.
- Gorczynski P, Sims-Schouten W. Evaluating mental health literacy amongst US college students: a cross sectional study. J Am Coll Health. 2024;72(3):676-9. [PubMed ID: 35471990]. https://doi.org/10.1080/07448481.2022.2063690.
- Kantaş Yılmaz F, ÜNKÜR P. Determination of Mental Health Literacy, Help-Seeking Behaviours and Psychological Distress Levels of Health Science Students. J Health Literacy. 2023;7(4):9-20.
- Mackenzie CS, Scott T, Mather A, Sareen J. Older adults' help-seeking attitudes and treatment beliefs concerning mental health problems. *Am J Geriatr Psychiatry*. 2008;**16**(12):1010-9. [PubMed ID: 19038900].
  [PubMed Central ID: PMC2735824]. https://doi.org/10.1097/JGP.0b013e31818cd3be.
- Kessler EM, Agines S, Bowen CE. Attitudes towards seeking mental health services among older adults: personal and contextual correlates. *Aging Ment Health.* 2015;19(2):182-91. [PubMed ID: 24898327]. https://doi.org/10.1080/13607863.2014.920300.