







The Relationship Between Metabolic Syndrome with Physical Activity and Eating Habits: The Result of the Tabari Cohort Study

Leyla Mansoori ¹, Zahra Kashi ¹, Amirhosein Khoshgoeian¹, Adele Bahar ¹, Mahmood Moosazadeh ², Atena Ramezani^{3,*}, Sanaz Soltani³

¹ Diabetes Research Center, Mazandaran University of Medical Sciences, Sari, Iran

² Gastrointestinal Cancer Research Center, Non-communicable Diseases Institute, Mazandaran University of Medical Sciences, Sari, Iran

³ Department of Nutrition Sciences, School of Health, Mazandaran University of Medical Sciences, Sari, Iran

*Corresponding Author: Department of Nutrition Sciences, School of Health, Mazandaran University of Medical Sciences, Sari, Iran. Email: ramezaniatena@yahoo.com

Received 2024 August 10: Revised 2024 August 21: Accepted 2024 August 21

Abstract

Background: Metabolic syndrome (MetS) is a group of metabolic issues that increase the risk of type 2 diabetes and heart disease.

Objectives: To reduce deaths from type 2 diabetes and heart disease, this study was conducted to determine the relationship between metabolic syndrome, physical activity, and eating habits in participants of the Tabari cohort study.

Methods: In this descriptive cross-sectional study, data from the Tabari cohort in an urban and mountain area in Sari, Mazandaran, was utilized from June 2015 to November 2017. Background information of participants was documented, and their eating habits were assessed using a food habit questionnaire. Additionally, the physical activity levels of the participants were recorded. All individuals were assessed for metabolic syndrome according to the third Adult Treatment Panel (ATP III) guidelines. The gathered data was then inputted into SPSS V.23 software for analysis. Qualitative variables, such as demographic and epidemiological data, were compared between the two groups using the chi-square test, while quantitative variables were compared using an independent *t*-test.

Results: A total of 10,255 adults between the ages of 35 and 70 participated in this study, and after applying entry and exit criteria, 9,976 people, including 4,083 with metabolic syndrome and 5,893 healthy individuals, were included in the final analysis. There was no significant difference in the average Body Mass Index (BMI) between individuals with metabolic syndrome (30.6 ± 4.5 kg/m²) and healthy individuals (26.8 ± 4.4 kg/m²). However, individuals with metabolic syndrome exhibited significantly higher values for height, weight, waist circumference, abdominal circumference, and wrist circumference compared to the control group ($P = 0.001$). The average Metabolic Equivalent of Task (MET) for the entire population was 43.03 ± 9.27 , with a median of 40.68 (range: Twenty-four-one hundred three) hours/day. A greater percentage of individuals with metabolic syndrome (57.0%) and healthy individuals (45.2%) had MET values below the average ($P = 0.001$). Regarding energy intake, individuals with metabolic syndrome consumed an average of 2189.74 ± 721.90 calories, while healthy individuals consumed 2354.92 ± 4.04 calories, showing a significant difference ($P = 0.006$). Moreover, healthy individuals demonstrated significantly higher consumption of added salt in food, fried foods, fried potatoes, fried onions, fried vegetables, and various types of cooking oils compared to those with metabolic syndrome ($P = 0.001$).

Conclusions: Based on the results of the Tabari cohort study, the prevalence of metabolic syndrome in Mazandaran was 41.10%, highlighting the need for prevention and screening programs. In this study, individuals with metabolic syndrome consumed fewer fried foods, trans fats, and daily calories compared to the healthy group. Considering that this study is cross-sectional and utilized data from the Tabari cohort, it is assumed that the difference is due to dietary compliance and lifestyle changes in patients with metabolic syndrome. It is recommended that prospective studies be designed to evaluate the causes of differences in metabolic syndrome components between the two groups.

Keywords: Metabolic Syndrome, Physical Conditioning, Food, Habits

1. Background

Metabolic syndrome (MetS) is a cluster of metabolic abnormalities linked to an increased risk of developing type 2 diabetes (T2D) and cardiovascular disease (CVD). It is characterized by high blood sugar levels, elevated blood pressure, raised triglyceride levels, low levels of HDL cholesterol, and obesity, particularly central obesity (1). Among the various guidelines for metabolic syndrome, the national cholesterol education program (NCEP) adult treatment panel (ATP III) is the most relevant (2). Obesity is widely recognized as the primary risk factor for metabolic syndrome (3). According to the World Health Organization, global obesity rates have nearly tripled since 1975 (4). This surge in obesity has contributed to approximately 34% of adults being affected by metabolic syndrome, indicating a rising prevalence (5). An analysis of NHANES data from 2011 to 2016 revealed that 34.7% of participants met the ATP III criteria for metabolic syndrome, a significant increase compared to the 22% observed in the 1988-1994 NHANES data (6, 7).

Obesity, particularly central obesity, has been linked to a reduced responsiveness to insulin, resulting in insulin resistance, which often leads to type 2 diabetes. Insulin resistance, along with elevated levels of insulin, glucose, and adipose cytokines, can negatively affect blood vessel function, lipid levels, and lead to conditions like high blood pressure, inflammation, and atherosclerotic cardiovascular disease (8, 9). As a result, metabolic syndrome is a crucial risk factor for both type 2 diabetes and cardiovascular diseases. Studies, including systematic reviews and meta-analyses, demonstrate the preventive and therapeutic benefits of exercise (10, 11) and leisure-time physical activity (12) in combatting major chronic diseases like metabolic syndrome. According to WHO guidelines on physical activity and sedentary behavior, the hazards of inactivity (13) and the importance of reducing sedentary behavior (14) are emphasized alongside the advantages of regular exercise. Recent research indicates that engaging in moderate to vigorous physical activity (MVPA) can lower the likelihood of developing metabolic syndrome, regardless of aerobic fitness or obesity (15-17). This type of physical activity also helps in preventing chronic diseases like type 2 diabetes and MetS (18). Additionally, a beneficial change in blood pressure indicates a link between physical activity and hypertension, as exercise impacts blood pressure and prehypertension (19). Engaging in 150 minutes of physical activity per week is

associated with reduced diastolic blood pressure, increased HDL cholesterol levels, and lower overall cholesterol (20).

Dietary behaviors, defined as intentional and recurring actions influenced by social and cultural factors, play a significant role in selecting, consuming, and utilizing particular foods or diets (21). These behaviors are shaped by cultural, social, economic, and environmental influences, along with individual preferences and beliefs (22). Optimal dietary habits typically involve a balanced selection of nutrient-rich foods, including fruits, vegetables, whole grains, lean proteins, and healthy fats. Consistent meal schedules, mindful eating practices, and proper hydration also contribute to healthy eating. On the contrary, poor dietary habits often involve excessive consumption of calorie-laden, nutritionally deficient foods rich in sugar, salt, and unhealthy fats, along with erratic eating patterns (23). Such unhealthy dietary habits are linked to a variety of health issues, including obesity, metabolic disorders, heart disease, and diabetes (24). The development and progression of metabolic syndrome involve a complex interplay of genetic and environmental factors (25), with diet being a key modifiable environmental factor (26). Research indicates that poor dietary choices contribute to the onset and worsening of metabolic syndrome (27, 28). Individuals with metabolic syndrome tend to consume high amounts of fatty and sugary foods, exhibit unhealthy eating behaviors, and have inadequate intake of vegetables. Additionally, diets high in salt, meat, fried foods, and typical Western dietary patterns are linked to an increased risk of metabolic syndrome (28, 29).

Given the rising prevalence of heart disease and diabetes in recent years, and the high prevalence of metabolic syndrome in Iran, particularly in Mazandaran province with its high rates of hypertension, hyperglycemia, and obesity, it is crucial to explore the connection between metabolic syndrome, dietary patterns, and physical activity levels among individuals enrolled in the Tabari cohort study.

2. Objectives

To reduce deaths from type 2 diabetes and heart disease, this study was conducted to determine the relationship between metabolic syndrome, physical activity, and eating habits in participants of the Tabari cohort study.

3. Methods

Data from the Tabari Cohort study (TCS) was utilized in this research. The purpose and structure of the Tabari Cohort have been described in previous publications (30). Details of the PERSIAN cohort study can also be found elsewhere (31, 32). The TCS registration period began in June 2015 and concluded in November 2017. Sampling was conducted through a census approach. Health volunteers in urban areas and medical personnel in mountainous regions recruited eligible individuals to the cohort center. Of the 12,191 qualified urban residents, 7,012 were included in the Tabari cohort. In mountainous areas, 4,417 individuals were eligible, with 3,243 being admitted into the cohort. Therefore, a total of 10,255 participants aged 35 to 70 were enrolled during the initial phase of the TCS (33).

3.1. Research Community

This study utilized data from the first cohort study of Mazandaran's urban and mountain population (the first phase took place between June 1, 2015, and November 9, 2017).

3.2. Research Sample

Sampling was based on the census method, with 10,255 adults participating in the Tabari Cohort study.

3.3. Inclusion and Exclusion Criteria

Inclusion criteria: All participants in the Tabari cohort study. Exclusion criteria: individuals with autoimmune, malignant, or endocrine diseases; those taking corticosteroids or antipsychotic drugs; pregnant and lactating women; and individuals who had taken nutritional supplements or herbal medicines in the last six months.

3.4. Data Collection

Data collection was carried out using a standardized questionnaire at the Tabari Cohort center. Information such as underlying diseases, height, weight, waist circumference, Body Mass Index, blood pressure, blood sugar, triglycerides, and HDL cholesterol were measured and recorded. Participants' eating habits were assessed using the food habit questionnaire (34), and their physical activity was measured using the Cohort Physical Activity Questionnaire. Physical activity was measured using the metabolic equivalent rates (METs) of self-reported daily activities in the PERSIAN cohort (30-32). Anthropometric measurements such as height, weight, and waist circumference were assessed following established guidelines (30, 32, 35). The height

of patients was measured using the SECA 226 (SECA, Hamburg, Germany), and their weight was measured using the SECA 755 (SECA, Hamburg, Germany). All patients were asked to stand still with minimal clothing and no shoes, maintaining balance in the center of the scale until the measurement was taken. Waist circumference was measured transversely at the upper border of the iliac crest using the NIH index (30, 32). In the first step, all participants were examined for metabolic syndrome based on the ATP III criteria. The nutritional patterns and physical activity levels of individuals with metabolic syndrome were investigated using data from the Tabari cohort study, which was collected and recorded by experts using standardized questionnaires.

Metabolic Equivalent of Task Index (MET): The MET Index quantifies the energy expenditure rate during a specific physical activity relative to energy expenditure at rest, typically set at 3.5 mL/kg per minute. This index is a valuable tool for assessing physical activity levels in the general population, particularly in epidemiological studies (36). Extensive information on MET values for various activities is available in online compendiums (37).

Historically, MET was defined as a multiple of an individual's resting metabolic rate (RMR), which can be determined through gas exchange measurement, heat output assessment, or steady-state diet assessment in sedentary conditions. It can also be estimated based on age, gender, height, weight, and fitness level. In essence, 1 MET represents the ratio between the metabolic rate during activity and at rest, with 1 MET being equivalent to the energy expenditure of sitting still. Additionally, 1 MET corresponds to an oxygen consumption rate of 3.5 mL/kg/min, which is the oxygen cost of sitting quietly. Originally developed for epidemiological surveys, the MET allows participants to report the duration of specific physical activities they engage in.

3.5. Data Analysis

The data were analyzed using SPSS V.23 software. Qualitative variables were described using frequency and percentage, while quantitative variables were summarized using mean and standard deviation. Demographic and epidemiological variables were compared between two groups using the chi-square test, and quantitative variables were compared using the independent *t*-test. A P-value of less than 0.05 was considered statistically significant.

3.6. Ethical Considerations

This descriptive cross-sectional study was conducted with ethical approval from the research ethics committee of Mazandaran University of Medical Sciences (code: [IR.MAZUMS.IMAMHOSPITAL.REC.1402.062](https://doi.org/10.2196/IR.MAZUMS.IMAMHOSPITAL.REC.1402.062)).

All participants provided written consent, were informed about the study, and were assured of confidentiality. No fees were charged to participants for their involvement in the study.

4. Results

4.1. Profile of Study Participants

A total of 10,255 adults (4,149 males and 6,106 females) between the ages of 35 and 70 participated in this study. After applying the inclusion and exclusion criteria, 9,976 participants were included in the final analysis, including 4,083 individuals with metabolic syndrome and 5,893 healthy individuals. The average Body Mass Index of individuals with metabolic syndrome was 30.6 ± 4.5 kg/m², while healthy individuals had an average BMI of 26.8 ± 4.4 kg/m². No significant difference in BMI was observed between the two groups. However, individuals with metabolic syndrome exhibited significantly higher values for height, weight, and abdominal circumference compared to the control group ($P = 0.001$) (Table 1).

The study's participants were found to engage in an average of 43.03 ± 9.27 MET hours/day, with a median of 40.68 (range: 24 - 103) hours/day. It was noted that 57.0% of individuals with metabolic syndrome and 45.2% of those without had MET values lower than the median ($P = 0.001$). Regarding dietary habits, the metabolic syndrome group consumed an average total energy intake of 2189.74 ± 721.90 kcal, while the healthy group consumed 2354.92 ± 4.04 kcal, showing a significant difference ($P = 0.006$). Significant variations were also observed between the groups in the intake levels of carbohydrates, alcohol, caffeine, fiber, potassium, beta-carotene, trans fatty acids, total saturated fats ($P = 0.001$), and omega-3 (Table 2). However, there were no notable differences in the intake of other nutrients analyzed.

Based on this research, significant differences were found between the two groups in terms of salt usage in food, consumption of fried foods like potatoes, onions, vegetables, and the type of oil used ($P = 0.001$). The study revealed that individuals with metabolic syndrome had lower rates of constantly (8.7%) or sometimes (16.4%) adding salt to their meals compared to the healthy group, where continuous (12.6%) and occasional (21.6%) salt addition was more common. Additionally, the

consumption of fried foods 1 - 3 times a week was notably higher in the metabolic syndrome group (49.4%) compared to the healthy group (46.5%) ($P = 0.001$) (Table 3).

5. Discussion

Considering the increase in the prevalence of heart diseases and diabetes in recent years, and the high prevalence of metabolic syndrome in Iran, especially in Mazandaran province—with its high rates of hypertension, hyperglycemia, and obesity—the study aimed to explore the connection between metabolic syndrome, dietary habits, and physical activity levels among individuals enrolled in the Tabari cohort study. Bread and rice are among the main food staples in Mazandaran, and reducing the prevalence of metabolic syndrome is essential for significantly reducing deaths caused by heart diseases and diabetes.

The study indicated a direct correlation between lower physical activity levels and increased susceptibility to coronary heart disease. Engaging in moderate physical activities, such as walking, was linked to a notable reduction in the risk of coronary heart disease. Vigorous activities with MET levels in the range of 5 - 9 were associated with a significant reduction in risk factors, such as high BMI, heart rate, systolic and diastolic blood pressure, and low HDL cholesterol levels among individuals with metabolic syndrome. Similar trends were observed in individuals with low physical activity levels and limited cardiorespiratory fitness (38). This aligns with the present study's findings, where patients diagnosed with metabolic syndrome exhibited a significantly lower MET Index compared to healthy individuals (38).

Zhang et al.'s meta-analysis revealed a direct correlation between leisure-time physical activity (LTPA) and the prevalence of metabolic syndrome. For every additional 10 hours per week of increased MET in LTPA, there was an 8% decrease in the risk of developing metabolic syndrome. Furthermore, meeting the recommended guidelines of 150 minutes of moderate physical activity per week (equivalent to 10 MET hours per week) resulted in a 10% decrease in the risk of metabolic syndrome onset compared to being inactive. Exceeding the recommended physical activity levels led to further reductions in the risk of metabolic syndrome, consistent with the findings of this study (12).

In another study by Sayon-Orea et al. (39), the relationship between metabolic syndrome and the consumption of fried foods was examined. Regular consumption of fried foods did not show a significant link to the development of metabolic syndrome (HR =

Table 1. Biometric Characteristics of the People Included in the Study

Variables	Healthy (N = 5893)	Metabolic Syndrome (N = 4083)	P-Value
Height (cm)	162.5 ± 9.0	159.9 ± 8.8	0.001
Weight (kg)	78.6 ± 13.5	70.92 ± 12.7	0.002
BMI (kg/m ²)	26.8 ± 4.4	30.6 ± 4.5	0.138
Waist circumference (cm)	100.5 ± 8.4	106.1 ± 9.3	0.001
Abdominal circumference (cm)	89.4 ± 10.4	99.4 ± 9.9	0.001
Wrist circumference (cm)	1.2 ± 16.7	1.3 ± 17.1	0.081

Abbreviation: BMI: Body Mass Index.

Table 2. Amount of Micronutrients and Macronutrients Received per Day in the Study Groups

Amount Received	Healthy (n = 5893)	Metabolic Syndrome (n = 4083)	P-Value
Energy (calories per day)	2354.92 ± 754.04	721.90 ± 2189.74	0.006
Protein (grams per day)	25.43 ± 74.98	25.40 ± 71.07	0.862
Fat (g/day)	60.77 ± 23.99	55.87 ± 23.86	0.100
Carbohydrates (grams per day)	134.91 ± 390.20	126.17 ± 364.97	0.001
Fructose (grams per day)	32.41 ± 18.86	34.06 ± 18.90	0.143
Alcohol (grams per day)	0.11 ± 0.03	0.10 ± 0.02	0.001
Caffeine (mg per day)	124.52 ± 164.54	110.54 ± 152.35	0.001
Sugar (grams per day)	62.74 ± 137.78	59.81 ± 130.23	0.109
Fiber (grams per day)	27.91 ± 11.46	28.68 ± 11.86	0.007
Magnesium (mg per day)	351.53 ± 116.98	344.36 ± 120.31	0.097
Potassium (mg/day)	3787.92 ± 1396.36	1461.14 ± 3829.92	0.001
Sodium (mg/day)	4140.39 ± 1756.23	1757.85 ± 3941.61	0.438
Selenium (mg per day)	56.86 ± 29.79	53.34 ± 29.91	0.832
Beta-carotene (micrograms per day)	4961.50 ± 3538.17	3759.06 ± 5412.55	0.001
Vitamin E (mg per day)	3.75 ± 7.27	3.81 ± 7.19	0.782
Vitamin D (micrograms per day)	1.44 ± 0.90	1.35 ± 0.96	0.432
Vitamin C (mg per day)	112.91 ± 163.08	116.24 ± 172.00	0.041
Cholesterol (mg/day)	264.92 ± 134.57	235.94 ± 137.42	0.051
Trans fatty acid (grams per day)	0.29 ± 0.29	0.27 ± 0.24	0.001
Saturated fatty acid (g/day)	22.67 ± 10.33	9.82 ± 20.49	0.001
Monounsaturated fatty acid (g/day)	18.84 ± 8.04	8.31 ± 17.26	0.471
Polyunsaturated fatty acid (g/day)	12.71 ± 5.76	5.81 ± 11.94	0.914
Three-Total Omega (grams per day)	0.033 ± 0.044	0.037 ± 0.043	0.027
Docosahexaenoic (acid grams per day)	0.024 ± 0.029	0.025 ± 0.032	0.001
Cozapentanoic acid (grams per day)	0.49 ± 0.35	0.45 ± 0.33	0.167
Alphalinolic acid (grams per day)	11.86 ± 11.17	11.15 ± 12.37	0.767
Omega 6 (grams per day)	5.22 ± 3.73	3.70 ± 4.78	0.068

0.98, 95% CI 0.77-1.26, P = 0.862 for trend). However, central obesity and hypertension demonstrated a positive correlation with fried food intake. Individuals who fried foods using oils or fats other than olive oil had a higher risk of metabolic syndrome, though these associations were not statistically significant.

A study conducted by Jusuf et al. found that the consumption of fatty, cooked, and salty foods was

significantly linked to metabolic syndrome (40). Similarly, research by Lutsey et al. found that the intake of red meat, fried foods, and diet soda was positively correlated with metabolic syndrome, whereas consuming dairy products had a protective effect (41). However, in the present study, the consumption of fried foods in the metabolic syndrome group was significantly lower than in the healthy control group.

Table 3. Eating Habits of the Studied Groups

Variables	Healthy (N = 5893)	Metabolic Syndrome (N = 4083)	P-Value
Number of meals per day			0.163
Three meals (breakfast, lunch and dinner)	720 (12.2)	436 (10.7)	
Four meals (three meals plus a snack)	2292 (38.9)	1607 (39.4)	
Five to six meals (three meals plus 2-3 snacks)	2548 (43.2)	1821 (44.6)	
More than 6 servings	272 (4.6)	181 (4.4)	
Less than 3 servings	61 (1.0)	38 (0.9)	
Adding salt to food			0.001
Yes	741 (12.6)	354 (8.7)	
Sometimes	1270 (21.6)	670 (16.4)	
No	3881 (65.9)	3059 (74.9)	
Eating grilled foods			0.595
Never	125 (2.1)	106 (2.6)	
Less than once a month	1525 (25.9)	1068 (26.2)	
One to three times a month	2946 (50.0)	2031 (49.7)	
One to three times a week	1284 (21.8)	870 (21.3)	
Daily	13 (0.2)	8 (0.2)	
Eating fried foods			0.001
Never	32 (0.5)	50 (1.2)	
Less than once a month	43 (0.7)	41 (1.0)	
One to three times a month	279 (4.7)	299 (7.3)	
One to three times a week	2743 (46.5)	2018 (49.4)	
Daily	2796 (47.4)	1675 (41.0)	
Consumption of french fries			0.001
Slightly fried	417 (7.1)	353 (8.6)	
Semi-fried	4165 (70.7)	2762 (67.6)	
Fully fried	1231 (20.9)	830 (20.3)	
No consumption	80 (1.4)	138 (3.4)	
Eating fried onions			
Slightly fried	1028 (17.4)	805 (19.7)	
Semi-fried	4211 (71.5)	2814 (68.9)	
Fully fried	614 (10.4)	403 (9.9)	
No consumption	40 (0.7)	61 (1.5)	
Eating fried vegetables			0.001
Slightly fried	4190 (71.1)	2742 (67.2)	
Semi-fried	998 (16.9)	737 (18.1)	
Fully fried	502 (8.5)	407 (10.0)	
No consumption	203 (3.4)	197 (4.8)	
Consumable oil			0.001
Solid	1360 (23.1)	844 (20.7)	
Semi solid	191 (3.2)	104 (2.5)	
Liquid	382 (6.5)	333 (8.2)	
Fried	3840 (65.2)	2664 (65.2)	
Other types of oil	111 (1.9)	111 (2.7)	
No oil consumption	9 (0.2)	27 (0.7)	
Used oil consumption			0.270
Yes	4049 (68.7)	2874 (69.7)	
No	1844 (31.3)	1235 (30.3)	
Consumption of smoked foods			0.769
Never	5652 (95.9)	3895 (95.4)	
Less than once a month	208 (3.5)	161 (3.9)	
One to three times a month	23 (0.4)	20 (0.5)	
One to three times a week	5 (0.1)	3 (0.1)	
Daily	5 (0.1)	4 (0.1)	
Eating fresh vegetables			0.544
Yes	302 (5.1)	198 (4.8)	
No	5590 (94.9)	3885 (95.2)	
The temperature of tea consumed			0.068
Very hot	1984 (33.7)	1282 (31.4)	
Hot	3617 (61.4)	2579 (63.2)	
Cold	249 (4.2)	196 (4.8)	
Not drinking tea	43 (0.7)	26 (0.6)	
The temperature of the soup			0.280
Very hot	1673 (28.4)	1130 (27.7)	
Hot	3936 (66.8)	2781 (68.1)	
Cold	259 (4.4)	161 (3.9)	
Not drinking tea	25 (0.4)	11 (0.3)	

One reason for this contradiction could be that the metabolic syndrome group was undergoing treatment with medication and dietary adjustments. It appears that the participants had recently been informed about

their condition and had begun to correct some of their eating habits.

Previous studies show that reducing the intake of saturated fatty acids (SFA) may be more effective in

preventing cardiometabolic risk factors (42, 43). However, recent studies have reported conflicting results regarding the effect of SFA consumption on cardiometabolic risk factors—some report no effect (44, 45), while others report a beneficial effect (46). For example, a prospective study examining nutrient interactions and coronary atherosclerosis risk found that SFA intake was associated with less progression of atherosclerosis, but only in those with lower MUFA intake (46).

A systematic review by Julibert et al. found that SFA intake was associated with hypertension and higher fasting triglyceride, HDL, and serum glucose levels (47). Studies have explored the impact of various fat types, showing that a higher unsaturated-to-saturated fat ratio (P/S ratio) positively influenced HDL cholesterol levels (48). However, polyunsaturated fatty acid (PUFA) intake was inversely related to HDL cholesterol and waist circumference (48, 49). Consumption of PUFA n-3 was linked to improved blood pressure, triglyceride levels, glucose tolerance, and reductions in insulin resistance, waist circumference, and obesity prevalence (49, 50). Some studies suggest that higher fish intake may reduce the risk of metabolic syndrome by improving triglyceride and HDL cholesterol levels (49, 51).

However, the present study found that the metabolic syndrome group had lower consumption of trans and saturated fatty acids compared to the healthy group. This discrepancy may be attributed to the fact that some participants were undergoing dietary interventions for metabolic syndrome. The study was conducted during the first phase of the Tabari cohort, which included patients following dietary changes, potentially impacting the outcomes. Further investigation in the cohort's second phase is necessary for more comprehensive insights.

In this study, fructose consumption was higher among healthy individuals compared to the metabolic syndrome group. Although fructose is a natural sugar found in fruits, excessive consumption has been linked to hyperlipidemia (52), diabetes, obesity, and certain cancers (53). While moderate consumption of fructose does not pose a problem, long-term overconsumption has been shown to increase the risk of obesity and conditions related to metabolic syndrome. Excessive fructose intake can also lead to fat accumulation, insulin resistance, and inflammation (54). Further investigation in future cohort phases is recommended to assess the prevalence of metabolic syndrome and refine dietary recommendations.

5.1. Limitations

Several limitations should be considered when interpreting the study's findings. Eating habits were evaluated only based on food intake, without accounting for dietary behaviors such as meal patterns, snack frequency, and meal timing (55). The study's cross-sectional nature also limits the ability to establish causality. Future studies should assess the relationship between dietary habits and metabolic syndrome in a prospective manner. Additionally, the findings cannot be generalized to the entire country, as eating habits are influenced by socio-economic factors. Nonetheless, participants from different regions with varying socio-economic statuses were included. Lastly, although we controlled for many confounding variables related to lifestyle, genetic factors were not accounted for, and they may confound the relationship between diet and metabolic syndrome.

5.2. Conclusions

The Tabari cohort study highlights the high prevalence of metabolic syndrome in Mazandaran, underscoring the importance of prevention and screening programs. Metabolic syndrome is a significant risk factor for diabetes and cardiovascular diseases, which have a high economic burden on individuals and healthcare systems. Contrary to previous studies, the present research found that the consumption of fried foods, trans fatty acids, and calories, as well as fructose intake, were lower in the metabolic syndrome group compared to the healthy group, indicating the positive effects of treatment programs in Mazandaran province. Based on the findings of this cross-sectional descriptive study, conducted during the first phase of the Tabari cohort, it is recommended that further studies be carried out to verify the types of food ingredients consumed in the current period (e.g., frying ingredients, trans fats, fructose intake, etc.). Future research should also evaluate the effects of dietary and lifestyle changes among individuals with metabolic syndrome, and their impact on preventing metabolic diseases such as diabetes and cardiovascular diseases. This should be explored in the second phase of the Tabari cohort to help reduce household healthcare expenses and alleviate the health burden on the country's healthcare system. Additionally, prospective studies are suggested to investigate the influence of these factors on reducing the incidence of related diseases and the associated costs.

Acknowledgements

We sincerely thank the esteemed professors who provided invaluable support in writing this project. We are also grateful to our colleagues who were responsible for collecting and organizing data across various healthcare departments. Lastly, we express our heartfelt thanks to God for granting us the strength to think, analyze, and complete this work.

Footnotes

Authors' Contribution: A. R. and Z. K.: Developed the original idea and the protocol; M. M. and A. R.: Analysis and interpretation of data; L. M., A. B. and A. H. Kh.: Wrote the manuscript; A. R., Z. K., A. B., L. M., M. M. and A. H. Kh.: Contributed to the development of the protocol, abstracted data, and prepared the manuscript.

Conflict of Interests Statement: The authors declared that they have no conflict of interest.

Data Availability: The dataset presented in the study is available on request from the corresponding author during submission or after publication.

Ethical Approval: The ethical approval code is IR.MAZUMS.IMAMHOSPITAL.REC.1402.062.

Funding/Support: This study was supported in part by grant and design number 18405 from the Mazandaran University of Medical Sciences, Sari, Iran.

Informed Consent: Informed consent was obtained from the participants.

References

- Geloneze B, Repetto EM, Geloneze SR, Tambascia MA, Ermetice MN. The threshold value for insulin resistance (HOMA-IR) in an admixed population IR in the Brazilian Metabolic Syndrome Study. *Diabetes Res Clin Pract.* 2006;**72**(2):219-20. [PubMed ID: 16310881]. <https://doi.org/10.1016/j.diabres.2005.10.017>.
- Alberti KG, Eckel RH, Grundy SM, Zimmet PZ, Cleeman JI, Donato KA, et al. Harmonizing the metabolic syndrome: a joint interim statement of the International Diabetes Federation Task Force on Epidemiology and Prevention; National Heart, Lung, and Blood Institute; American Heart Association; World Heart Federation; International Atherosclerosis Society; and International Association for the Study of Obesity. *Circulation.* 2009;**120**(16):1640-5. [PubMed ID: 19805654]. <https://doi.org/10.1161/CIRCULATIONAHA.109.192644>.
- Park YW, Zhu S, Palaniappan L, Heshka S, Carnethon MR, Heymsfield SB. The metabolic syndrome: prevalence and associated risk factor findings in the US population from the Third National Health and Nutrition Examination Survey, 1988-1994. *Arch Intern Med.* 2003;**163**(4):427-36. [PubMed ID: 12588201]. [PubMed Central ID: PMC3146257]. <https://doi.org/10.1001/archinte.163.4.427>.
- Ralston J, Brinsden H, Buse K, Candeias V, Caterson I, Hassell T, et al. Time for a new obesity narrative. *Lancet.* 2018;**392**(10156):1384-6. [PubMed ID: 30316458]. [https://doi.org/10.1016/S0140-6736\(18\)32537-6](https://doi.org/10.1016/S0140-6736(18)32537-6).
- Aguilar M, Bhuket T, Torres S, Liu B, Wong RJ. Prevalence of the metabolic syndrome in the United States, 2003-2012. *JAMA.* 2015;**313**(19):1973-4. [PubMed ID: 25988468]. <https://doi.org/10.1001/jama.2015.4260>.
- Ford ES, Giles WH, Dietz WH. Prevalence of the metabolic syndrome among US adults: findings from the third National Health and Nutrition Examination Survey. *JAMA.* 2002;**287**(3):356-9. [PubMed ID: 11790215]. <https://doi.org/10.1001/jama.287.3.356>.
- Hirode G, Wong RJ. Trends in the Prevalence of Metabolic Syndrome in the United States, 2011-2016. *JAMA.* 2020;**323**(24):2526-8. [PubMed ID: 32573660]. [PubMed Central ID: PMC7312413]. <https://doi.org/10.1001/jama.2020.4501>.
- Lindsay RS, Howard BV. Cardiovascular risk associated with the metabolic syndrome. *Curr Diab Rep.* 2004;**4**(1):63-8. [PubMed ID: 14764282]. <https://doi.org/10.1007/s11892-004-0013-9>.
- Koh KK, Han SH, Quon MJ. Inflammatory markers and the metabolic syndrome: insights from therapeutic interventions. *J Am Coll Cardiol.* 2005;**46**(11):1978-85. [PubMed ID: 16325028]. <https://doi.org/10.1016/j.jacc.2005.06.082>.
- van Namen M, Prendergast L, Peiris C. Supervised lifestyle intervention for people with metabolic syndrome improves outcomes and reduces individual risk factors of metabolic syndrome: A systematic review and meta-analysis. *Metabolism.* 2019;**101**:153988. [PubMed ID: 31672441]. <https://doi.org/10.1016/j.metabol.2019.153988>.
- Wewege MA, Thom JM, Rye KA, Parmenter BJ. Aerobic, resistance or combined training: A systematic review and meta-analysis of exercise to reduce cardiovascular risk in adults with metabolic syndrome. *Atherosclerosis.* 2018;**274**:162-71. [PubMed ID: 29783064]. <https://doi.org/10.1016/j.atherosclerosis.2018.05.002>.
- Zhang D, Liu X, Liu Y, Sun X, Wang B, Ren Y, et al. Leisure-time physical activity and incident metabolic syndrome: a systematic review and dose-response meta-analysis of cohort studies. *Metabolism.* 2017;**75**:36-44. [PubMed ID: 28927737]. <https://doi.org/10.1016/j.metabol.2017.08.001>.
- World Health Organization. *WHO guidelines on physical activity and sedentary behaviour: at a glance, 4 May 2021.* Geneva, Switzerland: World Health Organization; 2021. Available from: <https://www.who.int/europe/publications/item/9789240014886#:~:text=For%20health%20and%20wellbeing%2C%20WHO,All%20physical%20activity%20counts..>
- Katzmarzyk PT, Powell KE, Jakicic JM, Troiano RP, Piercy K, Tennant B. Sedentary Behavior and Health: Update from the 2018 Physical Activity Guidelines Advisory Committee. *Med Sci Sports Exerc.* 2019;**51**(6):1227-41. [PubMed ID: 31095080]. [PubMed Central ID: PMC6527341]. <https://doi.org/10.1249/mss.0000000000001935>.
- Brien SE, Katzmarzyk PT. Physical activity and the metabolic syndrome in Canada. *Appl Physiol Nutr Metab.* 2006;**31**(1):40-7. [PubMed ID: 16604140]. <https://doi.org/10.1139/h05-024>.
- Lakka TA, Laaksonen DE. Physical activity in prevention and treatment of the metabolic syndrome. *Appl Physiol Nutr Metab.* 2007;**32**(1):76-88. [PubMed ID: 17332786]. <https://doi.org/10.1139/h06-113>.
- Yang X, Telama R, Hirvensalo M, Mattsson N, Viikari JS, Raitakari OT. The longitudinal effects of physical activity history on metabolic syndrome. *Med Sci Sports Exerc.* 2008;**40**(8):1424-31. [PubMed ID: 18614950]. <https://doi.org/10.1249/MSS.0b013e318172ced4>.
- Laaksonen DE, Lindstrom J, Lakka TA, Eriksson JG, Niskanen L, Wikstrom K, et al. Physical activity in the prevention of type 2 diabetes: the Finnish diabetes prevention study. *Diabetes.* 2005;**54**(1):158-65. [PubMed ID: 15616024]. <https://doi.org/10.2337/diabetes.54.1.158>.

19. Diaz KM, Shimbo D. Physical activity and the prevention of hypertension. *Curr Hypertens Rep.* 2013;**15**(6):659-68. [PubMed ID: 24052212]. [PubMed Central ID: PMC3901083]. <https://doi.org/10.1007/s11906-013-0386-8>.
20. Burns RD. Public Health Implications of the Dose-Response Association Between Physical Activity and Cardiometabolic Health in Young Adults. *J Adolesc Health.* 2020;**67**(2):155-6. [PubMed ID: 32448598]. [PubMed Central ID: PMC7241332]. <https://doi.org/10.1016/j.jadohealth.2020.05.008>.
21. Rivera Medina C, Briones Urbano M, de Jesus Espinosa A, Toledo Lopez A. Eating Habits Associated with Nutrition-Related Knowledge among University Students Enrolled in Academic Programs Related to Nutrition and Culinary Arts in Puerto Rico. *Nutrients.* 2020;**12**(5):1408. [PubMed ID: 32422880]. [PubMed Central ID: PMC7285020]. <https://doi.org/10.3390/nu12051408>.
22. Lallukka T, Laaksonen M, Rahkonen O, Roos E, Lahelma E. Multiple socio-economic circumstances and healthy food habits. *Eur J Clin Nutr.* 2007;**61**(6):701-10. [PubMed ID: 17180154]. <https://doi.org/10.1038/sj.ejcn.1602583>.
23. Johansson L, Thelle DS, Solvoll K, Bjorneboe GE, Drevon CA. Healthy dietary habits in relation to social determinants and lifestyle factors. *Br J Nutr.* 1999;**81**(3):211-20. [PubMed ID: 10434847]. <https://doi.org/10.1017/s0007114599000409>.
24. Adeva-Andany MM, Rana-Muino E, Vila-Altesor M, Fernandez-Fernandez C, Funcasta-Calderon R, Castro-Quintela E. Dietary habits contribute to define the risk of type 2 diabetes in humans. *Clin Nutr ESPEN.* 2019;**34**:8-17. [PubMed ID: 31677716]. <https://doi.org/10.1016/j.clnesp.2019.08.002>.
25. McCracken E, Monaghan M, Sreenivasan S. Pathophysiology of the metabolic syndrome. *Clin Dermatol.* 2018;**36**(1):14-20. [PubMed ID: 29241747]. <https://doi.org/10.1016/j.clindermatol.2017.09.004>.
26. Yoshida J, Eguchi E, Nagaoka K, Ito T, Ogino K. Association of night eating habits with metabolic syndrome and its components: a longitudinal study. *BMC Public Health.* 2018;**18**(1):1366. [PubMed ID: 30537972]. [PubMed Central ID: PMC6288903]. <https://doi.org/10.1186/s12889-018-6262-3>.
27. Shin A, Lim SY, Sung J, Shin HR, Kim J. Dietary intake, eating habits, and metabolic syndrome in Korean men. *J Am Diet Assoc.* 2009;**109**(4):633-40. [PubMed ID: 19328258]. <https://doi.org/10.1016/j.jada.2008.12.015>.
28. Sarebanhassanabadi M, Mirhosseini SJ, Mirzaei M, Namayandeh SM, Soltani MH, Pakeresht M, et al. Effect of dietary habits on the risk of metabolic syndrome: Yazd Healthy Heart Project. *Public Health Nutr.* 2018;**21**(6):1139-46. [PubMed ID: 29233205]. [PubMed Central ID: PMC10261386]. <https://doi.org/10.1017/S1368980017003627>.
29. Lorzadeh E, Sangsefidi ZS, Mirzaei M, Hosseinzadeh M. Dietary Habits and their Association with Metabolic Syndrome in a sample of Iranian adults: A population-based study. *Food Sci Nutr.* 2020;**8**(11):6217-25. [PubMed ID: 33282272]. [PubMed Central ID: PMC7684605]. <https://doi.org/10.1002/fsn3.1918>.
30. Kheradmand M, Moosazadeh M, Saeedi M, Poustchi H, Eghtesad S, Esmaeili R, et al. Tabari Cohort Profile and Preliminary Results in Urban Areas and Mountainous Regions of Mazandaran, Iran. *Arch Iran Med.* 2019;**22**(6):279-85. [PubMed ID: 31356093].
31. Poustchi H, Eghtesad S, Kamangar F, Etemadi A, Keshtkar AA, Hekmatdoost A, et al. Prospective Epidemiological Research Studies in Iran (the PERSIAN Cohort Study): Rationale, Objectives, and Design. *Am J Epidemiol.* 2018;**187**(4):647-55. [PubMed ID: 29145581]. [PubMed Central ID: PMC6279089]. <https://doi.org/10.1093/aje/kwx314>.
32. Eghtesad S, Mohammadi Z, Shayanrad A, Faramarzi E, Joukar F, Hamzeh B, et al. The PERSIAN Cohort: Providing the Evidence Needed for Healthcare Reform. *Arch Iran Med.* 2017;**20**(11):691-5. [PubMed ID: 29480734].
33. Bahar A, Kashi Z, Kheradmand M, Hedayatizadeh-Omran A, Moradinazar M, Ramezani F, et al. Prevalence of metabolic syndrome using international diabetes federation, National Cholesterol Education Panel- Adult Treatment Panel III and Iranian criteria: results of Tabari cohort study. *J Diabetes Metab Disord.* 2020;**19**(1):205-11. [PubMed ID: 32550169]. [PubMed Central ID: PMC7270474]. <https://doi.org/10.1007/s40200-020-00492-6>.
34. Eghtesad S, Hekmatdoost A, Faramarzi E, Homayounfar R, Sharafkhan M, Hakimi H, et al. Validity and reproducibility of a food frequency questionnaire assessing food group intake in the PERSIAN Cohort Study. *Front Nutr.* 2023;**10**:1059870. [PubMed ID: 37599697]. [PubMed Central ID: PMC10436288]. <https://doi.org/10.3389/fnut.2023.1059870>.
35. Klein S, Allison DB, Heymsfield SB, Kelley DE, Leibel RL, Nonas C, et al. Waist circumference and cardiometabolic risk: a consensus statement from Shaping America's Health: Association for Weight Management and Obesity Prevention; NAAO, The Obesity Society; the American Society for Nutrition; and the American Diabetes Association. *Am J Clin Nutr.* 2007;**85**(5):1197-202. [PubMed ID: 17490953]. <https://doi.org/10.1093/ajcn/85.5.1197>.
36. Ainsworth BE, Haskell WL, Herrmann SD, Meckes N, Bassett DJ, Tudor-Locke C, et al. 2011 Compendium of Physical Activities: a second update of codes and MET values. *Med Sci Sports Exerc.* 2011;**43**(8):1575-81. [PubMed ID: 21681120]. <https://doi.org/10.1249/MSS.0b013e31821ecce2>.
37. Ainsworth BE, Haskell WL, Leon AS, Jacobs DR, Montoye H, Sallis JF, et al. Compendium of physical activities: classification of energy costs of human physical activities. *Med Sci Sports Exerc.* 1993;**25**(1):71-80. [PubMed ID: 8292105]. <https://doi.org/10.1249/00005768-199301000-00011>.
38. Rennie KL, McCarthy N, Yazdgerdi S, Marmot M, Brunner E. Association of the metabolic syndrome with both vigorous and moderate physical activity. *Int J Epidemiol.* 2003;**32**(4):600-6. [PubMed ID: 12913036]. <https://doi.org/10.1093/ije/dygi79>.
39. Sayon-Orea C, Martinez-Gonzalez MA, Gea A, Flores-Gomez E, Basterra-Gortari FJ, Bes-Rastrollo M. Consumption of fried foods and risk of metabolic syndrome: the SUN cohort study. *Clin Nutr.* 2014;**33**(3):545-9. [PubMed ID: 23954218]. <https://doi.org/10.1016/j.clnu.2013.07.014>.
40. Jusuf H, Rahma S, Monayo ER. Food consumption behavior and their association with metabolic syndrome: A cross-sectional study of adult in Gorontalo Province, Indonesia. *Syst Rev Pharm.* 2020;**11**(5):556-61.
41. Lutsey PL, Steffen LM, Stevens J. Dietary intake and the development of the metabolic syndrome: the Atherosclerosis Risk in Communities study. *Circulation.* 2008;**117**(6):754-61. [PubMed ID: 18212291]. <https://doi.org/10.1161/CIRCULATIONAHA.107.716159>.
42. Chiuev SE, Rimm EB, Sandhu RK, Bernstein AM, Rexrode KM, Manson JE, et al. Dietary fat quality and risk of sudden cardiac death in women. *Am J Clin Nutr.* 2012;**96**(3):498-507. [PubMed ID: 22854398]. [PubMed Central ID: PMC3417213]. <https://doi.org/10.3945/ajcn.112.040287>.
43. Trichopoulou A, Psaltopoulou T, Orfanos P, Trichopoulos D. Diet and physical activity in relation to overall mortality amongst adult diabetics in a general population cohort. *J Intern Med.* 2006;**259**(6):583-91. [PubMed ID: 16704559]. <https://doi.org/10.1111/j.1365-2796.2006.01638.x>.
44. Harding AH, Day NE, Khaw KT, Bingham S, Luben R, Welsh A, et al. Dietary fat and the risk of clinical type 2 diabetes: the European prospective investigation of Cancer-Norfolk study. *Am J Epidemiol.* 2004;**159**(1):73-82. [PubMed ID: 14693662]. <https://doi.org/10.1093/aje/kwh004>.
45. Meyer KA, Kushi LH, Jacobs DR, Folsom AR. Dietary fat and incidence of type 2 diabetes in older Iowa women. *Diabetes Care.*

- 2001;**24**(9):1528-35. [PubMed ID: [11522694](#)]. <https://doi.org/10.2337/diacare.24.9.1528>.
46. Mozaffarian D, Rimm EB, Herrington DM. Dietary fats, carbohydrate, and progression of coronary atherosclerosis in postmenopausal women. *Am J Clin Nutr*. 2004;**80**(5):1175-84. [PubMed ID: [15531663](#)]. [PubMed Central ID: [PMC1270002](#)]. <https://doi.org/10.1093/ajcn/80.5.1175>.
47. Julibert A, Bibiloni MDM, Tur JA. Dietary fat intake and metabolic syndrome in adults: A systematic review. *Nutr Metab Cardiovasc Dis*. 2019;**29**(9):887-905. [PubMed ID: [31377181](#)]. <https://doi.org/10.1016/j.numecd.2019.05.055>.
48. Shab-Bidar S, Hosseini-Esfahani F, Mirmiran P, Hosseinpour-Niazi S, Azizi F. Metabolic syndrome profiles, obesity measures and intake of dietary fatty acids in adults: Tehran Lipid and Glucose Study. *J Hum Nutr Diet*. 2014;**27** Suppl 2:98-108. [PubMed ID: [23731333](#)]. <https://doi.org/10.1111/jhn.12117>.
49. Noel SE, Newby PK, Ordovas JM, Tucker KL. Adherence to an (n-3) fatty acid/fish intake pattern is inversely associated with metabolic syndrome among Puerto Rican adults in the Greater Boston area. *J Nutr*. 2010;**140**(10):1846-54. [PubMed ID: [20702744](#)]. [PubMed Central ID: [PMC2937577](#)]. <https://doi.org/10.3945/jn.110.124297>.
50. Ebbesson SO, Tejero ME, Nobmann ED, Lopez-Alvarenga JC, Ebbesson L, Romenesko T, et al. Fatty acid consumption and metabolic syndrome components: the GOCADAN study. *J Cardiometab Syndr*. 2007;**2**(4):244-9. [PubMed ID: [18059206](#)]. <https://doi.org/10.1111/j.1559-4564.2007.07393.x>.
51. Park S, Ahn J, Kim NS, Lee BK. High carbohydrate diets are positively associated with the risk of metabolic syndrome irrespective to fatty acid composition in women: the KNHANES 2007-2014. *Int J Food Sci Nutr*. 2017;**68**(4):479-87. [PubMed ID: [27822963](#)]. <https://doi.org/10.1080/09637486.2016.1252318>.
52. Dolan LC, Potter SM, Burdock GA. Evidence-based review on the effect of normal dietary consumption of fructose on development of hyperlipidemia and obesity in healthy, normal weight individuals. *Crit Rev Food Sci Nutr*. 2010;**50**(1):53-84. [PubMed ID: [20047139](#)]. <https://doi.org/10.1080/10408390903461426>.
53. Jung S, Bae H, Song WS, Jang C. Dietary Fructose and Fructose-Induced Pathologies. *Annu Rev Nutr*. 2022;**42**:45-66. [PubMed ID: [35995049](#)]. [PubMed Central ID: [PMC9904196](#)]. <https://doi.org/10.1146/annurev-nutr-062220-025831>.
54. Legeza B, Marcolongo P, Gamberucci A, Varga V, Bánhegyi G, Benedetti A, et al. Fructose, Glucocorticoids and Adipose Tissue: Implications for the Metabolic Syndrome. *Nutrients*. 2017;**9**(5):422. [PubMed ID: [28445389](#)]. [PubMed Central ID: [PMC5452156](#)]. <https://doi.org/10.3390/nu9050426>.
55. Tseng M. Validation of dietary patterns assessed with a food-frequency questionnaire. *Am J Clin Nutr*. 1999;**70**(3):422. [PubMed ID: [10479207](#)]. <https://doi.org/10.1093/ajcn/70.3.422>.