Published online: 2024 April 24.

**Research Article** 

# Cardioprotective Effect of Hydroalcoholic Extract of *Lepidium draba* on Isoproterenol-Induced Acute Myocardial Infarction in Rats

Maryam Kahyaei-Aghdam <sup>1, 2</sup>, Vahid Shafiei-Irannejad 🔟 <sup>3</sup>, Sanaz Hamedeyazdan <sup>4</sup>, Hamid Soraya 🔟 <sup>1,\*</sup>

<sup>1</sup> Department of Pharmacology and Toxicology, School of Pharmacy, Urmia University of Medical Sciences, Urmia, Iran

<sup>2</sup> Student Research Committee, Urmia University of Medical Sciences, Urmia, Iran

<sup>3</sup> Cellular and Molecular Research Center, Cellular and Molecular Medicine Institute, Urmia University of Medical Sciences, Urmia, Iran

<sup>4</sup> Department of Pharmacognosy, Faculty of Pharmacy, Tabriz University of Medical Sciences, Tabriz, Iran

<sup>\*</sup>Corresponding author: Department of Pharmacology and Toxicology, School of Pharmacy, Urmia University of Medical Sciences, 571571441, Urmia, Iran. Tel: +98-4432754996; Fax: +98-4432754990; Email: soraya.h@umsu.ac.ir

Received 2024 February 18; Accepted 2024 April 24.

# Abstract

**Background:** Recently, there has been increased interest in medicinal plants for the discovery of new compounds to treat various disorders. A few studies have highlighted the antioxidant and anti-inflammatory effects of *Lepidium draba*.

**Objectives:** This study explores the potential cardioprotective effects of *L. draba* against myocardial infarction (MI) induced by isoproterenol.

**Methods:** Isoproterenol (ISO) (100 mg/kg, injected subcutaneously for two consecutive days) was utilized to induce acute myocardial infarction in rats. The treated groups were administered hydroalcoholic extracts of *L. draba intraperitoneally (ip)* at doses of 50, 250, and 500 mg/kg for 2 consecutive days. Subsequently, the electrocardiogram pattern, cardiodynamic parameters, cardiac remodeling, and lactate dehydrogenase levels were assessed.

**Results:** The findings revealed that isoproterenol administration elevated the ST-segment and diminished the R-amplitude on the electrocardiogram, whereas treatment with *L. draba* extracts significantly ameliorated the ECG pattern. Isoproterenol injections lowered the mean arterial pressure (MAP), which was significantly increased by *L. draba* extract at all doses (P < 0.05). Furthermore, isoproterenol injections elevated lactate dehydrogenase (LDH) levels (P < 0.05) and cardiac fibrosis (P < 0.001), and treatment with the hydroalcoholic extract of *L. draba* significantly reduced both indices (P < 0.001 and P < 0.05, respectively).

**Conclusions:** This study is the first to demonstrate that *L. draba* administration protects myocardial tissue from isoproterenolinduced infarction. Consequently, this plant could be recommended as an adjunct treatment for patients with acute MI.

Keywords: Lepidium draba, Myocardial Infarction, Isoproterenol, Cardiac Fibrosis, Hemodynamic

# 1. Background

Despite significant progress in the treatment and management of myocardial infarction (MI), the mortality rate remains high (1). Myocardial infarction occurs due to an imbalance between cardiac oxygen demand and the heart's inability to supply oxygen. In this situation, metabolic and ionic imbalances in myocardial cells and mitochondrial dysfunction play a key role in apoptosis and cardiac necrosis (2). Myocardial infarction can lead to significant changes in the patient's hemodynamics, and death may eventually occur (3). Some compensatory pathways are activated to improve heart function and reduce MI-induced damage. One of these pathways involves the replacement of damaged myocardial tissue with fibrotic tissue due to the presence of fibroblasts and myofibroblasts. Although this change is initially beneficial and prevents ventricular wall rupture, its continuation causes structural changes in the heart that eventually lead to heart dysfunction and the appearance of arrhythmias. Consequently, heart failure (HF) may occur (4, 5). Alterations in blood pressure are essential factors in myocardial infarction. Studies show that MI reduces

Copyright © 2024, Kahyaei-Aghdam et al. This open-access article is available under the Creative Commons Attribution 4.0 (CC BY 4.0) International License (https://creativecommons.org/licenses/by/4.0/), which allows for unrestricted use, distribution, and reproduction in any medium, provided that the original work is properly cited.

cardiac output and blood pressure. Although it improves over time, the average blood pressure level in individuals with MI is generally lower than that in healthy individuals (6, 7).

In general, MI can be divided into STEMI (ST elevation) and NSTEMI (without ST elevation). STsegment elevation in the patient's ECG can indicate myocardial infarction, but this is not true for all patients (2). In addition to these changes, R wave amplitude can also have a diagnostic aspect. A decrease in the R wave indicates acute myocardial infarction (8). Therefore, the use of electrocardiogram as a tool to diagnose myocardial infarction is helpful.

Today, the use of herbal medicines has grown significantly compared to synthetic medicines. Herbal medicines seem to be beneficial to patients both economically and therapeutically (9). Therefore, medicinal plants can be used as adjunctive therapy in cardiovascular diseases. Lepidium draba L. (Cruciferae), a native plant to southwestern and Central Asia, contains high amounts of sulforaphane (7), which has various effects, including anticancer effect, through the induction of phase 2 protein, and the antimicrobial application of sulforaphane is also reported (6). In addition, the antioxidant and anti-inflammatory effects of L. draba extract have been reported because of its flavonoids and phenolic acids, which exist in great amounts in this plant (10, 11). According to the reports in scientific resources, aerial parts of L. draba contain different classes of compounds, such as alkaloids, terpenoids, tannins, saponins, Leuco anthocyanins, triterpenoids, and flavonoids (12). Previously in a published article in 2018 carried out by Mahomoodally et al., LC-ESI-MS/MS analysis of the methanol extracts of Cardaria draba (synonym: Lepidium draba L.) presence of some major phenolic compounds known verbascoside, chlorogenic acid, apigenin (7)- glucoside, and 4-hydroxybenzoic acid, together with hesperidin, hyperoside, and chlorogenic acid in fewer content value plus vanillin and rosmarinic acid were reported in the plant (13). Elsewhere in another study Najim et al. reported L. draba contained many types of flavonoids including rutin, qurecetin, kampferol and luteolin (14).

# 2. Objectives

This is the inaugural study to investigate the effects of *L. draba* on cardiovascular diseases, including MI, thereby aiming to elucidate the potential protective effects of *L. draba* against isoproterenol-induced myocardial infarction.

#### 3. Methods

# 3.1. Plant

*Lepidium draba* was harvested from the north of West Azerbaijan Province in June 2020. Following the drying of plants in the shade, they were pulverized and used for further experimental analysis. The collected plant was authenticated by a pharmacognosist, and the sample was kept in the faculty's Herbarium (No: TBZFPH1989).

#### 3.2. Preparation of Plant Extract

The dried powdered flowering branches of *L. draba* were defatted with 5 L n-hexane, followed by maceration with 70% ethanol for 5 consecutive days, and then filtered with Whatman filter paper. The yielded hydroalcoholic extracts were solvent-evaporated up to complete dryness via a rotary evaporator (Heidolph, Germany) at 40°C under a vacuum.

# 3.3. Assessment of Phenolic Profile

#### 3.3.1. Total Phenolic Content

To determine the total phenolic content of the *L. draba* extract, the Folin-Ciocalteu (FC) method was used (15). Briefly, 5 mL of Folin-Ciocalteu reagent 10% (v/v) was mixed with 4 mL of Na<sub>2</sub>CO<sub>3</sub> solution, and 0.5 mL of plant extract (1 mg/mL) was placed at 20 - 22°C for 15 min in darkness. Then, the absorbance was recorded at  $\lambda$ max 760 nm. Finally, the calibration curve was depicted using gallic acid (10 - 100 µg/mL) in the same process, and the obtained values were calculated as gallic acid equivalents (GAE)/g of the extract.

#### 3.3.2. Total Flavonoid Content

The  $AlCl_3$  reagent was used to determine the total flavonoid content of the *L. draba* extract in a colorimetric approach (16). For this purpose, 1 mg/mL of hydroalcoholic extract of *L. draba* was blended with 2 mL

of  $AlCl_3$ , and then the absorbance was measured spectrophotometrically at  $\lambda$ max 415 nm. Finally, the calibration curve was depicted using quercetin (10 - 100  $\mu$ g/mL) in the same process, and the obtained values were calculated as quercetin equivalents (QE)/g of the extract.

# 3.4. Determination of in Vitro Antioxidant Activity

## 3.4.1. DPPH Radical Scavenging Assay

This assay was performed as previously reported (17-19). Two mL of 0.08% freshly prepared DPPH solution was mixed individually with 2 mL of varied concentrations of L. draba extract. The ensuing solutions were kept for 30 min at 25°C in darkness. The same procedure was applied for quercetin as a positive reference compound. A spectrophotometer (Shimadzu, Japan) was used to measure the absorbance of each test solution at 517 nm. Ultimately, the inhibition percentage (I %) of DPPH free radical scavenging potential of the extract and quercetin were both computed via the subsequent formula: I % = {(absorbance of the blank - absorbance of the sample)/absorbance of the blank}  $\times$  100. Overall, the IC50 values, indicating the sample concentration that inhibits the DPPH free radicals at 50%, were estimated for extract and quercetin, correspondingly.

# 3.4.2. FRAP Assay

The Ferric reducing antioxidant power assay was carried out as stated by the method reported by Bardakci et al. (20). 10  $\mu$ L of different concentrations of *L. draba* extract were individually mixed with distilled water (30  $\mu$ L) and FRAP reagent (260  $\mu$ L) comprising acetate buffer and FeCl<sub>3</sub>. The final solutions were kept for 30 min at 37°C. Later on, the achieved color absorbance was read at  $\lambda$ max 593 nm. The calibration curve was depicted using FeSO<sub>4</sub> in the same process, and the obtained values were calculated as FeSO<sub>4</sub> conforming to 1g of the extract.

# 3.5. Animal

Thirty male Wistar rats were used in this study. The rats weighing 250  $\pm$  20 g were kept under standard circumstances (22  $\pm$  2°C and a 12 - h light/12 - h dark cycle)

in the Animal House of Urmia Medical Sciences University. All protocols were done in accordance with the Ethics Committee guidelines of Urmia University of Medical Sciences (code: IR.UMSU.REC.1399.312).

#### 3.6. Experimental Protocol

The study was conducted in 2 days, and 5 groups were used in this study as follows: Group I (control) was left untreated during the study. Rats in group II (MI) received normal saline intraperitoneally (IP) and also isoproterenol (100 mg/kg, sc, for 2 consecutive days) to induce MI. Groups III-V were given hydroalcoholic extract of *L. draba* (50, 250, and 500 mg/kg respectively) for 2 days and were subcutaneously (sc) injected with isoproterenol at an interval of 24 h for 2 consecutive days.

#### 3.7. Electrocardiogram and Hemodynamic Measurement

A Ketamine (60 mg/kg)/ xylazine (10 mg/kg) mixture was used for the induction of anesthesia. Then, lead II was used to record electrocardiographic parameters. For hemodynamic measurement, an incision was made in the midline of the animal's neck, and after isolating the left carotid, systolic blood pressure (SBP), diastolic blood pressure (DBP), heart rate (HR), mean arterial pressure (MAP), and arterial Blood pressure (ABP) were measured using a polyethylene cannula. (Powerlab system, AD Instruments, Australia) (21).

# 3.8. Histopathological Study

For histopathological examination, cardiac apex samples were placed in 10% formalin and then stained with trichrome gomori. Finally, they were evaluated by 2 trained persons (22).

#### 3.9. Hypertrophy

The rats were euthanized by an overdose of anesthetics, and the hearts were rapidly excised and weighed. The degree of hypertrophy was measured by the heart-weight-to-body weight ratio (g/kg)(22).

# 3.10. LDH Measurement

At the end of the experiment, blood samples were collected from the portal vein, and serum was used to

measure LDH levels. The levels of serum LDH as a marker of tissue injury during MI were determined using enzymatic kits from Bayerpaul, Iran. The results were expressed as units of LDH present per liter (U/L).

#### 3.11. Statistical Analysis

All data were provided as means  $\pm$  standard error of the mean. To compare the mean values between groups, one-way ANOVA and the Tukey post-test were used. P < 0.05 was regarded as statistically significant.

# 4. Results

# 4.1. Phytochemical Analysis

The results of the phytochemical analysis for the hydroalcoholic extract of *L. draba* are shown in Table 1. The total flavonoid and phenolic contents of the *L. draba* extract were determined to be 92.30 mg GAE/g and 60.68 mg QE/g of the extract, respectively. Furthermore, the IC50 values in the DPPH assay for the extract and quercetin were found to be 304.45  $\mu$ g/mL and 4.5  $\mu$ g/mL, respectively. In the ferric-reducing antioxidant power assay, the extract exhibited a value of 0.33 mg FeSO<sub>4</sub>/g of the extract.

GAE/g)	Gentent (amg	Scavenging	(amg
	QE/g)	Activity (µg/mL)	FeSO <sub>4</sub> /g)
<i>L draba</i> Extract 92.30 ± 1.84	60.68±1.86	304.45±10.19 <sup>a</sup>	0.33±0.008

# 4.2. Hydroalcoholic Extract of L. draba Enhances Electrocardiogram Pattern

The electrocardiogram revealed a normal pattern in the control group, while the Iso (MI) group exhibited an ST-segment elevation (P < 0.01), indicative of myocardial infarction. Additionally, the R amplitude in the group receiving only isoproterenol injections was significantly reduced (P < 0.001) compared to the control group. The administration of the hydroalcoholic extract of *L. draba* led to an improvement in the ECG pattern, approaching normal levels (Figure 1A). As depicted in Figure 1B, the plant extract at doses of 50 and 500 mg/kg effectively reduced ST-segment elevation (P < 0.01 and P < 0.05, respectively) and significantly increased R-amplitude (P < 0.05) in comparison to the Iso group.



Figure 1. Effect of *L. draba* hydroalcoholic extract on ECG pattern in control and treated groups. Iso: Isoproterenol; LD: *Lepidium draba*.

# 4.3. Hydroalcoholic Extract of L. draba Improved the Hemodynamic Parameters

According to hemodynamic results, the MAP reduced from 138 ± 4 mmHg in the control group to 91 ± 8 mmHg in the Iso group (P < 0.001). Hydroalcoholic extract of *L. draba* at all three doses significantly improved MAP (P < 0.05). Furthermore, the heart rate of rats in the Iso group was higher than the control group (P < 0.001), while treatment with hydroalcoholic extract of *L. draba* at all three doses of 50, 250, and 500 mg/kg decreased (P < 0.001) as compared to Iso group (Table 2).

Table 2. Effects of *Lepidium draba* Hydroalcoholic Extract on Hemodynamic Parameters

Groups (N = 6)	ABP (mmHg)	Systolic Pressure (mmHg)	Diastolic Pressure (mmHg)	Mean Arterial Pressure (mmHg)	Heart rate (bpm)
Control	$155\pm1$	$138\pm4$	$129\pm4$	$138\pm4$	$247\pm10$
Iso	$95\pm 8$ <sup>b</sup>	$91\pm8 \ ^{b}$	$89\pm9^{\rm b}$	91 ± 8 <sup>b</sup>	400 ± 17 b
Iso+LD 50mg/kg	$118 \pm 4$ <sup>C</sup>	116 ± 3 <sup>c</sup>	113 $\pm$ 3 <sup>c</sup>	116 ± 3 <sup>c</sup>	$287\pm17^{\rm }{\rm d}$
Iso+LD 250mg/kg	116 ± 1 <sup>c</sup>	116 ± 1 <sup>c</sup>	$114 \pm 1^{\circ}$	116 ± 1 <sup>c</sup>	298±15 d
Iso+LD 500mg/kg	$124 \pm 12^{\text{C}}$	114 ± 12 <sup>c</sup>	$109\pm10$ <sup>c</sup>	114 ± 12 <sup>c</sup>	238 ± 16 d

 $^a$  Values are reported as mean  $\pm$  SEM.

<sup>b</sup> P < 0.001 vs. control group

<sup>c</sup> P < 0.05 vs. Iso group.

 $^{\rm d}$  P < 0.001 vs. Iso group.

# 4.4. Effects of the Hydroalcoholic Extract of L. draba on Cardiac Hypertrophy

The results indicated that the heart weight to body weight (HW/BW) ratio was significantly higher in the Iso group (P < 0.001) compared to the control group. Rats treated with the hydroalcoholic extract of *L. draba* at a dose of 500 mg/kg during the treatment period experienced a significant reduction in the HW/BW ratio (P < 0.01) compared to the group injected with isoproterenol (Figure 2).



**Figure 2.** Heart weight (HW) to body weight (BW) ratio between different groups. Data are presented as Mean  $\pm$  S.E.M (N = 6); # P < 0.001 from respective control value, \*\* P < 0.01 as compared with isoproterenol treated group.

#### 4.5. Histopathological Examination

Histological sections of the hearts from rats administered isoproterenol injections, stained with Gomori's Trichrome, showed significant subendocardial fibroblastic hyperplasia and severe fibrosis (Figure 3). The results indicated that treatment with the hydroalcoholic extract of *L. draba* (500 mg/kg) significantly reduced fibrosis (P < 0.001) induced by isoproterenol in heart tissue.



**Figure 3.** Histopathological fibrosis in rat cardiac apex tissues is graded. Pathological alterations are classified as mild, moderate, high, or severe in grades 1-4, respectively.

# 4.6. Effects of L. draba on Lactate Dehydrogenase Level

Isoproterenol injection increased LDH level from  $154.95 \pm 21$  U/L in the control group to  $461.66 \pm 183$  U/L in the Iso group (P < 0.05). However, *L. draba* administration significantly reduced the level of the serum LDH compared to the infarcted rats, who received isoproterenol only (Figure 4).



**Figure 4.** The Effect of Treatment with *Lepidium draba* (LD) extract on lactate dehydrogenase (LDH) level in the myocardial infarction. Iso: isoproterenol; LD: *Lepidium draba*. Values are mean  $\pm$  S.E.M (N = 6). #P < 0.05 from respective control value; "P < 0.05, as compared with iso group.

#### 5. Discussion

The present research aimed to elucidate the potential protective effects of the *L. draba* extract on hemodynamic and ECG parameters, as well as heart tissue fibrosis and LDH levels following MI in rats. Our findings demonstrate that acute administration of a hydroalcoholic extract of *L. draba* could protect against isoproterenol-induced myocardial infarction.

Lepidium draba is a plant of the Brassicaceae family several compounds, that contains such as glucosinolates and other compounds resulting from its degradation, with beneficial effects in preventing several diseases (23). Phytochemical studies have shown that aerial parts of L. draba contain sulforaphane and erysoline (13). It has been reported that glucosinolate can reduce the size of the infarcted area in myocardial tissue and also reduce apoptosis of cardiomyocytes (24). Our results show that L. draba extract reduced histopathological changes such as fibrosis, which can be partially attributed to glucosinolate in the plant. Additionally, the presence of compounds such as alkaloids and glucosinolates can also be detected in Lepidium meyenii, another member of the Brassicaceae family (25). Overall, there are few studies with *L. draba*, but studies on other plants of this family, including Lepidium sativum, indicate that polyphenols are present in large quantities in this plant (26). A study by Cheng et al. (27) demonstrated that polyphenols can improve the diastolic and endothelial function of the heart in people who have experienced myocardial infarction. Polyphenols also have protective effects in cardiac ischemia through their capability of decreasing oxygen free radicals (28). The findings of the phytochemical analysis of the *L. draba* hydroalcoholic extract revealed moderate antioxidant potential in DPPH radical scavenging activity, along with measurable ferric ion-reducing activity, which is mostly connected to the polyphenols and flavonoids content in the hydroalcoholic extract.

Several studies have indicated that isoproterenol, a synthetic β-adrenoceptor agonist, is used to induce MI in rats (29, 30). Studies show that isoproterenol-induced MI can be a valid and reproducible way to evaluate new cardioprotective drugs (31). We found that hemodynamic factors, such as mean arterial pressure, decreased in the MI (Iso) group, and administration of L. draba extract improved these parameters. It was also observed that heart rate improved and decreased in the group treated with L. draba extract. These positive effects on hemodynamic parameters can be partially attributed to the polyphenols present in the plant.

A decrease in R wave amplitude was observed in rats receiving isoproterenol. ECG analysis showed reduced R wave amplitude in the Iso group, but acute administration of L. draba extract in high doses increased the R wave amplitude, which can be related to the reduction of edema in the heart tissue (32, 33). Consistent with our results, Rajadurai and Stanely Mainzen Prince (34) reported that isoproterenolinduced MI can lead to ST-segment elevation on the ECG, which is an important sign of the presence of myocardial ischemia. Acute administration of L. draba extract at high doses reduced ST-segment elevation. Elevation of ST-segment on ECG pattern occurs due to a potential imbalance in the boundary between ischemic and non-ischemic regions, while R-amplitude decreases because of the presence of edema in myocardial tissue as a result of infarction induced by isoproterenol injection (35). It was also proved that elevation of the STsegment may be a sign of necrosis of the myocardial tissue (36). Hypertrophy caused by isoproterenol can be due to increased cell death. fibrosis. and abnormalities normal heart function. Treatment with in hydroalcoholic extract of L. draba reduces cardiac which hypertrophy, can be considered а cardioprotective mechanism (37, 38).

Metabolic damage or myocardial infarction causes a large number of diagnostic factors to leak into the extracellular space (39). The presence of cytosolic enzymes such as lactate dehydrogenase (LDH) is considered a diagnostic marker when tissue is damaged. The amount of these enzymes in serum can vary due to changes in membrane permeability or rupture (35). *Lepidium draba* extract-treated groups demonstrated less activity of LDH in the serum, indicating that the extract maintains cell permeability and prevents the leakage of these enzymes. It was reported that sulforaphane present in the plant can reduce the level of serum LDH activity caused by tissue damage (40), which is in line with our findings.

# 5.1. Conclusions

Acute administration of the hydroalcoholic extract of *L. draba* plays a protective role in myocardial infarction and could be considered an adjunctive treatment option for myocardial infarction.

#### Footnotes

**Authors' Contribution:** HS designed the experiments; MKA, VSI, and SH performed experiments and collected data; MKA, VSI, SH, and HS discussed the results and strategy; HS supervised, directed, and managed the study; MKA, VSI, SH, and HS, final approval of the version to be published.

**Conflict of Interests:** The authors declare no conflict of interest.

**Data Reproducibility:** Data will be provided upon request.

**Ethical Approval:** All protocols were approved by the Ethics Committee of Urmia University of Medical Sciences (code: IR.UMSU.REC.1399.312 ).

**Funding/Support:** The authors declared no funding/support for this study.

#### References

 Yu LJ, Zhang KJ, Zhu JZ, Zheng Q, Bao XY, Thapa S, et al. Salvianolic Acid Exerts Cardioprotection through Promoting Angiogenesis in Animal Models of Acute Myocardial Infarction: Preclinical Evidence. *Oxid Med Cell Longev.* 2017;2017:8192383. [PubMed ID: 28713492]. [PubMed Central ID: PMC5497657]. https://doi.org/10.1155/2017/8192383.

- Fu R, Song CX, Dou KF, Yang JG, Xu HY, Gao XJ, et al. Differences in symptoms and pre-hospital delay among acute myocardial infarction patients according to ST-segment elevation on electrocardiogram: an analysis of China Acute Myocardial Infarction (CAMI) registry. *Chin Med J (Engl)*. 2019;**132**(5):519-24. [PubMed ID: 30807351]. [PubMed Central ID: PMC6416090]. https://doi.org/10.1097/CM9.000000000000122.
- Thygesen K, Alpert JS, White HD, Joint EAFFTROMI. Universal definition of myocardial infarction. *Eur Heart J.* 2007;28(20):2525-38. [PubMed ID: 17951287]. https://doi.org/10.1093/eurheartj/ehm355.
- Talman V, Ruskoaho H. Cardiac fibrosis in myocardial infarctionfrom repair and remodeling to regeneration. *Cell Tissue Res.* 2016;**365**(3):563-81. [PubMed ID: 27324127]. [PubMed Central ID: PMC5010608]. https://doi.org/10.1007/s00441-016-2431-9.
- Bouchareb R, Katz M, Saadallah N, Sassi Y, Ali S, Lebeche D. Boron improves cardiac contractility and fibrotic remodeling following myocardial infarction injury. *Sci Rep.* 2020;**10**(1):17138. [PubMed ID: 33051505]. [PubMed Central ID: PMC7553911]. https://doi.org/10.1038/s41598-020-73864-w.
- Fahey JW, Haristoy X, Dolan PM, Kensler TW, Scholtus I, Stephenson KK, et al. Sulforaphane inhibits extracellular, intracellular, and antibiotic-resistant strains of Helicobacter pylori and prevents benzo[a]pyrene-induced stomach tumors. *Proc Natl Acad Sci U S A*. 2002;**99**(11):7610-5. [PubMed ID: 12032331]. [PubMed Central ID: PMC124299]. https://doi.org/10.1073/pnas.112203099.
- Fahey JW, Talalay P. Antioxidant functions of sulforaphane: a potent inducer of Phase II detoxication enzymes. *Food Chem Toxicol*. 1999;**37**(9-10):973-9. [PubMed ID: 10541453]. https://doi.org/10.1016/s0278-6915(99)00082-4.
- Sun X, Cai J, Fan X, Han P, Xie Y, Chen J, et al. Decreases in electrocardiographic R-wave amplitude and QT interval predict myocardial ischemic infarction in Rhesus monkeys with left anterior descending artery ligation. *PLoS One.* 2013;8(8). e71876. [PubMed ID: 23967258]. [PubMed Central ID: PMC3742514]. https://doi.org/10.1371/journal.pone.0071876.
- Shaito A, Thuan DTB, Phu HT, Nguyen THD, Hasan H, Halabi S, et al. Herbal Medicine for Cardiovascular Diseases: Efficacy, Mechanisms, and Safety. Front Pharmacol. 2020;11:422. [PubMed ID: 32317975]. [PubMed Central ID: PMC7155419]. https://doi.org/10.3389/fphar.2020.00422.
- Ouissem BS, Sabrina B, Lotfi B, Khellaf R, Chawki B, Ibrahim D, et al. HPLC Analysis and Antioxidant Properties of Algerian Lepidium draba Ethyl acetate Extract. J Bio Act Prod Nat. 2018;8(4):265-71. https://doi.org/10.1080/22311866.2018.1511381.
- Chyad AH. Evaluation of Anticancer, Analgesic and Anti-Inflammatory Activities of the Ethanolic Extract of Lepidium draba Linn. Leaves. Advances in Animal and Veterinary Sciences. 2017;5(1):7-13. https://doi.org/10.14737/journal.aavs/2017/5.1.7.13.
- 12. Bicha S, Benmekhebi L, Boubekri N, Khellaf R, Brouard I, Zama D, et al. Compositional study, antibacterial and antioxidant potential of Lepidium draba L.(Brascicaceae). *Res J Pharm Biol Chem Sci.* 2016;**7**.
- Mahomoodally MF, Zengin G, Aumeeruddy MZ, Sezgin M, Aktumsek
  A. Phytochemical profile and antioxidant properties of two Brassicaceae species: Cardaria draba subsp. draba and Descurainia

sophia. Biocatalysis and Agricultural Biotechnology. 2018;**16**:453-8. https://doi.org/10.1016/j.bcab.2018.09.020.

- Najim RS, Hasan ZYM, Al-Chalabi R. Study The Antimicrobial Activity of Ethanolic Extract of Lepidium draba on Some Skin Infectious Agents. J Biotechnol Res Center. 2020;14(1):10-9. https://doi.org/10.24126/jobrc.2020.14.1.583.
- Singleton VL, Rossi JA. Colorimetry of Total Phenolics with Phosphomolybdic-Phosphotungstic Acid Reagents. *Am J Enol Vitic*. 1965;16(3):144-58. https://doi.org/10.5344/ajev.1965.16.3.144.
- Chemsa AE, Zellagui A, Ozturk M, Erol E, Ceylan O, Duru ME, et al. Chemical composition, antioxidant, anticholinesterase, antimicrobial and antibiofilm activities of essential oil and methanolic extract of Anthemis stiparum subsp. sabulicola (Pomel) Oberpr. *Microb Pathog.* 2018;119:233-40. [PubMed ID: 29684540]. https://doi.org/10.1016/j.micpath.2018.04.033.
- Brand-Williams W, Cuvelier ME, Berset C. Use of a free radical method to evaluate antioxidant activity. *LWT - Food Science and Technology*. 1995;28(1):25-30. https://doi.org/10.1016/s0023-6438(95)80008-5.
- Hamedeyazdan S, Niroumand F, Fathiazad F. Phytochemical analysis and antioxidative properties of Centaurea albonitens. *Research Journal of Pharmacognosy*. 2017;4(4):57-64.
- Soraya H, Moloudizargari M, Aghajanshakeri S, Javaherypour S, Mokarizadeh A, Hamedeyazdan S, et al. Angiogenic effect of the aqueous extract of Cynodon dactylon on human umbilical vein endothelial cells and granulation tissue in rat. *Daru.* 2015;23(1):10. [PubMed ID: 25630338]. [PubMed Central ID: PMC4316609]. https://doi.org/10.1186/s40199-015-0093-x.
- Bardakci H, Celep E, Gözet T, Kurt-Celep I, Deniz I, Şen-Utsukarci B, et al. A comparative investigation on phenolic composition, antioxidant and antimicrobial potentials of Salvia heldreichiana Boiss. ex Bentham extracts. South African Journal of Botany. 2019;125:72-80. https://doi.org/10.1016/j.sajb.2019.07.010.
- Abbaszadeh S, Javidmehr A, Askari B, Janssen PML, Soraya H. Memantine, an NMDA receptor antagonist, attenuates cardiac remodeling, lipid peroxidation and neutrophil recruitment in heart failure: A cardioprotective agent? *Biomed Pharmacother*. 2018;**108**:1237-43. [PubMed ID: 30372825]. https://doi.org/10.1016/j.biopha.2018.09.153.
- Soraya H, Khorrami A, Garjani A, Maleki-Dizaji N, Garjani A. Acute treatment with metformin improves cardiac function following isoproterenol induced myocardial infarction in rats. *Pharmacol Rep.* 2012;64(6):1476-84. [PubMed ID: 23406758]. https://doi.org/10.1016/s1734-1140(12)70945-3.
- Radonic A, Blazevic I, Mastelic J, Zekic M, Skocibusic M, Maravic A. Phytochemical analysis and antimicrobial activity of Cardaria draba (L.) Desv. volatiles. *Chem Biodivers*. 2011;8(6):1170-81. [PubMed ID: 21674789]. https://doi.org/10.1002/cbdv.201000370.
- Mukherjee S, Lekli I, Ray D, Gangopadhyay H, Raychaudhuri U, Das DK. Comparison of the protective effects of steamed and cooked broccolis on ischaemia-reperfusion-induced cardiac injury. *Br J Nutr.* 2010;103(6):815-23. [PubMed ID: 19857366]. https://doi.org/10.1017/S0007114509992492.
- Tafuri S, Cocchia N, Carotenuto D, Vassetti A, Staropoli A, Mastellone V, et al. Chemical Analysis of Lepidium meyenii (Maca) and Its Effects on Redox Status and on Reproductive Biology in Stallions (dagger).

*Molecules*. 2019;**24**(10). [PubMed ID: 31126050]. [PubMed Central ID: PMC6571579]. https://doi.org/10.3390/molecules24101981.

- Ullah MA, Tungmunnithum D, Garros L, Drouet S, Hano C, Abbasi BH. Effect of Ultraviolet-C Radiation and Melatonin Stress on Biosynthesis of Antioxidant and Antidiabetic Metabolites Produced in In Vitro Callus Cultures of Lepidium sativum L. Int J Mol Sci. 2019;20(7). [PubMed ID: 30978911]. [PubMed Central ID: PMC6479895]. https://doi.org/10.3390/ijms20071787.
- Cheng YC, Sheen JM, Hu WL, Hung YC. Polyphenols and Oxidative Stress in Atherosclerosis-Related Ischemic Heart Disease and Stroke. Oxid Med Cell Longev. 2017;2017:8526438. [PubMed ID: 29317985]. [PubMed Central ID: PMC5727797]. https://doi.org/10.1155/2017/8526438.
- Cebova M, Pechanova O. Protective Effects of Polyphenols against Ischemia/Reperfusion Injury. *Molecules*. 2020;**25**(15). [PubMed ID: 32751587]. [PubMed Central ID: PMC7435883]. https://doi.org/10.3390/molecules25153469.
- Kurian GA, Philip S, Varghese T. Effect of aqueous extract of the Desmodium gangeticum DC root in the severity of myocardial infarction. *J Ethnopharmacol.* 2005;97(3):457-61. [PubMed ID: 15740881]. https://doi.org/10.1016/j.jep.2004.11.028.
- Fan Y. Cardioprotective Effect of Rhapontigenin in Isoproterenol-Induced Myocardial Infarction in a Rat Model. *Pharmacology*. 2019;**103**(5-6):291-302. [PubMed ID: 30820014]. https://doi.org/10.1159/000496800.
- Kannan MM, Quine SD. Ellagic acid ameliorates isoproterenol induced oxidative stress: Evidence from electrocardiological, biochemical and histological study. *Eur J Pharmacol*. 2011;659(1):45-52.
   [PubMed ID: 21385579]. https://doi.org/10.1016/j.ejphar.2011.02.037.
- 32. Moradi-Arzeloo M, Farshid AA, Tamaddonfard E, Asri-Rezaei S. Effects of histidine and vitamin C on isoproterenol-induced acute myocardial infarction in rats. *Vet Res Forum*. 2016;7(1):47. [PubMed ID: 27226887].
- Boarescu PM, Boarescu I, Bocsan IC, Pop RM, Gheban D, Bulboaca AE, et al. Curcumin Nanoparticles Protect against Isoproterenol Induced Myocardial Infarction by Alleviating Myocardial Tissue Oxidative Stress, Electrocardiogram, and Biological Changes. *Molecules*. 2019;24(15). [PubMed ID: 31374848]. [PubMed Central ID: PMC6696485]. https://doi.org/10.3390/molecules24152802.
- Rajadurai M, Stanely Mainzen Prince P. Preventive effect of naringin on cardiac markers, electrocardiographic patterns and lysosomal hydrolases in normal and isoproterenol-induced myocardial infarction in Wistar rats. *Toxicology*. 2007;230(2-3):178-88. [PubMed ID: 17188415]. https://doi.org/10.1016/j.tox.2006.11.053.
- Patel V, Upaganlawar A, Zalawadia R, Balaraman R. Cardioprotective effect of melatonin against isoproterenol induced myocardial infarction in rats: A biochemical, electrocardiographic and histoarchitectural evaluation. *Eur J Pharmacol.* 2010;644(1-3):160-8. [PubMed ID: 20624385]. https://doi.org/10.1016/j.ejphar.2010.06.065.
- 36. Prabhu S, Devi CS. Protective effect of mangiferin an active phytochemical and cardiotonic from Mangifera indica Linn on isoproterenol induced myocardial infarcted rats-an electrocardiographic, electrophoretic and biochemical evidences. *Pharmacogn Mag.* 2006;**2**(7).
- 37. Hemalatha KL, Stanely Mainzen Prince P. Antihyperlipidaemic, antihypertrophic, and reducing effects of zingerone on

experimentally induced myocardial infarcted rats. J Biochem Mol Toxicol. 2015;**29**(4):182-8. [PubMed ID: 25558849]. https://doi.org/10.1002/jbt.21683.

- Paulino ET, Barros Ferreira AK, da Silva JCG, Ferreira Costa CD, Smaniotto S, de Araujo-Junior JX, et al. Cardioprotective effects induced by hydroalcoholic extract of leaves of Alpinia zerumbet on myocardial infarction in rats. *J Ethnopharmacol.* 2019;**242**:112037. [PubMed ID: 31247239]. https://doi.org/10.1016/j.jep.2019.112037.
- Upaganlawar A, Gandhi C, Balaraman R. Effect of green tea and vitamin E combination in isoproterenol induced myocardial infarction in rats. *Plant Foods Hum Nutr.* 2009;64(1):75-80. [PubMed ID: 19058010]. https://doi.org/10.1007/s11130-008-0105-9.
- Kyto V, Sipila J, Rautava P. Likelihood and predictors of ST-elevation in patients hospitalized for myocardial infarction. *PLoS One*. 2014;9(9). e108440. [PubMed ID: 25254632]. [PubMed Central ID: PMC4177929]. https://doi.org/10.1371/journal.pone.0108440.