



# Effects of 810 nm and 940 nm Diode Laser on ROS Production in Macrophages: The Role of Interferon-Gamma

Milad Sadeghi<sup>1</sup>, Reyhaneh Shoorgashti <sup>1</sup>, Simin Lesan <sup>1</sup>, Sadegh Hasannia <sup>1</sup>, Taraneh Farrokhnia <sup>1,\*</sup>

<sup>1</sup>Department of Oral Medicine, TeMS. C., Islamic Azad University, Tehran, Iran

\*Corresponding Author: Department of Oral Medicine, TeMS. C., Islamic Azad University, Tehran, Iran. Email: taranehfars3@yahoo.com

Received: 5 February, 2025; Revised: 29 June, 2025; Accepted: 2 August, 2025

## Abstract

**Background:** Reactive oxygen species (ROS) are critical molecules produced during cellular metabolism. Macrophages, as essential immune cells, play a pivotal role in the immune response by producing inflammatory mediators and releasing ROS. Interferon-gamma (IFN- $\gamma$ ) is a key stimulator of macrophage activity, enhancing ROS production.

**Objectives:** With the increasing application of low-level laser therapy (LLLT) in medicine, this study aims to compare the effects of 810 nm and 940 nm diode laser irradiation on ROS production in macrophages.

**Methods:** This in vitro study utilized a monocyte/macrophage cell line. Cells were seeded into microwell plates and exposed to diode laser irradiation at 810 nm and 940 nm, with power settings of 0.4 watts (energy density 7-15 J/cm<sup>2</sup>) and irradiation times of 15 and 30 seconds. The IFN- $\gamma$  was added post-irradiation. The ROS production was assessed using a fluorometric method, measuring nitric oxide (NO) as an indirect indicator.

**Results:** The comparison between 810 nm and 940 nm wavelengths showed no significant difference in ROS production ( $P > 0.05$ ). However, higher energy density (4 J vs. 2 J,  $P = 0.001$ ), longer irradiation time (30 s vs. 15 s,  $P = 0.001$ ), and the presence of IFN- $\gamma$  ( $P < 0.001$ ) significantly increased ROS levels.

**Conclusions:** This study showed that both 810 nm and 940 nm diode lasers significantly increase ROS production in macrophage cells compared to a control group, despite no significant differences between the two wavelengths.

**Keywords:** Free Radicals, Interferon-gamma, Lasers, Macrophages, Phagocytosis, Reactive Oxygen Species

## 1. Background

Photobiomodulation (PBM), also known as low-level laser therapy (LLLT), was first identified nearly 50 years ago (1). In recent years, LLLT has gained significant traction in both medical and dental fields, primarily due to its ability to modulate biological processes without causing substantial cellular damage (2-4). Emerging evidence indicates that LLLT can modulate immune responses, including the regulation of reactive oxygen species (ROS) levels, thereby facilitating tissue repair and reducing inflammation (5-8).

The ROS are highly reactive oxygen-containing molecules that play a dual role in cellular physiology (9-

11). On one hand, they are essential for cellular signaling and defense mechanisms, particularly within the immune system. On the other hand, excessive ROS production can lead to oxidative stress, contributing to cellular damage and the pathogenesis of various diseases, including cardiovascular and neurodegenerative disorders (9, 12-16).

Hamblin indicates that LLLT can lead to a temporary and modest increase in ROS production in healthy cells, resulting from an elevation of mitochondrial membrane potential (MMP) beyond normal levels (17). Nonetheless, when applied as a treatment for tissue injury or muscle damage, LLLT has been shown to effectively lower markers of oxidative stress (17-19).

Macrophages, key players in the immune response, are central to the production of ROS, which they use to combat pathogens and mediate inflammation (20, 21). Interferon-gamma (IFN- $\gamma$ ), a critical cytokine in both innate and adaptive immunity, primes macrophages by enhancing their capacity to produce ROS and other inflammatory mediators. As a potent macrophage activator, IFN- $\gamma$  promotes the release of higher ROS levels (22, 23). However, the interaction between IFN- $\gamma$  and laser irradiation, particularly its influence on ROS production, remains relatively underexplored in the current literature.

Diode lasers emitting wavelengths of 810 nm and 940 nm have been extensively studied for their capacity to penetrate tissues and influence cellular activities (3, 5, 24, 25). Given the increasing relevance of laser technology in modern dentistry and medicine, this study aims to investigate a critical and underexplored question: Do 810 nm and 940 nm diode laser irradiations differentially affect the quantitative release of ROS in macrophages, in both the presence and absence of IFN- $\gamma$ ? If significant differences are observed, the findings could inform the selection of optimal laser wavelengths for therapeutic applications.

This study addresses a gap in the current research landscape by examining the specific effects of 810 nm and 940 nm diode lasers on ROS production in macrophages. While previous research has demonstrated that LLLT can enhance cellular repair mechanisms, such as in the treatment of recurrent oral ulcers and scar reduction (24, 26), the direct impact of these laser wavelengths on ROS release in macrophages remains insufficiently understood and necessitates further investigation.

## 2. Objectives

This study aimed to compare the effects of 810 nm and 940 nm diode laser irradiation on ROS production in macrophages.

## 3. Methods

### 3.1. Cell Line and Culture Conditions

The study utilized the murine monocyte/macrophage cell line J774A.1 (NCBIC 483), acquired from the Pasteur Institute of Iran. Cells were cultured in a flask containing Dulbecco's Modified Eagle

Medium (DMEM) enriched with 10% fetal bovine serum, 100  $\mu$ g/mL penicillin, 100  $\mu$ g/mL streptomycin, and 2 mM L-glutamine (27, 28). The culture was maintained at 37°C in a CO<sub>2</sub> incubator.

### 3.2. Laser Parameters

Two types of lasers were employed in this study: The 940 nm Epic X diode laser, featuring an adjustable power range from 0.1 to 10 W, manufactured by Biolase in the USA, and a 400  $\mu$ m tip; and the 810 nm Pulsar diode laser, made in Iran, with an adjustable power range of 0.5 to 8 W, operated at 0.4 W in this experiment. Both lasers were calibrated before use by measuring their output power with a power meter.

### 3.3. Experimental Groups

The release of ROS by macrophage cells was evaluated in 18 different groups as presented in Table 1.

### 3.4. Experimental Procedure

A volume of 100  $\mu$ L of the cultured J774A.1 cell line (containing 10<sup>4</sup> cells) was transferred into each well of a microplate (5 wells per group). Samples were allowed to stabilize for one hour before laser irradiation. The 940 nm and 810 nm diode lasers were employed at a 45-degree angle relative to the cell surface in the wells, using a 400  $\mu$ m tip. Laser output was set at 0.4 W, with exposure durations of 15 or 30 seconds, divided into intervals with 10-second pauses in a dark environment to prevent interference from external light sources. The effective energy per well was calibrated to approximately 2 J and 4 J, corresponding to calculated energy densities of ~7 - 15 J/cm<sup>2</sup>, based on the irradiated area. These values are cumulative and not per-pulse.

Following laser irradiation, 2  $\mu$ L of IFN- $\gamma$  was added to the wells to activate macrophages (29), and 4 hours post-irradiation, the ROS production level was assessed by measuring nitric oxide (NO) levels using a fluorometric method with DCFH-DA as the fluorescent probe (30, 31). Although DCFH-DA is a widely used fluorescent probe for general ROS detection, in this study, it was employed to assess NO levels as an indirect marker of ROS production. Previous research has demonstrated that DCFH-DA can respond to both ROS and reactive nitrogen species (RNS), including NO and peroxynitrite, making it suitable for detecting oxidative and nitrosative stress under specific conditions (27, 30-33).

**Table 1.** The Experimental Groups <sup>a</sup>

Groups	Laser Wavelength (nm)	Power (W)	Energy (J)	Irradiation Time (s)	With/Without IFN- $\gamma$
1	810	0.4	2	15	With and without
2	810	0.4	4	15	With and without
3	810	0.4	2	30	With and without
4	810	0.4	4	30	With and without
5	940	0.4	2	15	With and without
6	940	0.4	4	15	With and without
7	940	0.4	2	30	With and without
8	940	0.4	4	30	With and without
9	Control (no laser)	N/A	N/A	N/A	With and without

Abbreviation: N/A, not applicable.

<sup>a</sup> Each group was tested twice: Once with the application of IFN- $\gamma$  and once without.

The use of NO as a proxy for ROS has been validated in macrophage models where inducible nitric oxide synthase (iNOS) activity contributes to oxidative responses (31, 33). Nevertheless, we acknowledge that NO production can also occur independently of ROS generation, particularly via iNOS activation pathways stimulated by IFN- $\gamma$ . Therefore, while our measurements reflect total oxidative activity, they may not exclusively quantify ROS.

### 3.5. Data Analysis

The collected data were analyzed using SPSS 24. The Kolmogorov-Smirnov test was utilized to analyze the normal distribution of data, the *t*-test to assess the effects of IFN- $\gamma$  in each group, and the two-way analysis of variance (ANOVA) to compare ROS levels in the examined groups. A P-value < 0.05 was considered statistically significant.

## 4. Results

Table 2 presents a consolidated summary of NO levels under different experimental conditions, including laser wavelength, energy, exposure time, and the presence or absence of IFN- $\gamma$ . The values are expressed as means  $\pm$  standard deviations (SD) in micromoles ( $\mu$ mol). In addition to descriptive statistics, the table includes corresponding P-values and Cohen's *d* effect sizes for IFN- $\gamma$  comparisons to show the magnitude of biological effects.

Cohen's *d* values were calculated for each condition to quantify the magnitude of IFN- $\gamma$ 's effect on NO levels. The results demonstrate consistently large effect sizes (*d* > 0.8), with several conditions exceeding *d* = 2.0,

indicating a very large stimulatory effect of IFN- $\gamma$  on ROS production as measured by NO.

Based on the Kolmogorov-Smirnov test, data were normally distributed (*P* = 0.390), and Levene's test for equality of variances resulted in a P-value of 0.82, confirming that the variances are homogeneous.

One-way ANOVA revealed that irradiation energy (*F* = 19.00, *P* = 0.001), irradiation time (*F* = 12.20, *P* = 0.001), and the presence of IFN- $\gamma$  (*F* = 40.60, *P* < 0.001) all had significant main effects on NO production. In contrast, laser wavelength (810 nm vs. 940 nm) showed no significant effect (*F* = 0.22, *P* = 0.63). Additionally, no two-way, three-way, or four-way interaction terms were statistically significant (all *P* > 0.05), suggesting that these factors act independently rather than synergistically in modulating NO levels.

## 5. Discussion

This study aimed to evaluate the effect of two diode laser wavelengths, 810 nm and 940 nm, on ROS in macrophage cells, using NO levels as an indirect but widely accepted proxy for total oxidative activity. We hypothesized that the two wavelengths might influence ROS production differently due to their distinct absorption profiles. Additionally, we assessed how variations in irradiation energy, irradiation time, and the presence of IFN- $\gamma$  as a stimulant influenced ROS release. A control group, not exposed to laser irradiation, was included to provide a baseline for comparison.

Our results demonstrated that both 810 nm and 940 nm diode lasers can significantly increase ROS production compared to the control group. However, no

**Table 2.** Mean  $\pm$  Standard Deviations of Nitric Oxide Levels Under Different Experimental Conditions <sup>a,b</sup>

Laser Wavelength (nm)	Energy (J)	Irradiation time (s)	IFN- $\gamma$ Presence	Mean $\pm$ SD	Cohen's d	P-Value
810 nm	2	15	With	18 $\pm$ 2.73	3.33	0.001 <sup>c</sup>
			Without	11 $\pm$ 1.18		
	2	30	With	21 $\pm$ 2.64	2.55	0.001 <sup>c</sup>
			Without	13.8 $\pm$ 3.00		
	4	15	With	22.6 $\pm$ 3.7	1.51	0.005 <sup>d</sup>
			Without	15 $\pm$ 6.1		
940 nm	2	15	With	25.2 $\pm$ 5.2	0.75	0.01 <sup>d</sup>
			Without	21 $\pm$ 6.0		
	2	30	With	17.8 $\pm$ 3.1	1.85	0.002 <sup>d</sup>
			Without	11.4 $\pm$ 3.8		
	4	30	With	21 $\pm$ 3.2	2.23	0.001 <sup>c</sup>
			Without	14.4 $\pm$ 2.7		
Control (no laser)	N/A	N/A	With	20.8 $\pm$ 4.49	1.05	0.006 <sup>d</sup>
			Without	15.2 $\pm$ 6.1		
			With	24.6 $\pm$ 5.0	1.14	0.008 <sup>d</sup>
			Without	18.6 $\pm$ 5.5		
Control (no laser)	N/A	N/A	With	9.2 $\pm$ 0.83	0.66	0.05 <sup>e</sup>
			Without	8.4 $\pm$ 1.5		

Abbreviation: SD, standard deviations.

<sup>a</sup> The mean values indicate that nitric oxide (NO) levels were consistently higher at 4 J compared to 2 J, and with 30 seconds of irradiation compared to 15 seconds. Across all conditions, the presence of IFN- $\gamma$  resulted in substantially increased NO production compared to IFN- $\gamma$ -absent groups. These differences were statistically significant across all laser settings ( $P \leq 0.01$ ) and were also reflected by Cohen's d effect sizes ranging from 0.66 to 3.33, indicating moderate to very large biological effects of IFN- $\gamma$ .

<sup>b</sup> Laser treatment at both 810 nm and 940 nm, with varying energy levels and exposure times, significantly enhanced NO production when combined with IFN- $\gamma$  stimulation compared to both IFN- $\gamma$ -absent groups and untreated controls. The control group also demonstrated a significant difference between IFN- $\gamma$ -stimulated and unstimulated conditions ( $P = 0.05$ , Cohen's d = 0.66), supporting the role of IFN- $\gamma$  in promoting oxidative activity even in the absence of photostimulation.

<sup>c</sup>  $P \leq 0.001$ .

<sup>d</sup>  $P \leq 0.01$ .

<sup>e</sup>  $P \leq 0.05$ .

significant difference was observed in ROS release between the two wavelengths, leading to the rejection of the null hypothesis. Additionally, we observed that higher irradiation energy (4 J versus 2 J) and longer irradiation times (30 seconds versus 15 seconds) substantially elevated NO levels, indicating a corresponding increase in ROS production.

These results align with previous studies like Cerdeira (34) and Ivanov et al. (31). Cerdeira (34) found that LLLT can increase ROS production in neutrophils. Ivanov et al. (31) identified a positive correlation between ROS production and irradiation time but did not examine energy density (31). Our study builds on this by demonstrating that higher energy density also leads to increased ROS production, aligning with the results of Chen et al. (35).

In contrast to our findings, da Silva Neto Trajano et al. (36) observed no significant difference in ROS

production between laser-irradiated groups and the control group. Similarly, Yin et al. (37) reported a reduction in ROS levels with a 660 nm diode laser compared to their control, which is contrary to our results showing increased ROS levels with both 810 nm and 940 nm lasers. These discrepancies may be attributed to differences in experimental conditions, such as laser wavelength, power, energy density, and the use of IFN- $\gamma$  as a stimulant in our study.

Our study extends existing research by comparing two different diode laser wavelengths (810 nm and 940 nm). Previous studies, such as those by Yin et al. (37) and Chen et al. (35), primarily focused on single wavelengths without directly comparing different wavelengths as we did. The results of the current study also revealed that IFN- $\gamma$  led to a significant increase in ROS levels, indicating its efficacy in stimulating macrophage activation. The IFN- $\gamma$  is an important cytokine in both innate and adaptive immune responses, and it enhances

the secretion of anti-inflammatory cytokines by macrophages (29, 30).

While no other studies have used IFN- $\gamma$  to stimulate ROS production, our findings show a significant increase in ROS levels with its use. Compared to other priming agents such as LPS (31, 37), IFN- $\gamma$  appears to elicit a distinct and possibly more sustained activation of ROS-generating pathways. Ivanov et al. (31) demonstrated that LPS-stimulated serum proteins generate long-lived ROS species following laser exposure, whereas IFN- $\gamma$  may more directly and robustly activate intracellular enzymatic machinery. In fact, the observed synergistic effect between IFN- $\gamma$  and laser irradiation in enhancing ROS production may be attributed to the activation of multiple ROS-generating pathways. The IFN- $\gamma$  is known to prime macrophages through the upregulation of NADPH oxidase, an enzyme complex responsible for producing superoxide radicals during the respiratory burst (20, 29, 38). Simultaneously, laser irradiation, particularly in the near-infrared range, can stimulate mitochondrial ROS production via cytochrome c oxidase (CCO) activation. The combined stimulation of mitochondrial and NADPH oxidase pathways likely accounts for the significantly elevated ROS levels observed in IFN- $\gamma$ -treated, laser-irradiated groups. This dual-pathway activation supports the hypothesis that IFN- $\gamma$  sensitizes macrophages to LLLT by amplifying oxidative responses.

The absorption of light by mitochondrial chromophores, particularly CCO, is significant due to its prominent absorption peaks in the near-infrared region, especially around 810 nm (12). While we hypothesized that 810 nm would stimulate ROS production more effectively due to stronger CCO absorption, our findings showed no significant difference between the 810 nm and 940 nm wavelengths. This suggests that, although 940 nm is outside CCO's peak absorption range, it may still exert biological effects via secondary pathways such as water absorption or less well-characterized mitochondrial components. These results highlight the complexity of wavelength-specific interactions and underscore the need for further mechanistic studies.

While NO was used as a surrogate marker of ROS in this study, it is important to acknowledge that NO production can occur independently via iNOS activity and may not fully capture the complexity of ROS. Nonetheless, DCFH-DA is a validated probe for detecting general oxidative and nitrosative stress, including

species such as NO, hydrogen peroxide, and peroxynitrite (27, 31). To improve specificity, future studies should incorporate direct ROS detection methods, such as dihydroethidium (DHE) for superoxide and Amplex Red for hydrogen peroxide, to obtain a more nuanced profile of oxidative changes in macrophages post-irradiation.

The ROS are central to many physiological and pathological processes, ranging from microbial defense and wound healing to chronic inflammation and carcinogenesis (39, 40). The ability to modulate ROS levels via laser therapy has potential clinical applications in tissue regeneration, immune modulation, and inflammation control (39-41). Given ROS's dual role, being both beneficial and potentially harmful, precise control of laser parameters (wavelength, energy density, duration) is essential to ensure therapeutic safety and efficacy (42, 43).

While our study provides valuable insights, it is important to acknowledge its limitations. This study was conducted on a specific type of macrophage cell, and the results may not be generalizable to other cell types or in vivo conditions. The J774A.1 murine macrophage cell line was selected due to its reproducibility, availability, and established use in oxidative stress and immunomodulation studies. While this model provides consistency for controlled in vitro experimentation, it may not fully reflect the complexity of primary macrophages derived from human tissue. Differences in receptor expression, mitochondrial function, and cytokine responsiveness between murine cell lines and primary human macrophages could influence ROS production. Therefore, caution should be exercised when extrapolating these results to human physiology, and future studies should validate these findings using primary macrophages or human-derived cell models.

Future studies should aim to replicate these findings in different cell lines and explore the effects of other laser wavelengths and energy densities. Specifically, future studies should investigate the relationship between LLLT, ROS production, and clinical outcomes. Additionally, more extensive cellular studies are suggested to better understand the underlying mechanisms of laser-induced ROS production and its therapeutic potential.

Temperature changes during laser exposure were not continuously monitored in this study. However, the low-level irradiation parameters used (0.4 W for a maximum



of 30 seconds) fall within established non-thermal LLLT thresholds (3, 25). Previous studies have shown that under such conditions, cellular temperature elevations are minimal and unlikely to contribute significantly to the observed biological effects. Nevertheless, future work should incorporate direct temperature monitoring to definitively exclude thermal contributions to ROS generation.

In addition, future work should investigate how the mode of laser emission, pulsed versus continuous wave, modulates ROS production. Pulsed lasers may reduce heat buildup while still stimulating mitochondrial chromophores efficiently, potentially resulting in distinct oxidative responses compared to continuous wave modes. Some studies suggest that pulsed irradiation may produce lower peak ROS levels or induce different signaling cascades. Exploring this variable will be critical for optimizing LLLT protocols for specific clinical applications.

### 5.1. Conclusions

In conclusion, our study demonstrates that while the specific diode laser wavelengths of 810 nm and 940 nm do not significantly differ in ROS release, both significantly increase ROS production compared to a control group. Furthermore, irradiation time, energy density, and the presence of IFN- $\gamma$  play crucial roles in modulating ROS production in macrophage cells.

### Footnotes

**Authors' Contribution:** Study concept and design: T. F., S. L., and S. H.; Analysis and interpretation of data: M. S. and R. Sh.; Drafting of the manuscript: R. Sh.; Critical revision of the manuscript for important intellectual content: T. F., S. L., and S. H.; Statistical analysis: R. Sh.

**Conflict of Interests Statement:** The authors declare no conflict of interest.

**Data Availability:** All data used for this project, including fluorescence data, is available upon reasonable request from the corresponding author.

**Ethical Approval:** The study adhered to ethical guidelines and was approved by the relevant Institutional Review Board. This research is registered under the ethics code [IR.IAU.DENTAL.REC.1399.094](https://www.iaudental.ir/REC/1399.094).

**Funding/Support:** The present research received no funding/support.

### References

1. Hamblin MR, Agrawal T, de Sousa M. *Handbook of Low-Level Laser Therapy*. Singapore, Asia: Pan Stanford Publishing: CRC Press; 2016. <https://doi.org/10.1201/9781315364827>.
2. Miao L, Xu Z, Sui J, Meng X, Huo S, Liu S, et al. A New Nanoplatform Under NIR Released ROS Enhanced Photodynamic Therapy and Low Temperature Photothermal Therapy for Antibacterial and Wound Repair. *Int J Nanomedicine*. 2024;**19**:7509-27. [PubMed ID: 39071503]. [PubMed Central ID: PMC11283834]. <https://doi.org/10.2147/IJN.S471623>.
3. Ebrahimi H, Kargar M, Shoorgashti R. The Comparison of 940nm and 810nm Diode Laser Effects on the Repair of Inferior Alveolar Sensory Nerve Injury: A Clinical Trial. *J Dent (Shiraz)*. 2024;**25**(3):215-22. [PubMed ID: 39371960]. [PubMed Central ID: PMC11452597]. <https://doi.org/10.30476/dentjods.2023.97393.2013>.
4. Yeganehfard H, Azizi A, Shoorgashti R, Shokouhi Mostafavi SK, Kharazi-Fard M, Farrokhnia T. Evaluating Photodynamic Therapy at Two Wavelengths and Cold Plasma as Alternatives to Chlorhexidine for Targeting *Streptococcus mutans*: An In-vitro Study. *Archives of Clinical Infectious Diseases*. 2025;**20**(4). <https://doi.org/10.5812/archcid-160774>.
5. Escobar LM, Grajales M, Bendahan Z, Jaimes S, Baldion P. Osteoblastic differentiation and changes in the redox state in pulp stem cells by laser treatment. *Lasers Med Sci*. 2024;**39**(1):87. [PubMed ID: 38443654]. [PubMed Central ID: PMC10914891]. <https://doi.org/10.1007/s10103-024-04016-z>.
6. Yang J, Fu Q, Jiang H, Zhong H, Qin HK, Miao X, et al. Blue light photobiomodulation induced osteosarcoma cell death by facilitating ferroptosis and eliciting an incomplete tumor cell stress response. *J Photochem Photobiol B*. 2024;**258**:113003. [PubMed ID: 39121719]. <https://doi.org/10.1016/j.jphotobiol.2024.113003>.
7. Moradi A, Ghaffari Novin M, Bayat M. A Comprehensive Systematic Review of the Effects of Photobiomodulation Therapy in Different Light Wavelength Ranges (Blue, Green, Red, and Near-Infrared) on Sperm Cell Characteristics in Vitro and in Vivo. *Reprod Sci*. 2024;**31**(11):3275-302. [PubMed ID: 39095677]. <https://doi.org/10.1007/s43032-024-01657-x>.
8. Shoorgashti R, Moshiri A, Lesan S. Evaluation of Oral Mucosal Lesions in Iranian Smokers and Non-smokers. *Niger J Clin Pract*. 2024;**27**(4):467-74. [PubMed ID: 38679769]. [https://doi.org/10.4103/njcp.njcp\\_702\\_23](https://doi.org/10.4103/njcp.njcp_702_23).
9. Hong Y, Boiti A, Vallone D, Foulkes NS. Reactive Oxygen Species Signaling and Oxidative Stress: Transcriptional Regulation and Evolution. *Antioxidants (Basel)*. 2024;**13**(3). [PubMed ID: 38539845]. [PubMed Central ID: PMC10967436]. <https://doi.org/10.3390/antiox13030312>.
10. Checa J, Aran JM. Reactive Oxygen Species: Drivers of Physiological and Pathological Processes. *J Inflamm Res*. 2020;**13**:1057-73. [PubMed ID: 33293849]. [PubMed Central ID: PMC7719303]. <https://doi.org/10.2147/IJR.S275595>.
11. Shoorgashti R, Etemadi A, Mirzaee S, Shoorgashti R. Correlation of Periodontitis and In Vitro Fertilization Outcome: A Literature Review. *Journal of Research in Dental and Maxillofacial Sciences*. 2023;**8**(4):295-301. <https://doi.org/10.61186/jrdms.8.4.295>.

12. Shoorgashti R, Nikmaram R, Azimi Y, Rouientan A, Ebrahimi H, Lesan S. Effectiveness of cold plasma application in oral wound healing process: A scoping review. *Oral Dis.* 2024;**30**(8):5062-81. [PubMed ID: 39224064]. <https://doi.org/10.1111/odi.15119>.
13. Najafi M, Lotfali E, Shoorgashti R, Ebrahimi H, Lesan S. Evaluation of the Effects of Dielectric Barrier Discharge Plasma on Candida albicans Strains: An in vitro Study. *Jundishapur Journal of Microbiology.* 2024;**16**(11). <https://doi.org/10.5812/jjm-142236>.
14. Mirhosseini N, Shoorgashti R, Lesan S. The evaluation of clinical factors affecting oral health impacts on the quality of life of Iranian elderly patients visiting dental clinics: A cross-sectional study. *Spec Care Dentist.* 2024;**44**(4):1219-27. [PubMed ID: 38430466]. <https://doi.org/10.1111/scd.12980>.
15. Ghasemian Gorji R, Shoorgashti R, Lotfali E, Farhadi S. MIC Determination of Silver Nanoparticles on Nystatin-Resistant Candida albicans in Patients with Denture Stomatitis: An In Vitro Study. *Middle East Journal of Rehabilitation and Health Studies.* 2023;**11**(1). <https://doi.org/10.5812/mejrh-138030>.
16. Shoorgashti R, Shahrzad H, Nowroozi S, Ghadamgahi B, Mehrara R, Oroojalian F. Evaluation of the antibacterial and cytotoxic activities of Ag/ZnO nanoparticles loaded polycaprolactone/chitosan composites for dental applications. *Nanomedicine J.* 2023;**10**(1). <https://doi.org/10.22038/NMJ.2022.69755.1744>.
17. Hamblin MR. Mechanisms and applications of the anti-inflammatory effects of photobiomodulation. *AIMS Biophys.* 2017;**4**(3):337-61. [PubMed ID: 28748217]. [PubMed Central ID: PMC5523874]. <https://doi.org/10.3934/biophy.2017.3.337>.
18. He K, Zhou X, Zheng F, Ju X, Fu SN, Wong AYL. Histological, Physiological and Biomechanical Effects of Low-Level Laser Therapy on Tendon Healing in Animals and Humans: A Systematic Review. *Ann Biomed Eng.* 2023;**51**(12):2659-707. [PubMed ID: 37899380]. <https://doi.org/10.1007/s10439-023-03364-1>.
19. Mayahi M, Shoorgashti R, Lesan S, Rahimi MK, Ebrahimi H. Comparing the Antibacterial Effects of 660 nm Laser Therapy on *Staphylococcus aureus* Using Phycocyanin and Methylene Blue: An In-Vitro Study. *Middle East Journal of Rehabilitation and Health Studies.* 2024;**12**(2). <https://doi.org/10.5812/mejrh-155074>.
20. Herb M, Schramm M. Functions of ROS in Macrophages and Antimicrobial Immunity. *Antioxidants (Basel).* 2021;**10**(2). [PubMed ID: 33669824]. [PubMed Central ID: PMC7923022]. <https://doi.org/10.3390/antiox10020313>.
21. Virag L, Jaen RI, Regdon Z, Bosca L, Prieto P. Self-defense of macrophages against oxidative injury: Fighting for their own survival. *Redox Biol.* 2019;**26**:101261. [PubMed ID: 31279985]. [PubMed Central ID: PMC6614175]. <https://doi.org/10.1016/j.redox.2019.101261>.
22. Alspach E, Lussier DM, Schreiber RD. Interferon gamma and Its Important Roles in Promoting and Inhibiting Spontaneous and Therapeutic Cancer Immunity. *Cold Spring Harb Perspect Biol.* 2019;**11**(3). [PubMed ID: 29661791]. [PubMed Central ID: PMC6396335]. <https://doi.org/10.1101/cshperspect.a028480>.
23. Kopitar-Jerala N. The Role of Interferons in Inflammation and Inflammasome Activation. *Front Immunol.* 2017;**8**:873. [PubMed ID: 28791024]. [PubMed Central ID: PMC5525294]. <https://doi.org/10.3389/fimmu.2017.00873>.
24. Ebrahimi H, Darvish F, Alaeddini M, Etemad-Moghadam S. Comparison between the Effect of 810 nm and 940 nm Diode Laser Irradiation on Histopathological Changes in Iatrogenic Oral Ulcers: an Animal Study. *J Dent (Shiraz).* 2021;**22**(4):267-72. [PubMed ID: 34904123]. [PubMed Central ID: PMC8665437]. <https://doi.org/10.30476/DENTJODS.2021.86623.1202>.
25. Mohammadi S, Shoorgashti R, Lotfali E, Lesan S, Ebrahimi H. The Evaluation of 660, 810, and 940nm Laser Wavelengths on Nystatin-Resistant Candida albicans: An In-Vitro Study. *Jundishapur Journal of Microbiology.* 2024;**17**(7). <https://doi.org/10.5812/jjm-144680>.
26. Mansouri V, Arjmand B, Rezaei Tavirani M, Razzaghi M, Rostami-Nejad M, Hamdiah M. Evaluation of Efficacy of Low-Level Laser Therapy. *J Lasers Med Sci.* 2020;**11**(4):369-80. [PubMed ID: 33425286]. [PubMed Central ID: PMC7736953]. <https://doi.org/10.34172/jlms.2020.60>.
27. Taraszkievicz A, Szewczyk G, Sarna T, Bielawski KP, Nakonieczna J. Photodynamic Inactivation of Candida albicans with Imidazoacridinones: Influence of Irradiance, Photosensitizer Uptake and Reactive Oxygen Species Generation. *PLoS One.* 2015;**10**(6). e0129301. [PubMed ID: 26053272]. [PubMed Central ID: PMC4459871]. <https://doi.org/10.1371/journal.pone.0129301>.
28. Tamaki N, Tomofuji T, Ekuni D, Yamanaka R, Yamamoto T, Morita M. Short-term effects of non-surgical periodontal treatment on plasma level of reactive oxygen metabolites in patients with chronic periodontitis. *J Periodontol.* 2009;**80**(6):901-6. [PubMed ID: 19485819]. <https://doi.org/10.1902/jop.2009.080640>.
29. Wu C, Xue Y, Wang P, Lin L, Liu Q, Li N, et al. IFN-gamma primes macrophage activation by increasing phosphatase and tensin homolog via downregulation of miR-3473b. *J Immunol.* 2014;**193**(6):3036-44. [PubMed ID: 25092892]. <https://doi.org/10.4049/jimmunol.1302379>.
30. Tekautz TM, Zhu K, Grenet J, Kaushal D, Kidd VJ, Lahti JM. Evaluation of IFN-gamma effects on apoptosis and gene expression in neuroblastoma-preclinical studies. *Biochim Biophys Acta.* 2006;**1763**(10):1000-10. [PubMed ID: 16979248]. <https://doi.org/10.1016/j.bbamcr.2006.06.014>.
31. Ivanov VE, Usacheva AM, Chernikov AV, Bruskov VI, Gudkov SV. Formation of long-lived reactive species of blood serum proteins induced by low-intensity irradiation of helium-neon laser and their involvement in the generation of reactive oxygen species. *J Photochem Photobiol B.* 2017;**176**:36-43. [PubMed ID: 28941776]. <https://doi.org/10.1016/j.jphotobiol.2017.09.012>.
32. Gardiner B, Dougherty JA, Ponnalagu D, Singh H, Angelos M, Chen CA, et al. Measurement of Oxidative Stress Markers In Vitro Using Commercially Available Kits. In: Berliner LJ, Parinandi NL, editors. *Measuring Oxidants and Oxidative Stress in Biological Systems*. Cham (CH), Switzerland; 2020. p. 39-60. eng. [https://doi.org/10.1007/978-3-030-47318-1\\_4](https://doi.org/10.1007/978-3-030-47318-1_4).
33. Hardy M, Zielonka J, Karoui H, Sikora A, Michalski R, Podsiadly R, et al. Detection and Characterization of Reactive Oxygen and Nitrogen Species in Biological Systems by Monitoring Species-Specific Products. *Antioxid Redox Signal.* 2018;**28**(15):1416-32. [PubMed ID: 29037049]. [PubMed Central ID: PMC5910052]. <https://doi.org/10.1089/ars.2017.7398>.
34. Cerdeira CD, Lima Brigagao MR, Carli ML, de Souza Ferreira C, de Oliveira Isac Moraes G, Hadad H, et al. Low-level laser therapy stimulates the oxidative burst in human neutrophils and increases their fungicidal capacity. *J Biophotonics.* 2016;**9**(11-12):1180-8. [PubMed ID: 27243910]. [PubMed Central ID: PMC5133186]. <https://doi.org/10.1002/jbio.201600035>.
35. Chen ZY, Li DL, Duan XD, Peng DZ. [Effect of low-energy 633 nm red light stimulation on proliferation and reactive oxygen species level of human epidermal cell line HaCaT]. *Zhonghua Shao Shang Za Zhi.* 2016;**32**(9):560-5. [PubMed ID: 27647074]. <https://doi.org/10.3760/cma.j.issn.1009-2587.2016.09.010>.

36. da Silva Neto Trajano AL, da Silva CL, de Carvalho SN, Cortez E, Mencialha AL, de Souza da Fonseca A, et al. Cell viability, reactive oxygen species, apoptosis, and necrosis in myoblast cultures exposed to low-level infrared laser. *Lasers Med Sci.* 2016;**31**(5):841-8. [PubMed ID: 26886589]. <https://doi.org/10.1007/s10103-016-1909-8>.
37. Yin K, Zhu R, Wang S, Zhao RC. Low level laser (LLL) attenuate LPS-induced inflammatory responses in mesenchymal stem cells via the suppression of NF-kappaB signaling pathway in vitro. *PLoS One.* 2017;**12**(6). e0179175. [PubMed ID: 28594941]. [PubMed Central ID: PMC5464618]. <https://doi.org/10.1371/journal.pone.0179175>.
38. Voloshyna I, Littlefield MJ, Reiss AB. Atherosclerosis and interferon-gamma: new insights and therapeutic targets. *Trends Cardiovasc Med.* 2014;**24**(1):45-51. [PubMed ID: 23916809]. [PubMed Central ID: PMC3844070]. <https://doi.org/10.1016/j.tcm.2013.06.003>.
39. Cai Y, Zhang X, Yang C, Jiang Y, Chen Y. Melatonin alleviates high-fat-diet-induced dry eye by regulating macrophage polarization via IFT27 and lowering ERK/JNK phosphorylation. *iScience.* 2024;**27**(7):110367. [PubMed ID: 39100927]. [PubMed Central ID: PMC11294704]. <https://doi.org/10.1016/j.isci.2024.110367>.
40. Shoorgashti R, Khalililrad M, Haji Molla Hoseini M, Lesan S. [Evaluation of the relationship between nitric oxide and candida albicans-associated denture stomatitis: A cross-sectional study]. *Caspian J Dental Res.* 2023;**12**(2):82-8. FA. <https://doi.org/10.22088/cjdr.12.2.82>.
41. Hasturk H, Kantarci A, Van Dyke TE. Oral inflammatory diseases and systemic inflammation: role of the macrophage. *Front Immunol.* 2012;**3**:118. [PubMed ID: 22623923]. [PubMed Central ID: PMC3353263]. <https://doi.org/10.3389/fimmu.2012.00118>.
42. Dunnill C, Patton T, Brennan J, Barrett J, Dryden M, Cooke J, et al. Reactive oxygen species (ROS) and wound healing: the functional role of ROS and emerging ROS-modulating technologies for augmentation of the healing process. *Int Wound J.* 2017;**14**(1):89-96. [PubMed ID: 26688157]. [PubMed Central ID: PMC7950185]. <https://doi.org/10.1111/iwj.12557>.
43. Beirami AD, Akhtari N, Noroozi R, Hatamabadi D, Hasan SMF, Ayatollahi SA, et al. Bringing back Galium aparine L. from forgotten corners of traditional wound treatment procedures: an antimicrobial, antioxidant, and in-vitro wound healing assay along with HPTLC fingerprinting study. *BMC Complement Med Ther.* 2024;**24**(1):279. [PubMed ID: 39044256]. [PubMed Central ID: PMC11267910]. <https://doi.org/10.1186/s12906-024-04355-y>.