Published online: 2024 July 2.



Research Article

Integration of Lipofilling with Blepharoplasty to Optimize Periorbital Rejuvenation: A Retrospective Study

Daniele Bollero¹, Sabrina Donzelli (b^{2,*}, Bruno Bovani³, Matteo Tretti Clementoni⁴, Fabrizio Melfa⁵, Domenico Piccolo⁶, Melinda Keener⁷

¹ Plastic Surgery and Burn Unit, CTO Hospital, Torino, Italy

² National Security Council (NSC), Washington, USA

³ Centro Medico Esculapio Perugia, Perugia PG, Italy

⁴ Laserplast, Milano, Italy

⁵ Mediaging Clinic Center Palermo Italy, Italy

⁶ Novea Skin Center Avezzano, Avezzano AQ, Italy

⁷ Private Practice, Jacksonville, USA

 * *Corresponding author:* National Security Council (NSC), Washington, USA. Email: sab.donzelli@gmail.com

Received 2024 March 25; Revised 2024 April 27; Accepted 2024 April 29.

Abstract

Background: Surgical blepharoplasty has traditionally been used to rejuvenate the periorbita by removing excess tissue. While this often enhances appearance, it does not address the volume loss associated with aging. Hyaluronic acid (HA) fillers have been used to address volume loss; however, issues such as the Tyndall effect, chronic edema, and the cost of repeated injections make them less than ideal. Advances in contemporary surgery have made autologous fat a natural alternative for filling volume loss in the periorbita. Fat grafting, whether using macrofat, microfat, or nanofat, can provide excellent results, but a separate procedure is required to harvest the fat from the patient.

Objectives: The aim of this study was to analyze and describe complications and patient satisfaction by comparing blepharoplasty alone with blepharoplasty combined with lipofilling.

Methods: A retrospective review of a single surgeon's private practice cases was performed on 45 consecutive patients who underwent surgical blepharoplasty alone and 64 patients who had surgical blepharoplasty combined with lipofilling from 2002 to 2021. Complications, pain scores, and patient satisfaction reports were assessed.

Results: Upon comparing the blepharoplasty alone group to the blepharoplasty with lipofilling group, the 109 patients did not differ significantly in age. Both groups were predominantly female. There were no major complications, only minor ones, and though the blepharoplasty alone group had a higher complication rate, this was not statistically significant. Lipofilling added approximately 30 minutes to the 60-minute blepharoplasty procedure, but pain scores were higher in the combination group. However, pain dissipated quickly and, while statistically significant, was not clinically significant. Patient satisfaction with the additional lipofilling was high, which resulted in lipofilling being added to all blepharoplasty procedures.

Conclusions: Traditional surgical blepharoplasty can be combined with fat grafting techniques to address tissue loss and descent. Lipofilling integrated with blepharoplasty is a safe procedure. It is well tolerated by patients and provides high satisfaction with good aesthetic outcomes. The use of autologous fat offers an economical and effective advantage for patients.

Keywords: Blepharoplasty, Eye Lift, Lipofilling, Fat Transfer, Fat Graft, Surgery, Eyelid

1. Background

For most individuals, the first signs of aging are noted around the eyes. Due to the complexity of periorbital anatomy and the consequent aging process, which includes volume loss, skin laxity, and tissue descent, a variety of surgical and non-surgical methods can be combined to rejuvenate the periorbita (1). Surgical blepharoplasty has traditionally been used to rejuvenate the upper and lower eyelids by removing excess and lax soft tissue that has descended with aging (1). Overall, blepharoplasty has a low complication rate

Copyright @ 2024, Journal of Skin and Stem Cell. This open-access article is available under the Creative Commons Attribution-NonCommercial 4.0 (CC BY-NC 4.0) International License (https://creativecommons.org/licenses/by-nc/4.0/), which allows for the copying and redistribution of the material only for noncommercial purposes, provided that the original work is properly cited.

and is well tolerated by most patients; however, it does not address the issue of volume loss. It is important to find the best strategy to maximize the rejuvenation effect while minimizing complications. In-depth discussions with the patient to determine goals, risk factors, and alternatives are required to decide on the best combination of modalities.

A variety of modalities have been used with blepharoplasty to enhance the aesthetic rejuvenation of the periorbita. Botulinum toxin, hyaluronic acid fillers, and platelet-rich plasma (PRP) have all been used. Botulinum toxin can be costly and is required frequently. Hyaluronic acid fillers carry an increased risk of vessel occlusion, prolonged edema, scarring of the lymphatics of the periorbita, and discoloration around the eyes. PRP also requires more than one treatment and provides short-term results. In the last 10 years, advances in fat grafting have made autologous fat a very reasonable option to combine with surgical blepharoplasty (2). Recently, Marten and Elyassnia (2) reviewed the role of fat grafting in surgical procedures to rejuvenate the periorbital area by treating the problem of volume loss. Most agree that fat grafting provides a more youthful and natural aesthetic appearance than blepharoplasty alone. The shift from macrofat of the past to microfat has allowed for more preservation of the stem cell component, providing both a regenerative therapy and an aesthetic effect. Nanofat has transformed into "lipofilling" with very small fat particles and a high concentration of growth factors and stem cells, making the fat highly regenerative. This is relevant to understanding the potential impact of adipose tissue adjunct to an already well-standardized intervention. Nanofat addresses volume loss and revitalizes tissues of the periorbita affected by aging (3, 4). Fat grafting also has the advantage of providing longer-lasting results compared to other modalities. However, fat grafting does require an additional harvest site, thus an additional procedure, raising concerns about creating undue burden on the patient and the postoperative course. This retrospective study aimed to investigate the short-term postoperative effects of lipofilling in combination with blepharoplasty for patients undergoing periorbital rejuvenation by an experienced plastic surgeon in a private practice setting.

2. Objectives

This retrospective study was aimed to investigate the short term post operative effects of lipofilling in combination with blepharoplasty for patients undergoing periorbita rejuvenation by an experienced plastic surgeon in a private practice setting.

3. Methods

3.1. Study Design

This retrospective study analyzes the short-term postoperative period of patients undergoing surgical blepharoplasty alone (Group 1) or blepharoplasty combined with lipofilling (Group 2) from 2002 to 2021.

3.2. Participants

Patients for the study were recruited from a private practice setting. All patients signed written informed consent forms for the procedures and photography.

3.3. Procedures

All procedures were performed by a single plastic surgeon. Meticulous marking was done while the patient was standing to preserve landmarks once the patient was supine and local anesthesia was infiltrated. Intravenous sedation was administered using Midazolam (2 mg), Fentanyl (50 - 100 micrograms), and Propofol (in small boluses). Additional local anesthesia was provided using 2% Mepivacaine with epinephrine for eyelids and infraorbital nerve blocks. Oxygen flow was supplied during the procedure to assist with the patient's spontaneous breathing. Amoxicillin-clavulanic acid was used intraoperatively for prophylaxis and postoperatively.

Ophthalmic Betadine was used to prep the periorbital area, while regular Betadine was used to prep the fat harvest site. For upper blepharoplasty, excess upper eyelid skin was sharply excised along with a small portion of the orbicularis muscle and the medial fat pad, if in excess. Lower blepharoplasty was performed with a subciliary incision, and a musculocutaneous flap was dissected up to the lower orbital edge. Excess adipose tissue was removed from the three fat pads, and a thin strip of skin and muscle was excised. Care was taken to ensure meticulous hemostasis in all phases of the surgery. The upper blepharoplasty incision was closed with a running intradermal Nylon 5 - 0 suture, while the lower blepharoplasty incision was closed with interrupted 4 -0 Silk sutures. When performing lipofilling, 2% Mepivacaine with epinephrine was infiltrated at the harvesting cannula entrance site. Fat was harvested using a Tonnard Tulip cannula. The harvest sites were the inner knee for females and the flanks for males. A total of 20 - 40 mL of fat was harvested from each side (5). The choice of harvest sites was based on the increased stability and survivability of the fat in these regions, ensuring that if weight loss occurred postoperatively, there would not be a drastic change. Local anesthesia was avoided in the harvesting area to preserve the adipocytes as much as possible. The cannula entry site was closed with a 5 - 0 silk suture. Fat was processed according to the Coleman protocol, with centrifugation for 3 minutes at 3000 rpm to separate the nonviable components, which were then drained. Approximately 10 cc of the harvested fat was processed with the Tulip system as nanofat (6-8). Nanofat was produced by mechanical shuffling and filtration of microfat. Because the nanofat particles are extremely small, they do not provide notable volume effects. Studies indicate that nanofat contains abundant stromal cells and adipose-derived stem cells, which help reconstruct dermal support structures, such as collagen, and regenerate healthier, younger-appearing skin (9). In Italy, only mechanically treated stem cells are permitted, while enzymatic separation is prohibited. Microfat was injected using 25G cannulas for volume restoration, while nanofat was injected with 27 - 30G needles for fine rhytid filling and overall regenerative effects. Recently, a high-precision injection pen (Lipopen) has been used to provide greater uniformity of distribution; traditionally, injections were performed manually using syringes. Injections were performed slowly, with small boluses administered while withdrawing the cannula to avoid vascular occlusion and uniformly distribute fat. Microfat was placed with a microcannula in the tear trough and supraperiosteal in the lower eyelid. In other areas, such as the zygomatic region, microfat was placed subdermally into the subcutaneous tissue. A total of 4 -15 mL was placed bilaterally to adequately restore volume. Nanofat was injected into the intradermal plane to form a wheal, which resorbed within a few days postoperatively.Surgical blepharoplasty typically took an average of 60 minutes to perform, and lipofilling

added an average of 30 minutes to the procedure. Postoperatively, patients were observed for 30 minutes in recovery and then discharged home. Figures 1 and 2 illustrate two cases preoperatively and postoperatively.



Figure 1. Female patient, 40 years old. Upper and lower blepharoplasty and lipofiling micro and nano for a better harmonization of periorbital area



Figure 2. Female patient, 42 years old. Upper and lower blepharoplasty and lipofiling micro and nano

3.4. Postsurgical Care

A dressing was applied over the incisions, ensuring the eyes could open and vision was not obstructed. An NSAID was administered postoperatively for 3 days to address pain, and antibiotics were also given. Immediately after surgery, a Pulsed Short Wave Therapy (PSWT) device (RecoveryRX, BioElectroninc, Frederick, MD, USA) was placed on each side to help reduce pain and inflammation. These devices were activated and checked to ensure proper functioning before placement. The device was secured to the head and face for 6 days until suture removal.Botox, 50 units, was injected one month postoperatively to decrease periorbital rhytids and enhance the aesthetic result. Botox therapy also helped reduce tension on the scar.

3.5. Outcome Measures

Table 1. Demographic Description of The Groups and Evaluation of Pain: Chi-Square Test and P-Value are Reported and Show no Statistically Significant Differences Be	tween
Groups ^{a, b}	

Variables	Group 1	Group 2	P-Value	Chi2
variables	Blepharoplasty (N = 45)	Blepharoplasty + lipofilling (N = 64)		
Age (y)	54.3	54.1		
Gender			0.65	0.21
Females	40 (89)	55 (86)		
Pain control: How would you judge it?			3.04	0.22
Insufficient	0(0)	0(0)		
Sufficient	0(0)	3 (5)		
Good	9 (20)	17 (26)		
Optimal	36 (80)	44 (69)		
Satisfaction: Would you recommend it?			1.74	0.42
No	0(0)	2 (3)		
Yes	42 (93)	56 (88)		
Maybe	3 (7)	6 (9)		

^a Statistically Significant Difference, Alpha level of significant set below 0.05.

^b Values are presented as mean SD or No (%).

The outcome measures considered for comparison between the two groups were postoperative complications, patient-reported satisfaction levels, and postoperative pain on days 1 - 4 using a Numeric Rating Scale (NRS) (10, 11).

3.6. Data Analysis

Descriptive statistics were performed with proportions for categorical variables and means and standard deviations for normally distributed continuous variables. After assessing normal distribution, parametric tests (t-test, ANOVA) for repeated measures were applied to compare the differences between the two groups and analyze whether the type of procedure affected pain over time.

4. Results

This retrospective analysis included 109 patients with a mean age of 54.2 (SD \pm 10.1). Of the 109 patients, 45 were in Group 1 (blepharoplasty alone) and 64 were in Group 2 (combined blepharoplasty-lipofilling). The average age was 53 years for both groups. In Group 1, 86% were female, and in Group 2, 89% were female. Demographic details are reported in Table 1. Pain control was rated as good to optimal in the majority of the sample in both groups. Satisfaction was rated as good in both groups, but dissatisfaction was reported by only two patients, with no statistically significant differences between the groups.

The decision to add lipofilling to blepharoplasty was made by the patient during the preoperative consultation. Both groups were comprised of similar patients, with no significant differences in age, and females were predominant. The two groups showed similar characteristics at baseline, with no differences in age and an equal distribution of females (chi2 0.21, P = 0.65): 86% in the blepharoplasty alone group and 89% in the blepharoplasty plus lipofilling group.

As reported in Table 2, pain scores were low for both groups. Higher scores were reported in the first two days in the blepharoplasty plus lipofilling group, but by days 3 and 4, the difference was no longer statistically significant (Figure 3). Postoperative complications reported by patients are shown in Table 3. Overall, 20% of patients (9 out of 45) undergoing blepharoplasty alone and 12% of patients (8 out of 64) undergoing blepharoplasty with lipofilling reported mild and transient problems.



Figure 3. Trend of pain average values over time in days after surgery in the two intervention groups

Table 2. Postoperative subjective evaluation of pain comparing two intervention procedures statistically significant difference $^{\rm a}$

	Group 1 Group 2				
Pain	Bleph 45)	naroplasty (N =	Blepharoplasty + lipofilling (N = 64)		P-Value
Day					
1	t	0.5 (1.1)	1.2 (1.7)	9.51	0.0022 b
1	2	0.3 (0.9)	0.8(1.4)	5.28	0.022 ^b
3	3	0.1(0.8)	0.3 (0.9)	0.83	0.37
	4	0.1(0.5)	0.2 (0.7)	0.41	0.52

^a Values are presented as mean SD.

^bAlpha level of significant set below 0.05.

Table 3. Postoperative Complications Reported by Patients Group 1 Group 2 Variables Blepharoplasty (N = 45) Blepharoplasty + lipofilling (N = 64) 8/64 9/45 Itching 3 2 Nausea 3 3 Hypotension 2 1 Headache 0 2 Bleeding 1 0

5. Discussion

Retrospective analysis revealed that the integration of lipofilling with blepharoplasty is very successful and well tolerated by patients. The combination of the two procedures does not increase complications and enhances the aesthetics of the periorbita. The focus of this study was on postoperative pain, patient complications, and patient satisfaction. Although pain

J Skin Stem Cell. 2024; 11(2): e147228.

was reportedly higher in the blepharoplasty-lipofilling group, it was brief and described as a "mild ache." Despite being statistically significant, the differences in pain between the groups were not clinically significant. The fact that patients in both groups stated they would undergo the surgery again is evidence that they tolerated the procedures well (12).

The best candidates for lipofilling combined with blepharoplasty are those who have had no prior surgery of the periorbita and, as suggested by Cohen (3), those with more advanced aging requiring more than 5 syringes of fillers. It is important to note that both groups included in the analysis had an average age of 53 years, which is in line with previous studies on blepharoplasty (10). The numerical rating scale (NRS) has been utilized to evaluate pain levels in a variety of disorders. It is widely recognized and accepted in all fields of medicine that a pain level below 3 is considered acceptable. Therefore, the average pain level reported by the patients in both groups in the present study can be interpreted as mild pain, an ache by definition (11, 13).

All our patients showed good compliance with the dressing and use of the devices, with no nuisances reported in the provided sheets. Postoperative bruising and ecchymosis are expected in the early postoperative period and are usually minimized by the application of cold compresses for 48 hours (14). However, the PSWT device has replaced the use of cold application. Ice packs are difficult to manage in the postoperative period as they obstruct vision and cause cold-related headaches. The sensitivity of fat cells to low temperatures is also well known (15). In our experience, the PSWT device for postoperative management allowed excellent control of pain and edema and improved wound healing, confirming the initial experience reported by Nicolle (16) and further supported by additional publications (17). Clearly, the addition of fat and stem cells provides a benefit in healing, as evidenced by the literature and patient postoperative courses. **Postoperative** management was the same for both groups, and patients in this study demonstrated compliance. Previously, ice packs were used, but they were difficult for the patient to manage, so the PSWT device was used to decrease pain and inflammation. In this surgeon's experience, the PSWT has allowed for excellent postoperative pain control, decreased edema, and improved healing (18, 19). The current results are in agreement with the scientific literature. A recently

published review with meta-analysis reveals that overall patient satisfaction is relatively high, ranging from 86.4% to 94.0%. Most complications were minor and could be easily treated or may disappear spontaneously. There was no high rate of severe complications (20). The level of evidence exploring these surgical techniques is still too low to provide a better understanding of the pros and cons of different surgical choices. Observational cohort studies comparing pain and complications outcomes in larger cohorts and with longer follow-up would enrich the current body of knowledge (21). This study did not demonstrate a significant increase in pain or complications for the patient; however, it did increase overall satisfaction and aesthetics. The authors are planning a long-term followup for the included patients to provide more insights on lipofilling integrated with blepharoplasty in the long term. Because of these results, a paradigm shift occurred during the 2002-2021 period, integrating lipofilling with all blepharoplasty surgeries. It is important to inform patients that even though the rejuvenation effects of fat grafting or lipofilling persist for long periods, they are not permanent, and the aging process will continue. Therefore, patients should understand their options regarding future modalities for the correction of rhytids (22, 23). Fat banks in the future could allow subsequent correction as aging and volume loss continue.

Footnotes

Authors' Contribution: Study concept and design: DB and SD, acquisition of data: .DB and SD, analysis and interpretation of data: DB SD and MK, drafting of the manuscript: DB MK BB MTC FM and DP, critical revision of the manuscript for important intellectual content: .MK BB MTC FM and DP, statistical analysis: SD, administrative, technical, and material support: .SD, Study supervision: SD

Conflict of Interests Statement: All the authors declared they have nothing to disclose.

Data Availability: Data are available upon request.

Funding/Support: This study was run with independent resources and without any fundings.

Informed Consent: All the patients included in the present study signed an informed consent to participate and for publication.

References

- Trepsat F. Periorbital rejuvenation combining fat grafting and blepharoplasties. *Aesthetic Plast Surg.* 2003;27(4):243-53. [PubMed ID: 15058544]. https://doi.org/10.1007/s00266-003-2126-y.
- Marten T, Elyassnia D. Periorbital Fat Grafting: A New Paradigm for Rejuvenation of the Eyelids. *Facial Plast Surg Clin North Am.* 2021;29(2):243-73. [PubMed ID: 33906759]. https://doi.org/10.1016/j.fsc.2021.02.003.
- Cohen SR, Womack H, Ghanem A. Fat Grafting for Facial Rejuvenation through Injectable Tissue Replacement and Regeneration: A Differential, Standardized, Anatomic Approach. *Clin Plast Surg.* 2020;47(1):31-41. [PubMed ID: 31739895]. https://doi.org/10.1016/j.cps.2019.08.005.
- Crowley JS, Kream E, Fabi S, Cohen SR. Facial Rejuvenation With Fat Grafting and Fillers. *Aesthet Surg J.* 2021;41(Suppl 1):S31-8. [PubMed ID: 34002771]. https://doi.org/10.1093/asj/sjab014.
- Moore JJ, Kolaczynski JW, Morales LM, Considine RV, Pietrzkowski Z, Noto PF, et al. Viability of fat obtained by syringe suction lipectomy: effects of local anesthesia with lidocaine. *Aesthetic Plast Surg.* 1995;**19**(4):335-9. [PubMed ID: 7484471]. https://doi.org/10.1007/BF00451659.
- Coleman SR. Facial recontouring with lipostructure. *Clin Plast Surg.* 1997;**24**(2):347-67. [PubMed ID: 9142473].
- Azoury SC, Shakir S, Bucky LP, Percec I. Modern Fat Grafting Techniques to the Face and Neck. *Plast Reconstr Surg.* 2021;**148**(4):620-33. [PubMed ID: 34550946]. https://doi.org/10.1097/PRS.00000000008405.
- Tonnard P, Verpaele A, Peeters G, Hamdi M, Cornelissen M, Declercq H. Nanofat grafting: basic research and clinical applications. *Plast Reconstr Surg.* 2013;132(4):1017-26. [PubMed ID: 23783059]. https://doi.org/10.1097/PRS.0b013e31829fe1b0.
- Ding P, Lu E, Li G, Sun Y, Yang W, Zhao Z. Research Progress on Preparation, Mechanism, and Clinical Application of Nanofat. *J Burn Care Res.* 2022;43(5):1140-4. [PubMed ID: 35015870]. [PubMed Central ID: PMC9435497]. https://doi.org/10.1093/jbcr/irab250.
- Deliaert AE, van den Elzen ME, van den Kerckhove E, Fieuws S, van der Hulst RR. Smoking in relation to age in aesthetic facial surgery. *Aesthetic Plast Surg.* 2012;36(4):853-6. [PubMed ID: 22711217]. [PubMed Central ID: PMC3404290]. https://doi.org/10.1007/s00266-012-9913-2.
- Tashjian RZ, Deloach J, Porucznik CA, Powell AP. Minimal clinically important differences (MCID) and patient acceptable symptomatic state (PASS) for visual analog scales (VAS) measuring pain in patients treated for rotator cuff disease. J Shoulder Elbow Surg. 2009;18(6):927-32. [PubMed ID: 19535272]. https://doi.org/10.1016/j.jse.2009.03.021.
- Mack WP. Blepharoplasty complications. Facial Plast Surg. 2012;28(3):273-87. [PubMed ID: 22723228]. https://doi.org/10.1055/s-0032-1312705.
- Kelly AM. The minimum clinically significant difference in visual analogue scale pain score does not differ with severity of pain. *Emerg Med J.* 2001;18(3):205-7. [PubMed ID: 11354213]. [PubMed Central ID: PMC1725574]. https://doi.org/10.1136/emj.18.3.205.
- Pool SMW, van Exsel DCE, Melenhorst W, Cromheecke M, van der Lei B. The effect of eyelid cooling on pain, edema, erythema, and hematoma after upper blepharoplasty: a randomized, controlled, observer-blinded evaluation study. *Plast Reconstr Surg.* 2015;**135**(2):277-81. [PubMed ID: 25626811]. https://doi.org/10.1097/PRS.000000000000919.

- Svalgaard JD, Juul S, Vester-Glovinski PV, Haastrup EK, Ballesteros OR, Lynggaard CD, et al. Lipoaspirate Storage Time and Temperature: Effects on Stromal Vascular Fraction Quality and Cell Composition. *Cells Tissues Organs*. 2020;209(1):54-63. [PubMed ID: 32580198]. https://doi.org/10.1159/000507825.
- Nicolle FV, Bentall RM. Use of radio-frequency pulsed energy in the control of postoperative reaction in blepharoplasty. *Aesthetic Plast Surg.* 1982;6(3):169-71. [PubMed ID: 7180721]. https://doi.org/10.1007/BF01570638.
- Bollero D. Jawline reshape technique with suspension threads to improve the effect of blepharoplasty on face rejuvenation. *J Cosmet Dermatol.* 2021;20 Suppl 2:12-7. [PubMed ID: 34318987]. https://doi.org/10.1111/jocd.14283.
- Lambros V. Observations on periorbital and midface aging. *Plast Reconstr Surg.* 2007;**120**(5):1367-76. [PubMed ID: 17898614]. https://doi.org/10.1097/01.prs.0000279348.09156.c3.
- Kahn DM, Shaw RJ. Aging of the bony orbit: a three-dimensional computed tomographic study. *Aesthet Surg J.* 2008;**28**(3):258-64. [PubMed ID: 19083535]. https://doi.org/10.1016/j.asj.2008.02.007.

- Yang F, Ji Z, Peng L, Fu T, Liu K, Dou W, et al. Efficacy, safety and complications of autologous fat grafting to the eyelids and periorbital area: A systematic review and meta-analysis. *PLoS One.* 2021;**16**(4). e0248505. [PubMed ID: 33793573]. [PubMed Central ID: PMC8016360]. https://doi.org/10.1371/journal.pone.0248505.
- Boureaux E, Chaput B, Bannani S, Herlin C, De Runz A, Carloni R, et al. Eyelid fat grafting: Indications, operative technique and complications; a systematic review. *J Craniomaxillofac Surg.* 2016;44(4):374-80. [PubMed ID: 26880013]. https://doi.org/10.1016/j.jcms.2015.12.013.
- Mendelson B, Wong CH. Changes in the facial skeleton with aging: implications and clinical applications in facial rejuvenation. *Aesthetic Plast Surg.* 2012;36(4):753-60. [PubMed ID: 22580543]. [PubMed Central ID: PMC3404279]. https://doi.org/10.1007/s00266-012-9904-3.
- 23. Rohrich RJ, Pessa JE. The fat compartments of the face: anatomy and clinical implications for cosmetic surgery. *Plast Reconstr Surg.* 2007;**119**(7):2219-27. [PubMed ID: 17519724]. https://doi.org/10.1097/01.prs.0000265403.66886.54.