



Antibacterial and Wound Healing Activity of *Dipterocarpus alatus* Crude Extract Against Methicillin-Resistant *Staphylococcus aureus*-Induced Superficial Skin Infection in Mice

Waranya Chatuphonprasert¹, Nitima Tatiya-aphiradee², Sutthiwan Thammawat¹, Chawalit Yongram³, Ploenthip Puthongking³ and Kanokwan Jarukamjorn^{2,*}

¹Faculty of Medicine, Mahasarakham University, Maha Sarakham, Thailand

²Research Group for Pharmaceutical Activities of Natural Products Using Pharmaceutical Biotechnology (PANPB), Khon Kaen University, Khon Kaen, Thailand

³Faculty of Pharmaceutical Sciences, Khon Kaen University, Khon Kaen, Thailand

*Corresponding author: Faculty of Pharmaceutical Sciences, Khon Kaen University, 123 Mitrapaab Rd., Khon Kaen 40002, Thailand. Tel/Fax: +66-043202379, +66-043202305, Email: kanok_ja@kku.ac.th

Received 2019 November 14; Revised 2019 December 16; Accepted 2019 December 18.

Abstract

Background: The bark oil of *Dipterocarpus alatus* Roxb. ex G. Don (resin tree) has been used for the treatment of ulcerated wounds in Ayurvedic medicine. However, information is still scarce about *D. alatus* extract and its constituents.

Objectives: This study aimed to examine the antibacterial and wound healing effects of *D. alatus* crude extracts against methicillin-resistant *Staphylococcus aureus* (MRSA) in a superficial skin infection using a tape-stripping mouse model.

Methods: Different parts (bark, leaves, twig, wood, and oleo-resin) of *D. alatus* were used for extraction. Crude extracts with two pure compounds (α,β -gurjunene and dipterocarpol) were examined for antibacterial activity against methicillin-susceptible *S. aureus* (MSSA) and MRSA *in vitro*. The *in vivo* antibacterial activity against MRSA was examined in a tape-stripping mouse model of superficial skin infection by evaluating transepidermal water loss (TEWL), colony counts from wound culture swabs, and H&E staining.

Results: *Dipterocarpus alatus* twig extract showed the lowest MIC (250 $\mu\text{g}/\text{mL}$) and MBC (500 $\mu\text{g}/\text{mL}$) against MRSA among extracts from other *D. alatus* parts. In addition, α,β -gurjunene showed lower MIC (250 $\mu\text{g}/\text{mL}$) and MBC (250 $\mu\text{g}/\text{mL}$) against MRSA than dipterocarpol. *In vivo*, *D. alatus* twig extract significantly reduced the number of MRSA recovered from superficial wounds to levels comparable to the non-infected control group on the second day. Oleo-resin, *D. alatus* wood extract, α,β -gurjunene, and dipterocarpol also significantly reduced the number of MRSA. Moreover, *D. alatus* twig extract and dipterocarpol healed infected wounds at rates comparable to the non-infected control group. From H&E staining of wounds, fibrin accumulation and neutrophil infiltration were attenuated in mice treated with *D. alatus* twig extract and dipterocarpol.

Conclusions: *Dipterocarpus alatus* twig extract and dipterocarpol possess antibacterial and wound healing properties against MRSA-induced superficial skin infection in mice.

Keywords: Dipterocarpol, Oleo-resin, Resin Tree, MRSA, Wound, Antibacterial

1. Background

Staphylococcus aureus is a Gram-positive pathogen of the skin, soft tissues, and septic infections (1). The first-line drugs for *S. aureus* treatment are β -lactam antibiotics, such as penicillin and cloxacillin, but the incidence of β -lactam-resistant *S. aureus* strains (methicillin-resistant *S. aureus*, MRSA) is increasing (2). Hence, new effective antibacterial agents against MRSA are required.

Traditionally, the oleo-resin of *Dipterocarpus alatus* Roxb. ex G. Don was used for illumination and waterproofing baskets and boats (3). *Dipterocarpus alatus* bark oil has been applied in Ayurvedic medicine for external

wound healing (4). Most of the bioactive chemical constituents isolated from genus *Dipterocarpus* are from the bark, which is an important source of resins such as oleoresin, dammars, and camphor. The main chemical constituents are sesquiterpenes, triterpenes, and coumarin derivatives (5). A previous study showed that the oleo-resin of *D. alatus* was an effective treatment for skin ulcers (6). Moreover, crude extracts from several parts of *D. alatus* have been reported to have free radical scavenging and anti-cancer activities (7). Besides, an ethanolic leaf extract of *D. alatus* was shown to inhibit monoamine oxidase and reduce the symptoms of depression in unpredictable

chronic mild stress in mice (8). To date, there is no information about the antimicrobial effects of *D. alatus* extracts.

2. Objectives

We aimed to investigate the antibacterial and wound healing effects of *D. alatus* crude extracts in an MRSA-induced superficial skin infection mouse model.

3. Methods

3.1. Materials and Chemicals

Two strains of *S. aureus*, methicillin-susceptible *S. aureus* (MSSA, DMST-2933) and methicillin-resistant *S. aureus* (MRSA, DMST-20651), were identified and provided by the Department of Medical Sciences, Thailand Biodiversity Center, National Center for Genetic Engineering and Biotechnology, Nonthaburi, Thailand. Mannitol salt agar (MSA) and Mueller Hinton broth and agar (MHB and MHA) were prepared from Himedia (Mumbai, India). Mayer's Hematoxylin, Eosin Y, and other staining reagents were the products of Bio Optica (Milano, Italy). Oxacillin, erythromycin, and tetracycline were supplied by Sigma-Aldrich Chemical (St. Louis, MO). Other chemicals and materials were from suppliers with premium quality.

3.2. Extraction of *Dipterocarpus alatus*

Dipterocarpus alatus crude extracts were prepared from different parts, namely bark, leaves, twig, and wood. Dipterocarpol and α,β -gurjunene were isolated from the oleo-resin extract as described elsewhere (6). Briefly, bark, leaves, twig, and wood were macerated with methanol. After evaporation to eliminate methanol, the crude extracts were obtained in the yields of 4.24, 11.80, 6.07, and 6.01%, respectively. Oleo-resin was heated at 40 - 60°C to give clear oil for use in this study. Oleo-resin (25.04 g) was isolated using the open column chromatographic technique to achieve the two known compounds, dipterocarpol (1.01 g) and α,β -gurjunene (4.76 g).

3.3. Determination of Minimum Inhibitory Concentration and Minimum Bactericidal Concentration In Vitro

The minimum inhibitory concentration (MIC) was evaluated by the broth dilution method in MHB as described elsewhere (9). An inoculum of MSSA or MRSA (2×10^6 CFU/mL) was mixed with *D. alatus* extracts, α,β -gurjunene, dipterocarpol, oxacillin, erythromycin, and tetracycline, and incubated at 37°C for 24 hours. The growth of bacteria was determined at 625 nm by a spectrophotometer. The MIC value was identified from the well in which MSSA or MRSA growth was inhibited.

For minimum bactericidal concentration (MBC), all wells with no visible growth were cultured on MHA plates by simple streaking and the plates were incubated at 37°C for 18 - 24 hours. The MBC was determined from the lowest concentration of the test compound without colonies (9).

3.4. Tape-Stripping Mouse Model

Seven-week-old male ICR mice (30 - 35 g) were bred and provided by the National Laboratory Animal Center, Mahidol University, Thailand. All mice were housed in the Northeast Laboratory Animal Center, Khon Kaen University, Thailand. Animal-related protocols were adopted from the Animal Ethics Committee for Use and Care of Khon Kaen University (approval no. ACUC-KKU-38/2559) in agreement with the Guide for the Care and Use of Laboratory Animals prepared by the National Research Council of Thailand. After transportation, mice were acclimated for a week before being randomly divided into eight groups (n = 10 per group). The mice were anesthetized by intraperitoneal injection of pentobarbital sodium 50 μ g/kg and the back hair was shaved. An adhesive plaster was quickly applied to and immediately removed from a 2×2 cm² dorsal area for 15 - 20 times until the skin became swollen and ruddy without bleeding. The transepidermal water loss (TEWL) value was measured by DermaLab[®] Combo (Hadsund, Denmark) and standardized at the level of 70 - 75 g/m²h to normalize the skin damage of all mice. After that, a 10 μ L-MRSA inoculum (1×10^8 CFU/mL) was applied to wounds (9). A 100- μ L aliquot of each sample, i.e., 10% ethanol in propylene glycol (vehicle), 20 mg/mL of *D. alatus* twig/oleoresin/wood extract, 4 mg/mL of α,β -gurjunene, 1.1 mg/mL of dipterocarpol, or 160 μ g/mL of tetracycline, was applied to the wound area daily for nine consecutive days. The non-infected control group and non-treated control group (MRSA-infected group without treatment) were treated with a vehicle. The TEWL was determined on days 1, 2, 5, and 7. Swabs from infected-wounds were cultured on 6 μ g/mL oxacillin-contained MSA plates on days 2, 4, 6, and 8. The number of MRSA colonies was recorded after incubation for 18 - 24 hours (10). Wounds were photographed on days 1, 3, 5, 7, and 9.

3.5. Skin Staining by Hematoxylin and Eosin

Mice from each treatment group were sacrificed on days 1, 4, and 8. The skin from the dorsal area was collected and washed in iced isotonic phosphate-buffered saline for 30 minutes before soaking in a 10% formalin solution at 4°C for 24 hours. The skin sample was gradually dehydrated in ethanol (50 to 100% v/v) at room temperature, embedded in warm paraffin (60°C), and frozen at -20°C. The paraffin-embedded skin was cut into 5 μ m sections

and fixed on a glass slide before staining with hematoxylin and eosin (H&E) solution following the instructions of the supplier (10).

3.6. Statistical Analysis

The TEWL values and the number of MRSA colonies are presented as mean \pm SD. Statistical significance was analyzed by one-way ANOVA followed by Tukey's test using SPSS V. 22.0 program. A P value of < 0.05 was considered significant.

4. Results

4.1. Effects of *Dipterocarpus alatus* Extracts on MSSA and MRSA In Vitro

Table 1 shows the MIC and MBC of *D. alatus* extracts, α , β -gurjunene, and dipterocarpol, compared to three antibiotics, oxacillin, erythromycin, and tetracycline. Tetracycline showed the most anti-MSSA and MRSA activity, while erythromycin and oxacillin did not inhibit MRSA growth. Almost all *D. alatus* extracts showed less antibacterial activity than tetracycline, with MICs and MBCs of 500 μ g/mL or higher. The *D. alatus* twig extract had the lowest MIC and MBC against MRSA at 250 and 500 μ g/mL, respectively. Moreover, α , β -gurjunene showed lower MIC and MBC (250 μ g/mL) against MRSA than dipterocarpol. Thus, twig and wood extracts, α , β -gurjunene, and tetracycline (a positive control) were selected for *in vivo* experiments in the mouse superficial wound model. In addition, we studied the oleo-resin extract, which has been claimed to heal ulcerated wounds in Ayurvedic medicine (4), and its principal component dipterocarpol (6).

4.2. Antibacterial Effects of *Dipterocarpus alatus* Extracts on MRSA-Induced Superficial Skin Infection in Mice

Mice with MRSA-superficial skin infections had very high numbers of MRSA colonies on the second day of infection. Of these, the MRSA-non-treated and MRSA-tetracycline-treated groups showed higher numbers of MRSA colonies than the others (Table 2). The *D. alatus* twig extract significantly decreased the number of MRSA colonies on the second day of treatment to the same number of MRSA colonies found in the non-infected control. The extracts of oleo-resin and wood, α , β -gurjunene, and dipterocarpol significantly decreased the number of MRSA colonies compared to the MRSA-infected group without treatment. Therefore, the *D. alatus* extracts, α , β -gurjunene, and dipterocarpol possessed antibacterial activity against MRSA in this model.

Table 1. Minimum Inhibitory Concentration (MIC) and Minimum Bactericidal Concentration (MBC) of *Dipterocarpus alatus* Crude Extracts Against Methicillin-Susceptible *Staphylococcus aureus* (MSSA) and Methicillin-Resistant *Staphylococcus aureus* (MRSA)

Test Compounds	MIC (μ g/mL)	MBC (μ g/mL)
Bark crude extract		
MSSA	500	1000
MRSA	500	1000
Leaves crude extract		
MSSA	500	500
MRSA	500	1000
Twig crude extract		
MSSA	250	500
MRSA	250	500
Oleo-resin crude extract		
MSSA	> 1000	> 1000
MRSA	> 1000	> 1000
Wood crude extract		
MSSA	500	500
MRSA	500	500
Tetracycline		
MSSA	≤ 0.0625	2
MRSA	4	16
Erythromycin		
MSSA	≤ 0.0625	0.5
MRSA	≤ 64	> 64
Oxacillin		
MSSA	≤ 0.0625	2
MRSA	> 64	> 64
Dipterocarpol		
MSSA	500	500
MRSA	500	1000
α,β-Gurjunene		
MSSA	500	500
MRSA	250	250

4.3. Wound Healing Effects of *Dipterocarpus alatus* Extracts on MRSA-Induced Superficial Skin Infection in the Mouse Model

The daily examination revealed that the wounds of non-infected control mice completely healed nine days after tape stripping (Figure 1) while MRSA-infected non-treated mice were found to be ulcerated until the ninth day and remained infected according to the MRSA colony counts (Table 2). The *D. alatus* twig extract and dipterocarpol were effective in wound healing at a rate comparable to control mice. The oleo-resin extract demonstrated some wound healing effects though it was inferior to the twig extract and dipterocarpol. The wounds of the groups treated with wood extract, α , β -gurjunene, and tetracycline remained scabby on the ninth day. The TEWL indicated the permeability of the skin layer by mea-

Table 2. Numbers of MRSA Colonies Recovered from Infected Superficial Mouse Wounds

	Number of MRSA Colonies ^a			
	Day 2	Day 4	Day 6	Day 8
Control	0.00	0.00	0.00	0.00
MRSA-non-treatment	86.90 ± 79.79 ^b	14.63 ± 13.34 ^b	10.83 ± 12.25 ^b	9.75 ± 11.06 ^b
MRSA-twig	1.10 ± 1.29 ^c	0.00 ± 0.00 ^c	0.00 ± 0.00 ^c	0.00 ± 0.00 ^c
MRSA-oleo-resin	12.60 ± 20.13 ^{b, c}	3.88 ± 4.32 ^c	0.50 ± 0.84 ^c	0.00 ± 0.00 ^c
MRSA-wood	14.80 ± 14.02 ^{b, c}	0.25 ± 0.46 ^c	0.17 ± 0.41 ^c	0.00 ± 0.00 ^c
MRSA- α, β -gurjunene	27.10 ± 23.19 ^{b, c}	2.50 ± 3.12 ^c	2.83 ± 2.79 ^c	0.25 ± 0.50 ^c
MRSA-dipterocarpol	21.50 ± 27.23 ^{b, c}	0.25 ± 0.46 ^c	1.00 ± 1.26 ^c	0.00 ± 0.00 ^c
MRSA-tetracycline	131.70 ± 167.21 ^b	11.13 ± 10.34 ^b	24.00 ± 31.59 ^b	12.00 ± 18.74 ^b

^aValues are expressed as mean ± SD.

^bP < 0.05 vs. control on the same day.

^cP < 0.05 vs. MRSA-non-treatment on the same day.

asuring the rate of water loss. The TEWL value was the highest after the epidermal barrier removal by tape stripping. In the non-infected control group, TEWL recovered to pre-wound levels within seven days (Figure 2). The *D. alatus* twig extract and dipterocarpol treatments gradually decreased the TEWL to the same level as the non-infected control group, which was significantly lower than the MRSA-non-treated group. The improvement in wound appearance and a decrease in TEWL indicated that the *D. alatus* twig extract and dipterocarpol possessed wound healing properties.

4.4. Effects of *Dipterocarpus alatus* Extracts and Dipterocarpol on Wound Histology

The epidermis (E) of all mice was partially disrupted by tape stripping (Figure 3). The H&E staining showed fibrin accumulation (black arrows) on the fourth day. The *D. alatus* twig extract and dipterocarpol reduced fibrin accumulation comparable to the non-infected control. Neutrophil infiltration or accumulation of neutrophils (Figure 3, squares) was observed in the upper dermis of all mice from the first day. On the fourth day, the *D. alatus* twig extract and dipterocarpol were regenerating the epidermis and healing the wounds; the dermal-epidermal junction was clearly differentiated. In contrast, neutrophils remained in the MRSA-non-treated group, as well as α, β -gurjunene and tetracycline-treated groups. Therefore, the *D. alatus* twig extract and dipterocarpol extensively restored the integrity of the epidermis in MRSA-induced superficial skin infection.

5. Discussion

The number of skin and soft tissue infections commonly caused by *S. aureus* and community-acquired MRSA increased between 1998 to 2004 in North America (11). The frequency of *S. aureus*-induced skin infection was 44.6% and the rate of MRSA among isolates was 35.9% (12). Hence, there is a need for an effective compound for the treatment of MRSA infection. To date, there are limited reports on the antibacterial effects of *Dipterocarpus* species and dipterocarpol. *Dipterocarpus verrucosus* has been reported to have *in vitro* antibacterial effects (5). The oleo-resin of *Shorea robusta* Gaertn (containing dipterocarpol) is an antibacterial, astringent, analgesic, and wound healing agent (13). Here, we described the first experiments demonstrating the activity of *D. alatus* extracts and dipterocarpol against *S. aureus* both in *in vitro* and *in vivo*.

The MRSA colony counting from mouse wounds indicated that the number of MRSA colonies gradually reduced without treatment (Table 2); however, all *D. alatus* extracts accelerated the decrease in the colony number from the first day of treatment. The comparatively reduced antibacterial properties of *D. alatus* extracts and dipterocarpol *in vitro* may be due to their low water solubility (9). One study reported that the rate of antibacterial susceptibility to tetracycline in MRSA isolates was about 82% (11), which corresponded to our *in vitro* results. Hence, tetracycline was applied as a positive control in the mouse model. However, tetracycline showed low antibacterial activity against MRSA in this model, presumably due to its low permeability and inability to cross the skin membrane layer.

The tape stripping technique in combination with MRSA infection is a well-known mouse model for studying antibacterial and wound healing effects, representing




Figure 1. Mouse superficial wound appearance. Mouse superficial wounds were established by repeated tape stripping to remove the epidermis. Wounds were infected with 1×10^8 MRSA and treated daily with 10% ethanol in propylene glycol (non-treated control), 20 mg/mL *Dipterocarpus alatus* twig/oleo-resin/wood extract in 10% ethanol in propylene glycol, 4 mg/mL α,β -gurjunene, 1.1 mg/mL dipterocarpol, or 160 μ g/mL tetracycline for nine days along with a non-infected control group (eight groups, 9-10 mice in each group). The wounds of all mice were photographed on days 1, 3, 5, 7, and 9.




Figure 2. Transepidermal water loss (TEWL) of mouse superficial wounds. Mouse superficial wounds were established by repeated tape stripping to remove the epidermis. Wounds were infected with 1×10^8 MRSA and treated daily with 10% ethanol in propylene glycol (non-treated control), 20 mg/mL *Dipterocarpus alatus* twig/oleo-resin/wood extract in 10% ethanol in propylene glycol, 4 mg/mL α,β -gurjunene, 1.1 mg/mL dipterocarpol, or 160 μ g/mL tetracycline for nine days along with a non-infected control group (eight groups, 9–10 mice in each group). The TEWL from the superficial wounds was measured on days 1, 2, 5, and 7 and presented as mean \pm SD. # $p < 0.05$ vs. MRSA-non-treatment on the same day by one-way ANOVA followed by Tukey's post hoc test.

an abrasive wound in *S. aureus* skin-infected patients (10, 14). The TEWL indicates skin barrier damage (10) and it was used to normalize the same level of skin barrier damage in all mice on the first day of study. After the tape stripping, the application of MRSA inoculum to an abrasive wound allowed MRSA to grow, permeate the wound, and stimulate the systematic inflammatory response. Fibrin deposition in the skin is an eminent and compatible marker for type IV hypersensitivity skin reactions, contact dermatitis, and MRSA-infected skin (10, 15). Neutrophil infiltration is a process required for an effective immune response against *S. aureus* (16). Our observations showed both fibrin deposition and neutrophil infiltration after tape stripping and MRSA infection. Treatment with *D. alatus* twig extract and dipterocarpol completely reversed these conditions, comparable to the non-infection control group. The MRSA colony counting, TEWL, wound appearance, and H&E staining clearly indicated that the *D. alatus* twig extract and dipterocarpol had anti-MRSA activity and wound healing properties that would be suitable for further development as a topical formulation for superficial skin infection.

5.1. Conclusions

The *D. alatus* twig extract and dipterocarpol had antibacterial effects against MSSA and MRSA *in vitro*. The *D. alatus* twig extract and dipterocarpol possessed wound healing and antibacterial activities shown by decreasing the number of MRSA colonies, reducing fibrin accumulation and neutrophil infiltration, and restoring TEWL to pre-

wound levels. Therefore, *D. alatus* twig extract and dipterocarpol are candidates for MRSA treatment, particularly in superficial skin infection.

Footnotes

Authors' Contribution: Waranya Chatuphonprasert designed the study, conducted the experiments, analyzed and interpreted the data, conducted the statistical analysis, and wrote the manuscript. Nitima Tatiya-aphiradee took care of animal models and collected data from mice. Sutthiwan Thammawat collected and interpreted data from *in vitro* experiments. Chawalit Yonggram and Ploenthip Puthongking prepared plant extracts and pure compounds. Kanokwan Jarukamjorn contributed to the development of the protocol, abstracted the data, and prepared the manuscript.

Conflict of Interests: The authors declare that there is no conflict of interest in this study.

Ethical Approval: The research protocol was approved by the Animal Ethics Committee for Use and Care of Khon Kaen University, Thailand (approval no. ACUC-KKU-38/2559) in accordance with the Guide for the Care and Use of Laboratory Animals prepared by the National Academy of Sciences and published by the National Institutes of Health.

Funding/Support: The Plant Genetic Conservation Project under the royal initiative of Her Royal Highness Princess




Figure 3. Histology of wounds with H&E staining. Mice were sacrificed on days 1, 4, and 8 and tissues were cut into 5- μ m sections, stained with H&E, and mounted. E indicates epidermis, black arrows indicate fibrin accumulation, and squares indicate areas with neutrophil infiltration. The tissues were photographed at 400x magnification.

Maha Chakri Sirindhorn, Research Group for Pharmaceutical Activities of Natural Products using Pharmaceutical Biotechnology (PANPB), Faculty of Pharmaceutical Sciences, Khon Kaen University, and Faculty of Medicine, Mahasarakham University, Thailand, are acknowledged for financial support and facilities. The authors thank Dr. Glenn Borlace for English assistance.

References

1. Daum RS. Clinical practice. Skin and soft-tissue infections caused by methicillin-resistant *Staphylococcus aureus*. *N Engl J Med*. 2007;**357**(4):380–90. doi: [10.1056/NEJMc070747](https://doi.org/10.1056/NEJMc070747). [PubMed: [17652653](https://pubmed.ncbi.nlm.nih.gov/17652653/)].
2. Haddadin AS, Fappiano SA, Lipsett PA. Methicillin resistant *Staphylococcus aureus* (MRSA) in the intensive care unit. *Postgrad Med J*. 2002;**78**(921):385–92. doi: [10.1136/pmj.78.921.385](https://doi.org/10.1136/pmj.78.921.385). [PubMed: [12151652](https://pubmed.ncbi.nlm.nih.gov/12151652/)]. [PubMed Central: [PMC1742438](https://pubmed.ncbi.nlm.nih.gov/PMC1742438/)].
3. Ankarjard R, Kegl M. Tapping oleoresin from *Dipterocarpus alatus* (Dipterocarpaceae) in a Lao village. *Econ Bot*. 1998;**52**(1):7–14. doi: [10.1007/bf02861291](https://doi.org/10.1007/bf02861291).
4. Karnick CR, Hocking GM. Ethnobotanical records of drug plants described in Valmiki Ramayana and their uses in the Ayurvedic system of medicine. *Q J Crude Drug Res*. 2010;**13**(3-4):143–54. doi: [10.1080/13880207509162671](https://doi.org/10.1080/13880207509162671).
5. Aslam MS, Ahmad MS, Mamat AS. A phytochemical, ethnomedicinal and pharmacological review of genus *dipterocarpus*. *Int J Pharm Pharmaceut Sci*. 2015;**7**(4):27–38.
6. Chandra PPR. Ecological analysis of Dipterocarpaceae of North Andaman Forest, India. *J Plant Dev*. 2011;**18**:135–49.
7. Yongram C, Sungthong B, Puthongking P, Weerapreeyakul N. Chemical composition, antioxidant and cytotoxicity activities of leaves, bark, twigs and oleo-resin of *Dipterocarpus alatus*. *Molecules*. 2019;**24**(17). doi: [10.3390/molecules24173083](https://doi.org/10.3390/molecules24173083). [PubMed: [31450678](https://pubmed.ncbi.nlm.nih.gov/31450678/)]. [PubMed Central: [PMC6751497](https://pubmed.ncbi.nlm.nih.gov/PMC6751497/)].
8. Daodee S, Monthakantirat O, Ruengwinitwong K, Gatenakorn K, Maneenet J, Khamphukdee C, et al. Effects of the ethanol extract of *Dipterocarpus alatus* leaf on the unpredictable chronic mild stress-induced depression in ICR mice and its possible mechanism of action. *Molecules*. 2019;**24**(18). doi: [10.3390/molecules24183396](https://doi.org/10.3390/molecules24183396). [PubMed: [31540539](https://pubmed.ncbi.nlm.nih.gov/31540539/)]. [PubMed Central: [PMC6767234](https://pubmed.ncbi.nlm.nih.gov/PMC6767234/)].
9. Tatiya-aphiradee N, Chatuphonprasert W, Jarukamjorn K. In vivo antibacterial activity of *Garcinia mangostana* pericarp extract against methicillin-resistant *Staphylococcus aureus* in a mouse superficial skin infection model. *Pharm Biol*. 2016;**54**(11):2606–15. doi: [10.3109/13880209.2016.1172321](https://doi.org/10.3109/13880209.2016.1172321). [PubMed: [27180784](https://pubmed.ncbi.nlm.nih.gov/27180784/)].
10. Tatiya-aphiradee N, Chatuphonprasert W, Jarukamjorn K. Anti-inflammatory effect of *Garcinia mangostana* Linn. pericarp extract in methicillin-resistant *Staphylococcus aureus*-induced superficial skin infection in mice. *Biomed Pharmacother*. 2019;**111**:705–13. doi: [10.1016/j.biopha.2018.12.142](https://doi.org/10.1016/j.biopha.2018.12.142). [PubMed: [30611995](https://pubmed.ncbi.nlm.nih.gov/30611995/)].
11. Moran GJ, Amii RN, Abrahamian FM, Talan DA. Methicillin-resistant *Staphylococcus aureus* in community-acquired skin infections. *Emerg Infect Dis*. 2005;**11**(6):928–30. doi: [10.3201/eid1106.040641](https://doi.org/10.3201/eid1106.040641). [PubMed: [15963289](https://pubmed.ncbi.nlm.nih.gov/15963289/)]. [PubMed Central: [PMC3367577](https://pubmed.ncbi.nlm.nih.gov/PMC3367577/)].
12. Moet GJ, Jones RN, Biedenbach DJ, Stilwell MG, Fritsche TR. Contemporary causes of skin and soft tissue infections in North America, Latin America, and Europe: Report from the SENTRY Antimicrobial Surveillance Program (1998–2004). *Diagn Microbiol Infect Dis*. 2007;**57**(1):7–13. doi: [10.1016/j.diagmicrobio.2006.05.009](https://doi.org/10.1016/j.diagmicrobio.2006.05.009). [PubMed: [17059876](https://pubmed.ncbi.nlm.nih.gov/17059876/)].
13. Poornima B. Comparative phytochemical analysis of *Shorea robusta* Gaertn (oleoresin) WSR to its seasonal collection. *Anc Sci Life*. 2009;**29**(1):26–8. [PubMed: [22557341](https://pubmed.ncbi.nlm.nih.gov/22557341/)]. [PubMed Central: [PMC3336301](https://pubmed.ncbi.nlm.nih.gov/PMC3336301/)].
14. Martinez LR, Han G, Chacko M, Mihu MR, Jacobson M, Gialanella P, et al. Antimicrobial and healing efficacy of sustained release nitric oxide nanoparticles against *Staphylococcus aureus* skin infection. *J Invest Dermatol*. 2009;**129**(10):2463–9. doi: [10.1038/jid.2009.95](https://doi.org/10.1038/jid.2009.95). [PubMed: [19387479](https://pubmed.ncbi.nlm.nih.gov/19387479/)].
15. Colvin RB, Johnson RA, Mihm MJ, Dvorak HF. Role of the clotting system in cell-mediated hypersensitivity. I. Fibrin deposition in delayed skin reactions in man. *J Exp Med*. 1973;**138**(3):686–98. doi: [10.1084/jem.138.3.686](https://doi.org/10.1084/jem.138.3.686). [PubMed: [4580465](https://pubmed.ncbi.nlm.nih.gov/4580465/)]. [PubMed Central: [PMC2139425](https://pubmed.ncbi.nlm.nih.gov/PMC2139425/)].
16. Molne L, Verdrengh M, Tarkowski A. Role of neutrophil leukocytes in cutaneous infection caused by *Staphylococcus aureus*. *Infect Immun*. 2000;**68**(11):6162–7. doi: [10.1128/jai.68.11.6162-6167.2000](https://doi.org/10.1128/jai.68.11.6162-6167.2000). [PubMed: [11035720](https://pubmed.ncbi.nlm.nih.gov/11035720/)]. [PubMed Central: [PMC97694](https://pubmed.ncbi.nlm.nih.gov/PMC97694/)].