



Antioxidant Mechanisms in Myocardial Ischemia: Narrative Review

Najmeh Sadeghi ¹, Hassanali Abedi ^{2,*}, Shekoufeh Atashpour ³ and Soodabeh Etemadi¹

¹Sirjan School of Medical Sciences, Sirjan, Iran

²Department of Physiology, Jahrom University of Medical Sciences, Jahrom, Iran

³Department of Pharmacology, Jahrom University of Medical Sciences, Jahrom, Iran

*Corresponding author: Department of Physiology, Jahrom University of Medical Sciences, Jahrom, Iran. Email: hassanaliabedi@gmail.com

Received 2022 October 08; Revised 2023 July 29; Accepted 2023 August 28.

Abstract

Ischemia is a common condition in the population and has become one of the leading causes of mortality in recent years. It occurs due to a shortage of oxygen in the cell caused by inadequate blood flow, leading to the formation of free radicals. These free radicals can bind to the phospholipids in the cell membrane, resulting in proxy radicals and lipid hydroperoxidation. Malondialdehyde (MDA) is a cytotoxic byproduct of lipid peroxidation that causes cell damage and necrosis by rupturing the cell membrane. Subsequently, mitochondrial dysfunction related to fatty acid oxidation can develop, further exacerbating the damage caused by ischemia. Reactive oxygen species (ROS) produced in mitochondria have been shown to elevate oxidative phosphorylation disorder and disrupt the functioning of complexes I and IV. These enzyme activities can be restored to normal levels by activating antioxidant enzymes such as superoxide dismutase (Mn-SOD) and catalase, thereby strengthening the antioxidant defense system. Various antioxidants have been found to have protective effects. In this review, we compare the antioxidant effects of various plants and tannins that have been studied up to this point on myocardial ischemia. We also explain some of the cellular and molecular pathways that have been investigated to protect the myocardium against ischemia-reperfusion injury.

Keywords: Ischemia, Antioxidant, ROS (Reactive Oxygen Species), NO (Nitric Oxide), Cardioprotection, Tannins

1. Context

Pathologically, it can lead to cardiac myocyte death due to chronic ischemia. Reports indicate that coronary artery disease (CAD) causes approximately seven million deaths worldwide each year (1). Myocardial infarction (MI) imposes a substantial economic burden. Hospitalization costs alone in the United States amount to at least \$450 billion, and patients lose many years of healthy life (2).

To preserve myocardial health, it is crucial to restore blood flow quickly during ischemia (3). Reperfusion interventions such as primary percutaneous coronary intervention (PCI) and thrombolysis have been shown to reduce mortality and infarct size and improve left ventricular function. However, reperfusion can also lead to complications (4). Swift restoration of coronary flow can result in reversible myocardial contractility (myocardial stunning), ventricular arrhythmias, and microvascular dysfunction.

Myocardial ischemia/reperfusion (I/R) injury follows a pattern of damage, with the adverse effects of cardiac ischemia/reperfusion leading to myocyte necrosis and

heart failure (5).

One of the yet-to-be-discovered molecular mechanisms of IR damage in muscles is the production of reactive oxygen species (ROS). Excessive ROS accumulation activates the mitochondrial permeability transition pore, causing cellular oxidative stress, mitochondrial dysfunction, and cell death. Elevated ROS levels may reduce myofibrillar Ca²⁺ sensitivity, impairing muscle contractile function. Additionally, modest ROS levels can act as signaling molecules in protective ischemia preconditioning (IPC) pathways (6).

Antioxidant treatment has shown variable and inconsistent effects in experimental and clinical studies, highlighting the need for further research in this field. Cohort studies, however, consistently show a connection between antioxidants such as flavonoids and beta-carotene and reduced cardiovascular disease (CVD)-related mortality (1).

2. Evidence Acquisition

Novel cardioprotective strategies for enhancing myocardial salvage and cardiac function are of paramount importance. Recently, plant-based medicines with antioxidant properties, such as grape seed extract and its derivatives like epigallocatechin or Gallic acid, have gained attention (7). Consequently, we have embarked on investigating their molecular mechanisms as a therapeutic alternative to conventional pharmacological approaches, which often come with numerous side effects (8).

3. Results

3.1. Inorganic Nitrate or Nitrite Therapy

Inorganic nitrate or nitrite therapy has shown significant reductions in oxidative stress, heart hypertrophy, fibrosis, and blood pressure levels while also improving renal function. In unilaterally nephrectomized rats, treatment with a high dose of sodium nitrate (140 - 1400 mg/kg/day) resulted in decreased mean arterial pressure, cardiac hypertrophy/fibrosis, plasma malondialdehyde, isoprostane, and 8-oxo-2-deoxyguanosine levels, along with improved kidney morphology (9).

Using a lower dosage of sodium nitrite (15 mg/kg/day), spontaneously hypertensive rats exhibited lower blood pressure and improved endothelial function. These blood vessels produced more nitric oxide (NO), leading to increased cGMP production and enhanced endothelial nitric oxide synthase (eNOS) activity (10). Sodium nitrite at a dose of 7.5 mg/kg/day significantly lowered systolic blood pressure and reduced oxidative stress in the vasculature and kidneys in angiotensin-II-infused mice, a model of arterial hypertension (11).

The same study demonstrated that sodium nitrite (15 mg/kg/day) diminished systolic blood pressure, malondialdehyde levels, and vascular dihydroethidium in rats treated with deoxycorticosterone (DOCA) and salt (12).

Using inorganic nitrite/nitrates has been found effective in reducing inflammation and inhibiting oxidation, including the upregulation of arginase and pathways dependent on cyclic guanosine monophosphate (cGMP)/cyclic adenosine monophosphate (cAMP) (13). For example, supplementation with inorganic nitrite/nitrate prevented microvascular inflammation in rats and mice treated with non-steroidal anti-inflammatory drugs (NSAIDs), leading to a reduction in tumor necrosis factor-alpha (TNF- α) expression and leukocyte recruitment (14). In rats and mice, inorganic nitrite/nitrate supplementation reduced the production of tumor

necrosis factor-alpha (TNF- α) and adhesion molecules, as well as leukocyte recruitment, which was enhanced by NSAID treatment (15). In lipopolysaccharide (LPS)-stimulated phagocytes (macrophages), nitrite directly inhibited NADPH oxidase, mediated by xanthine oxidoreductase-dependent bioactivation to nitric oxide (NO), leading to the downregulation of protein-level cytosolic NADPH oxidase subunits p47phox and p67phox but not mRNA (16).

In Apo E knockout animals fed a high-fat diet, inorganic nitrate reduced leukocyte rolling, recruitment to the vascular wall, and myeloperoxidase activity (17). In active neutrophils, inorganic nitrite inhibited the production of proxy nitrite and hypochlorite. However, the specific mechanism by which inorganic nitrite and nitrate inhibit immune cell activation remains unknown (18).

3.2. Cardioprotective cGMP-PKG Pathway

The cGMP-PKG pathway is initiated by nitric oxide (NO) and natriuretic peptides (NPs). Nitric oxide initiates this signaling pathway by activating soluble guanylate cyclase (sGC), a heterodimeric enzyme composed of α - and β -subunits with prosthetic heme groups. Atrial NP (ANP), brain NP (BNP), and C-type NP (CNP) activate particulate GC (pGC) in the plasma membrane to produce cGMP.

The second messenger, cGMP, is generated by both sGC and pGC. However, the downstream functions of cGMP vary depending on its subcellular location. High intracellular levels of cGMP primarily mediate physiological actions through protein kinase G (PKG), which is cGMP-dependent. PKG-I, the original kinase in animals, plays a crucial role in transmitting physiological signals to the cardiovascular system. Notably, PKG-I and PKG-1 differ in their N-termini due to alternative splicing. Research indicates that PKG-I is ten times more sensitive to cGMP than PKG-1. Additionally, PKG-I can be activated independently of cGMP through thiol oxidation, thanks to its unique cysteine residue (Cys 42). This cGMP-independent activation of PKG-1 can reduce blood pressure by promoting vasodilation, though its effects on cardiomyocytes are not fully understood.

Numerous studies have identified multiple downstream effectors of cGMP-PKG-I in the cardiovascular system. For instance, in cardiomyocytes, PKG-I phosphorylates phospholamban at Ser16, activating the sarcoplasmic reticulum Ca²⁺-ATPase (SERCA). This activation enhances Ca²⁺ uptake into the sarcoplasmic reticulum during systole, reducing cytosolic Ca²⁺ levels. Additionally, cGMP-induced PKG-I activation has been linked to the opening of mitochondrial ATP-sensitive K⁺ (mitoKATP) channels in the inner mitochondrial membrane, leading to an increase in

K⁺ influx and mitochondrial matrix alkalinization. This process boosts H₂O₂ production from complex I and subsequently activates protein kinase C (PKC), protecting cardiomyocytes against cell death by inhibiting the opening of mitochondrial permeability transition pores (mPTP).

Furthermore, the cardiac myocyte Ca²⁺-Activated BK type K⁺ channels (CMBK) play a crucial role in cardiac remodeling following myocardial infarction (MI). CMBK deficiency has been associated with more severe myocardial injury after ischemia/reperfusion (I/R), characterized by increased reactive oxygen species (ROS) production. In addition, angiotensin II (ATII)-induced cardiac hypertrophy, mediated by ROS and mitogen-activated protein kinase (MAPK) activation, was found to be mitigated by antioxidant treatment.

Changes in ion flow, mediated by ROS, have also been linked to the regulation of muscle contractility in cardiomyocytes. Increased ROS production reduces myofibrillar calcium (Ca²⁺) sensitivity and inhibits sarcoplasmic reticulum Ca²⁺ ATPase 2 activity (SERCA2), potentially disrupting excitation-contraction coupling and accelerating heart failure progression.

While the detailed mechanistic interactions between downstream effectors and the cGMP-PKG pathway in maintaining heart health require further characterization, numerous preclinical studies have demonstrated the cardio-protective potential of this pathway through pharmacological interventions and gene manipulations. Based on the available evidence, strategies involving either enhanced cGMP synthesis (e.g., sGC activators or stimulators) or reduced cGMP degradation (e.g., PDE inhibitors) represent promising therapeutic approaches for targeting this system.

3.3. Ginsenoside Rb1

Ginsenoside Rb1 (G-Rb1) is a component of ginseng, an herbal remedy with a centuries-old history of use worldwide, particularly in Asian countries. Ginseng contains various compounds known as ginsenosides, which are triterpene saponins. To date, researchers have identified over 30 distinct types of ginsenosides.

Ginsenoside Rb1 (G-Rb₁) exhibits numerous beneficial effects within the cardiovascular system, likely attributable to its antioxidant, anti-inflammatory, and anti-apoptotic properties. These attributes enhance its angiogenesis-promoting and antiarrhythmic effects. In cases of acute myocardial infarction (MI), G-Rb₁ can prevent ventricular hypertrophy, as illustrated in [Figure 1](#).

The following are potential mechanisms by which G-Rb1 may mitigate myocardial ischemia/reperfusion (I/R) injury: (1) Inhibition of apoptosis: G-Rb1 achieves

this by down-regulating the expression of caspase-3 (19); (2) inhibition of the p38MAPK signaling pathway (20); (3) enhancement of antioxidant defenses: G-Rb1 elevates antioxidant levels, including glutathione (GSH) and superoxide dismutase (SOD), reducing malondialdehyde (MDA) levels—a marker of oxidative stress. This process is likely linked to chondrogenesis attenuation (21); (4) suppression of inflammation: G-Rb1 inhibits the expression of tumor necrosis factor-alpha (TNF-α) by preventing the activation of the PI3K/Akt signaling pathway and interfering with p38MAPK signaling. This crosstalk between the two pathways contributes to its anti-inflammatory effects (21); (5) promotion of angiogenesis: G-Rb1 increases the expression of vascular endothelial growth factor (VEGF) and enhances circulation by elevating nitric oxide (NO) levels through the activation of the PI3K/Akt signaling pathway (22, 23).

3.4. Redox Mechanisms of IPC in Cardiac Muscle

Ischemic preconditioning (IPC) has been demonstrated to protect both cardiac and skeletal muscles from ischemia-reperfusion (IR) injury by subjecting them to sub-lethal levels of IR for several cycles (24). However, the exact mechanism behind IPC remains unknown despite it being one of the most effective methods for protecting against ischemic injury in both experimental and partially clinical studies (25).

During prolonged exposure to IR, preconditioned myocardium exhibits reduced oxidative stress (26). The release of ATP-sensitive K⁺ (mitoKATP) channels and the limited production of reactive oxygen species (ROS) play a role in IPC's protective effect. Disruption of either mitoKATP channels or ROS production can counteract IPC's safety net (27). Activation of mitoKATP channels may result in the generation of mitochondrial ROS, which prevents cell death by inhibiting mitochondrial permeability transition pore (mPTP) opening through protein kinase C (PKC) activation (28). Protein kinase C serves as a signaling molecule that protects mitochondria throughout the IPC process. In the early stages of IPC, activation of opioid, bradykinin, and adenosine receptors leads to PKC activation and sensitizes the A2b adenosine receptor (A2bAR). This sensitization triggers a survival pathway involving PI3 kinase, Akt, and ERK, ultimately blocking mPTP opening (29). MitoKATP activation has also been observed to inhibit mitochondrial matrix contraction, potentially enhancing ATP production (6).

3.5. Molecular Mechanisms of H₂S-Induced Cardio Protection

With advances in scientific technology, researchers have uncovered the role of hydrogen sulfide (H₂S) in




Figure 1. Chemical structures of ginsenoside Rb1

various physiological and pathological processes in mammals. H_2S , akin to nitric oxide (NO) and carbon monoxide (CO), is now recognized as a novel gaseous signaling molecule (30). H_2S is endogenously produced by several enzymes, including cystathionine-synthase (CBS), cystathionine-lyase (CSE), 3-mercaptopyruvate sulfur-transferase (3-MST), and cysteine aminotransferase (CAT) (31). The expression of these enzymes varies depending on the tissue. CBS is a crucial enzyme for

H_2S production in the nervous system, while CSE is the primary H_2S -producing enzyme in the cardiovascular system (32).

Numerous studies have shown that H_2S is associated with various pathophysiological processes such as oxidative stress, inflammation, apoptosis, and angiogenesis (30). Increasing evidence suggests that H_2S plays a significant role in regulating cardiac functions and has a protective effect on the pathophysiology and

development of heart disorders.

In this context, we will elucidate the mechanisms underlying this protective role, particularly in the context of preventing arrhythmias, which are undesirable side effects of ischemia that H₂S has been shown to guard against.

3.6. Regulation of Ion Channels

Hydrogen sulfide (H₂S) has significant effects on heart electrophysiology. The myocardial membrane contains two different types of Ca²⁺ channels, namely L-type and T-type. L-type Ca²⁺ channels are essential for maintaining the electrophysiological basis for the plateau phase of action potentials and excitation-contraction (EC) coupling (33). Whole patch clamp experiments in rat cardiomyocytes have revealed that NaHS negatively regulates L-type Ca²⁺ channels composed of CaV1.2 subunits (34).

In various clinical conditions such as cardiac hypertrophy and heart failure, T-type Ca²⁺ channels can be observed in atrial and ventricular myocytes, contributing to aberrant electrical activity and EC coupling (35). Recent research has confirmed that NaHS selectively inhibits T-type Ca²⁺ channels heterologously expressed in HEK293 cells (10 μM - 1 mM) (36).

KATP channels are distributed on the cell membranes and mitochondria in the myocardium and play a crucial role in the principal endogenous cardioprotective mechanism involved in cardiac ischemia preconditioning-opening of KATP channels. Opening of KATP channels leads to hyperpolarization, reducing calcium influx via L-type Ca²⁺ channels and preventing Ca²⁺ overload. Tang and coworkers have reported that NaHS (100 μM) opens KATP channels in vascular smooth muscle cells (37). Additionally, it may indirectly activate KATP channels by causing intracellular acidosis (38). H₂S shortens action potential duration (APD) and exerts cardioprotective effects by activating KATP channels, while it does not affect the amplitude of action potentials or resting potential (39).

Studies have indicated that H₂S can modulate voltage-dependent Na⁺ channels (Nav). There are native Nav and recombinant Nav (NaV1.5) expressed in different tissues, including the heart. Strega et al. demonstrated that NaHS altered peak sodium current, voltage dependence of inactivation, and activation in Nav channels (40). Since NaV1.5 is responsible for the upstroke of the cardiac action potential, H₂S might have a similar impact on Nav channels expressed in the heart.

Chloride channels play a significant role in the normal physiological functions of myocardial cells. However, abnormal changes may occur in pathological conditions

such as myocardial ischemia and arrhythmias. Malenkov et al. proposed that H₂S affects single-channel currents of chloride channels, showing that NaHS inhibits chloride channels by reducing channel open probability in a concentration-dependent manner, as demonstrated by patch clamp experiments (41). The biological effects of H₂S in the heart may be linked to its inhibitory impact on chloride channels (42).

3.7. Pyruvate's Cardio-Protective Antioxidant Mechanisms

In addition to its ATP-producing activities, pyruvate exhibits potent antioxidant effects as a metabolic fuel. The antioxidant properties of pyruvate are mediated through various pathways, as illustrated in Figure 2 (43). Through direct, non-enzymatic processes, pyruvate-containing aliphatic carboxylates undergo oxidative decarboxylation, effectively reducing and detoxifying peroxides, yielding electrons, hydroxyl radicals, and nitrite substitutes (44). Consequently, pyruvate has several effects: (1) Reduction of lipid peroxides to their conjugated alcohols, thereby preventing membrane lipid peroxidation; (2) conversion of hydrogen peroxide (H₂O₂) and hydroxyl radicals (OH) into water; and (3) transformation of peroxynitrite into a benign form of nitrite (45).

Pyruvate decarboxylation also generates acetate, an oxidizable fuel. Glutathione, a tripeptide and the most abundant natural antioxidant in cells contains a sulfhydryl moiety capable of neutralizing peroxides, peroxynitrites, and hydroxyl radicals (46). During ischemia, particularly during efforts to restore circulation, such as revascularization and fluid resuscitation, large quantities of these hazardous metabolites are generated (47). The oxyradical load can deplete cellular glutathione (GSH), jeopardizing GSH-dependent antioxidant mechanisms. Pyruvate supports GSH in two ways, as depicted in Figure 2 (48).

Pyruvate safeguards cellular GSH levels through two primary mechanisms: first, by directly detoxifying RONS, it unburdens GSH-dependent antioxidant enzymes. Second, carboxylation of pyruvate increases the amount of Krebs cycle intermediates, such as citrate, which is a natural inhibitor of the glycolytic enzyme phosphofructokinase (49).

In isolated hearts and in vivo myocardium, pyruvate treatment significantly elevates citrate content (50). Subsequently, citrate is transported into the cytosol by tricarboxylate transporters in the inner mitochondrial membrane (Figure 2). This action impedes phosphofructokinase, leading to the accumulation of glucose 6-phosphate, which serves as a substrate for the hexose monophosphate pathway used to maintain GSH levels. Another source of NADPH is the NADP⁺-dependent

concentration of 0.5 mM) did not inhibit the suppression of RONS by pyruvate, whereas sarcolemma pyruvate uptake had no impact. Consequently, in this model, pyruvate's antioxidant actions are primarily cytosolic, unlike its energy-producing and inotropic activities that require mitochondrial metabolism (55).

Pyruvate's antioxidant properties are believed to contribute to its ability to enhance β -adrenergic pathways in shocked myocardium. Ischemia-reperfusion reduced β -adrenergic stimulation of contractile function in isolated guinea pig hearts, but pyruvate therapy largely restored it. The relationship between cardiac output and isoproterenol concentration was altered by ischemia-reperfusion, as evidenced by the drop in EC50 from 0.3 ± 0.06 nM pre-ischemia to 5.2 ± 1.9 nM in anesthetized hearts. However, when 5 mM pyruvate was administered after 15 minutes of reperfusion, the EC50 for isoproterenol decreased to 1.1 ± 0.3 nM. Consequently, 2 nM isoproterenol, which had a pronounced inotropic effect in pre-ischemic hearts, was ineffective in shocked myocardium. Still, 5 mM pyruvate substantially restored the inotropic response to isoproterenol (56).

However, while pyruvate alone increased GATP levels, GATP remained comparable in the absence vs. presence of 2 nM isoproterenol stimulation. The increase in GATP couldn't account for the fivefold greater power of the hearts receiving 2 nM isoproterenol and pyruvate compared to the hearts receiving 2 nM isoproterenol alone. In contrast, the antioxidant ratios GSH/GSSG and NADPH/NADP⁺ were significantly increased by pyruvate, similar to the effects of the sulfhydryl antioxidant N-acetylcysteine. This increase in antioxidant ratios accompanied the augmentation of isoproterenol-induced power and GSH/GSSG, similar to pyruvate, without altering Δ GATP levels (52). Therefore, pyruvate's antioxidant mechanisms strengthen β -adrenergic inotropic responses in shocked myocardium (57).

Other research demonstrated that inhibiting the activity of Pyruvate dehydrogenase kinase 4 (PDK4) altered the stimulation of glucose oxidation, potentially alleviating glucose oxidation, increasing glucose uptake, and reducing myocardial ischemia-reperfusion injury (58).

3.8. Tannins and Their Biological Effects

Tannins are a diverse group of water-soluble polyphenolic compounds with a high molecular weight (500 - 3000 Daltons) and up to 20 hydroxyl groups. They are commonly found in plants, foods, and beverages (59). Tannins are chemically reactive phenolic compounds that form both inter- and intramolecular hydrogen bonds, enabling them to interact with and precipitate

macromolecules such as proteins and carbohydrates. They are also responsible for the astringent taste found in many fruits and vegetables (60). Astringency tends to increase with the presence of one to five hydroxyl groups but decreases when there are seven or more, as steric hindrance begins to counteract the strength of hydrogen bonding (61).

Despite being abundant in human food (with an estimated daily intake of 0.1 - 0.5 g), tannins have received relatively little attention due to their polymeric structure and structural complexity (62). In recent years, there has been growing interest in the potential beneficial effects of proanthocyanidins and their monomers on human health. These effects include immunomodulation, anti-inflammatory properties, anticancer activity, antioxidant effects, cardio-protection, and anti-thrombotic properties (63).

For instance, sumac extracts and isolates containing tannins are known to enhance the quality and oxidative stability of animal products such as meat and milk (64).

However, more research is needed to fully understand the pharmacological and toxicological activities of tannins and their specific roles in human health. Despite the fact that several plant-derived proanthocyanidins are now widely consumed as dietary supplements, there is currently no conclusive evidence regarding their safety or potential for long-term harm (65). In this context, we have examined the most recent research and findings regarding the characteristics of tannins, with a particular focus on the antioxidant effects of proanthocyanidins on the heart (66).

3.9. Cardioprotective Properties

The relationship between proanthocyanidin consumption and its cardioprotective effects has been extensively investigated (67). Numerous in vitro and animal studies have been conducted to elucidate the mechanisms through which proanthocyanidins contribute to the prevention and improvement of various cardiac conditions, including atherosclerosis, blood pressure regulation, and lipid homeostasis (68).

Recent research has identified additional modes of action. For example, oligomeric proanthocyanidins (OPCs) derived from *Crataegus oxyacantha* L. have been shown to inhibit monocyte-to-macrophage differentiation in atherosclerosis, as demonstrated in both in vitro and in vivo studies. Oligomeric proanthocyanidins were found to reduce levels of vascular cell adhesion protein 1, chemokine CCL2, as well as inflammatory markers like MMP 2 and 9 and PPAR. These findings suggest that these phytochemicals may play a role in

modulating macrophage activity during the early stages of atherosclerosis and related conditions (69).

Another animal study investigated the cardioprotective effects of proanthocyanidins in rats with doxorubicin-induced heart damage. Proanthocyanidins significantly reduced doxorubicin-induced electrocardiographic abnormalities and corrected ventricular tachycardia induced by aconitine. Additionally, proanthocyanidins markedly reduced the biochemical changes induced by doxorubicin, including creatine kinase-myocardial band, LDH, MDA, SOD, and CAT levels (70). Moreover, GSPs have been observed to possess antioxidant, anti-inflammatory, and antiapoptotic effects, making them capable of protecting the liver against ischemia/reperfusion injury by attenuating endoplasmic reticulum stress (71).

Furthermore, researchers reported that intraperitoneal administration of fisetin (20 mg/kg) 15 minutes before surgery attenuated I/R-induced mitochondrial dysfunction, restored mitochondrial copy number, modulated Pgc1 gene expression, reduced mitochondrial fission, enhanced mitochondrial fusion, and promoted mitochondrial autophagy. These actions collectively counteracted the I/R-related reduction in mitochondrial mass in renal tissue (72).

There is also evidence suggesting that a diet rich in proanthocyanins can protect against acute ischemic brain injury in rats, leading to improved motor function, reduced cerebral infarction volume, and decreased levels of peroxidative markers such as 4-hydroxyl reduction and advanced glycation products. Additionally, inflammatory markers, including CCL2, ionized calcium-binding adapter molecule-1, and TNF- α , were reduced (73).

Furthermore, a study highlighted the bioactive components of *Tamarindus indica*, specifically Thymine and 4H-Pyran-4-one, 2,3-dihydro-3,5-dihydroxy-6-methyl-, which exhibited significant binding affinity with the human AT1 receptor antagonist via PPAR γ agonist. These antioxidant properties could potentially serve as a dietary supplement for mitigating cardiac toxicity (74).

Lastly, recent research explored the inhibitory effects of procyanidin B2 on the NOD-like receptor family. A significant downregulation of NLRP3 inflammasome in human umbilical vein endothelial cells (HUVECs) was observed, leading to the inhibition of caspase-1 activation and IL-1 β secretion in response to lipopolysaccharides (LPS). Furthermore, procyanidin B2 reduced the production of reactive oxygen species (ROS) induced by LPS and suppressed the transcriptional activity of activator protein-1 (75, 76).

4. Conclusions

This review has provided a comparative analysis of the antioxidant effects of various plants and tannins in the context of myocardial ischemia. It is evident that different antioxidants possess anti-ischemic and anti-reperfusion properties.

Footnotes

Authors' Contribution: Study concept and design, Najmeh Sadeghi; Acquisition of data, Najmeh Sadeghi, Hassanali Abedi; Analysis and interpretation of data, Najmeh Sadeghi; Drafting of the manuscript, Najmeh Sadeghi, Shekofe Atashpour; Critical revision of the manuscript for important intellectual content, Najmeh Sadeghi, Hassanali Abedi; Administrative, technical, and material support, Najmeh Sadeghi, Shekofe Atashpour; Study supervision, Sodabeh Etemadi.

Conflict of Interests: The authors disclose no conflicts of interest.

Funding/Support: This study did not receive any funding/support.

References

1. Mozaffarian D, Benjamin EJ, Go AS, Arnett DK, Blaha MJ, Cushman M, et al. Heart disease and stroke statistics—2016 update: A report from the American Heart Association. *Circulation*. 2016;**133**(4):e38–e360.
2. Weintraub WS, Daniels SR, Burke LE, Franklin BA, Goff DJ, Hayman LL, et al. Value of primordial and primary prevention for cardiovascular disease: A policy statement from the American Heart Association. *Circulation*. 2011;**124**(8):967–90. [PubMed ID: 21788592]. <https://doi.org/10.1161/CIR.0b013e3182285a81>.
3. Ribas N, Garcia-Garcia C, Merono O, Recasens L, Perez-Fernandez S, Bazan V, et al. Secondary prevention strategies after an acute ST-segment elevation myocardial infarction in the AMI code era: Beyond myocardial mechanical reperfusion. *BMC Cardiovasc Disord*. 2017;**17**(1):54. [PubMed ID: 28173757]. [PubMed Central ID: PMC5297147]. <https://doi.org/10.1186/s12872-017-0493-6>.
4. Heusch G, Gersh BJ. The pathophysiology of acute myocardial infarction and strategies of protection beyond reperfusion: A continual challenge. *Eur Heart J*. 2017;**38**(11):774–84. [PubMed ID: 27354052]. <https://doi.org/10.1093/eurheartj/ehw224>.
5. Seewald M, Coles JJ, Sigg DC, Iazzo PA. Featured Article: Pharmacological postconditioning with delta opioid attenuates myocardial reperfusion injury in isolated porcine hearts. *Exp Biol Med (Maywood)*. 2017;**242**(9):986–95. [PubMed ID: 28440739]. [PubMed Central ID: PMC5407584]. <https://doi.org/10.1177/1535370216684041>.
6. Zhou T, Prather ER, Garrison DE, Zuo L. Interplay between ROS and antioxidants during ischemia-reperfusion injuries in cardiac and skeletal muscle. *Int J Mol Sci*. 2018;**19**(2). [PubMed ID: 29385043]. [PubMed Central ID: PMC5855639]. <https://doi.org/10.3390/ijms19020417>.
7. Safaei N, Babaei H, Azarfarin R, Jodati AR, Yaghoubi A, Sheikhalizadeh MA. Comparative effect of grape seed extract (*Vitis vinifera*) and ascorbic acid in oxidative stress induced by on-pump coronary artery bypass surgery. *Ann Card Anaesth*. 2017;**20**(1):45–51.

- [PubMed ID: 28074795]. [PubMed Central ID: PMC5290695]. <https://doi.org/10.4103/0971-9784.197834>.
8. Booth J, Clayton T, Pepper J, Nugara F, Flather M, Sigwart U, et al. Randomized, controlled trial of coronary artery bypass surgery versus percutaneous coronary intervention in patients with multivessel coronary artery disease: Six-year follow-up from the Stent or Surgery Trial (SoS). *Circulation*. 2008;**118**(4):381-8. [PubMed ID: 18606919]. <https://doi.org/10.1161/CIRCULATIONAHA.107.739144>.
 9. Carlstrom M, Persson AE, Larsson E, Hezel M, Scheffer PG, Teerlink T, et al. Dietary nitrate attenuates oxidative stress, prevents cardiac and renal injuries, and reduces blood pressure in salt-induced hypertension. *Cardiovasc Res*. 2011;**89**(3):574-85. [PubMed ID: 21097806]. <https://doi.org/10.1093/cvr/cvq366>.
 10. Ling WC, Murugan DD, Lau YS, Vanhoutte PM, Mustafa MR. Sodium nitrite exerts an antihypertensive effect and improves endothelial function through activation of eNOS in the SHR. *Sci Rep*. 2016;**6**:33048. [PubMed ID: 27616322]. [PubMed Central ID: PMC5018809]. <https://doi.org/10.1038/srep33048>.
 11. Ling WC, Mustafa MR, Vanhoutte PM, Murugan DD. Chronic administration of sodium nitrite prevents hypertension and protects arterial endothelial function by reducing oxidative stress in angiotensin II-infused mice. *Vascul Pharmacol*. 2018;**102**:11-20. [PubMed ID: 28552746]. <https://doi.org/10.1016/j.vph.2017.05.003>.
 12. Amaral JH, Ferreira GC, Pinheiro LC, Montenegro MF, Tanus-Santos JE. Consistent antioxidant and antihypertensive effects of oral sodium nitrite in DOCA-salt hypertension. *Redox Biol*. 2015;**5**:340-6. [PubMed ID: 26119848]. [PubMed Central ID: PMC4491646]. <https://doi.org/10.1016/j.redox.2015.06.009>.
 13. Waltz P, Escobar D, Botero AM, Zuckerbraun BS. Nitrate/Nitrite as Critical Mediators to Limit Oxidative Injury and Inflammation. *Antioxid Redox Signal*. 2015;**23**(4):328-39. [PubMed ID: 26140517]. [PubMed Central ID: PMC4692126]. <https://doi.org/10.1089/ars.2015.6256>.
 14. Jadert C, Petersson J, Massena S, Ahl D, Grapensparr L, Holm L, et al. Decreased leukocyte recruitment by inorganic nitrate and nitrite in microvascular inflammation and NSAID-induced intestinal injury. *Free Radic Biol Med*. 2012;**52**(3):683-92. [PubMed ID: 22178413]. <https://doi.org/10.1016/j.freeradbiomed.2011.11.018>.
 15. Apostoli GL, Solomon A, Smallwood MJ, Winyard PG, Emerson M. Role of inorganic nitrate and nitrite in driving nitric oxide-cGMP-mediated inhibition of platelet aggregation in vitro and in vivo. *J Thromb Haemost*. 2014;**12**(11):1880-9. [PubMed ID: 25163536]. <https://doi.org/10.1111/jth.12711>.
 16. Zollbrecht C, Persson AE, Lundberg JO, Weitzberg E, Carlstrom M. Nitrite-mediated reduction of macrophage NADPH oxidase activity is dependent on xanthine oxidoreductase-derived nitric oxide but independent of S-nitrosation. *Redox Biol*. 2016;**10**:119-27. [PubMed ID: 27744114]. [PubMed Central ID: PMC5065649]. <https://doi.org/10.1016/j.redox.2016.09.015>.
 17. Khambata RS, Ghosh SM, Rathod KS, Thevathasan T, Filomena F, Xiao Q, et al. Antiinflammatory actions of inorganic nitrate stabilize the atherosclerotic plaque. *Proc Natl Acad Sci U S A*. 2017;**114**(4):E550-9. [PubMed ID: 28057862]. [PubMed Central ID: PMC5278489]. <https://doi.org/10.1073/pnas.1613063114>.
 18. Munzel T, Daiber A. Inorganic nitrite and nitrate in cardiovascular therapy: A better alternative to organic nitrates as nitric oxide donors? *Vascul Pharmacol*. 2018;**102**:1-10. [PubMed ID: 29174923]. <https://doi.org/10.1016/j.vph.2017.11.003>.
 19. Wu Y, Xia ZY, Dou J, Zhang L, Xu JJ, Zhao B, et al. Protective effect of ginsenoside Rb1 against myocardial ischemia/reperfusion injury in streptozotocin-induced diabetic rats. *Mol Biol Rep*. 2011;**38**(7):4327-35. [PubMed ID: 21136666]. <https://doi.org/10.1007/s11033-010-0558-4>.
 20. Bodiga S, Wang W, Oudit GY. Use of ginseng to reduce post-myocardial adverse myocardial remodeling: Applying scientific principles to the use of herbal therapies. *J Mol Med (Berl)*. 2011;**89**(4):317-20. [PubMed ID: 21340500]. <https://doi.org/10.1007/s00109-011-0736-4>.
 21. Leng X, Zhang LD, Jia LQ, Zang AY, Cao J, Li QF, et al. Effect of ginsenoside Rb1 on isoproterenol-induced acute myocardial ischemia in rats and its mechanism of action. *Chinese J Experimental Traditional Medical Formulae*. 2015;**21**(24):104-8.
 22. Zheng Q, Bao XY, Zhu PC, Tong Q, Zheng GQ, Wang Y. Ginsenoside Rb1 for myocardial ischemia/reperfusion injury: Preclinical evidence and possible mechanisms. *Oxid Med Cell Longev*. 2017;**2017**:6313625. [PubMed ID: 29430282]. [PubMed Central ID: PMC5753014]. <https://doi.org/10.1155/2017/6313625>.
 23. Liu YF, Liu SW, Liu ZX. Effects of ginsenoside Rb1 on blood vessel regeneration after ischemia and reperfusion in rats. *Chin J Histochem Cytochem*. 2008;**17**:39-44.
 24. Pang CY, Yang RZ, Zhong A, Xu N, Boyd B, Forrest CR. Acute ischaemic preconditioning protects against skeletal muscle infarction in the pig. *Cardiovasc Res*. 1995;**29**(6):782-8. [PubMed ID: 7656281].
 25. Liang J, Han R, Zhou B. Metabolic reprogramming: Strategy for ischemic stroke treatment by ischemic preconditioning. *Biology (Basel)*. 2021;**10**(5). [PubMed ID: 34064579]. [PubMed Central ID: PMC8151271]. <https://doi.org/10.3390/biology10050424>.
 26. Zhou T, Chuang CC, Zuo L. Molecular characterization of reactive oxygen species in myocardial ischemia-reperfusion injury. *Biomed Res Int*. 2015;**2015**:864946. [PubMed ID: 26509170]. [PubMed Central ID: PMC4609796]. <https://doi.org/10.1155/2015/864946>.
 27. Garlid KD, Dos Santos P, Xie ZJ, Costa AD, Paucek P. Mitochondrial potassium transport: The role of the mitochondrial ATP-sensitive K(+) channel in cardiac function and cardioprotection. *Biochim Biophys Acta*. 2003;**1606**(1-3):1-21. [PubMed ID: 14507424]. [https://doi.org/10.1016/s0005-2728\(03\)00109-9](https://doi.org/10.1016/s0005-2728(03)00109-9).
 28. Kalogeris T, Bao Y, Korhuis RJ. Mitochondrial reactive oxygen species: A double edged sword in ischemia/reperfusion vs preconditioning. *Redox Biol*. 2014;**2**:702-14. [PubMed ID: 24944913]. [PubMed Central ID: PMC4060303]. <https://doi.org/10.1016/j.redox.2014.05.006>.
 29. Yang X, Cohen MV, Downey JM. Mechanism of cardioprotection by early ischemic preconditioning. *Cardiovasc Drugs Ther*. 2010;**24**(3):225-34. [PubMed ID: 20505987]. [PubMed Central ID: PMC2932886]. <https://doi.org/10.1007/s10557-010-6236-x>.
 30. Wang R. Physiological implications of hydrogen sulfide: A whiff exploration that blossomed. *Physiol Rev*. 2012;**92**(2):791-896. [PubMed ID: 22535897]. <https://doi.org/10.1152/physrev.00017.2011>.
 31. Wagner F, Asfar P, Calzia E, Radermacher P, Szabo C. Bench-to bedside review: Hydrogen sulfide—the third gaseous transmitter: Applications for critical care. *Crit Care*. 2009;**13**(3):213. [PubMed ID: 19519960]. [PubMed Central ID: PMC2717401]. <https://doi.org/10.1186/cc7700>.
 32. Yang C, Yang Z, Zhang M, Dong Q, Wang X, Lan A, et al. Hydrogen sulfide protects against chemical hypoxia-induced cytotoxicity and inflammation in HaCaT cells through inhibition of ROS/NF-kappaB/COX-2 pathway. *PLoS One*. 2011;**6**(7). e21971. [PubMed ID: 21779360]. [PubMed Central ID: PMC3136491]. <https://doi.org/10.1371/journal.pone.0021971>.
 33. Bers DM. Calcium cycling and signaling in cardiac myocytes. *Annu Rev Physiol*. 2008;**70**:23-49. [PubMed ID: 17988210]. <https://doi.org/10.1146/annurev.physiol.70.113006.100455>.
 34. Zhang R, Sun Y, Tsai H, Tang C, Jin H, Du J. Hydrogen sulfide inhibits L-type calcium currents depending upon the protein sulphydryl state in rat cardiomyocytes. *PLoS One*. 2012;**7**(5). e37073. [PubMed ID: 22590646]. [PubMed Central ID: PMC3349658]. <https://doi.org/10.1371/journal.pone.0037073>.
 35. Vassort G, Talavera K, Alvarez JL. Role of T-type Ca²⁺ channels in the heart. *Cell Calcium*. 2006;**40**(2):205-20. [PubMed ID: 16766028]. <https://doi.org/10.1016/j.ceca.2006.04.025>.
 36. Elies J, Scragg JL, Huang S, Dallas ML, Huang D, MacDougall D, et al. Hydrogen sulfide inhibits Cav3.2 T-type Ca²⁺ channels. *FASEB J*.

- 2014;**28**(12):5376–87. [PubMed ID: 25183670]. <https://doi.org/10.1096/fj.14-257113>.
37. Tang G, Wu L, Liang W, Wang R. Direct stimulation of K(ATP) channels by exogenous and endogenous hydrogen sulfide in vascular smooth muscle cells. *Mol Pharmacol*. 2005;**68**(6):1757–64. [PubMed ID: 16150926]. <https://doi.org/10.1124/mol.105.017467>.
38. Lee SW, Cheng Y, Moore PK, Bian JS. Hydrogen sulphide regulates intracellular pH in vascular smooth muscle cells. *Biochem Biophys Res Commun*. 2007;**358**(4):1142–7. [PubMed ID: 17531202]. <https://doi.org/10.1016/j.bbrc.2007.05.063>.
39. Zhang Z, Huang H, Liu P, Tang C, Wang J. Hydrogen sulfide contributes to cardioprotection during ischemia-reperfusion injury by opening K ATP channels. *Can J Physiol Pharmacol*. 2007;**85**(12):1248–53. [PubMed ID: 18066126]. <https://doi.org/10.1139/Y07-120>.
40. Strege PR, Bernard CE, Kraichely RE, Mazzone A, Sha L, Beyder A, et al. Hydrogen sulfide is a partially redox-independent activator of the human jejunal Na⁺ channel, Nav1.5. *Am J Physiol Gastrointest Liver Physiol*. 2011;**300**(6):G105–14. [PubMed ID: 21393430]. [PubMed Central ID: PMC3119119]. <https://doi.org/10.1152/ajpgi.00556.2010>.
41. Malekova L, Krizanova O, Ondrias K. H(2)S and HS(-) donor NaHS inhibits intracellular chloride channels. *Gen Physiol Biophys*. 2009;**28**(2):190–4. [PubMed ID: 19592715]. <https://doi.org/10.4149/gpb.2009.02.190>.
42. Shen Y, Shen Z, Luo S, Guo W, Zhu YZ. The cardioprotective effects of hydrogen sulfide in heart diseases: From molecular mechanisms to therapeutic potential. *Oxid Med Cell Longev*. 2015;**2015**:925167. [PubMed ID: 26078822]. [PubMed Central ID: PMC4442295]. <https://doi.org/10.1155/2015/925167>.
43. Bunger R, Kang YH, Lee YJ, Mongan P, Sharma P. Redox and antioxidant mechanisms of pyruvate in cellular oxidative stress: Enhanced survival signaling and mitochondrial stability. *Recent Res Devel Physiol*. 2006;**4**:1–25.
44. Vasquez-Vivar J, Denicola A, Radi R, Augusto O. Peroxynitrite-mediated decarboxylation of pyruvate to both carbon dioxide and carbon dioxide radical anion. *Chem Res Toxicol*. 1997;**10**(7):786–94. [PubMed ID: 9250413]. <https://doi.org/10.1021/tx970031g>.
45. Raghavamenon A, Garelnabi M, Babu S, Aldrich A, Litvinov D, Parthasarathy S. Alpha-tocopherol is ineffective in preventing the decomposition of preformed lipid peroxides and may promote the accumulation of toxic aldehydes: A potential explanation for the failure of antioxidants to affect human atherosclerosis. *Antioxid Redox Signal*. 2009;**11**(6):1237–48. [PubMed ID: 19186999]. [PubMed Central ID: PMC2842134]. <https://doi.org/10.1089/ars.2008.2248>.
46. Cheung PY, Wang W, Schulz R. Glutathione protects against myocardial ischemia-reperfusion injury by detoxifying peroxynitrite. *J Mol Cell Cardiol*. 2000;**32**(9):1669–78. [PubMed ID: 10966829]. <https://doi.org/10.1006/jmcc.2000.1203>.
47. Zakkar M, Guida G, Suleiman MS, Angelini GD. Cardiopulmonary bypass and oxidative stress. *Oxid Med Cell Longev*. 2015;**2015**:189863. [PubMed ID: 25722792]. [PubMed Central ID: PMC4334937]. <https://doi.org/10.1155/2015/189863>.
48. Mallet RT, Sun J, Knott EM, Sharma AB, Olivencia-Yurvati AH. Metabolic cardioprotection by pyruvate: Recent progress. *Exp Biol Med (Maywood)*. 2005;**230**(7):435–43. [PubMed ID: 15985618]. <https://doi.org/10.1177/153537020523000701>.
49. Pound KM, Sorokina N, Ballal K, Berkich DA, Fasano M, Lanoue KF, et al. Substrate-enzyme competition attenuates upregulated anaplerotic flux through malic enzyme in hypertrophied rat heart and restores triacylglyceride content: Attenuating upregulated anaplerosis in hypertrophy. *Circ Res*. 2009;**104**(6):805–12. [PubMed ID: 19213957]. [PubMed Central ID: PMC2908318]. <https://doi.org/10.1161/CIRCRESAHA.108.189951>.
50. Gurji HA, White DW, Hoxha B, Sun J, Olivencia-Yurvati AH, Mallet RT. Pyruvate-fortified resuscitation stabilizes cardiac electrical activity and energy metabolism during hypovolemia. *World J Crit Care Med*. 2013;**2**(4):56–64. [PubMed ID: 24701417]. [PubMed Central ID: PMC3953871]. <https://doi.org/10.5492/wjccm.v2.i4.56>.
51. Sundqvist KE, Heikkila J, Hassinen IE, Hiltunen JK. Role of NADP⁺ (corrected)-linked malic enzymes as regulators of the pool size of tricarboxylic acid-cycle intermediates in the perfused rat heart. *Biochem J*. 1987;**243**(3):853–7. [PubMed ID: 3663104]. [PubMed Central ID: PMC1147935]. <https://doi.org/10.1042/bj2430853>.
52. Tejero-Taldo MI, Caffrey JL, Sun J, Mallet RT. Antioxidant properties of pyruvate mediate its potentiation of beta-adrenergic inotropism in stunned myocardium. *J Mol Cell Cardiol*. 1999;**31**(10):1863–72. [PubMed ID: 10525424]. <https://doi.org/10.1006/jmcc.1999.1020>.
53. Sharma AB, Sun J, Howard LL, Williams AJ, Mallet RT. Oxidative stress reversibly inactivates myocardial enzymes during cardiac arrest. *Am J Physiol Heart Circ Physiol*. 2007;**292**(1):H198–206. [PubMed ID: 16920803]. <https://doi.org/10.1152/ajpheart.00698.2006>.
54. Bassenge E, Sommer O, Schwemmer M, Bunger R. Antioxidant pyruvate inhibits cardiac formation of reactive oxygen species through changes in redox state. *Am J Physiol Heart Circ Physiol*. 2000;**279**(5):H2431–8. [PubMed ID: 11045981]. <https://doi.org/10.1152/ajpheart.2000.279.5.H2431>.
55. Mallet RT, Sun J. Mitochondrial metabolism of pyruvate is required for its enhancement of cardiac function and energetics. *Cardiovasc Res*. 1999;**42**(1):149–61. [PubMed ID: 10435006]. [https://doi.org/10.1016/S0008-6363\(98\)00300-9](https://doi.org/10.1016/S0008-6363(98)00300-9).
56. Tejero-Taldo MI, Sun J, Caffrey JL, Mallet RT. Pyruvate potentiates beta-adrenergic inotropism of stunned guinea-pig myocardium. *J Mol Cell Cardiol*. 1998;**30**(11):2327–39. [PubMed ID: 9925369]. <https://doi.org/10.1006/jmcc.1998.0792>.
57. Mallet RT, Olivencia-Yurvati AH, Bunger R. Pyruvate enhancement of cardiac performance: Cellular mechanisms and clinical application. *Exp Biol Med (Maywood)*. 2018;**243**(2):198–210. [PubMed ID: 29154687]. [PubMed Central ID: PMC5788146]. <https://doi.org/10.1177/1535370217743919>.
58. Wang X, Shen X, Yan Y, Li H. Pyruvate dehydrogenase kinases (PDKs): An overview toward clinical applications. *Biosci Rep*. 2021;**41**(4). [PubMed ID: 33739396]. [PubMed Central ID: PMC8026821]. <https://doi.org/10.1042/BSR202004402>.
59. De Jesus NZT, Falcao HS, Gomes IF, Leite TJA, Lima GRM, Barbosa-Filho JM, et al. Tannins, peptic ulcers and related mechanisms. *Int J Mol Sci*. 2012;**13**(3):3203–28. [PubMed ID: 22489149]. [PubMed Central ID: PMC3317710]. <https://doi.org/10.3390/ijms13033203>.
60. Lamy E, Pinheiro C, Rodrigues L, Capela-Silva F, Lopes O, Tavares S, et al. Determinants of tannin-rich food and beverage consumption: Oral perception vs. Psychosocial aspects. 2016. Available from: <https://dspace.uevora.pt/rdpc/handle/10174/18018>.
61. Zhou Q, Yin ZP, Ma L, Zhao W, Hao HW, Li HL. Free radical-scavenging activities of oligomeric proanthocyanidin from *Rhodiola rosea* L. and its antioxidant effects in vivo. *Nat Prod Res*. 2014;**28**(24):2301–3. [PubMed ID: 24896611]. <https://doi.org/10.1080/14786419.2014.921786>.
62. Serrano J, Puupponen-Pimia R, Dauer A, Aura AM, Saura-Calixto F. Tannins: current knowledge of food sources, intake, bioavailability and biological effects. *Mol Nutr Food Res*. 2009;**53** Suppl 2:S310–29. [PubMed ID: 19437486]. <https://doi.org/10.1002/mnfr.200900039>.
63. Sieniawska E. Activities of tannins—From in vitro studies to clinical trials. *Natural Product Communications*. 2015;**10**(11):1934578X150100118.
64. Batiha GE, Ogunyemi OM, Shaheen HM, Kutu FR, Olaiya CO, Sabatier JM, et al. *Rhus coriaria* L. (Sumac), a Versatile and Resourceful Food Spice with Cornucopia of Polyphenols. *Molecules*. 2022;**27**(16). [PubMed ID: 36014419]. [PubMed Central ID: PMC9414570]. <https://doi.org/10.3390/molecules27165179>.
65. Berry AC, Nakshabendi R, Abidali H, Atchaneeyasakul K, Dholaria K, Johnson C, et al. Adverse effects of grape seed extract supplement: A clinical case and long-term follow-up. *J Diet Suppl*. 2016;**13**(2):232–5. [PubMed ID: 25688637]. <https://doi.org/10.3109/19390211.2015.1008617>.

66. Smeriglio A, Barreca D, Bellocco E, Trombetta D. Proanthocyanidins and hydrolysable tannins: Occurrence, dietary intake and pharmacological effects. *Br J Pharmacol*. 2017;**174**(11):1244–62. [PubMed ID: 27646690]. [PubMed Central ID: PMC5429339]. <https://doi.org/10.1111/bph.13630>.
67. Blade C, Aragones G, Arola-Arnal A, Muguerza B, Bravo FI, Salvado MJ, et al. Proanthocyanidins in health and disease. *Biofactors*. 2016;**42**(1):5–12. [PubMed ID: 26762288]. <https://doi.org/10.1002/biof.1249>.
68. Quifer-Rada P, Choy YY, Calvert CC, Waterhouse AL, Lamuela-Raventos RM. Use of metabolomics and lipidomics to evaluate the hypocholesterolemic effect of Proanthocyanidins from grape seed in a pig model. *Mol Nutr Food Res*. 2016;**60**(10):2219–27. [PubMed ID: 27240545]. <https://doi.org/10.1002/mnfr.201600190>.
69. Mohana T, Navin AV, Jamuna S, Sakeena Sadullah MS, Niranjali Devaraj S. Inhibition of differentiation of monocyte to macrophages in atherosclerosis by oligomeric proanthocyanidins -In-vivo and in-vitro study. *Food Chem Toxicol*. 2015;**82**:96–105. [PubMed ID: 25981678]. <https://doi.org/10.1016/j.fct.2015.04.028>.
70. Ammar el SM, Said SA, El-Damarawy SL, Suddek GM. Cardioprotective effect of grape-seed proanthocyanidins on doxorubicin-induced cardiac toxicity in rats. *Pharm Biol*. 2013;**51**(3):339–44. [PubMed ID: 23134235]. <https://doi.org/10.3109/13880209.2012.729065>.
71. Xu ZC, Yin J, Zhou B, Liu YT, Yu Y, Li GQ. Grape seed proanthocyanidin protects liver against ischemia/reperfusion injury by attenuating endoplasmic reticulum stress. *World J Gastroenterol*. 2015;**21**(24):7468–77. [PubMed ID: 26139992]. [PubMed Central ID: PMC4481441]. <https://doi.org/10.3748/wjg.v21.i24.7468>.
72. Peng P, Zou J, Zhong B, Zhang G, Zou X, Xie T. Protective Effects and Mechanisms of Flavonoids in Renal Ischemia-Reperfusion Injury. *Pharmacology*. 2023;**108**(1):27–36. [PubMed ID: 36446333]. <https://doi.org/10.1159/000527262>.
73. Yunoki T, Deguchi K, Omote Y, Liu N, Liu W, Hishikawa N, et al. Anti-oxidative nutrient-rich diet protects against acute ischemic brain damage in rats. *Brain Res*. 2014;**1587**:33–9. [PubMed ID: 25175837]. <https://doi.org/10.1016/j.brainres.2014.08.056>.
74. Akter H, Rashid MM, Islam MS, Hossen MA, Rahman MA, Algheshairy RM, et al. Biometabolites of Tamarindus indica play a remarkable cardioprotective role as a functional food in doxorubicin-induced cardiotoxicity models. *J Functional Foods*. 2022;**96**:105212. <https://doi.org/10.1016/j.jbcp.2014.10.001>.
75. Yang H, Xiao L, Yuan Y, Luo X, Jiang M, Ni J, et al. Procyanidin B2 inhibits NLRP3 inflammasome activation in human vascular endothelial cells. *Biochem Pharmacol*. 2014;**92**(4):599–606. [PubMed ID: 25450671]. <https://doi.org/10.1016/j.bcp.2014.10.001>.
76. Yang L, Xian D, Xiong X, Lai R, Song J, Zhong J. Proanthocyanidins against oxidative stress: From molecular mechanisms to clinical applications. *Biomed Res Int*. 2018;**2018**:8584136. [PubMed ID: 29750172]. [PubMed Central ID: PMC5884402]. <https://doi.org/10.1155/2018/8584136>.