



The Effect of Nursing Care at Home on the Quality of Life and Movement Problems of Older Adults with Hip Fractures Operation

Sümeýra Koçyiğit ¹, Hatice Yorulmaz ^{2,*}

¹ Faculty of Health Sciences, Haliç University, İstanbul, Türkiye

² Faculty of Medicine, Haliç University, İstanbul, Türkiye

*Corresponding Author: Faculty of Medicine, Haliç University, İstanbul, Türkiye. Email: haticeyorulmaz@halic.edu.tr

Received: 9 November, 2024; Revised: 11 January, 2025; Accepted: 21 January, 2024

Abstract

Background: The most important factor affecting mortality after hip fracture is the patient's comorbidities and their overall health status.

Objectives: This study examined the effect of a home-based nursing care program, in addition to routine maintenance services, on the quality of life and mobility issues encountered in older adults who underwent hip fracture surgery.

Methods: This semi-experimental study was conducted in İstanbul. Patients were divided into two groups: (1) Control, and (2) experimental. The control group received only the standard care provided at the hospital or community clinic. In contrast, the experimental group received home care services, including educational training prepared by the researcher, at the first and third months post-discharge. The Oxford Hip Score (OHS) and the 'Quality of Life Scale' were administered to the patients.

Results: In this study, 30% of the control group were aged 65 - 69, and 56.7% were female. In the experimental group, 50% were aged 75 - 79, and 56.7% were female. We observed that scores for 'Hip Pain', 'Walking Distance', and 'Interference with General Work' at the first month, and 'Hip Pain', 'Washing', 'Transportation', and 'Wearing Socks' at the third month, decreased in the experimental group compared to the control group in the OHS. The experimental group also showed higher average scores in physical function, physical role, general health perception, and physical summary health score compared to the control one after the training ($P < 0.05$).

Conclusions: The postoperative discharge education program implemented for older adults resulted in significant improvements in their physical activities. Thorough preparation and execution of such educational programs can substantially alleviate the adverse outcomes associated with hip fractures in the elderly population.

Keywords: Hip Fracture, Home Care, Nursing, Older adults, Quality of Life

1. Background

Old age is a period characterized by a loss of status, increased risk of dependency and accidents, and a decline in physical abilities. Hip fractures remain a significant health issue, leading to mortality and disability, particularly in the older adult population. It is reported that approximately one-third of individuals over the age of 65 experience a fall at least once annually (1). Approximately one-third of women and one in twelve men will suffer a hip fracture in their lifetime. According to the World Health Organization, the number of hip fractures associated with osteoporosis is projected to triple over the next 50 years, increasing

from 1.7 million cases in 1990 to 6.3 million in 2050 worldwide (2). Hip fractures result in a 12 - 20% decrease in life expectancy and a 5 - 20% increase in disability rate within one year post-fracture (3). Patients with hip fractures are at high risk for age-related and fracture-related complications, even under optimal conditions (4). It has been determined that older adults who undergo hip fracture surgery face significant challenges in meeting their bathing, dressing, and toileting needs, often feeling dependent on others (5). Studies have confirmed that hip fractures have a strong and persistent negative impact on patients' quality of life (5, 6). Nursing care is emphasized as playing a crucial role in the recovery and return to normal life for individuals

in the postoperative period (7). Older adults with a high quality of life can maintain independence and meet their needs and daily living activities themselves (8). Therefore, nurses have the opportunity to evaluate the healthy lifestyle behaviors and quality of life of the patients they care for, and to plan and implement assistance and home care optimally according to the patient's needs (9).

2. Objectives

The present study aimed to investigate the effect of home-based nursing care, in addition to routine care, on the quality of life and other challenges faced by older adults undergoing hip fracture surgery.

3. Methods

3.1. Study Design

The study was conducted as a semi-experimental design.

3.2. Participants

The research population consisted of 300 patients who visited a health center and a state hospital within the borders of Istanbul in 2018. From this population, the sample comprised 60 patients who volunteered to participate in the study. Patients were selected using a simple random sampling method. The sample size was determined to be sufficient at a 90% confidence interval. The patients were divided into two groups: (1) Control (N = 30), and (2) experimental (N = 30). Both groups were composed of patients with similar sociodemographic characteristics. The research sample included patients aged 65 and over who had undergone hip fracture surgery, were literate, and were mentally competent. No exclusion criteria were applied. The experimental and control groups were formed with patients having similar sociodemographic characteristics.

3.3. Scales

Patients were asked to complete the patient information form, hip evaluation form, and Quality of Life Scale.

3.3.1. Patient Information Form

The patient information form included questions about age, gender, marital status, and other demographic details.

3.3.2. Short Form Health Survey

The Short Form Health Survey (SF-36) was developed by Ware and Sherbourne in 1992 (10). The reliability and validity of the Turkish version were evaluated by Kocyiğit et al. The scale includes 8 sub-dimensions (physical function, physical role, vitality/fatigue, pain, general health perception, social function, emotional role, mental health) and 2 main dimensions (physical dimension and mental dimension) (11). The SF-36 is scored such that higher scores in each health domain indicate a higher health-related quality of life. Scores for health-related life domains range from 0 to 100, with higher scores representing better health status.

3.3.3. The Oxford Hip Score

The Oxford Hip Score (OHS) was developed by Dawson et al. (12), and its validity and reliability for the Turkish population were established by Sendir and Babadag (13). The OHS consists of 12 statements covering aspects such as pain, washing, transportation, wearing socks, shopping, walking time, climbing stairs, getting up from a chair, limping, type of hip pain, night pain, and the effect of pain on daily activities. Each statement is scored on a Likert scale ranging from 1 to 5. Lower scores indicate less impact from the disease, while higher scores indicate greater impact.

3.4. Data Collection

In the procedure, patients in the control group received a brief training session upon discharge and were subsequently called for a routine check-up in the first month. The control group received routine care services at the hospital or health center. In contrast, the experimental group received home care services, including educational topics prepared by a specialist nurse, at the 1st and 3rd months post-discharge. The purpose of the research was explained to both the experimental and control groups, and data were collected using a face-to-face interview technique. The research sample consisted of patients aged 65 and over who had undergone hip fracture surgery, were literate, and mentally competent. The specialist nurse visited the homes of the patients in the experimental group during the first and third months after discharge to provide home care services and training. The specialist nurse administered the questionnaire to both the control and experimental groups at months 1 and 3.

3.4.1. Postoperative Discharge Education for Individuals with Hip Fracture Surgery

The training content included pain management, wound care, daily living activities, exercise, self-care, nutrition, monitoring for signs of complications, medication use, safety precautions at home, physical limitations, considerations for social and sexual life, use of assistive devices while walking, physical therapy as recommended by the doctor, and the importance of adhering to outpatient clinic follow-ups. Education was provided to patients by a specialist nurse and was supported with visual diagrams.

3.5. Data Analysis

The SPSS 22.0 statistical package program was used for data analysis. The Pearson's chi-square test and Fisher's exact test were employed to compare qualitative data. The Mann-Whitney U test and Wilcoxon signed-rank test were used to compare the parameters. Results were evaluated at a significance level of $P < 0.05$. The content validity ratio (CVR) for the items was above 0.76, and the Content Validity Index (CVI) was above 0.79 in the SF-36. The CVR for the items was above 0.81, and the CVI was above 0.74 in the OHS.

3.6. Ethical Consideration

Permission for the study was received from the Ministry of Health, Turkish Public Hospitals Institution (70794255-663.08/12.01.2018).

4. Results

Descriptive characteristics of the groups are presented in [Table 1](#). The distribution of the groups' responses to the OHS before (1st month) and after (3rd month) the training is shown in [Table 2](#). [Table 3](#) presents the distribution of the groups' OHS total score and question-based responses before (1st month) and after (3rd month) the training. After the training, the averages of physical function, pain, and mental role in the experimental group were higher than those in the control group ($P < 0.05$) ([Table 4](#)).

Before and after the training, patients in the experimental group had higher physical role, general health perception, and physical component summary (PCS) averages than those in the control group. The increase in physical role, pain, general health perception, vitality, social function, PCS, and mental

component summary (MCS) levels in the control group after the training was statistically significant ($P < 0.05$) ([Table 4](#)). The increase in physical function, physical role, pain, general health perception, vitality, social function, mental role, mental health, PCS, and MCS levels in the experimental group after the training was statistically significant ($P < 0.05$) ([Table 4](#)).

5. Discussion

Hip fractures in the geriatric population represent a significant public health concern (14). This study observed a higher prevalence of female patients. Raichandani et al. reported osteoporosis in 65% of females and 50% of males with hip fractures. The higher prevalence in females is attributed to increased bone loss during peri-menopause and post-menopause (15). As hip fracture surgeries are often performed at older ages, it is common for individuals in this age group to be married, with many being widowed. Marriage is considered a factor that facilitates and supports home care for individuals undergoing hip fracture surgery (16). The study observed that most individuals had lost their spouses and generally lived with their children and family members. Acute and chronic diseases, which increase with advancing age, lead to long-term and multiple drug use (17).

In the third month, the proportion of those experiencing severe pain in the control group decreased to 23.3%, whereas no one in the experimental group reported severe pain. Analysis of the OHS revealed that mean scores in the third month decreased in both groups compared to the first month. A study on hip replacement indicated that pain began to decrease in the third month, with optimal results observed in the sixth month (18). In the third month, the rate of patients unable to walk at all in the control group was 10%, which decreased to 3.3% in the experimental group. The OHS analysis showed that mean night pain scores in both the control and experimental groups decreased in the third month compared to the first month. Data collected from patients admitted to four New York city hospitals with hip fractures suggest that improved pain control can reduce the length of stay and enhance long-term functional outcomes (19).

Hip fracture is a critical condition that adversely affects the quality of life, particularly in older patients, where lost functionality due to physiological disorders and related problems, including disorders related to physiological and mental qualities, are observed (20).

Table 1. Descriptive Characteristics of the Groups^a

Variables	Control (n = 30)	Experimental (n = 30)	P; χ^2
Age			$\chi^2 = 4.273$; P = 0.233 ^b
65 - 69	6 (20.0)	3 (10.0)	
70 - 74	9 (30.0)	5 (16.7)	
75 - 79	8 (26.7)	15 (50.0)	
80 and over	7 (23.3)	7 (23.3)	
Gender			-
Female	17 (56.7)	17 (56.7)	
Male	13 (43.3)	13 (43.3)	
Education			$\chi^2 = 0.128$; P = 0.938 ^c
Illiterate	11 (36.7)	11 (36.7)	
Literate	14 (46.7)	13 (43.3)	
Primary school	5 (16.7)	6 (20.0)	
Marital status			$\chi^2 = 0.081$; P = 0.961 ^b
Married	11 (36.7)	12 (40.0)	
Single	5 (16.7)	5 (16.7)	
Widow	14 (46.7)	13 (43.3)	
Working status			$\chi^2 = 3.774$; P = 0.047 ^b
Yes	6 (20.0)	13 (43.3)	
No	24 (80.0)	17 (56.7)	
Income status			$\chi^2 = 3.826$; P = 0.148 ^c
Income less than expenses	11 (36.7)	5 (16.7)	
Income equals expenditure	13 (43.3)	20 (66.7)	
Income more than expenditure	6 (20.0)	5 (16.7)	
Shared life person(s)			$\chi^2 = 1.270$; P = 0.199 ^b
Family	19 (63.3)	23 (76.7)	
Retirement home and single	11 (36.7)	7 (23.3)	
Additional disease			$\chi^2 = 0.073$; P = 0.50 ^b
Yes	19 (63.3)	20 (66.7)	
No	11 (36.7)	10 (33.3)	
Previous hip fracture surgery			$\chi^2 = 1.763$; P = 0.144 ^b
Yes	14 (46.7)	9 (30.0)	
No	16 (53.3)	21 (70.0)	

^a Values are expressed as No (%).

^b Pearson's chi-square test.

^c Fisher's exact test.

Partial improvements have been demonstrated in the physical and psychosocial functions of patients three to four months post-fracture (21). The mean physical role scores of patients in the experimental group, both before and after training, were higher than those of the control group. Adachi et al. conducted similar studies and reported that hip fractures significantly decrease all dimensions of quality of life, particularly physical and social functions (22). The Functional Capacity Scale (FCS) averages of patients in the experimental group, both

before and after training, were higher than those of the control group. A study indicated that the quality of life for older patients with hip fractures was severely impaired one month post-fracture, with partial improvement by the fourth month (23).

In this study, the mean post-training pain levels of patients in the experimental group were higher than those in the control group. Poorly managed postoperative pain is associated with delayed ambulation and pulmonary complications (24). The

Table 3. Distribution of Oxford Hip Score Total Score and Question Based Responses of the Groups at Month 1 and then at Month 3^a

Variables	Control	Experimental	P-Value
Hip pain			
1st month	4.63 ± 0.490	4.07 ± 0.69	0.001
3rd month	4 ± 0.69	3.03 ± 0.71	0.000
Wilcoxon Z/P	-3.64/0.000	-3.77/0.000	
Washing			
1st month	4.30 ± 0.70	4.30 ± 0.65	0.942
3rd month	4.10 ± 0.66	2.63 ± 0.61	0.000
Wilcoxon Z/P	-1.269/0.20	-4.62/0.000	
Public transport			
1st month	4.53 ± 0.50	4.43 ± 0.50	0.442
3rd month	3.93 ± 0.69	2.47 ± 0.50	0.000
Wilcoxon Z/P	-3.175/0.001	-4.85/0.000	
Wearing socks			
1st month	4.43 ± 0.56	4.23 ± 0.62	0.217
3rd month	4.10 ± 0.60	2.80 ± 0.55	0.000
Wilcoxon Z/P	-2.23/0.025	-4.65/0.000	
Household shopping			
1st month	4.13 ± 0.73	4.27 ± 0.64	0.495
3rd month	4.03 ± 0.66	2.67 ± 0.54	0.000
Wilcoxon Z/P	-0.60/0.54	-4.82/0.000	
Walking			
1st month	3.80 ± 0.96	3.40 ± 0.72	0.083
3rd month	3.57 ± 0.81	2.90 ± 0.84	0.003
Wilcoxon Z/P	-1.09/0.27	-2.41/0.016	
Flat of stairs			
1st month	4.03 ± 0.80	3.93 ± 0.64	0.603
3rd month	3.57 ± 0.50	2.07 ± 0.64	0.000
Wilcoxon Z/P	-2.562/0.010	-4.769/0.000	
Stand up from a chair			
1st month	4.27 ± 0.691	4.07 ± 0.64	0.225
3rd month	3.47 ± 0.507	2.50 ± 0.50	0.000
Wilcoxon Z/P	-3.96/0.000	-4.815/0.000	
Limping			
1st month	4.43 ± 0.504	4.23 ± 0.72	0.343
3rd month	4.10 ± 0.71	2.33 ± 0.66	0.000
Wilcoxon Z/P	-2.041/0.041	-4.848/0.000	
Sudden/severe pain			
1st month	4.57 ± 0.50	4.5 ± 0.5	0.608
3rd month	4.40 ± 0.49	2.77 ± 0.62	0.000
Wilcoxon Z/P	-1.213/0.22	-4.86/0.000	
Usual work			
1st month	4.6 ± 0.49	4.1 ± 0.71	0.005
3rd month	4.5 ± 0.50	3.5 ± 0.68	0.000
Wilcoxon Z/P	-0.77/0.43	-2.99/0.003	
Hip pain in bed			
1st month	4.5 ± 0.50	3.97 ± 0.669	0.002
3rd month	3.9 ± 0.66	2.53 ± 0.507	0.000
Wilcoxon Z/P	-3.38/0.001	-4.696/0.000	
OHS toplam			
1st month	52.23 ± 4.38	49.5 ± 3.17	0.014
3rd month	47.56 ± 3.1	32.2 ± 2.52	0.000
Wilcoxon Z/P	-4.23/0.000	-4.786/0.000	

Abbreviation: OHS, Oxford Hip Score.

^a Values are expressed as mean ± SD.

mean mental role scores of patients in the experimental group, both before and after training, were higher than those of the control group. The lack of significant

differences, despite nursing care provided at home after an operation that caused more distress to patients, suggests that patients should receive mental support

Table 4. Distribution of Main and Sub-dimensions of Short Form Health Survey Before and After Training by Groups^a

Variables	Control	Experimental	P-Value
Physical function			
1st month	36.50 ± 15.54	35.83 ± 11.52	0.89
3rd month	40.33 ± 14.32	66.83 ± 11.48	0.000
Wilcoxon Z/P	-1.147/0.25	-4.74/0.000	
Physical role			
1st month	12.5 ± 12.7	35 ± 26.74	0.000
3rd month	50 ± 30.79	71.6 ± 24.33	0.004
Wilcoxon Z/P	-3.94/0.000	-4.037/0.000	
Pain			
1st month	21.96 ± 15.74	19.26 ± 9.91	0.471
3rd month	38.76 ± 12.51	54.9 ± 10.49	0.000
Wilcoxon Z/P	-3.601/0.000	-4.713/0.000	
General health			
1st month	32.76 ± 6.64	36.16 ± 5.03	0.034
3rd month	48.2 ± 7.08	65 ± 5.01	0.000
Wilcoxon Z/P	-4.79/0.000	-4.82/0.000	
Vitality			
1st month	25 ± 8.3	26.16 ± 7.15	0.67
3rd month	47.83 ± 7.15	50.33 ± 6.94	0.26
Wilcoxon Z/P	-4.72/0.000	-4.81/0.000	
Social function			
1st month	26.66 ± 11.24	22.91 ± 12.74	0.20
3rd month	50.41 ± 11.12	56.66 ± 10.24	0.06
Wilcoxon Z/P	-4.475/0.000	-4.820/0.000	
Role emotional			
1st month	33.33 ± 31.56	42.22 ± 38.09	0.38
3rd month	40 ± 28.23	68.88 ± 23.05	0.000
Wilcoxon Z/P	-1.06/0.28	-3.12/0.002	
Mental health			
1st month	49.06 ± 8.39	44.8 ± 8.75	0.055
3rd month	51.6 ± 8.36	49.73 ± 5.32	0.39
Wilcoxon Z/P	-1.35/0.17	-2.26/0.024	
PCS			
1st month	27.51 ± 4.82	29.82 ± 4.26	0.032
3rd month	35.83 ± 4.35	45.8 ± 4.01	0.000
Wilcoxon Z/P	-4.24/0.000	-4.78/0.000	
MCS			
1st month	35.68 ± 5.26	34.71 ± 5.34	0.47
3rd month	40.10 ± 4.3	40.18 ± 3.82	0.85
Wilcoxon Z/P	-3.38/0.001	-3.73/0.000	

Abbreviations: PCS, physical component summary; MCS, mental component summary.

^a Values are expressed as mean ± SD.

and nurses should be made aware of this issue (25). Su et al. demonstrated that a nurse-led care program improved health-related quality of life in older adults following hip fracture surgery (26). Similarly, Banappagoudar et al. showed that nursing intervention

was beneficial in improving physical and psychosocial functioning in elderly patients with hip fractures (27).

When responses to the OHS were examined, a significant decrease was observed in scores for hip pain, bathing, transportation, wearing socks, shopping,

walking distance, climbing stairs, pain when rising from a chair, limping while walking, sudden/serious pain, affecting general work, and pain in bed at night in the experimental group compared to the control group. Additionally, OHS analysis revealed that the mean scores for hip pain at the first month, the variable of general work being affected, and the total OHS score were higher in the control group than in the experimental group. In the experimental group, the mean total OHS score and responses to questions in the third month decreased significantly compared to the control group.

The small sample size is a limitation of this study. Furthermore, the number and duration of education provided to patients could not be monitored for at least six months.

5.1. Conclusions

These results indicate that the training program developed for patients made significant contributions to their healing process. Based on these findings, to reduce the frequency of hospitalization for patients undergoing hip fracture surgery, education programs for the patient and family should be initiated upon hospital admission, and discharge education should be comprehensively planned and implemented.

Acknowledgements

The authors are thankful for the participants who participated in the study.

Footnotes

Authors' Contribution: H. Y.: Study concept and design; S. K.: Analysis and interpretation of data; H. Y.: Drafting of the manuscript; S. K.: Critical revision of the manuscript for important intellectual content and H. Y.: Statistical analysis.

Conflict of Interests Statement: The authors declared no conflict of interest.

Data Availability: The data presented in this study are uploaded during submission as a supplementary file and are openly available for readers.

Ethical Approval: Permission for the study was received from the Ministry of Health, Turkish Public Hospitals Institution (70794255-663.08/12.01.2018).

Funding/Support: The present study received no funding/support.

Informed Consent: Informed consent was obtained from all participants.

References

- Veronese N, Maggi S. Epidemiology and social costs of hip fracture. *Injury*. 2018;**49**(8):1458-60. [PubMed ID: 29699731]. <https://doi.org/10.1016/j.injury.2018.04.015>.
- Dhanwal DK, Dennison EM, Harvey NC, Cooper C. Epidemiology of hip fracture: Worldwide geographic variation. *Indian J Orthop*. 2011;**45**(1):15-22. [PubMed ID: 21221218]. [PubMed Central ID: PMC3004072]. <https://doi.org/10.4103/0019-5413.73656>.
- Amarilla-Donoso FJ, Lopez-Espuela F, Roncero-Martin R, Leal-Hernandez O, Puerto-Parejo LM, Aliaga-Vera I, et al. Quality of life in elderly people after a hip fracture: a prospective study. *Health Qual Life Outcomes*. 2020;**18**(1):71. [PubMed ID: 32171319]. [PubMed Central ID: PMC7071575]. <https://doi.org/10.1186/s12955-020-01314-2>.
- Gonzalez-Zabaleta J, Pita-Fernandez S, Seoane-Pillado T, Lopez-Calvino B, Gonzalez-Zabaleta JL. Dependence for basic and instrumental activities of daily living after hip fractures. *Arch Gerontol Geriatr*. 2015;**60**(1):66-70. [PubMed ID: 25465506]. <https://doi.org/10.1016/j.archger.2014.10.020>.
- Handoll HH, Cameron ID, Mak JC, Panagoda CE, Finnegan TP. Multidisciplinary rehabilitation for older people with hip fractures. *Cochrane Database Syst Rev*. 2021;**11**(11). CD007125. [PubMed ID: 34766330]. [PubMed Central ID: PMC8586844]. <https://doi.org/10.1002/14651858.CD007125.pub3>.
- Campefeldt P, Ekstrom W, Al-Ani AN, Weibust E, Greve K, Hedstrom M. Health related quality of life and mortality 10 years after a femoral neck fracture in patients younger than 70 years. *Injury*. 2020;**51**(10):2283-8. [PubMed ID: 32620326]. <https://doi.org/10.1016/j.injury.2020.06.029>.
- MacDonald V, Maher AB, Mainz H, Meehan AJ, Brent L, Hommel A, et al. Developing and Testing an International Audit of Nursing Quality Indicators for Older Adults With Fragility Hip Fracture. *Orthop Nurs*. 2018;**37**(2):115-21. [PubMed ID: 29570544]. <https://doi.org/10.1097/NOR.0000000000000431>.
- Jazayeri E, Kazemipour S, Hosseini SR, Radfar M. Quality of life in the elderly: A community study. *Caspian J Int Med*. 2023;**14**(3):534-42. eng. <https://doi.org/10.22088/cjim.14.3.543>.
- Chang HK, Gil CR, Kim HJ, Bea HJ. Factors Affecting Quality of Life Among the Elderly in Long-Term Care Hospitals. *J Nurs Res*. 2020;**29**(1). e134. [PubMed ID: 33284135]. [PubMed Central ID: PMC7808359]. <https://doi.org/10.1097/jnr.0000000000000413>.
- Ware JJ, Sherbourne CD. The MOS 36-item short-form health survey (SF-36). I. Conceptual framework and item selection. *Med Care*. 1992;**30**(6):473-83. [PubMed ID: 1593914].
- Koçyiğit H, Aydemir Ö, Fişek G, Ölmez N, Memiş A. Reliability and validity of the Turkish version of short form 36 (SF-36) in patients with rheumatoid arthritis. *J Med and Therap*. 1999;**12**:102-6.
- Dawson J, Fitzpatrick R, Carr A, Murray D. Questionnaire on the Perceptions of Patients About Total Hip Replacement. *J Bone Joint Surg*. 1996;**78-B**(2):185-90. <https://doi.org/10.1302/0301-620X.78b2.0780185>.

13. Şendir M, Babadağ K. Effect of Preoperative Patient Education on Postoperative Physical Adaptation and Quality of Life after Total Hip Replacement Surgery. *Florence Nightingale J Nurs*. 2000;**12**(46):27-42.
14. Dimet-Wiley A, Golovko G, Watowich SJ. One-Year Postfracture Mortality Rate in Older Adults With Hip Fractures Relative to Other Lower Extremity Fractures: Retrospective Cohort Study. *JMIR Aging*. 2022;**5**(1). e32683. [PubMed ID: 35293865]. [PubMed Central ID: PMC8968577]. <https://doi.org/10.2196/32683>.
15. Raichandani K, Agarwal S, Jain H, Bharwani N. Mortality profile after 2 years of hip fractures in elderly patients treated with early surgery. *J Clin Orthop Trauma*. 2021;**18**:1-5. [PubMed ID: 33936949]. [PubMed Central ID: PMC8079428]. <https://doi.org/10.1016/j.jcot.2021.04.009>.
16. Hokby A, Reimers A, Laflamme L. Hip fractures among older people: do marital status and type of residence matter? *Public Health*. 2003;**117**(3):196-201. [PubMed ID: 12825470]. [https://doi.org/10.1016/S0033-3506\(03\)00033-7](https://doi.org/10.1016/S0033-3506(03)00033-7).
17. Nobili A, Garattini S, Mannucci PM. Multiple diseases and polypharmacy in the elderly: challenges for the internist of the third millennium. *J Comorb*. 2011;**1**:28-44. [PubMed ID: 29090134]. [PubMed Central ID: PMC5556419]. <https://doi.org/10.15256/joc.2011.1.4>.
18. Erlenwein J, Muller M, Falla D, Przemek M, Pflingsten M, Budde S, et al. Clinical relevance of persistent postoperative pain after total hip replacement - a prospective observational cohort study. *J Pain Res*. 2017;**10**:2183-93. [PubMed ID: 28919814]. [PubMed Central ID: PMC5593409]. <https://doi.org/10.2147/JPR.S137892>.
19. Morrison SR, Magaziner J, McLaughlin MA, Orosz G, Silberzweig SB, Koval KJ, et al. The impact of post-operative pain on outcomes following hip fracture. *Pain*. 2003;**103**(3):303-11. [PubMed ID: 12791436]. [https://doi.org/10.1016/S0304-3959\(02\)00458-X](https://doi.org/10.1016/S0304-3959(02)00458-X).
20. Peeters CM, Visser E, Van de Ree CL, Gosens T, Den Ouden BL, De Vries J. Quality of life after hip fracture in the elderly: A systematic literature review. *Injury*. 2016;**47**(7):1369-82. [PubMed ID: 27178770]. <https://doi.org/10.1016/j.injury.2016.04.018>.
21. Shyu YI, Chen MC, Liang J, Lu JF, Wu CC, Su JY. Changes in quality of life among elderly patients with hip fracture in Taiwan. *Osteoporos Int*. 2004;**15**(2):95-102. [PubMed ID: 14605800]. <https://doi.org/10.1007/s00198-003-1533-x>.
22. Adachi JD, Adami S, Gehlbach S, Anderson Jr FA, Boonen S, Chapurlat RD, et al. Impact of prevalent fractures on quality of life: baseline results from the global longitudinal study of osteoporosis in women. *Mayo Clinic Proceedings*. Elsevier; 2010. p. 806-13.
23. Mendonca TM, Silva CH, Canto RS, Morales NM, Pinto RM, Morales RR. Evaluation of the health-related quality of life in elderly patients according to the type of hip fracture: femoral neck or trochanteric. *Clinics (Sao Paulo)*. 2008;**63**(5):607-12. [PubMed ID: 18925319]. [PubMed Central ID: PMC2664717]. <https://doi.org/10.1590/s1807-59322008000500007>.
24. Nussenzweig TC. Pain management after total joint replacement and its impact on patient outcomes. *AORN J*. 1999;**70**(6):1060-2. [PubMed ID: 10635429]. [https://doi.org/10.1016/s0001-2092\(06\)62213-8](https://doi.org/10.1016/s0001-2092(06)62213-8).
25. Romagnoli KM, Handler SM, Hochheiser H. Home care: more than just a visiting nurse. *BMJ Qual Saf*. 2013;**22**(12):972-4. [PubMed ID: 23940375]. [PubMed Central ID: PMC4120108]. <https://doi.org/10.1136/bmjqs-2013-002339>.
26. Su SF, Lin SN, Chen CS. Self-Efficacy Care Program for Older Adults Receiving Hip-Fracture Surgery. *Clin Nurs Res*. 2021;**30**(6):911-20. [PubMed ID: 33736482]. <https://doi.org/10.1177/10547738211001486>.
27. Banappagoudar SB, Parveen A, Prabhudas S, Madhappan A, James S. Nursing Intervention for Improving Physical and Psychosocial Functioning after Hip Fracture in Older People. *J ReAttach Ther Dev Diversities*. 2023;**6**(10s):452-7.

Table 2. Distribution of the Groups' Responses to Oxford Hip Score Before (1st Month) and After (3rd Month) Training ^{a, b}

Variables	Control	Experimental	Text Values
Hip pain			
1st month			$\chi^2 = 11.407; P = 0.003$
Mild	0 (0.0)	6 (20.0)	
Moderate	11 (36.7)	16 (53.3)	
Severe	19 (63.3)	8 (26.7)	
3rd month			$\chi^2 = 19.576; P = 0.000$
Very mild	0 (0.0)	7 (23.3)	
Mild	7 (23.3)	15 (50.0)	
Moderate	16 (53.3)	8 (26.7)	
Severe	7 (23.3)	0 (0.0)	
Washing			
1st month			$\chi^2 = 0.326; P = 0.850$
Moderate trouble	4 (13.3)	3 (10.0)	
Extreme difficulty	13 (43.3)	15 (50.0)	
Impossible to do	13 (43.3)	12 (40.0)	
3rd month			$\chi^2 = 37.842; P = 0.000$
Very little trouble	0 (0.0)	13 (43.3)	
Moderate trouble	5 (16.7)	15 (50.0)	
Extreme difficulty	17 (56.7)	2 (6.7)	
Impossible to do	8 (26.7)	0 (0.0)	
Public transport			
1st month			$\chi^2 = 0.601; P = 0.303$
Extreme difficulty	14 (46.7)	17 (56.7)	
Impossible to do	16 (53.3)	13 (43.3)	
3rd month			$\chi^2 = 39.636; P = 0.000$
With little trouble	0 (0.0)	16 (53.3)	
Moderate trouble	8 (26.7)	14 (46.7)	
Extreme difficulty	16 (53.3)	0 (0.0)	
Impossible to do	6 (20.0)	0 (0.0)	
Wearing socks			
1st month			$\chi^2 = 1.792; P = 0.408$
With moderate difficulty	1 (3.3)	3 (10.0)	
With extreme difficulty	15 (50.0)	17 (56.7)	
Not impossible	14 (46.7)	10 (33.3)	
3rd month			$\chi^2 = 39.429; P = 0.000$
With little difficulty	0 (0.0)	8 (26.7)	
With moderate difficulty	4 (13.3)	20 (66.7)	
With extreme difficulty	19 (63.3)	2 (6.7)	
Not impossible	7 (23.3)	0 (0.0)	
Household shopping			
1st month			$\chi^2 = 1.181; P = 0.554$
With moderate difficulty	6 (20.0)	3 (10.0)	
With extreme difficulty	14 (46.7)	16 (53.3)	
Not impossible	10 (33.3)	11 (36.7)	
3rd month			$\chi^2 = 38.222; P = 0.000$
With little difficulty	0 (0.0)	11 (36.7)	
With moderate difficulty	6 (20.0)	18 (60.0)	
With extreme difficulty	17 (56.7)	1 (3.3)	
Not impossible	7 (23.3)	0 (0.0)	
Walking			
1st month			$\chi^2 = 9.291; P = 0.026$
16 - 30 mins	3 (10.0)	4 (13.3)	
5 - 15 mins	8 (26.7)	10 (33.3)	
Around the house only	11 (36.7)	16 (53.3)	
Not at all	8 (26.7)	0 (0.0)	
3rd month			$\chi^2 = 8.953; P = 0.030$
16 - 30 mins	3 (10.0)	11 (36.7)	
5 - 15 mins	10 (33.3)	12 (40.0)	
Around the house only	14 (46.7)	6 (20.0)	
Not at all	3 (10.0)	1 (3.3)	
Flat of stairs			
1st month			$\chi^2 = 3.606; P = 0.165$
With moderate difficulty	9 (30.0)	7 (23.3)	
With extreme difficulty	11 (36.7)	18 (60.0)	
Not impossible	10 (33.3)	5 (16.7)	
3rd month			$\chi^2 = 41.800; P = 0.000$
Yes, easily	0 (0.0)	5 (16.7)	
With little difficulty	0 (0.0)	18 (60.0)	
With moderate difficulty	13 (43.3)	7 (23.3)	
With extreme difficulty	17 (56.7)	0 (0.0)	
Standup from a chair			
1st month			$\chi^2 = 1.927; P = 0.382$
Moderately painful	4 (13.3)	5 (16.7)	
Very painful	14 (46.7)	18 (60.0)	
Unbearable	12 (40.0)	7 (23.3)	
3rd month			$\chi^2 = 29.032; P = 0.000$
Slightly painful	0 (0.0)	15 (50.0)	
Moderately painful	16 (53.3)	15 (50.0)	

Variables	Control	Experimental	Text Values
Very painful	14 (46.7)	0 (0.0)	
Limping			
1st month			$\chi^2 = 5.573; P = 0.062$
Often not just at first	0 (0.0)	5 (16.7)	
Most of the time	17 (56.7)	13 (43.3)	
All of the time	13 (43.3)	12 (40.0)	
3rd month			$\chi^2 = 43.579; P = 0.000$
Rarely/never	0 (0.0)	3 (10.0)	
Sometimes or just at first	0 (0.0)	14 (46.7)	
Often not just at first	6 (20.0)	13 (43.3)	
Most of the time	15 (50.0)	0 (0.0)	
All of the time	9 (30.0)	0 (0.0)	
Sudden, severe pain			
1st month			$\chi^2 = 0.268; P = 0.398$
Most days	13 (43.3)	15 (50.0)	
Every day	17 (56.7)	15 (50.0)	
3rd month			$\chi^2 = 49.714; P = 0.000$
Only 1 or 2 days	0 (0.0)	10 (33.3)	
Some days	0 (0.0)	17 (56.7)	
Most days	18 (60.0)	3 (10.0)	
Every day	12 (40.0)	0 (0.0)	
Usual work			
1st month			$\chi^2 = 9.333; P = 0.009$
Moderately	0 (0.0)	6 (20.0)	
Greatly	12 (40.0)	15 (50.0)	
Totally	18 (60.0)	9 (30.0)	
3rd month			$\chi^2 = 26.275; P = 0.000$
A little bit	0 (0.0)	1 (3.3)	
Moderately	0 (0.0)	15 (50.0)	
Greatly	15 (50.0)	12 (40.0)	
Totally	15 (50.0)	2 (6.7)	
Hip pain in bed			
1st month			$\chi^2 = 10.982; P = 0.004$
Some nights	0 (0.0)	7 (23.3)	
Most nights	15 (50.0)	17 (56.7)	
Every night	15 (50.0)	6 (20.0)	
3rd month			$\chi^2 = 38.667; P = 0.000$
Only 1 or 2 nights	0 (0.0)	14 (46.7)	
Some nights	8 (26.7)	16 (53.3)	

^a Values are expressed as No (%).

^b Man Whitney U test.