Published online 2023 August 26.

Research Article

Relief of Morphine Withdrawal Symptoms by Japanese Sake Yeast Supplement (*Saccharomyces cerevisiae sake*) Through Adenosine A₁ Receptor Activation in Mice

Hooman Bozorgi ¹, Ali Rashidy Pour ^{2,*}, Nasrollah Moradikor ³, Hamed Darbanian ⁴, Amirmehdi Tayyebi Sereshki ⁴, Seyed Ali Seyedinia ⁵, Parnia Tarahomi ⁵, Mahdi Saghiri Ganjeh ⁵, Azita Alasvand Zarasvand ⁶ and Parham Zamani ^{7,**}

¹Department of Pharmacology, Research Center of Physiology, Semnan University of Medical Sciences, Semnan, Iran

²Research Center of Physiology, Semnan University of Medical Sciences, Semnan, Iran

³International Center for Neuroscience Research, Institute for Intelligent Research, Tbilisi, Georgia

⁴School of Paramedical Sciences, Semnan University of Medical Sciences, Semnan, Iran
⁵School of Medicine, Semnan University of Medical Sciences, Semnan, Iran

⁶Department of Bioinformatics and Genomics, University of North Carolina, Charlotte, USA

⁷School of Pharmacy, Tehran University of Medical Sciences, Tehran, Iran

, Corresponding author: Research Center of Physiology, Semnan University of Medical Sciences, Semnan, Iran. Email: arashidy_pour44@yahoo.com

Received 2023 March 01; **Revised** 2023 July 16; **Accepted** 2023 July 17.

Abstract

Background: Morphine withdrawal syndrome is often treated with chemical drugs; however, these drugs can have concerning side effects. Some natural substances might offer safer alternatives for managing withdrawal symptoms. The neuroprotective activity of the adenosine A₁ receptor (A₁R) in the central nervous system (CNS) has been mentioned before. As a novel natural agent, Japanese sake yeast is enriched with adenosine analogs.

Objectives: As the first report, the present study was designed to evaluate the effects of the Japanese sake yeast supplement against the excitatory signs of morphine withdrawal syndrome in mice.

Methods: Mice were treated orally (gavage) with sake yeast at doses of 100, 200, and 300 mg kg⁻¹ once daily for a week. Furthermore, during the last 3 days of receiving sake yeast, the animals were made physically dependent on morphine by being given twice-daily subcutaneous injections of increasing doses of morphine (30 - 90 mg/kg) for 3 consecutive days. The withdrawal syndrome was induced in animals through the intraperitoneal (i.p.) administration of naloxone (3 mg kg⁻¹) or was elicited spontaneously. After the examination of withdrawal signs, the animals were decapitated, and their brains were prepared for histopathology.

Results: Sake yeast alleviated the intensity of ptosis, piloerection, diarrhea, irritability, and tremor during the two withdrawal syndrome protocols (P < 0.05) and diminished the number of jumping in naloxone-induced withdrawal syndrome (P < 0.01). The withdrawal symptom-lowering effect of sake yeast was reversed by the i.p. injection of an AIR-selective antagonist, 8-cyclopentyltheophylline (10 mg kg⁻¹), and maintained following the i.p. injection of ZM241385 (15 mg kg⁻¹), a selective adenosine A2AR antagonist. Furthermore, histopathological findings demonstrated that sake yeast at doses of 200 and 300 mg kg⁻¹ inhibited the CA1 hippocampal neuronal damage induced by withdrawal syndrome (P < 0.05).

Conclusions: Sake yeast suppresses the morphine withdrawal syndrome via the activation of A₁R.

Keywords: Japanese Sake Yeast, Adenosine A₁ Receptor, Morphine Withdrawal Syndrome, Pharmacoresistance, Mice

1. Background

Long-term administration and subsequent discontinuation of opioids, such as morphine, cause physical dependence and withdrawal syndrome, respectively. Withdrawal syndrome leads to neuronal excitotoxicity-related symptoms, such as agitation and seizure, induced by the primary excitatory amino acid neurotransmitter glutamate (i.e., one of the crucial molecular mechanisms involved in the pathophysiology of opioid withdrawal syndrome) (1). N-methyl D-aspartate (NMDA), as the main glutamate receptor, inhibitors have been shown to alleviate the physical dependence and

Copyright © 2023, Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CC BY 4.0) (https://creativecommons.org/licenses/by/4.0/) which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

the symptoms of opioid withdrawal (2), implicating that the maintenance of opioid dependence and withdrawal syndrome is established by glutamatergic transmission, especially via NMDA receptors (3). N-methyl D-aspartate-motivated calcium influx leads to the launching of neuronal injury induced by a biochemical cascade and worsens the cell's conditions (4).

Adenosine (i.e., an organic and purinergic compound that exerts physiological and pharmacological roles via the stimulation of four known adenosine receptor subtypes [A1, A2A, A2B, and A3]), an endogenous inhibitory neuromodulator in the central nervous system (CNS), displays neuroprotective activity that is mainly based on the prevention of glutamate-induced neurotoxicity by the inhibition of its release through the activation of adenosine receptors particularly adenosine A1 receptor $(A_1R)(5, 6)$. Adenosine induces neuroprotection at least via two pathways: (1) The inhibition of the release of excitatory neurotransmitters, such as glutamate-mediated, by the presynaptic activation of A₁Rs linked via G proteins to both Ca^{2+} and K^{+} ion channels (7-9); and (2) the activation of postsynaptic receptors that promote homeostatic potentiation of intracellular Ca^{2+} (10).

Unlike existing common neuroprotective medications, the body's natural production of adenosine is not hampered by mechanisms in the brain that lead to drug resistance. Drug resistance can arise through structural or metabolic changes in the brain of unknown or genetic origin. Increased levels of transporter proteins in the blood-brain barrier and increased levels of proteins, such as P-glycoprotein (P-gp), in astrocytes and blood vessel cells might be nonspecific or adaptive mechanisms that lead to drug resistance. However, these mechanisms do not impact endogenous adenosine (11). Instead, this organic molecule still shows therapeutic effects against pharmacoresistant conditions, such as epilepsy, anxiety, and depression (12).

Japanese sake yeast (*Saccharomyces cerevisiae* or Japanese rice wine) has long been part of Japanese culture. Sake is a unique brewed alcoholic beverage made through a more complex fermentation process than most other alcoholic beverages (13, 14). Orally administered sake yeast contains adenosine analogs that can activate adenosine receptors and show potential neuroprotective effects. For instance, studies have shown that sake yeast supplements can improve sleep quality and promote non-rapid eye movement (NREM) sleep through the activation of adenosine A₂A receptors in both clinical and animal experiments (15, 16). The most significant advantage of Japanese sake yeast supplement as an organic and natural agent, compared to the routine drugs available to treat withdrawal syndrome on the market (e.g., benzodiazepines), is that sake yeast shows no evidence of tolerance, dependence, addiction, or withdrawal symptoms. This suggests that it might have a lower risk of developing pharmacological resistance and more tolerability in patients.

Furthermore, orally administering sake yeast and its active ingredient, S-adenosyl methionine (SAM), might have advantages over injecting synthetic adenosine or adenosine A₁R agonists. The ability to take sake yeast in oral powder or tablet form provides a non-invasive dosage option compared to injections. Not requiring intravenous delivery expands the practicality and convenience of sake yeast as a potential intervention. Finally, this study aimed at the oral administration of this novel natural agent to evaluate its neuroprotective effects on a mouse model of morphine withdrawal syndrome. The present study is the first to investigate the possible neurobehavioral protective effects of Japanese sake yeast against morphine withdrawal syndrome.

2. Objectives

The present study aimed at the neurobehavioral and histopathological evaluation of oral adenosine-enriched Japanese sake yeast in a mouse model of morphine withdrawal syndrome.

3. Methods

3.1. Animals

Male NMRI mice weighing 20 to 30 g were used for the present study. Standard conditions, including temperature, light/dark (LD), food, and water, were maintained the same for all the animals. The adaptation phase to the vivarium was exerted in animals 1 week before starting the experiments. All the experiments were performed in accordance with the European Communities Council Directive of 22 September 2010 (2010/63/EU) and the local FUM Committee for Human and Animal Ethics.

3.2. Drugs

The dried sake yeast powder (GSP6), provided by Dr. Yuki Nagamori from Lion Corporation in Odawara-shi, Japan, was used. The powder was dispersed in the water shortly before use and then administered to the animals by oral gavage at doses of 100, 200, or 300 mg/kg. The sake yeast doses were chosen based on previously published studies (16). Methylcellulose, used as a vehicle for the sake of yeast, and 8-cyclopentyltheophylline (CPT-8) were purchased from Sigma-Aldrich (St Louis, MO, USA). ZM241385 was purchased from Tocris Bioscience (Bristol, UK). Morphine sulfate was purchased from Darou Pakhsh Pharmaceutical MFG Co. (Tehran, Iran) and administered subcutaneously (s.c.). Diazepam was purchased from Sobhan Darou Pharmaceutical Co. (Tehran, Iran). Additionally, CPT-8 and ZM241385 were injected intraperitoneally. Sake, methylcellulose, and diazepam were administered orally.

3.3. Experimental Design and Pharmacological Phase

The mice were administered orally with sake yeast for a week and concurrently in the last 3 days of sake yeast administration. The animals were rendered dependent by gradually increasing doses of morphine sulfate (s.c. twice daily with 12-hour intervals for 3 consecutive days). Administered doses were 30 and 45 mg/kg on the first day, 60 and 90 mg/kg on the second day, and 90 mg/kg on the last day (17). After the continuous intraperitoneal (i.p.) injections of morphine, the withdrawal signs were observed and monitored following the two experimental protocols; firstly, it was by the i.p. administration of naloxone (3 mg/kg), 3 hours after the last dose of morphine (18). Immediately after the injection of naloxone, the animals were housed in a confined wooden box with high vertical walls. The number of jumps was counted, and the other signs, including chattering teeth, irritability, tremors, diarrhea, ptosis, piloerection, genital grooming, squirming posture, and wet dog shaking, were monitored over the 30-minute period (310). Each character was shown with a point given for 10 minutes (scoring range: 0 - 3) (17). Secondly, it was spontaneously, immediately after the sudden cessation of morphine, and the mentioned withdrawal signs were checked in mice for 24 hours after the last dose of morphine. Immediately after recording behavioral data, the brains were removed, dissected, and prepared for in vitro test.

A total of 100 mice (in general, 200 mice were used for both methods of inducing naloxone and spontaneous withdrawal syndrome) were randomly divided into 10 groups. The animals of all the following groups were exposed to 3-day increasing doses of morphine (in all groups equal to n = 10):

(1) Control group (CONT)

(2) Mice were orally administered (VEH) once daily with 0.5% methylcellulose in water (vehicle) for 1 week.

(3) Mice were orally administered diazepam (2.5 mg/kg) daily for 1 week (DZP).

(4), (5), and (6) Three treatment groups of mice were orally administered sake yeast powder in water (100, 200, and 300 mg/kg) (SAKE).

(7) Mice were orally administered 0.5% methylcellulose in water (vehicle) once daily for 1 week and finally injected with a single dose of

8-cyclopentyltheophylline (a selective adenosine A_1R antagonist, 10 mg/kg, i.p.) on day 7 (VEH + CPT).

(8) Mice were orally administered sake yeast powder in water (200 mg/kg) once a day for 1 week and finally injected with a single dose of 8-cyclopentyltheophylline (10 mg/kg, i.p.) on day 7 (SAKE + CPT).

(9) Mice were orally administered 0.5% methylcellulose in water (vehicle) once daily for 1 week and finally injected with a single dose of ZM24138 (a selective adenosine A_2AR antagonist, 15 mg/kg, i.p.) on day 7 (VEH + ZM).

(10) Mice were orally administered sake yeast powder in water (200 mg/kg) once a day for 1 week and finally injected with a single dose of ZM24138 (15 mg/kg, i.p.) (12).

3.4. Histopathological Evaluation

The histopathological changes in the hippocampal CA1 region were examined using Nissl staining. After recording behavioral data from spontaneous morphine-withdrawn animals, the mice were anesthetized using the i.p. administration of chloral hydrate (35 mg/mL) and then perfused with 0.9% saline solution plus 4% paraformaldehyde. Immediately, the brain tissue was dissected, removed, and postfixed at 4°C in 4% paraformaldehyde for 12 hours. The tissue was then placed in paraffin. The paraffin-embedded tissues were washed by running water, underwent graded ethanol dehydration, and then placed in paraffin again. Tissue samples were prepared for Nissl staining after sectioning (4 μ m thick), deparaffinized in xylene, and rehydrated in ethanol (100 - 70%). The sections were washed with phosphate-buffered saline (PBS), then dried at 70°C for 2 hours, immersed in 0.9% cresyl violet (Sigma-Aldrich) for 1 hour at 56°C, rinsed with distilled water, differentiated in 95% alcohol, dehydrated with 70%, 80%, 90%, and 100% ethanol, cleared in xylene, and then mounted with neutral balsam. Eventually, the number of surviving pyramidal cells in the hippocampal CA1 region (per 1 mm length of each section) was counted in a blind manner using a microscope (\times 400).

3.5. Data Analysis

GraphPad Prism software (GraphPad Software Inc., San Diego, CA, USA) was applied for statistical analysis. For counting jumps, the data were expressed as the mean \pm standard error of the mean (SEM), and the statistical analysis was performed using the one-way analysis of variance (ANOVA) followed by Tukey's posthoc test. A P-value < 0.05 was considered a significant result. The Mann-Whitney U test was applied for the analysis of the qualitative withdrawal signs.

4. Results

As shown in Table 1, morphine-treated mice demonstrated a significant increase in the number of jumping after the induction of withdrawal syndrome compared to the intact mice. As shown in Figure 1, similar to 2.5 mg/kg of diazepam (P < 0.001, compared to the control group), sake yeast led to a significant reduction of the number of jumps in naloxone-precipitated withdrawal syndrome in morphine-dependent mice (one-way ANOVA: F 3,36 = 22.83, P < 0.001). For 100, 200, and 300 mg/kg of supplement, this reduction was 6.45%, 38.7%, and 70.96%, with statistical significances reported as P > 0.05, P < 0.01, and P < 0.001, compared to the control, respectively. Moreover, 100 mg/kg of supplement did not show a significant difference compared to the control.

Additionally, 100 mg/kg of sake yeast significantly decreased the intensity of naloxone-precipitated tremor compared to the control group (U-value = 25, P < 0.05). The intensity attenuation of irritability (agitation, U-value = 25), tremor (U-value = 5), diarrhea (U-value = 5), ptosis (U-value = 5), piloerection (U-value = 5), genital grooming (U-value = 0), writhing posture (U-value = 0), and the wet dog shake (U-value = 0) was observed with the administration of 200 mg/kg of sake yeast (P < 0.05), and all of the mentioned withdrawal signs (except for teeth chattering, U-value = 50) were attenuated with 300 mg/kg of the supplement (P < 0.05)(Table 2). Teeth chattering did not show any significant difference between the control and all the sake yeast-treated (lonely) groups (P > 0.05).

In this study, 100 mg/kg of sake yeast showed no significant differences in spontaneous-precipitated withdrawal signs compared to the control group (P > 0.05).The intensity attenuation of irritability (U-value = 0), tremor (U-value = 0), diarrhea (U-value = 0), ptosis (U-value = 0), and piloerection (U-value = 0) was observed by 200 mg/kg (P < 0.05) or 300 mg/kg (P < 0.01) of sake yeast (Table 3). Teeth chattering, genital grooming, writhing posture, and wet dog shake showed no significant difference between the control and the sake yeast-treated groups (P > 0.05). Furthermore, the withdrawal symptoms-attenuating effect of sake yeast was inhibited by the administration of CPT-8 (lonely or in combination with 200 mg/kg of supplement) and maintained following the administration of ZM241385 (Figure 1, Tables 2, and 3).

In the diazepam group (i.e., diazepam-treated morphine-withdrawn mice), the neurons in the hippocampal CA1 region were demonstrated round-shaped, and cytoplasmic intact cells were shown without any distorted morphology. In the control group (i.e., non-treated morphine-withdrawn mice), masses

of dysplastic neurons were scattered; disappearance, irregularity, swelling, and cracking of the neurons were obvious, and the number of Nissl bodies was diminished in the cytoplasm and nuclear pyknosis. The aforementioned signs were attenuated in mice treated with 200 and 300 mg/kg of sake yeast. The number of surviving cells was quantitatively analyzed in each group and is statistically shown in the histogram (Figure 2). The number of surviving neurons increased significantly in the diazepam group (P < 0.01) and by 200 and 300 mg/kg of sake yeast (P < 0.05) compared to the control group. Similar to the previously mentioned results, these effects were also inhibited by the administration of CPT-8 (in combination with 200 mg/kg of the supplement) and maintained following the administration of ZM241385.

5. Discussion

Adenosine receptors and their functions have been shown to be adjusted by chronic opiate treatment (19, 20). The result of the present study revealed that sake yeast supplement (enriched by high concentrations of oral adenosine analogs) exerts neuroprotection, including behavioral and morphological improvement, spontaneous and naloxone-induced morphine in withdrawal syndrome. Reversing and maintaining this effect following CPT-8 and ZM241385 injections, respectively, demonstrated that sake yeast exerted a withdrawal symptoms-lowering effect predominantly by the activation of A₁Rs that is consistent with the relevant previous findings; the adenosine A₁R selective agonists, N6-cyclohexyladenosine (CHA) and N6-phenylisopropyladenosine decrease jumping diarrhea in naloxone-precipitated morphine and withdrawal symptoms in mice (19). Subsequently, jumping increased; nevertheless, diarrhea decreased with the administration of the adenosine A₁R antagonist, 8-cyclopentyl-1,3-dipropylxanthine (DPCPX).

The adenosine A₂R agonist, 5'-(N-cyclopropyl)-carboxamidoadenosine (CPCA), decreases jumping and diarrhea, and another adenosine antagonist, 3,7-dimethyl-1-propargylxanthine A_2R (DMPX), did not elicit any response in this respect. 8-Cyclopentyl-1,3-dipropylxanthine decreases the inhibition of jumping and diarrhea induced by CHA; however, DMPX decreases the inhibition of diarrhea induced by CPCA (19). The aforementioned findings indicate that jumping induced by naloxone in morphine-dependent mice might be modified by the adenosine A₁R mechanisms, and diarrhea induced by the opioid receptor antagonist could be mediated by both adenosine A1Rs and A2Rs. The local administration of a

Table 1. A Pilot Study for Precision of 3-Day Period of Addiction Protocol in Mice ^{a, b}						
Groups/Parameters	Weight (g)	Number of Jumping				
Intact	23 ± 3.22	0				
Morphine-treated mice and withdrawn by naloxone	23.4 ± 3.1	19 ± 2.34 **				
Morphine-treated mice and withdrawn spontaneously	22.8 ± 2.61	3± 0.74 *				

^a The data were expressed as mean ± standard error of the mean (SEM). Statistically significant differences are reported from the intact mice (*P < 0.05, **P < 0.01). ^b In all groups, n = 10.



Figure 1. Effect of sake yeast (mg/kg) on the number of jumps during the naloxone-precipitated morphine withdrawal in mice. The data were expressed as mean \pm standard error of the mean (SEM). Statistically significant differences are reported from the control group (** P < 0.01, ***P < 0.001) or from the diazepam group (#P < 0.05, ##P < 0.01). In all groups, n = 10, CPT-8: 8-cyclopentyltheophylline; ZM: ZM241385

selective adenosine A_1R agonist, N6-cyclopentyladenosine (CPA), into the nucleus accumbens decreased the Gellert-Holtzman withdrawal scale and attenuated opioid withdrawal symptoms in a dose-dependent manner (21). A_2ARs are required to induce quasi-morphine withdrawal syndrome by the co-administration of caffeine and naloxone (22). The adenosine A_1R agonist R-N6(phenylisopropyl) adenosine significantly reduces wet dog shakes and diarrhea; nonetheless, the adenosine A_1R antagonist DPCPX significantly increases weight loss (19).

The inhibition of the N-type calcium channels via the opioid receptors is considered the underlying mechanism of morphine (23). The long-term consumption of opioids, such as morphine, leads to a decrease in calcium influx

into the neurons; following that, the release of glutamate would be diminished, and its NMDA receptor would be upregulated. When morphine consumption is ceased, the calcium channels change their status to the open state. The excessive calcium influx releases a large amount of the main excitatory neurotransmitter (glutamate). Eventually, neuronal cell death occurs and causes a withdrawal syndrome (24, 25).

On the other hand, the neurological effects of sake can be induced by its adenosine analogs. Adenosine, known as a neuromodulator, has biphasic excitatory and inhibitory effects on neurotransmission, acting via the inhibitory adenosine A_1R and the less common facilitating A_2AR . It is well known that adenosine receptors play an important role in neuroprotection as they inhibit



Figure 2. Nissl staining of the hippocampal CA1 (A) pyramidal neurons with cresyl violet (\times 400 Magnification); (a), control; (b), diazepam; (c), sake 200; (d), sake 300; (e), CPT-8 + sake 200; (f), ZM + sake 200; Bar 20 μ m; quantitative analysis of the expression of surviving neurons in the hippocampal CA1 (B). The data were expressed as mean \pm standard error of the mean (SEM). Statistically significant differences are reported from the control group (* P < 0.05, **P < 0.01) or from the diazepam group (#P < 0.05). In all groups, n = 10. CPT-8: 8-cyclopentyltheophylline; ZM: ZM241385

Groups/Paramete	ers Teeth Chattering	Ptosis	Piloerection	Diarrhea	Irritability	Tremor	Genital Grooming	Writhing Posture	Wet Dog Shake
Control	1(0-3)	2(0-3)	2(0-3)	2 (0 - 3)	3(0-3)	3 (0 - 3)	1(0-3)	1(0-3)	1(0-3)
Vehicle	1(0-3)	3(0-3)	2(0-3)	2(0-3)	3(0-3)	2(0-3)	1(0-3)	1(0-3)	1(0-3)
Diazepam	0 (0 - 3)*	0 (0 - 3) **	0 (0 - 3) **	0 (0 - 3) **	1(0-3)**	0 (0 - 3) **	0(0-3)*	0(0-3)*	0 (0 - 3)*
Sake 100	1(0-3)	2(0-3)	2(0-3)	2(0-3)	3(0-3)	2(0-3)*	1(0-3)	1(0-3)	1(0-3)
Sake 200	1(0-3)	0 (0 - 3) **	0 (0 - 3) **	0 (0 - 3) **	2(0-3)*	1(0-3)**	0(0-3)*	0(0-3)*	0 (0 - 3)*
Sake 300	1(0-3)	0 (0 - 3) **	0 (0 - 3) **	0 (0 - 3) **	0 (0 - 3) **	1(0-3)**	0(0-3)*	0 (0 - 3)*	0 (0 - 3) *
CPT-8 + Sake 200	1(0-3)	2(0-3)	2 (0 - 3)	2 (0 - 3)	2(0-3)*	3(0-3)	1(0-3)	1(0-3)	1(0-3)
ZM + Sake 200	0 (0 - 3)*	1(0-3)*	0 (0 - 3) **	1(0-3)*	1(0-3)**	1(0-3)**	0(0-3)*	0(0-3)*	0 (0 - 3)*
CPT-8	1(0-3)	2(0-3)	2(0-3)	2(0-3)	3(0-3)	2(0-3)*	1(0-3)	1(0-3)	1(0-3)
ZM	0 (0 - 3)*	0 (0 - 3) **	0 (0 - 3) **	1(0-3)*	1(0-3)**	1(0-3)**	0(0-3)*	1(0-3)	0 (0 - 3) *

 a Withdrawal signs were monitored over 3 imes 10 min periods, and the scores were given to the animals (0 - 3). The data were expressed as mean \pm standard error of the mean (SEM). Statistically significant differences are reported from the control group (*P < 0.05, **P < 0.01) using the Mann-Whitney U test.

^b In all groups, n = 10.

^c CPT-8: 8-cyclopentyltheophylline; ZM: ZM241385

Table 3. Effects of Sake Yeast on Spontaneous Morphine Withdrawal Signs in Mice^{a,b,c}

Groups/Paramete	ers Teeth Chattering	Ptosis	Piloerection	Diarrhea	Irritability	Tremor	Genital Grooming	Writhing Posture	Wet Dog Shake
Control	1(0-3)	2(0-3)	2(0-3)	2 (0 - 3)	2(0-3)	2(0-3)	0 (0 - 3)	0 (0 - 3)	0 (0 - 3)
Vehicle	1(0-3)	2(0-3)	2(0-3)	2 (0 - 3)	2(0-3)	2(0-3)	0 (0 - 3)	0(0-3)	0 (0 - 3)
Diazepam	0 (0 - 3)*	0 (0 - 3) **	0 (0 - 3) **	0 (0 - 3) **	0(0-3)**	0 (0-3)**	0(0-3)	0(0-3)	0 (0 - 3)
Sake 100	1(0-3)	2(0-3)	2(0-3)	2 (0 - 3)	2(0-3)	2(0-3)	0(0-3)	0 (0 - 3)	0 (0 - 3)
Sake 200	1(0-3)	1(0-3)*	1(0-3)*	1(0-3)*	1(0-3)*	1(0-3)*	0 (0 - 3)	0(0-3)	0 (0 - 3)
Sake 300	1(0-3)	0 (0 - 3) **	0 (0 - 3) **	0 (0 - 3) **	0 (0 - 3) **	0 (0 - 3) **	0(0-3)	0 (0 - 3)	0 (0 - 3)
CPT-8 + Sake 200	1(0-3)	2(0-3)	2(0-3)	2 (0 - 3)	2(0-3)	2(0-3)	0 (0 - 3)	0 (0 - 3)	0 (0 - 3)
ZM + Sake 200	1(0-3)	1(0-3)*	1(0-3)*	1(0-3)*	1(0-3)*	1(0-3)*	0 (0 - 3)	0(0-3)	0 (0 - 3)
CPT-8	1(0-3)	2(0-3)	2(0-3)	2 (0 - 3)	2(0-3)	2(0-3)	0 (0 - 3)	0(0-3)	0 (0 - 3)
ZM	1(0-3)	1(0-3)*	1(0-3)*	1(0-3)*	1(0-3)*	1(0-3)*	0(0-3)	0(0-3)	0 (0 - 3

^a The signs were evaluated 24 hours after the last dose of morphine, and the scores were given to the animals (0 - 3). The data were expressed as mean ± standard error 100 spin rest valuated at hours are that use of the mean (SEM). Statistically significant differences are reported from the control group (*P < 0.05, **P < 0.01) using the Mann-Whitney U test. In all groups, n = 10.

^c CPT-8: 8-cyclopentyltheophylline; ZM: ZM241385

glutamate release and hyperpolarize neurons (26). Adenosine mainly inhibits intraamygdalar circuits through A_1R activation. The activation of A_1R by the selective A1R agonist CPA decreased NMDAR-mediated currents, and the activation of A2AR with a selective A2AR agonist, CGS21680 ([2-p-[2 carboxyethyl] phenethylamino-5¢-N-ethylcarboxamidoadenosine

hydrochloride hydrate [CGS21680]) also decreased NMDAR-mediated synaptic currents in the amygdala. This effect inhibited by the A₁R antagonist, DPCPX, rather than by the selective A2AR antagonist, 2-(2-furanyl)-7-(2-phenylethyl)-7H-pyrazolo[4,3-e] [1,2,4] triazolo[1,5-c]pyrimidin-5-amine (SCH58261) (27). Therefore, sake yeast-induced neurobehavioral protection can be attributed to activating the adenosine receptors (mainly A₁Rs) and inhibiting glutamatergic transmission in a withdrawal syndrome.

5.1. Conclusions

Eventually, this study concluded that both adenosine A₁R and A₂AR are involved in opioid dependency and withdrawal syndrome. The activation of A1Rs and blockade of A_2ARs elicit a withdrawal syndrome-attenuating effect. Oral adenosine-enriched sake yeast appears to primarily stimulate A_1Rs . The key benefit of using Japanese sake yeast and its active component SAM, compared to injectable synthetic adenosine and adenosine A_1R agonists, is the availability of non-invasive oral forms, such as powders or tablets. Additionally, this natural compound might have lower drug resistance and improved tolerance in patients.

Further research is needed to determine the optimal dosage of sake yeast that provides maximum neurobehavioral protection with minimal CNS depression. Although the authors of this study previously reported the memory-enhancing (28), anxiolytic, and antidepressant-like effects (29) of sake yeast, the present preclinical data provide basic evidence for the clinical use of sake yeast in patients suffering from morphine withdrawal syndrome. Therefore, additional physical activity (30), the stimulation of adenosine receptors by natural compounds, such as sake yeast, would be useful for neurobehavioral problems.

Footnotes

Authors' Contribution: Study concept and design: H. B., A. R., and P. Z. Acquisition of the data: SAS, PT, MSG, and AR. Analysis and interpretation of the data: H. B., N. M., and P. Z. Drafting of the manuscript: A. T. S. and H. D. Critical revision of the manuscript for important intellectual content: H. B. and A. A. Z. Statistical analysis: HB, A. A. Z., and P. Z. Administrative, technical, and material support: H. B. Study supervision: H. B. and H. D.

Conflict of Interests: We declared that two of our authors (Hooman Bozorgi and Ali Rashidy Pour) are of the editorial board. The journal confirmed that the authors with CoI were excluded from all review processes.

Data Reproducibility: The dataset presented in the study is available on request from the corresponding author during submission or after publication. The data are not publicly available.

Ethical Approval: This study was approved under the ethical approval code of IR (approval ID: IR.SEMUMS.REC.1400.090.

Funding/Support: The present study did not receive any grants from funding agencies in the public or commercial sectors. The authors are grateful to Dr. Yuki Nagamori (Lion Corporation, Odawara-shi, Japan) and the Research Center of Physiology of Semnan University of Medical Sciences for their cooperation.

References

1. O'Brien CP. Drug addiction and drug abuse. *Goodman and Gilman's the pharmacological basis of therapeutics*. 2006;**11**:607-27.

- Tanganelli S, Antonelli T, Morari M, Bianchi C, Beani L. Glutamate antagonists prevent morphine withdrawal in mice and guinea pigs. *Neurosci Lett.* 1991;**122**(2):270–2. [PubMed ID: 2027527]. https://doi.org/ 10.1016/0304-3940(91)90875-t.
- Bisaga A, Comer SD, Ward AS, Popik P, Kleber HD, Fischman MW. The NMDA antagonist memantine attenuates the expression of opioid physical dependence in humans. *Psychopharmacology* (*Berl*). 2001;157(1):1–10. [PubMed ID: 11512037]. https://doi.org/10.1007/ s002130100739.
- Rubio M, Villain H, Docagne F, Roussel BD, Ramos JA, Vivien D, et al. Pharmacological activation/inhibition of the cannabinoid system affects alcohol withdrawal-induced neuronal hypersensitivity to excitotoxic insults. *PLoS One*. 2011;6(8). e23690. [PubMed ID: 21886913]. [PubMed Central ID: PMC3158793]. https://doi.org/10.1371/journal. pone.0023690.
- Barrie AP, Nicholls DG. Adenosine AI receptor inhibition of glutamate exocytosis and protein kinase C-mediated decoupling. J Neurochem. 1993;60(3):1081-6. [PubMed ID: 8094742]. https://doi.org/10.1111/j.1471-4159.1993.tb03257.x.
- Masino SA, Kawamura M, Ruskin DN. Adenosine receptors and epilepsy: Current evidence and future potential. *Int Rev Neurobiol.* 2014;**119**:233–55. [PubMed ID: 25175969]. [PubMed Central ID: PMC6026023]. https://doi.org/10.1016/B978-0-12-801022-8.00011-8.
- Wheeler DB, Randall A, Tsien RW. Roles of N-type and Q-type Ca2+ channels in supporting hippocampal synaptic transmission. *Science*. 1994;264(5155):107-11. [PubMed ID: 7832825]. https://doi.org/ 10.1126/science.7832825.
- Palmer TM, Stiles GL. Adenosine receptors. *Neuropharmacology*. 1995;34(7):683–94. [PubMed ID: 8532135]. https://doi.org/10.1016/0028-3908(95)00044-7.
- Fredholm BB. Adenosine and neuroprotection. Int Rev Neurobiol. 1997;40:259–80. [PubMed ID: 8989624].
- Rudolphi KA, Schubert P, Parkinson FE, Fredholm BB. Neuroprotective role of adenosine in cerebral ischaemia. *Trends Pharmacol Sci.* 1992;**13**(12):439–45. [PubMed ID: 1293870]. https://doi.org/10.1016/0165-6147(92)90141-r.
- Fattorusso A, Matricardi S, Mencaroni E, Dell'Isola GB, Di Cara G, Striano P, et al. The pharmacoresistant epilepsy: An overview on existant and new emerging therapies. *Front Neurol*. 2021;12:674483. [PubMed ID: 34239494]. [PubMed Central ID: PMC8258148]. https:// doi.org/10.3389/fneur.2021.674483.
- Klaft ZJ, Hollnagel JO, Salar S, Caliskan G, Schulz SB, Schneider UC, et al. Adenosine A1 receptor-mediated suppression of carbamazepine-resistant seizure-like events in human neocortical slices. *Epilepsia*. 2016;**57**(5):746–56. [PubMed ID: 27087530]. https://doi.org/10.1111/epi.13360.
- Tadenuma M. [Concerning the ingredients of Shea seeds. Modern Science (In Japan Nese)] [Sheishu no seibun wo megutte. Gendai Kagaku (in Japanese)]. Gendai Kagaku. 1987;1:26–31. Japannese.
- Nakahara M, Mishima T, Hayakawa T. Effect of a sake concentrate on the epidermis of aged mice and confirmation of ethyl alpha-D-glucoside as its active component. *Biosci Biotechnol Biochem*. 2007;71(2):427–34. [PubMed ID: 17284832]. https://doi.org/10.1271/bbb. 60489.
- Monoi N, Matsuno A, Nagamori Y, Kimura E, Nakamura Y, Oka K, et al. Japanese sake yeast supplementation improves the quality of sleep: A double-blind randomised controlled clinical trial. J Sleep Res. 2016;25(1):116–23. [PubMed ID: 26354605]. https://doi.org/10.1111/ jsr.12336.
- 16. Nakamura Y, Midorikawa T, Monoi N, Kimura E, Murata-Matsuno A, Sano T, et al. Oral administration of Japanese sake yeast (Saccharomyces cerevisiae sake) promotes non-rapid eye movement sleep in mice via adenosine A(2A) receptors. J

Sleep Res. 2016;**25**(6):746–53. [PubMed ID: 27338238]. https: //doi.org/10.1111/jsr.12434.

- HAJ HV, RABANI MOHAMMAD, ASGHARI GR, KARAMI SZ. Effects of Otostegia persica (Burm.) Boiss on morphine withdrawal syndrome in mice. *Iran J Pharm Res.* 2004.
- Zelena D, Barna I, Mlynarik M, Gupta OP, Jezova D, Makara GB. Stress symptoms induced by repeated morphine withdrawal in comparison to other chronic stress models in mice. *Neuroendocrinology*. 2005;81(3):205–15. [PubMed ID: 16020930]. https://doi.org/10.1159/000087034.
- Kaplan GB, Sears MT. Adenosine receptor agonists attenuate and adenosine receptor antagonists exacerbate opiate withdrawal signs. *Psychopharmacology (Berl)*. 1996;123(1):64–70. [PubMed ID: 8741956]. https://doi.org/10.1007/BF02246282.
- Zarrindast MR, Naghipour B, Roushan-zamir F, Shafaghi B. Effects of adenosine receptor agents on the expression of morphine withdrawal in mice. *Eur J Pharmacol.* 1999;**369**(1):17-22. [PubMed ID: 10204676]. https://doi.org/10.1016/s0014-2999(99) 00021-7.
- Jafarova Demirkapu M, Yananli HR, Kaleli M, Sakalli HE, Goren MZ, Topkara B. The role of adenosine A1 receptors in the nucleus accumbens during morphine withdrawal. *Clin Exp Pharmacol Physiol.* 2020;47(4):553–60. [PubMed ID: 31808190]. https://doi.org/10.1111/1440-1681.13224.
- Bilbao A, Cippitelli A, Martin AB, Granado N, Ortiz O, Bezard E, et al. Absence of quasi-morphine withdrawal syndrome in adenosine A2A receptor knockout mice. *Psychopharmacology (Berl)*. 2006;**185**(2):160-8. [PubMed ID: 16470403]. https://doi.org/10.1007/s00213-005-0284-0.
- 23. Chahl LA. Opioids-mechanisms of action. *Australian Prescriber*. 1996;**19**(3).
- 24. Bozorgi H, Jahanian-Najafabadi A, Rabbani M. Effect of N-type calcium channel blocker Ziconotide on memory in rats following morphine

withdrawal. Online J Veterinary Res. 2016;20(5):335-44.

- Bozorgi H, Motaghi E, Zamani M, Ghavimi R. Neuronal calcium channels blocker, ziconotide (Omega—conotoxin MVIIA), reverses morphine withdrawal-induced memory impairments via alteration in hippocampal NMDA receptor expression in rats. *Toxin Reviews*. 2018;**38**:210–20. https://doi.org/10.1080/15569543.2018.1525402.
- Cunha RA. Neuroprotection by adenosine in the brain: From A(1) receptor activation to A (2A) receptor blockade. *Purinergic Signal.* 2005;1(2):111–34. [PubMed ID: 18404497]. [PubMed Central ID: PMC2096528]. https://doi.org/10.1007/s11302-005-0649-1.
- 27. Pliássova A, Henriques M, Silva HB, Agostinho P, Cunha RA, Ferreira SG. Control of NMDA receptor-mediated currents by adenosine A1 and A2A receptors within the basolateral amygdala. *J Caffeine Adenosine Res.* 2020;**10**(2):61–70.
- Bozorgi H, Rashidy-Pour A, Moradikor N, Motaghi E, Zamani M, Budde T, et al. Reversal of chronic restraint stress-induced memory impairment by Japanese sake yeast supplement in mice: Role of adenosine A(1) and A(2A) receptors. *J Psychiatr Res.* 2023;**161**:123–31. [PubMed ID: 36921500]. https://doi.org/10.1016/j.jpsychires.2023.02. 033.
- 29. Bozorgi H, Rashidy-Pour A, Moradikor N, Zamani M, Motaghi E. Neurobehavioral protective effects of Japanese sake yeast supplement (Saccharomyces cerevisiae) against chronic stress-induced anxiety and depression-like symptoms in mice: Validation of central adenosine receptors as novel therapeutic targets for psychiatric disorders. *Psychopharmacology (Berl)*. 2023;Forthcoming.
- Sani MM, Dambele MY, Raheem S, Akindele MO. The Effect of Physical Activity Level and Body Adiposity on Fatty Infiltration of the Lumbar Multifidus in Apparently Healthy Individuals From an Urban African Setting. *Middle East j rehabil health stud.* 2021;8(1). https://doi.org/10. 5812/mejrh.105506.