Published Online: 2025 June 17

Systematic Review



A Comparative Study of The Efficacy of Photobiomodulation and Electrical Stimulation on Healing of Diabetic Foot Ulcer: A Systematic Review and Meta-Analysis

Parsa Owliaee (b) ^{1, 2}, Soheila Saeedi (b) ^{3, 4}, Mohammadtaha Kouchaki (b) ^{1, 2}, Mohammad Reza Asadi (b) ^{2,*}

¹ Student Research Committee, Hamadan University of Medical Sciences, Hamadan, Iran

- ² Department of Physiotherapy, School of Rehabilitation Sciences, Hamadan University of Medical Sciences, Hamadan, Iran
- ³ Department of Health Information Technology, School of Allied Medical Sciences, Hamadan University of Medical Sciences, Hamadan, Iran
- ⁴ Clinical Research Development Unit of Farshchian Hospital, Hamadan University of Medical Sciences, Hamadan, Iran

* Corresponding Author: Department of Physiotherapy, School of Rehabilitation Sciences, Hamadan University of Medical Sciences, Hamadan, Iran. Email: reza.asadi21@yahoo.com

Received: 31 December, 2024; Revised: 21 May, 2025; Accepted: 25 May, 2025

Abstract

Context: Diabetic foot ulcers (DFUs) represent a prevalent complication of uncontrolled diabetes. Physical modalities like electrical stimulation (ES) and photobiomodulation (PBM) demonstrate significant potential in treating DFUs.

Objectives: This study aimed to systematically review the literature and perform a meta-analysis comparing PBM and ES for reducing DFU size.

Data Sources: We conducted an extensive search for relevant studies by systematically exploring various international databases, including Medline (through PubMed), Web of Science, Cochrane, Scopus, and Pedro. The search was not limited by time, extending until April 29th, 2024, and was restricted to studies published in English.

Study Selection: Two independent researchers participated in the search and screening of sources, and a third researcher resolved any discrepancies in case of conflicting opinions. The initial search yielded 478 articles, of which 225 remained after removing duplicates. After screening titles and abstracts, 31 full-text articles were assessed for eligibility. Ten studies were excluded due to inadequate data, lack of relevant intervention, or being non-randomized or pilot studies.

Data Extraction: Twenty-one studies were included in the systematic review and meta-analysis.

Results: Twenty-one studies were included. The standardized mean change for PBM was 1.79 (95% CI 0.18 - 3.39), and for ES, it was 0.36 (95% CI -0.33 - 1.06). Both PBM and ES significantly reduced DFU size compared to baseline. Although the difference between treatments was not statistically significant (P = 0.095), PBM demonstrated a clinically relevant larger effect size than ES. No adverse effects were reported for PBM, while ES was associated with minor skin irritation or burns in some cases.

Conclusions: Both PBM and ES, as adjuvant treatments, significantly decrease DFU size compared to standard care alone. The PBM's larger effect size suggests potentially greater clinical efficacy, despite non-significant statistical differences. However, further high-quality studies are needed to confirm these findings.

Keywords: Diabetic Foot Ulcer, Electrical Stimulation, Photobiomodulation, Low-Level Laser Therapy, Ulcer Size, Wound Healing

1. Context

Diabetes mellitus poses a significant public health challenge, characterized by its high prevalence, morbidity, and mortality (1). The rising prevalence of diabetes has led to increased complications, including foot ulcers, affecting roughly 15% of diabetic patients, and subsequent amputations, observed in 15 to 20% of patients with such ulcers (2). Diabetic foot ulcers (DFUs) represent a prevalent complication of uncontrolled

Copyright © 2025, Owliaee et al. This open-access article is available under the Creative Commons Attribution 4.0 (CC BY 4.0) International License (https://creativecommons.org/licenses/by/4.0/), which allows for unrestricted use, distribution, and reproduction in any medium, provided that the original work is properly cited.

How to Cite: Owliaee P, Saeedi S, Kouchaki M, Asadi M R. A Comparative Study of The Efficacy of Photobiomodulation and Electrical Stimulation on Healing of Diabetic Foot Ulcer: A Systematic Review and Meta-Analysis. Middle East J Rehabil Health Stud. 2025; 12 (3): e159265. https://doi.org/10.5812/mejrh-159265.

diabetes, often stemming from diabetic peripheral neuropathy, heightened pressure on the foot, vascular issues, and traumatic injuries (3). The DFUs are divided into three categories: Neuropathic (commonly occurring in weight-bearing areas of the foot, such as metatarsal heads and the heel), ischemic (often located in areas around the lateral fifth metatarsal head, medial first metatarsal head, and tips of toes), and neuroischemic (a combination of the first two) (4, 5). The DFUs lead to disability, distress, activity disruption, and significant healthcare costs (6).

While standard treatments for DFUs include various dressings, debridement, offloading, revascularization surgery, and compression therapy (7), many DFUs do not respond to these methods. This lack of response highlights the need for more effective therapeutic approaches to expedite wound healing and reduce the burden of DFUs. Consequently, many studies are exploring therapeutic approaches to expedite wound healing. Modalities like electrical stimulation (ES), ultrasound therapy, low-level laser therapy (LLLT) or photobiomodulation (PBM), and shockwave therapy are being investigated as adjunct treatments for improving chronic wound healing, including DFUs (8, 9). Electrical stimulation and LLLT are the most commonly employed methods, demonstrating significant potential in treating DFUs (10). These modalities offer promising alternatives to standard care, particularly for patients who do not respond to conventional treatments.

It is important to note that patient responses vary depending on ulcer severity, comorbidities, and treatment adherence. Additionally, adjunctive therapies like physical therapy are crucial in managing DFUs by addressing functional limitations, improving mobility, and enhancing patient compliance with offloading strategies. These therapies, when combined with standard care, can further optimize outcomes and reduce the risk of ulcer recurrence. The ES is a safe and economical physiotherapy method that has shown effectiveness in healing DFUs. However, its clinical implementation necessitates staff training for correct electrode positioning and may face obstacles in resource-limited environments due to equipment needs. Similarly, PBM, while non-invasive, demands precise dosing protocols and has higher initial costs for laser devices. Both treatments generally require several sessions, potentially posing challenges for patient adherence.

While the neurophysiological mechanisms behind ES's wound-healing effects remain unclear, research suggests ES enhances wound healing through antibacterial effects (11), increases blood circulation (12),

and promotes cell migration by replicating the body's natural electric currents at the wound site (10, 13). A recent systematic review and meta-analysis by Zheng et al. (14) found that combining ES with standard care leads to accelerated healing and reduced ulcer size in DFUs compared to treatment with routine care alone.

The LLLT, recently known as PBM, is a non-invasive physiotherapy technique that utilizes low-power, monochromatic, and coherent light, typically ranging between 10 and 90 mW/cm², with an intensity of 1 - 4 I/cm^2 when applied to treatment areas (15). Various systematic reviews and network meta-analyses have investigated the effectiveness of LLLT in healing ulcers among individuals with DFUs (16-18). These studies have consistently highlighted the therapeutic advantages of LLLT in diabetic wound healing, including increased ulcer healing rates, reduced ulcer size, enhanced granulation tissue formation, and relief from DFUrelated pain. The precise mechanisms driving the healing effects of LLLT for DFUs remain elusive. However, reports suggest that these effects may be attributed to increased collagen and extracellular matrix synthesis (19, 20), improved local blood circulation (21), heightened ATP production (20), increased growth factor secretion (22), reduced levels of matrix metalloproteinases (MMPs), and decreased activity of inflammatory cytokines (23).

Numerous clinical studies have demonstrated that ES and LLLT expedite the healing of DFUs and exhibit superiority over placebos and standard care. For over five decades, ES devices have been used to facilitate wound healing by utilizing Direct Current and mono and biphasic pulsed current waveforms. In recent years, newer technologies, including bioelectric dressing-like devices, have emerged as potential tools in this endeavor (24).

The LLLT was discovered by Mester et al. in 1968 (25-27). Mester's research showed that LLLT has promising effects on cell function and tissue repair, particularly in enhancing wound healing. Since then, advancements in laser technology, such as light-emitting diodes (LEDs), broadband light using filters, and high-power lasers, have further expanded the potential applications of LLLT in wound management.

2. Objectives

However, despite the extensive use of ES and LLLT in chronic wounds, their effectiveness in treating DFUs remains relatively unexplored. Notably, no clinical trials, published reviews, or meta-analyses have yet compared the effectiveness of ES and LLLT in DFU treatment. This gap in the literature underscores the need for a comprehensive evaluation of these therapies to guide clinical decision-making. Furthermore, the variability in patient responses to these therapies, influenced by factors such as ulcer severity, patient comorbidities, and adherence to treatment protocols, underscores the need for a comprehensive evaluation. This variability highlights the ongoing challenges in the field and emphasizes the importance of synthesizing existing evidence to guide clinical decision-making. Therefore, this systematic review aims to evaluate the relative efficacy of these two therapies for managing DFUs and provide an evidence-based foundation for clinical decision-making in wound management.

3. Data Sources

3.1. Study Design and Registration

This literature review, conducted until April 29, 2024, followed a structured approach involving a systematic review of scientific articles and a meta-analysis, adhering to the Preferred Reporting Items for Systematic Review and Meta-analysis (PRISMA) guidelines (28). This systematic review and metaanalysis was not registered in a public trial registry because it is a secondary analysis of existing studies, and registration is not typically required for systematic reviews and meta-analyses. The research process consisted of several distinct phases. Initially, we formulated the research topic and devised a comprehensive search strategy. Subsequently, we systematically screened and curated relevant articles from various databases. We rigorously assessed these articles based on predefined inclusion and exclusion criteria. Additionally, a qualitative evaluation was performed, and the review culminated in a comprehensive statistical analysis.

3.2. Search Strategy

We conducted an extensive search for relevant studies by systematically exploring various international databases, including Medline (through PubMed), Web of Science, Cochrane, Scopus, and Pedro. Our search was not limited by time, extending until April 29, 2024. Furthermore, an extensive examination of pertinent studies' reference lists was undertaken to identify all extant articles within this domain. The search strategy employed a combination of MeSH terms and keywords such as "DFUs," "Electrical Stimulation," "Photobiomodulation," "Low-Level Laser Therapy," "Low-Power Laser Therapy," "Electric Stimulation Therapy,"

and "Low-Level Laser Therapy." Boolean operators (AND/OR) were used to refine the search results.

3.3. Eligibility Criteria

In this review, we considered studies that explored the impact of PBM and ES on the treatment of DFUs. To identify these relevant studies, we initially assessed their titles and abstracts. After this initial screening, we thoroughly reviewed the full texts of the selected studies. For inclusion in this systematic review and meta-analysis, studies had to meet specific criteria: (1) Randomized clinical trials (RCTs), (2) original articles, (3) participants with DFUs, (4) the treatment of DFUs using PBM or ES, and (5) providing specific data on DFU healing, including initial ulcer size and final ulcer size. We excluded studies that did not meet these criteria, including those with insufficient data, a lack of quantitative reporting on initial or final ulcer size, articles without full texts, animal studies, non-original articles, pilot studies, in vitro studies, letters to editors, protocols, review articles, monographs, and theses/dissertations.

3.4. Data Extraction

In the initial phase of this study, two authors independently reviewed the titles and abstracts of all the retrieved articles. Subsequently, articles that appeared relevant were selected, and their full texts were carefully examined. Pertinent information was then extracted from these articles. Furthermore, the references of these relevant articles were investigated to identify any additional relevant sources. Any discrepancies or differences in judgment between the authors were resolved through discussion with the involvement of the corresponding author. To systematically gather and organize the information from the selected articles, an Excel spreadsheet was designed. The following details were extracted and entered into this spreadsheet: Author names, publication years, study countries, information about the intervention and control groups, initial ulcer sizes, final ulcer sizes, Wagner ulcer grade, average duration of ulcers, the number of treatment sessions, PBM/ES parameters, and the study outcomes. Additionally, a separate Excel sheet was used to assess the quality of the articles based on predefined quality assessment criteria.

3.5. Quality Assessment

The Effective Public Health Practice Project (EPHPP) tool was employed to appraise the methodology of the

selected articles (29). This tool allows for the assessment of six key components.

(1) Study design: Evaluating the structure and design of the study.

(2) Withdrawals and dropouts: Examining the handling of participant withdrawals and dropouts.

(3) Data collection practices: Assessing the methods used for data collection.

(4) Selection bias: Investigating any bias in the selection of participants.

(5) Blinding in controlled trials: Examining whether blinding was appropriately implemented in controlled trials.

(6) Confounders: Analyzing the identification and management of potential confounding variables.

The EPHPP tool proves valuable for evaluating the quality of quantitative studies. Based on the aforementioned criteria, studies are categorized into one of three levels.

(1) Strong: No weak ratings in any of the six aspects.

(2) Moderate: One weak rating in any of the six aspects.

(3) Weak: Two or more weak ratings across the six aspects.

Two authors conducted the quality assessment of the selected articles. After independently completing their assessments, the results were compared, and any discrepancies between the authors were resolved through discussions with SS.

3.6. Data Analysis

Standardized mean change using change score standardization (SMCC Index) was analyzed to assess the changes in wound size before and after the intervention in the two groups. In this study, analyses related to metaanalysis were conducted using a model with random effects. To account for the heterogeneity of the studies, we used a random effects model. The random effects model in meta-analysis is used when the studies included in the analysis are assumed to have different underlying effect sizes, rather than a single common effect size. This model is particularly useful when there is significant heterogeneity among the studies, as it provides a more realistic estimate of the overall effect size by acknowledging differences between studies. In the second step, the multilevel linear mixed-effect model was used, which considered dataset variabilities and heterogeneities in the form of covariances and error terms (30-32). The values of I^2 , τ^2 , and H^2 in the two groups were used to assess heterogeneity between studies. Additionally, we employed Graphical Display of Study Heterogeneity (GOSH) plots to visually inspect study heterogeneity and identify potential outliers within the included studies (33). EndNote X9, a reference management software, was utilized for resource management, and the meta-analysis was conducted using R 4.2.2 software and the metafor package.

4. Results

4.1. The Literature Search

The search results and the number of retrieved articles based on PRISMA are shown in Figure 1. The search resulted in retrieving 478 articles from five databases; after removing duplicates, 225 remained for title and abstract screening. After removing 200 articles that did not meet the inclusion criteria, the full text of 31 articles was examined. Among these, ten references were excluded due to inadequate data, lack of related intervention, or being non-randomized or pilot studies. Finally, 21 articles were included in this systematic review, and data were extracted from them. It is necessary to explain that, considering the keywords used for the search were in English, all retrieved articles had at least English titles. After checking the full text of the articles included in this systematic review, only one article had the full text in a non-English language. Considering that the authors were proficient in that non-English language, this article was included in the systematic review.

4.2. General Characteristics of the Included Studies

The general characteristics of the articles included in this systematic review and meta-analysis are given in Table 1 (Full study characteristics are provided in Appendix 1 in Supplementary File). Out of the 21 articles included in this systematic review, 20 studies were published in journals, and one study was presented at a conference. Of the 16 journals that accepted articles in this field, two journals, "Lasers in Medical Science" and "Photomedicine and Laser Surgery," had the highest acceptance rate of articles in this field with three articles each. The oldest article was published in 1992, and the most recent article was published in 2023. The highest number of published articles was related to 2017 and 2018, with four articles each. The three countries of Iran, India, and Brazil accounted for 61.9% of the published studies, with five, four, and four published articles, respectively. The therapy method was PBM in 15 studies (71.4%), ES in five studies (23.8%), and both ES and PBM in one study (4.8%). Wagner grade was not mentioned in four studies, and in most studies (n = 6, 28.6%), it was



Figure 1. Flow diagram of the literature search and study selection

grade II. Six studies did not mention challenges or limitations. The most mentioned limitations in the studies were "small sample size" (12 studies) and "short study period" (six studies).

4.3. Effects of Photobiomodulation and ES on Wound Size

Meta-analysis was used to investigate changes in wound size before and after the intervention. The characteristics of the studies included in the metaanalysis are given in Table 2.

In each study, the SMCC Index was analyzed to assess the changes in wound size before and after the intervention in the two groups. The overall SMCC value, along with the homogeneity indices of the studies, is reported in Table 3. Based on the Q-test results (P <0.0001 for both groups), significant heterogeneity was observed between the effect sizes of studies in both the PBM and ES groups. Therefore, a random-effects model (RE model) was used to calculate the effect sizes. In the PBM group, the SMCC was 1.79 (95% CI: 0.18, 3.39), indicating a statistically significant reduction in wound size after the intervention (Z statistic = 2.18, P = 0.029). Similarly, the SMCC in the ES group was 0.36 (95% CI: -0.33, 1.06), suggesting a reduction in wound size, though this was not statistically significant (Z statistic = 1.02, P = 0.308).

The values of I^2 , τ^2 , and H^2 in the two groups indicate the presence of high dispersion and heterogeneity between studies, and therefore the obtained SMCC values are not valid. Also, forest diagrams for the PBM and ES groups are presented in Figures 2 and 3.

The GOSH analysis was used to check for outliers in the included studies. After removing studies 1 and 4 from the PBM group and study 4 from the ES group, the

habie i. General characterist	ies of the m	eraded studies					
Authors, Y/Country	Sample Size	Groups	Initial Size (cm ²)	Final Size (cm ²)	Wagner Grade	PBM/ES Parameters	Treatment Time
Zulbaran-Rojas et al, (34) 2023/USA	16;17	ES + PLA	$\begin{array}{c} 7.4 \pm 8.5; 3.1 \pm \\ 5.6 \end{array}$	$5.8 \pm 8.0; 3.2 \pm 8.7$	Not clear I-II (converted)	HVPC; 150 - 250V; ankle electrodes	Home daily for 1 hour for 4 weeks
Wadee et al, (35) 2021/Egypt	25; 25; 25	HBOT + TT + LLLT + TT + TT	$5.48 \pm 0.70; 5.58 \pm 0.72; 5.61 \pm 0.75$	$0.45 \pm 0.16; 0.75 \pm 0.23;$ 5.50 ± 0.75	Ш	GaAlAs laser; 850 nm; 4 J/cm ²	3 sessions (8 min) /week for 6 weeks
Vitoriano et al, (36) 2019/Brazil	6;6	LLLT+TT + LED+TT	$\begin{array}{c} 1.76 \pm 1.69 ; 1.45 \pm \\ 1.52 \end{array}$	$0.36 \pm 0.50; 0.64 \pm 0.81$	I (converted)	830 nm laser; 30 mW; 7 J/cm ²	10 sessions/ 2 weeks
Tantawy et al, (37) 2018/Egypt	33; 32	LLLT + TT + infrared laser therapy + TT	$\begin{array}{c} 10.2\pm 5.6; 9.5\pm \\ 4.2\end{array}$	3.7±1.2; 4.1±1.3	I-II	632 nm laser; 20 mW	90s application/cm ² and the dose of 5 J/cm ² for 8 weeks
Frangez et al, (38) 2018/Slovenia	30;30	LLLT + TT + PLA + TT	13.15; 15.84	7.364;10.296	NA	Wavelengths: 625/660/850 nm; 2.4 J/cm ²	3 sessions/week for 8 weeks (5 min)
Priyadarshini et al, (39) 2018/India	50; 50	LLLT + TT	13.74 ± 11.88; 19.09 ± 15.03	3.97±5.41; 18.80± 17.70	I-II	660nm laser; 4 - 8 J/cm ²	15 day (20 min)
de Alencar Fonseca Santos et al, (40) 2018/Brazil	8;8	LLLT + TT	$\begin{array}{c} 1.83 \pm 1.08; 2.97 \\ \pm 1.66 \end{array}$	0.32±0.26; 1.63±1.57	II-III (converted)	660 nm laser; 6 J/cm ²	4 sessions /week for 4 weeks
Srilestari et al, (41) 2017/Indonesia	18; 18	LLLT + TT + SHAM	4.75(0.10 - 9.94); 2.33 (0.90 - 9.88)	0.24 (0.00 -2.54); 1.25 (0.00 - 8.11)	I-III (converted)	630 nm laser; 100 mW; 4 J/point	2 sessions/week for 4 weeks
Asadi et al, (42)2017/Iran	13; 11	ES + TT + PLA + TT	$\begin{array}{c} 4.19 \pm 2.2; 3.82 \pm \\ 1.7 \end{array}$	$1.83 \pm 1.63; 2.88 \pm 1.51$	II	Cathodal DC; sensory threshold	3 sessions/week for 4 weeks
Mathur et al, (21) 2017/India	15;15	LLLT + TT + TT	14.84; 13.52	9.30; 11.46	I	660 nm laser; 3 J/cm ²	15 day (1 min)
El Rasheed et al, (43) 2017/India	15; 15	PEMF + TT + LLLT+ TT	13.096 ± 5.93; 17.55 ± 12.1	$5.84 \pm 3.63; 2.033 \pm 2.01$	Ш	904 nm laser; 10 J/cm ²	3 sessions (10 min) /week for 4 weeks
Hoseini Sanati et al, (44) 2016/Iran	15;12	LLLT + PLA	4.94 ± 2.91; 4.63 ± 2.34	$1.25 \pm 1.04; 3.30 \pm 1.25$	II	904 nm laser; 2 J/cm ²	3 sessions /week for 4 weeks
Asadi et al, (45) 2015/Iran	10;10	ES + PLA	$\begin{array}{c} 4.05 \pm 2.01; 4.27 \\ \pm 3.2 \end{array}$	1.01± 0.8; 2.6 ±1.1	II	Cathodal DC; sensory threshold	3 sessions/week for 4 weeks
Feitosa et al, (46) 2015/Brazil	8;8	LLLT+ TT + TT	$\begin{array}{c} 7.98 \pm 2.06 ; 2.55 \\ \pm \ 0.77 \end{array}$	$2.39 \pm 1.26; 8.43 \pm 1.84$	NA	632.8 nm laser; 4 J/cm ²	3 sessions (80s) /week for 4 weeks
Mohajeri-Tehrani et al, (47) 2014/Iran	10;10	ESP + LA	$2.48 \pm 0.97; 2.43 \\ \pm 0.39$	1.71 ± 0.66 not clear; 2.18 ± 0.35 not clear	II	Cathodal DC; sensory threshold	3 sessions (1 hour) /week for 4 weeks
Sandova Ortíz et al, (48) 2014/Colombia	9;10;9	ES + LLLT + TT	NA	NA	I-II	685 nm laser; 30 mW / 100 pps ES	3 sessions (45min) /week for 16 weeks or until the wound closed
Kajagar et al, (49) 2012/India	34;34	LLLT+ TT + TT	$26.08 \pm 68.31;$ 27.47 ± 6.03	$\begin{array}{c} 15.64 \pm 43.73 ; 24.24 \pm \\ 5.51 \end{array}$	Ι	Pulsed laser; 5 kHz; 2 - 4 J/cm ²	Daily for 15 days
Kaviani et al, (50) 2011/Iran	13;10	LLLT + TTPLA + TT	10.7 ± 25.7; 7.8 ± 11	73.7±10.2 (reduction %); 47.3±15.4 (reduction %)	I-II	685 nm laser; 10 J/cm ²	6 sessions/week
Minatel et al, (51) 2009/Brazil	7;7	LLLT + TTPLA + TT	$\begin{array}{c} 11.8 \pm 20.46 ; 3.8 \\ \pm 4.13 \end{array}$	NA	NA	Wavelengths: (660 + 890 nm); 3 J/cm ²	2 sessions/week for 90 days
Naidu et al, (52) 2005/Malaysia	8; 8	LLLT + TT	5.28; 4.21	1.31; 3.59	I	He-Ne laser; 1 - 5 J/cm ²	5 sessions/week for 6 weeks
Lundeberg et al, (53) 1992/Sweden	32; 32	ES + TTPLA + TT	$24.2 \pm 12.6; 22 \pm 9.6$	39 ± 14 (remaining %); 59 ± 11 (remaining %)	NA	80 Hz AC; muscle contraction	14 sessions/week for 12 weeks

Table 1. General Characteristics of the Included Studies^a

Abbreviations: PBM, photobiomodulation; ES, electrical stimulation; HVPC, high voltage pulsed current; HBOT, hyperbaric oxygen therapy; TT, traditional treatment; LLLT, low-level laser therapy; DFU, diabetic foot ulcer; LED, light-emitting diodes.
^a Full intervention details are provided in Supplementary Table 1.

results were extracted again and reported in Table 4. The obtained values of I^2 , τ^2 , and H^2 in the two groups indicate the existence of moderate dispersion between the studies, and therefore the obtained SMCC values were valid. Intervention in both the PBM and ES groups

reduced the wound size, and this size change was statistically significant. Forest diagrams for the PBM and ES groups are presented in Figures 4 and 5.

The effect of the BMI variable on the wound size in the groups is reported in Table 5. In the PBM group, although the effect of the BMI variable was not

Fable 2. Studies Included in the Meta-Analysis	
Groups	Number
PBM; author (y)	
Wadee et al. (2021) (35)	25
Hoseini Sanati et al. (2016) (44)	15
Feitosa et al. (2015) (46)	8
Kajagar et al. (2012) (49)	34
Priyadarshini et al. (2018) (39)	50
Santos et al. (2018) (16)	8
ES; author (y)	
Zulbaran-Rojas et al. (2023) (34)	16
Asadi et al. (2017) (42)	13
Asadi et al. (2015) (45)	10
Lundeberg et al. (1992) (53)	32
Mohajeri-Tehrani et al. (2014) (47)	10

Table 3. Standardized Mean Change Using Change Score Standardization Value, Q Statistics, I ² , H ² , and τ^2 Indices in Photobiomodulation and Electrical Stimulation Groups								
Groups	SMCC (95% CI)	Z Statistics (P-Value)	Q Statistics (P-Value)	I^2	H ²	τ^2		
PBM	1.79 (0.18, 3.39)	2.18 (0.029)	56.13 (< 0.0001)	97.84	46.20	3.78		
ES	0.36 (-0.33, 1.06)	1.02 (0.308)	33.12 (< 0.0001)	86.39	7.35	0.53		

Abbreviations: SMCC, standardized mean change using change score standardization; CI, confidence interval; PBM, photobiomodulation; ES, electrical stimulation.

significant (P = 0.937), adding this variable to the model reduced I², τ^2 , and H² values. In the ES group, the effect of the BMI variable was significant (P = 0.017), and adding this variable to the model caused a significant decrease in I², τ^2 , and H² values. To investigate the effect of time (before and after intervention), BMI, and group (PBM and ES) on the wound size mean, multilevel metaanalysis using mixed-effect models was applied.

The results of this analysis are reported in Table 6. According to the results, the wound size mean after the intervention was smaller than before the intervention, and it was statistically significant (P = 0.002). Also, the wound size mean in the PBM group was smaller than in the ES group but wasn't statistically significant (P = 0.095).

4.4. Quality Assessment Results

The quality of the included studies was assessed using the EPHPP tool, which evaluates six key components: Selection bias, study design, confounders, blinding, data collection, and drop-outs. Each component was rated as 1 = strong, 2 = moderate, or 3 = weak, and an overall global rating was assigned to each study. The detailed results of the quality assessment are presented in Table 7.

The results showed that 100% of the articles included in this systematic review were strong in terms of data collection and study design. However, 28.6% of the articles were classified as weak in terms of blinding. Global ratings also showed that 52.4% of the articles were moderate and 38.1% were strong. These findings are summarized in Figure 6.

5. Discussion

5.1. Summary of Findings

This study aimed to compare the effects of PBM and ES on the healing of DFUs. An extensive literature search was conducted across five reputable databases, initially yielding 478 articles. After screening, only 21 studies met the predefined inclusion criteria and were included in this review, highlighting the rigorous methodology utilized. Our analysis found that PBM was the predominant therapeutic intervention, utilized in 15 studies, representing 71.4% of the total. ES was employed in five studies, accounting for 23.8%. One study, comprising 4.8% of the sample, used both PBM and ES.



Figure 2. Forest plot of the photobiomodulation (PBM) group. SMCC, standardized mean change using change score standardization; SD, standard deviation; CI, confidence interval; RE, random effects; df, degrees of freedom (16, 35, 39, 44, 46, 49).

The peak in publications in 2017 and 2018 may reflect a surge in research interest or critical developments during this period. Further investigation into the historical events and breakthroughs in those years that stimulated research output would provide insight. Tracking the progression of research historically can identify emerging areas of focus. The geographical concentration of studies in Iran, India, and Brazil suggests that these countries have substantially contributed to the field. Further investigation into the research ecosystems and collaborations in these regions can shed light on the reasons behind their high research output. This could include examining government funding initiatives, academic institutions, and partnerships with international researchers.

5.2. Efficacy of Photobiomodulation and Electrical Stimulation

The meta-analysis demonstrated that both PBM and ES significantly reduced DFU size compared to baseline. However, there was no statistically significant difference between the treatment effects of PBM and ES for decreasing ulcer area. This indicates that neither modality showed superior efficacy over the other. It should be noted, however, that PBM demonstrated a

more significant effect size and standardized mean change than ES. Although this larger effect was not statistically significant, it may suggest potentially enhanced efficacy of PBM over ES. The PBM can impact various biological pathways linked to wound healing by enhancing cellular activity, improving blood circulation, and boosting growth factor levels, potentially leading to faster reductions in ulcer size (54). additional However, high-quality comparative controlled trials with larger sample sizes are required to further compare the effects of PBM and ES on reducing DFU size over time. The current findings, while promising, must be interpreted cautiously due to heterogeneity in study designs and intervention protocols across the included trials. Nevertheless, the current analysis provides evidence supporting both modalities as effective adjuvant therapies that should be considered along with standard care to enhance wound closure in this population.

5.3. Comparison with Previous Studies

The results of this study align with previous research showing the benefits of LLLT and ES for treating DFUs. A recent systematic review and meta-analysis by Santos et al. found that LLLT significantly reduced ulcer size and



Figure 3. Forest plot of the ES group. SMCC, standardized mean change using change score standardization; SD, standard deviation; CI, confidence interval; RE, random effects; df, degrees of freedom (34, 42, 45, 47, 53).

Groups	SMCC (95% CI)	Z Statistics (P-Value)	Q Statistics (P-Value)	I^2	H^2	τ^2
PBM	1.06 (0.62, 1.50)	4.69 (< 0.0001)	5.39 (0.145)	42.53	1.74	0.09
ES	0.64 (0.17, 1.10)	2.67 (0.008)	6.42 (0.093)	52.64	2.11	0.12

Abbreviations: SMCC, standardized mean change using change score standardization; CI, confidence interval; PBM, photobiomodulation; ES, electrical stimulation.

promoted greater healing rates than control treatments, with a 22.96% greater reduction in ulcer size with LLLT (16). Our findings on the efficacy of ES are supported by a recent meta-analysis by Chen et al. on ES for DFUs. In their analysis of seven randomized controlled trials with 274 patients, ES as an adjunct to standard wound care resulted in a significantly greater decrease in ulcer area at four weeks (standardized mean difference 1.09, 95% CI 0.62 - 1.57, P < 0.001) and a higher ulcer healing rate at 12 weeks (risk difference 0.19, 95% CI 0.06 - 0.32, P = 0.005) compared to standard wound care alone (55).

A previous study by Sandoval Ortiz et al. did not demonstrate additional effects of LLLT or high voltage pulsed current (HVPC) compared to standard wound care alone on the healing of DFUs (48). Their randomized controlled trial found no significant differences between the LLLT, HVPC, and control groups in ulcer healing rates, nerve conduction studies, protective sensation, or quality of life after 16 weeks of treatment. These negative results contradict our findings, which show improved diabetic ulcer healing with LLLT and ES. However, Sandoval Ortiz et al. had a small sample size (n = 28), which may have limited the power to detect group differences. They recommend more extensive trials to be conducted. The advanced neuropathy state of their patients (mean 11 years diabetes duration) may have also reduced treatment effects on the assessed neurological outcomes. Overall, this well-conducted study provides substantial evidence that LLLT and HVPC may not provide added benefit over standard care in treating diabetic ulcers, particularly in



Figure 4. Forest plot of the photobiomodulation (PBM) group after removing outliers. SMCC, standardized mean change using change score standardization; SD, standard deviation; CI, confidence interval; RE, random effects; df, degrees of freedom (16,39,44,46).



Figure 5. Forest plot of the ES group after removing outliers. SMCC, standardized mean change using change score standardization; SD, standard deviation; CI, confidence interval; RE, random effects; df, degrees of freedom (34, 42, 45, 47).

those with severe neuropathy. However, Sandoval Ortiz et al. did not report the initial and final ulcer sizes, precluding the inclusion of their data in a meta-analysis.

5.4. Implications for Clinical Practice

We observed a significant decrease in the size of DFUs when either PBM or ES was used in conjunction with standard care. This suggests that employing these therapies alongside conventional treatment enhances wound area reduction. There was no statistically

Table	F able 5. Meta-Analysis via Random Effects Model with BMI Variable								
Groups PBM		Coefficient (95% CI)	Z Statistics (P-Value)	I² 42.37	H ²	τ^2			
					1.74	0.05			
	Intercept	1.68 (-17.08, 20.44)	0.17 (0.861)						
	BMI	-0.03 (-0.72, 0.664)	-0.08 (0.937)						
ES				0.00	1.00	0.00			
	Intercept	3.68 (1.09, 6.29)	2.78 (0.006)						
	BMI	-0.11 (-0.19, -0.02)	-2.39 (0.017)						

Abbreviations: CI, confidence interval; BMI, Body Mass Index; PBM, photobiomodulation; ES, electrical stimulation.

Fable 6. Meta-Analysis via Multilevel Linear Mixed-Effect Models					
Variables	Coefficient (95% CI)	Z Statistics (P-Value)			
Intercept	-5.84 (-20.48,8.98)	-0.78 (0.433)			
Time (Final)	-3.53 (-5.79, -1.25)	-3.05 (0.002)			
Group (PBM)	-1.78 (-3.85, 0.31)	-1.67(0.095)			
BMI	0.45 (-0.11, 1.01)	1.56 (0.118)			

Abbreviations: CI, confidence interval; BMI, Body Mass Index; PBM, photobiomodulation; ES, electrical stimulation.

Studies	Selection Bias	Study Design	Confounders	Blinding	Data Collection	Drop-outs	Global Rating ^b
Zulbaran-Rojas et al, (34) 2023/USA	1	1	1	1	1	1	1
Wadee et al, (35) 2021/Egypt	3	1	1	2	1	1	2
Vitoriano et al, (36) 2019/Brazil	2	1	2	3	1	1	2
Tantawy et al, (37) 2018/Egypt	3	1	1	2	1	1	2
de Alencar Fonseca Santos et al, (40) 2018/Brazil	2	1	1	3	1	1	2
Priyadarshini et al, (39) 2018/India	1	1	1	3	1	1	2
Frangez et al, (38) 2018/Slovenia	2	1	1	1	1	3	2
Srilestari et al, (41) 2017/Indonesia	2	1	1	1	1	1	1
El Rasheed et al, (43) 2017/India	2	1	1	3	1	1	2
Mathur et al, (21) 2017/India	2	1	1	3	1	1	2
Asadi et al, (42) 2017/Iran	2	1	1	2	1	1	1
Hoseini Sanati et al, (44) 2016/Iran	2	1	3	3	1	2	3
Feitosa et al, (46) 2015/Brazil	3	1	1	2	1	1	2
Asadi et al, (45) 2015/Iran	2	1	1	2	1	1	1
Sandoval Ortíz et al, (48) 2014/Colombia	1	1	1	1	1	3	2
Mohajeri-Tehrani et al, (47) 2014/Iran	2	1	1	1	1	1	1
Kajagar et al, (49) 2012/India	2	1	1	1	1	1	1
Kaviani et al, (50) 2011/Iran	1	1	1	1	1	1	1
Minatel et al, (51) 2009/Brazil	3	1	3	1	1	3	3
Naidu et al, (52) 2005/Malaysia	2	1	3	2	1	1	2
Lundeberg et al, (53) 1992/Sweden	2	1	1	2	1	1	1

 $^{\rm a}$ Quality ratings are based on the Effective Public Health Practice Project (EPHPP) tool.

^b The Global Rating is an overall assessment of study quality based on the EPHPP tool.

significant difference between the treatment effects of

PBM and ES for decreasing ulcer size. The side effect



Figure 6. Results of quality assessment of included articles with Effective Public Health Practice Project (EPHPP) check

profile and risks should be considered when selecting between modalities. None of the studies reported adverse events with PBM treatment, likely due to low power densities avoiding tissue damage. In contrast, ES carries risks of skin irritation or burns with improper parameters. Additionally, PBM typically requires shorter treatment sessions than ES. Therefore, the improved safety profile and shorter treatment durations of PBM should be considered when choosing adjuvant therapies with standard care to improve wound closure.

Our meta-analysis comprehensively assessed the impact of BMI on wound size reduction across PBM and ES treatment groups. The findings revealed intriguing distinctions in treatment efficacy based on patients' BMI. For PBM, our analysis demonstrated that BMI did not significantly affect ulcer area reduction. This suggests that PBM demonstrates consistent therapeutic efficacy across different body weight categories. In contrast, ES exhibited a significant BMI effect, indicating reduced treatment efficacy specifically in obese diabetic patients.

The differential impact of BMI on these therapies highlights the importance of considering patient body composition when selecting treatment protocols. For ES, the significant BMI influence suggests that clinicians should carefully stratify patients and potentially develop tailored interventions for individuals with higher body weight. Supporting this observation, a study by Doheny et al. investigated the impact of subcutaneous fat thickness on ES, revealing that increased fat thickness reduced activation function and necessitated higher electrical currents for effective stimulation (56). This finding aligns closely with our research, where we observed correlations between slower wound healing in diabetic patients with higher BMI. Our results suggest that while PBM appears to be a more universally applicable adjuvant therapy for DFUs, ES may require more nuanced, patient-specific approaches. Further research is warranted to explore the underlying mechanisms driving the differential impact of BMI on these therapies and to optimize treatment protocols based on individual patient characteristics.

Individual factors such as age, sex, duration of diabetes, and neuropathy can negatively influence DFU healing (10). While these variables are known to affect wound healing, they were not always consistently controlled for in the studies included in this metaanalysis. We acknowledge that variables such as age, sex, and neuropathy were not fully addressed in every study. In the cases where these factors were reported, we integrated them as much as possible into our analysis.

The PBM and ES encounter significant challenges for real-world adoption, such as (1) high equipment costs (especially for PBM); (2) the requirement for specialized staff training; and (3) demanding treatment schedules that can impact patient adherence. Although PBM offers safety benefits, ES might present a more cost-efficient option. Future studies should tackle these obstacles by exploring subsidized programs or streamlined protocols.

5.5. Characteristics of Interventions

5.5.1. Photobiomodulation

Based on the included RCTs in our work, the PBM parameters for DFU were as follows: Wavelength of 630 - 904 nm, power density of 20 - 300 mW, and dosage of 1 - 6]/cm², with an irradiation time of 30 - 60 seconds at

each point at a distance of 1 cm away from the wound, and at least 3 times a week for 4 - 6 weeks.

5.5.2. Electrical Stimulation

The parameters used in most trials for ES include the HVPC and DC with negative polarity, the intensity of sensory threshold, 30 - 60 minutes per day, 3 - 5 sessions per week, and electrodes placed around the wound. Yet, further evidence from higher-quality RCTs is imperative to determine the most appropriate parameters for healing DFUs.

5.6. Limitations and Potential Biases

This systematic review and meta-analysis have several important limitations that should be considered when interpreting the results. First, many of the included studies had fewer than 30 participants, which may limit the generalizability of the findings and increase the risk of random error. Additionally, many studies had short follow-up periods, often around four weeks, which provides only a snapshot of the ulcer healing trajectory and limits the ability to assess the long-term sustainability of treatment effects. Another limitation is the heterogeneity of the included studies in terms of intervention parameters, such as wavelength, power density, and treatment duration for PBM, as well as waveform, intensity, and electrode placement for ES. Finally, the quality assessment of the included studies revealed that 28.6% of the studies were weak in terms of blinding, which may have introduced performance or detection bias. These limitations underscore the need for larger, more standardized trials with longer followup periods to confirm these preliminary findings.

5.7. Future Research Directions

Given the limitations of current evidence, future RCTs should prioritize larger sample sizes, standardized protocols, and longer follow-up periods to validate these preliminary findings. These studies would provide more definitive, high-quality evidence regarding the impact of PBM and ES on reducing diabetic ulcer size over time. By incorporating these elements, researchers can enhance our understanding of these treatments and their long-term effects on DFUs.

6. Conclusions

This systematic review and meta-analysis found that both PBM and ES are effective adjuvant therapies for enhancing DFU healing when combined with standard care. Although PBM showed a larger effect size, the difference between the two treatments was not statistically significant. Overall, healthcare providers may consider incorporating PBM or ES to improve DFU healing. Given this, treatment selection should be guided by practical considerations such as equipment availability, cost, treatment duration, ease of use, side effect profile, and patient preference. Notably, most included studies (6 of 11) focused on Wagner grade II ulcers, indicating that these findings are generalizable primarily to moderate-severity DFUs. Future studies should investigate the efficacy of these treatments across different Wagner grades to further establish their applicability in varying ulcer severities.

Acknowledgements

This study was derived from research project No. 14030121355, approved by Hamadan University of Medical Sciences. The authors appreciate the financial and non-financial support of the university's Vice Chancellor for Research and Technology and the patients who participated in the study. The funding source had no role in the study design, data collection, analysis, interpretation, or writing of the manuscript. The authors would also like to thank Dr. Niloofar Rabiei for assisting with the statistical analysis.

Supplementary Material

Supplementary material(s) is available here [To read supplementary materials, please refer to the journal website and open PDF/HTML].

Footnotes

Authors' Contribution: Study concept and design: M. R. M; Acquisition of data: P. O. and M. T. K; Analysis and interpretation of data: S. S. and P. O.; Drafting of the manuscript: P. O. and M. R. A; Critical revision of the manuscript for important intellectual content: P. O., M. T. K., and M. R. A.; Statistical analysis: S. S.

Conflict of Interests Statement: The authors declare no conflict of interest.

Data Availability: The datasets used and/or analyzed during the current study are available from the corresponding author upon reasonable request.

Ethical Approval: This study was approved by the ethical code IR.UMSHA.REC.1403.025 at Hamadan University of Medical Sciences.

Funding/Support: This study was derived from research project No. 14030121355 approved by Hamadan University of Medical Sciences.

References

- 1. Diabetes Atlas IDF. International diabetes federation. *Executive summary*. 2017:8-11.
- Pendsey SP. Understanding diabetic foot. Int J Diabetes Dev Ctries. 2010;30(2):75-9. [PubMed ID: 20535310]. [PubMed Central ID: PMC2878694]. https://doi.org/10.4103/0973-3930.62596.
- 3. Oliver TI, Mutluoglu M. Diabetic Foot Ulcer (Archived). *StatPearls*. Treasure Island (FL): StatPearls Publishing Copyright © 2025, StatPearls Publishing LLC; 2025.
- Brocklehurst JD. The Validity and Reliability of the SINBAD Classification System for Diabetic Foot Ulcers. Adv Skin Wound Care. 2023;36(11):1-5. [PubMed ID: 37861669]. https://doi.org/10.1097/asw.000000000000050.
- 5. Habtemariam S, Habtemariam S. Pathophysiology of type 2 diabetes complications. *Medicinal Foods as Potential Therapies for Type-2 Diabetes Ass Dis.* 2019:69-88. https://doi.org/10.1016/B978-0-08-102922-0.00004-3.
- Guest JF, Fuller GW, Vowden P. Diabetic foot ulcer management in clinical practice in the UK: costs and outcomes. *Int Wound J.* 2018;15(1):43-52. [PubMed ID: 29243399]. [PubMed Central ID: PMC7950039]. https://doi.org/10.1111/iwj.12816.
- Perez-Favila A, Martinez-Fierro ML, Rodriguez-Lazalde JG, Cid-Baez MA, Zamudio-Osuna MJ, Martinez-Blanco MDR, et al. Current Therapeutic Strategies in Diabetic Foot Ulcers. *Medicina (Kaunas)*. 2019;55(11). [PubMed ID: 31731539]. [PubMed Central ID: PMC6915664]. https://doi.org/10.3390/medicina55110714.
- Medeiros S, Rodrigues A, Costa R. Physiotherapeutic interventions in the treatment of patients with diabetic foot ulcers: a systematic literature review. *Physiotherapy*. 2023;**118**:79-87. [PubMed ID: 36244842]. https://doi.org/10.1016/j.physio.2022.09.006.
- Kwan RL, Cheing GL, Vong SK, Lo SK. Electrophysical therapy for managing diabetic foot ulcers: a systematic review. *Int Wound J.* 2013;10(2):121-31. [PubMed ID: 22958779]. [PubMed Central ID: PMC7950529]. https://doi.org/10.1111/j.1742-481X.2012.01085.x.
- Huang H, Xin R, Li X, Zhang X, Chen Z, Zhu Q, et al. Physical therapy in diabetic foot ulcer: Research progress and clinical application. *Int Wound J.* 2023;**20**(8):3417-34. [PubMed ID: 37095726]. [PubMed Central ID: PMC10502280]. https://doi.org/10.1111/iwj.14196.
- Asadi MR, Torkaman G. Bacterial Inhibition by Electrical Stimulation. *Adv Wound Care (New Rochelle)*. 2014;3(2):91-7. [PubMed ID: 24761349]. [PubMed Central ID: PMC3929208]. https://doi.org/10.1089/wound.2012.0410.
- Polak A, Franek A, Taradaj J. High-Voltage Pulsed Current Electrical Stimulation in Wound Treatment. *Adv Wound Care (New Rochelle)*. 2014;3(2):104-17. [PubMed ID: 24761351]. [PubMed Central ID: PMC3928985]. https://doi.org/10.1089/wound.2013.0445.
- Melotto G, Tunprasert T, Forss JR. The effects of electrical stimulation on diabetic ulcers of foot and lower limb: A systematic review. *Int Wound J.* 2022;**19**(7):1911-33. [PubMed ID: 35112496]. [PubMed Central ID: PMC9615295]. https://doi.org/10.1111/iwj.13762.
- Zheng Y, Du X, Yin L, Liu H. Effect of electrical stimulation on patients with diabetes-related ulcers: a systematic review and meta-analysis. *BMC Endocr Disord*. 2022;**22**(1):112. [PubMed ID: 35477391]. [PubMed Central ID: PMC9044601]. https://doi.org/10.1186/s12902-022-01029-z.
- 15. Hawkins D, Houreld N, Abrahamse H. Low level laser therapy (LLLT) as an effective therapeutic modality for delayed wound healing. *Ann*

N Y Acad Sci. 2005;**1056**:486-93. [PubMed ID: 16387711]. https://doi.org/10.1196/annals.1352.040.

- Santos CMD, Rocha RBD, Hazime FA, Cardoso VS. A Systematic Review and Meta-Analysis of the Effects of Low-Level Laser Therapy in the Treatment of Diabetic Foot Ulcers. Int J Low Extrem Wounds. 2021;20(3):198-207. [PubMed ID: 32394760]. https://doi.org/10.1177/1534734620914439.
- Huang J, Chen J, Xiong S, Huang J, Liu Z. The effect of low-level laser therapy on diabetic foot ulcers: A meta-analysis of randomised controlled trials. *Int Wound J.* 2021;**18**(6):763-76. [PubMed ID: 33751853]. [PubMed Central ID: PMC8613395]. https://doi.org/10.1111/iwj.13577.
- Li S, Wang C, Wang B, Liu L, Tang L, Liu D, et al. Efficacy of low-level light therapy for treatment of diabetic foot ulcer: A systematic review and meta-analysis of randomized controlled trials. *Diabetes Res Clin Pract.* 2018;143:215-24. [PubMed ID: 30009935]. https://doi.org/10.1016/j.diabres.2018.07.014.
- Kailashdan AA, Arjunbhai BR. Effect of low-level laser therapy on diabetic foot ulcers. Int J Med Biomed Studies. 2019;3(9):125-9. https://doi.org/10.32553/ijmbs.v3i9.545.
- Maiya AG, Kumar AS, Hazari A, Jadhav R, Ramachandra L, Hande HM, et al. Photobiomodulation therapy in neuroischaemic diabetic foot ulcers: a novel method of limb salvage. J Wound Care. 2018;27(12):837-42. [PubMed ID: 30557112]. https://doi.org/10.12968/jowc.2018.27.12.837.
- 21. Mathur RK, Sahu K, Saraf S, Patheja P, Khan F, Gupta PK. Low-level laser therapy as an adjunct to conventional therapy in the treatment of diabetic foot ulcers. *Lasers Med Sci.* 2017;**32**(2):275-82. [PubMed ID: 27896528]. https://doi.org/10.1007/s10103-016-2109-2.
- Rahbar Layegh E, Fadaei Fathabadi F, Lotfinia M, Zare F, Mohammadi Tofigh A, Abrishami S, et al. Photobiomodulation therapy improves the growth factor and cytokine secretory profile in human type 2 diabetic fibroblasts. *J Photochem Photobiol B.* 2020;**210**:111962.
 [PubMed ID: 32712344]. https://doi.org/10.1016/j.jphotobiol.2020.111962.
- 23. Houreld N. Review healing effects of photobiomodulation on diabetic wounds. *Applied Sci.* 2019;**9**(23):81-90. https://doi.org/10.3390/app9235114.
- 24. Kloth LC. Electrical stimulation technologies for wound healing. *Advances Wound Care.* 2014;3(2):81-90. https://doi.org/10.1089/wound.2013.0459.
- 25. Mester E, Szende B, Gärtner P. [The effect of laser beams on the growth of hair in mice]. *Radiobiol Radiother (Berl)*. 1968;**9**(5):621-6. Germany. [PubMed ID: 5732466].
- 26. Mester E, Spiry T, Szende B, Tota JG. Effect of laser rays on wound healing. *The American J Surgery*. 1971;**122**(4):532-5. https://doi.org/10.1016/0002-9610(71)90482-X.
- Mester E. [Clinical results of laser stimulation and experimental studies on its mechanism of action]. *Minerva Med.* 1981;72(33):2195-9. IT. [PubMed ID: 7290441].
- Moher D, Liberati A, Tetzlaff J, Altman DG. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Med.* 2009;6(7). e1000097. [PubMed ID: 19621072]. [PubMed Central ID: PMC2707599]. https://doi.org/10.1371/journal.pmed.1000097.
- Armijo-Olivo S, Stiles CR, Hagen NA, Biondo PD, Cummings GG. Assessment of study quality for systematic reviews: a comparison of the Cochrane Collaboration Risk of Bias Tool and the Effective Public Health Practice Project Quality Assessment Tool: methodological research. J Eval Clin Pract. 2012;18(1):12-8. [PubMed ID: 20698919]. https://doi.org/10.1111/j.1365-2753.2010.01516.x.
- Hedges LV, Vevea JL. Fixed-and random-effects models in metaanalysis. Psychol Methods. 1998;3(4):486. https://doi.org/10.1037/1082-989X.3.4.486.

- 31. Kanters S. Fixed-and random-effects models. *Meta-research: Methods and protocols*. New York, USA: Springer; 2021. p. 41-65. https://doi.org/10.1007/978-1-0716-1566-9_3.
- Riley RD, Higgins JP, Deeks JJ. Interpretation of random effects metaanalyses. *BMJ*. 2011;342:d549. [PubMed ID: 21310794]. https://doi.org/10.1136/bmj.d549.
- Olkin I, Dahabreh IJ, Trikalinos TA. GOSH a graphical display of study heterogeneity. *Res Synth Methods*. 2012;3(3):214-23. [PubMed ID: 26062164]. https://doi.org/10.1002/jrsm.1053.
- Zulbaran-Rojas A, Park C, El-Refaei N, Lepow B, Najafi B. Home-Based Electrical Stimulation to Accelerate Wound Healing-A Double-Blinded Randomized Control Trial. J Diabetes Sci Technol. 2023;17(1):15-24. [PubMed ID: 34328024]. [PubMed Central ID: PMC9846397]. https://doi.org/10.1177/19322968211035128.
- Wadee AN, Fahmy SM, Bahey El-Deen HA. Low-level laser therapy (photobiomodulation) versus hyperbaric oxygen therapy on healing of chronic diabetic foot ulcers: a controlled randomized trial. *Physical Therapy Rev.* 2021;26(1):73-80. https://doi.org/10.1080/10833196.2021.1876380.
- 36. Vitoriano NAM, Mont'Alverne DGB, Martins MIS, Silva PS, Martins CA, Teixeira HD, et al. Comparative study on laser and LED influence on tissue repair and improvement of neuropathic symptoms during the treatment of diabetic ulcers. *Lasers Med Sci.* 2019;**34**(7):1365-71. [PubMed ID: 30715637]. https://doi.org/10.1007/s10103-019-02724-5.
- Tantawy SA, Abdelbasset WK, Kamel DM, Alrawaili SM. A randomized controlled trial comparing helium-neon laser therapy and infrared laser therapy in patients with diabetic foot ulcer. *Lasers Med Sci.* 2018;33(9):1901-6. [PubMed ID: 29846832]. https://doi.org/10.1007/s10103-018-2553-2.
- Frangež I, Nizič-Kos T, Frangež HB. Phototherapy with LED Shows Promising Results in Healing Chronic Wounds in Diabetes Mellitus Patients: A Prospective Randomized Double-Blind Study. *Photomed Laser Surg.* 2018;**36**(7):377-82. [PubMed ID: 29668397]. https://doi.org/10.1089/pho.2017.4382.
- Priyadarshini LMJ, Babu KEP, Thariq IA. Effect of low level laser therapy on diabetic foot ulcers: a randomized control trial. *Int Surg J.* 2018;5:1008-15. https://doi.org/10.18203/2349-2902.isj20180821.
- 40. de Alencar Fonseca Santos J, Campelo MBD, de Oliveira RA, Nicolau RA, Rezende VEA, Arisawa EÂ L. Effects of Low-Power Light Therapy on the Tissue Repair Process of Chronic Wounds in Diabetic Feet. *Photomed Laser Surg.* 2018;**36**(6):298-304. [PubMed ID: 29882738]. https://doi.org/10.1089/pho.2018.4455.
- 41. Srilestari A, Nareswari I, Simadibrata C, Tarigan TJ. Effectiveness of combined laser-puncture and conventional wound care to accelerate diabetic foot ulcer healing. *Med J Indonesia*. 2017;**26**(1):26-34.
- Asadi MR, Torkaman G, Hedayati M, Mohajeri-Tehrani MR, Ahmadi M, Gohardani RF. Angiogenic effects of low-intensity cathodal direct current on ischemic diabetic foot ulcers: A randomized controlled trial. *Diabetes Res Clin Pract.* 2017;**127**:147-55. [PubMed ID: 28371685]. https://doi.org/10.1016/j.diabres.2017.03.012.
- 43. El Rasheed NAA, Mahmoud NF, Hamada HA, El Khatib A. Pulsed electromagnetic fields versus laser therapy on enhancing recovery of diabetic foot ulcer: A single blind randomized controlled trial. *Biomed Res (0970-938X).* 2017;**28**(19).

- Hoseini Sanati M, Torkaman G, Hedayati M, Iranparvar Alamdari M.
 [Effect of Ga-As laser on decrease of wound surface area and ABI value in diabetic foot ulcers]. *J Advances Med Biomed Res.* 2016;24(103):20-1.
 FA.
- 45. Asadi MR, Torkaman G, Mohajeri-Tehrani MR, Hedayati M. [Effects of electrical stimulation on the management of ischemic diabetic foot ulcers]. *J Babol Univ Med Sci.* 2015;**17**(7):7-14. FA.
- 46. Feitosa MC, Carvalho AF, Feitosa VC, Coelho IM, Oliveira RA, Arisawa EÂ. Effects of the Low-Level Laser Therapy (LLLT) in the process of healing diabetic foot ulcers. *Acta Cir Bras.* 2015;**30**(12):852-7. [PubMed ID: 26735058]. https://doi.org/10.1590/s0102-865020150120000010.
- Mohajeri-Tehrani MR, Nasiripoor F, Torkaman G, Hedayati M, Annabestani Z, Asadi MR. Effect of low-intensity direct current on expression of vascular endothelial growth factor and nitric oxide in diabetic foot ulcers. J Rehabil Res Dev. 2014;51(5):815-24. [PubMed ID: 25509057]. https://doi.org/10.1682/jrrd.2013.08.0174.
- 48. Sandoval Ortíz MC, Herrera Villabona E, Camargo Lemos DM, Castellanos R. Effects of low level laser therapy and high voltage stimulation on diabetic wound healing. *Revista de la Universidad Industrial de Santander Salud*. 2014;46(2):107-17.
- Kajagar BM, Godhi AS, Pandit A, Khatri S. Efficacy of low level laser therapy on wound healing in patients with chronic diabetic foot ulcers-a randomised control trial. *Indian J Surg.* 2012;74(5):359-63. [PubMed ID: 24082586]. [PubMed Central ID: PMC3477409]. https://doi.org/10.1007/s12262-011-0393-4.
- 50. Kaviani A, Djavid GE, Ataie-Fashtami L, Fateh M, Ghodsi M, Salami M, et al. A randomized clinical trial on the effect of low-level laser therapy on chronic diabetic foot wound healing: a preliminary report. *Photomed Laser Surg.* 2011;**29**(2):109-14. [PubMed ID: 21214368]. https://doi.org/10.1089/pho.2009.2680.
- Minatel DG, Frade MA, França SC, Enwemeka CS. Phototherapy promotes healing of chronic diabetic leg ulcers that failed to respond to other therapies. *Lasers Surg Med.* 2009;41(6):433-41. [PubMed ID: 19588536]. https://doi.org/10.1002/lsm.20789.
- 52. Naidu SVLG, Subapriya S, Yeoh CN, Soosai S, Shalini V, Harwant S. The use of low output laser therapy to accelerate healing of diabetic foot ulcers-a randomized prospective controlled trial. *Laser Florence 2004: A Window on the Laser Medicine World*. SPIE; 2005. p. 51-7.
- Lundeberg TC, Eriksson SV, Malm M. Electrical nerve stimulation improves healing of diabetic ulcers. *Ann Plast Surg.* 1992;**29**(4):328-31. [PubMed ID: 1466529]. https://doi.org/10.1097/00000637-199210000-00009.
- Avci P, Gupta A, Sadasivam M, Vecchio D, Pam Z, Pam N, et al. Low-level laser (light) therapy (LLLT) in skin: stimulating, healing, restoring. *Semin Cutan Med Surg.* 2013;**32**(1):41-52. [PubMed ID: 24049929]. [PubMed Central ID: PMC4126803].
- Chen Z, Chen ZY, Liu WH, Li GS. Electric Stimulation as an Effective Adjunctive Therapy for Diabetic Foot Ulcer: A Meta-analysis of Randomized Controlled Trials. *Adv Skin Wound Care*. 2020;**33**(11):608-12. [PubMed ID: 33065683]. https://doi.org/10.1097/01.ASW.0000695784.82605.1e.
- Doheny EP, Caulfield BM, Minogue CM, Lowery MM. Effect of subcutaneous fat thickness and surface electrode configuration during neuromuscular electrical stimulation. *Med Eng Phys.* 2010;**32**(5):468-74. [PubMed ID: 20417145]. https://doi.org/10.1016/j.medengphy.2010.03.004.