



Effect of Family-centered Empowerment Model on Quality of Life of Patients with Hypertension

Mahsa Asadollahi Hamedani ¹, Alireza Salar ^{2,*} and Fatihe Kermansaravi ²

¹Zahedan University of Medical Sciences, Zahedan, Iran

²Community Nursing Research Center, Zahedan University of Medical Sciences, Zahedan, Iran

*Corresponding author: Community Nursing Research Center, Zahedan University of Medical Sciences, Zahedan, Iran. Email: salar293@gmail.com

Received 2021 June 20; Accepted 2021 June 20.

Abstract

Background: Nowadays, the high prevalence of hypertension and its serious complications for affected patients have made this disease a major health issue worldwide. Hypertension, like other chronic diseases, reduces the patients' quality of life. Maintaining and improving their lifestyles requires the serious support of families.

Objectives: This study aimed to explore the effect of the family-centered empowerment model on the quality of life of patients with hypertension.

Methods: This clinical trial was performed on 70 hypertensive patients who were referred to two comprehensive health centers in Zahedan University of Medical Sciences in Zahedan, southeastern Iran, in 2020. The participants were selected and randomly allocated based on the inclusion criteria to the intervention (n = 35) and control (n = 35) groups using the consort. The participants in the intervention group received training in four 60-min sessions individually based on the family empowerment model. To collect data, a personal information form and the Quality of Life Questionnaire for hypertensive patients were used. The questionnaire was completed before the intervention and one and three months after the intervention by a self-report method. The collected data were analyzed by SPSS-22 using the independent samples *t* test, chi-square test, and repeated-measures analysis of variance.

Results: The results of repeated-measures analysis of variance regarding the effect of time and group on the quality of life showed a statistically significant difference in the second and third phases between the two groups. In other words, the two variables of intervention and time affected the mean score of quality of life, and a statistically significant difference was observed in the measured time intervals ($P < 0.001$).

Conclusions: This study showed that implementing the family-centered empowerment model improves the quality of life of patients with hypertension. Therefore, we recommend employing this method by nurses to engage the patients with chronic diseases and their families in the process of treatment.

Keywords: Family-centered Empowerment, Quality of Life, Hypertension

1. Background

Today, chronic diseases account for a large proportion of health problems, not only in developing countries but also in many developed countries (1). It is estimated that more than 1.5 billion people worldwide will have high blood pressure by 2025, which will increase the risk of heart disease by 50% and the risk of stroke by 75% (2). With the high prevalence and serious complications on body organs, high blood pressure is a disease, making major health issues worldwide. The presence of one billion hypertensive people and the annual occurrence of four million deaths all over the world are the direct results of this disease (3). Blood pressure can be controlled only in 20% of

patients suffering from hypertension in Iran (4). In 2019, 21,000 people were diagnosed with high blood pressure in Sistan and Baluchestan province, and according to the reports in this area, about 700 people died because of high blood pressure in 1996. Besides, a total of 29,549 patients with hypertension have been recently identified in a medical center affiliated with Zahedan University of Medical Sciences (5, 6).

Complications such as myocardial infarction, stroke, heart failure, atrial fibrillation, aortic rupture, peripheral artery disease, and cognitive problems can be prevented by the early detection of hypertension. Indeed, hypertension is a preventable risk factor for these complications (7, 8), which can be controlled by measures like lifestyle

modification such as restricting sodium in the diet, weight loss, increasing activity and exercise, eating more fruits and vegetables, not smoking, reducing alcohol consumption, and adhering to the prescribed medication regimen (9). Therefore, self-efficacy is an important prerequisite in the process of changing behavior and improving the quality of life (10).

The incidence of any chronic disease has a great impact on the patient's quality of life. Physical, psychological, social, and economic dimensions of the quality of life undergo many changes due to the duration and severity of chronic diseases (11). Numerous studies have shown that the quality of life of those patients is low (12, 13), which, in turn, reduces life expectancy and increases the risk of depression, social isolation, reduced daily activities, dependence, and economic burden (14). It is worth mentioning that nowadays, one of the most important goals of the Iranian health care system is to improve the quality of life (15). Thus, educational interventions (16), empowerment models (17), self-care education (18), and integrated strategies for health services delivery have received growing attention (19).

Currently, given the growing rate of chronic diseases and lack of physical space in treatment centers, attention to the family-centered empowerment model is a key element in medical and nursing education (20). Family-centered empowerment reflects professional interactions with families so that family members can gain a sense of control over their lives (21). The family empowerment model considers the beliefs and cultural values of family members, takes their needs into account, and tries to reduce their independence from the hospital (22). The term family-centered empowerment has played an important role in nursing and medical studies in recent decades, and this has been considered a necessity of the nursing process (23). Family-centered care is defined as an innovative approach to health care planning, delivery, and evaluation, based on beneficial and reciprocal collaboration between patients, families, and caregivers (24). Educating family members about disease control and even prevention can be important (25) because there is a strong relationship between family and members' health status (26). The family-centered empowerment model can empower individuals by emphasizing perceived threats (through perceived severity and sensitivity), promoting self-efficacy (by fostering necessary skills), and increasing self-esteem (through self-help, self-control, and self-efficacy) (27).

The family-centered empowerment model consists of

four stages of identification, sensitization, control, and evaluation (28). The mentioned pattern is a native model and concepts of qualitative research of grounded theory. This model can increase the knowledge, beliefs, and skills of the patient and the family by emphasizing the important role of the family to play in terms of motivation, psychology, self-efficacy, and self-esteem. The main goal of the family-centered empowerment model is to empower the patients and their families to improve their level of health (29). A review study showed that this model affects the self-esteem (30) and the lifestyle of patients with hypertension (31), self-efficacy of patients on hemodialysis (32), and the quality of life of patients with myocardial infarction (33). Since the nurse is the best source for transferring information among the health team, patients, and their families, many studies have highlighted the effects of this model on patients with other diagnoses and different variables (23).

2. Objectives

The present study aimed to investigate the effect of the family-centered empowerment model on the quality of life of patients with hypertension.

3. Methods

The present clinical trial study was performed with registration number IRCT20201016049043N1 based on a permit from the Ethics Committee of Zahedan University of Medical Sciences (IR.ZAUMS.REC.1399.171) in Sistan and Baluchestan province. This province is located in the southeast of Iran, Central Asia. The participants were patients with hypertension who were referred to the comprehensive health centers in Zahedan in 2020.

The inclusion criteria included having a history of hypertension diagnosed in medical centers, having a blood pressure of higher than 140/90 for at least six months, having minimum literacy, being 25-65-years-old, and not participating in the same training program in the past six months. The inclusion criteria for the caregiver included having minimum literacy and living in the patient's home. The exclusion criteria for the patients were having an acute illness, hospitalization during the study, the unwillingness of the patient and his family to continue participating in the study, and not taking part in the posttest.

Sampling was performed in several steps. First, several urban districts were considered clusters. Then, two districts and one center from each district were randomly selected, and a list of eligible patients was prepared based on

the inclusion criteria from the SIB system (The integrated health system) and the medical files available in the two centers. Out of 100 eligible patients, 70 patients were selected from two centers using convenience sampling. The selected patients were randomly assigned to the intervention and control groups (Figure 1).

Following a similar study (Noori et al. (17) in 2015), taking a 95% confidence interval and a 95% statistical test power, the sample size was calculated as 32 patients using the following formula: $n = \frac{(Z_{1-\frac{\alpha}{2}} + Z_{1-\beta})^2 (S_1^2 + S_2^2)}{(\bar{X}_1 - \bar{X}_2)^2} =$

31.68

$Z_{1-\frac{\alpha}{2}} = 1.96$, $Z_{1-\beta} = 1.64$, $S_1 = 10.92$, $S_2 = 6.75$, $X_1 = 51.95$, $X_2 = 43.74$

With a possible dropouts, 35 patients in each group and a total of 70 patients were evaluated (17).

The instruments used in the present study were a personal information form and the Quality of Life Questionnaire for patients with hypertension. The personal information form assessed the patients' age, education (for the patient and caregiver), body mass index, duration of illness, underlying disease(s), gender, ethnicity, employment status, primary caregiver, marriage, smoking, a family history of hypertension, and the disease control method. The Quality of Life Questionnaire for Patients with Hypertension was designed and validated by Shamsi (2016). This questionnaire contains 42 items to assess the quality of life of Iranian patients with hypertension and was piloted in Tehran in 2015-2016. The items are scored on a five-point Likert scale: Never (5), rarely (4), sometimes (3), often (2), and always (1). The total score obtained varies from 42 to 210, and a higher score means a higher quality of life. The face, content, and construct validity indices of the questionnaire were confirmed, and its reliability with Cronbach's alpha was reported to be 0.89 (34).

After obtaining the necessary permits, the researcher referred to the selected health centers and provided some information about the objectives of the study and the procedure taken to conduct it. Then, he invited the patients to participate in the study. First, the personal information form and the questionnaire were completed by conducting interviews with the participants in both groups. Afterward, the intervention was conducted for the participants in the intervention group using the empowerment model in four 60-min sessions for four weeks. The training sessions were held individually for each patient and a member of his family as the main caregiver (Table 1).

The sessions were held in the form of discussions, ver-

bal exchanges, questions, and answers by the researcher in health centers using educational aids such as PowerPoint presentations, videos, and booklets based on valid scientific texts (35) under the supervision of experts in this field. The participants in the control group did not receive any training. During the training, caregivers were asked to practice the given training content in the presence of a nurse to remove any possible ambiguities. The family-centered empowerment model was implemented through the following steps:

3.1. Step 1 (Perceiving the Threats)

The first step in the family-centered empowerment model was to increase the intensity and sensitivity perceived by the participants by lecturing and discussing the nature and complications of the disease, aggravating factors, nutritional factors, exercise, and other effective factors in disease control in two sessions. At the end of the training session, the participants' comprehension was ensured through questions and answers.

3.2. Step 2 (Self-efficacy)

The focus was on improving the problem-solving skills. For this purpose, a problem-solving session was held individually for patients and their families. In these sessions, patients and families were faced with problems and the problem-solving process. They were asked to discuss concrete examples of their situation and what they had done to solve the problem. Besides, the practical methods of measuring blood pressure and weight and the consequences of not controlling these two variables and their normal levels were explained in detail. Then, the patients and families were asked to practice the skills. At this stage, the best way to solve the patients' problems was selected with a focus on the clients' skills.

3.3. Step 3 (Self-confidence)

At this stage, the patient was asked to teach the family members the issues covered in the previous sessions with the help of the researcher. The purpose of this step was to increase the patients' self-confidence concerning their ability to provide information to family members and family support. To ensure and follow up on this step, several questions related to the educational issues were given to the patient to be answered with the help of family members and be delivered to the researcher.




Figure 1. The consort flow diagram

Table 1. Training Sessions

Educational Content	
Step one, sessions 1 and 2	Familiarity with the objectives of the intervention program, the generality of the disease, the importance of a healthy lifestyle to the individual's health and prevention of chronic diseases, increasing awareness and improving the attitude of the participants about healthy nutrition; raising their awareness and improving their attitudes toward physical activity and its importance in the prevention and control of hypertension, improving the mental health of the participants with stories about ways to manage stress, improving the spiritual health of the participants with stories about the impact of religion on health.
Step two, session 3	Familiarity with devices, principles, and objective methods of measuring blood pressure, how to solve problems, providing suggestions for solving problems, emphasizing the client's skills
Step three, session 4	Reviewing content by the client accompanied by relevant interventions and support by the researcher, questions and answers
Step four	Evaluation

3.4. Step 4 (Evaluation)

The evaluation was performed one and three months after the intervention in the two groups by completing the quality of life questionnaire. Furthermore, during the study, the participants were in contact with the researcher by phone to solve possible problems and answer possible

questions.

To comply with ethical principles, a training booklet was provided to the participants in both intervention and control groups after completing the intervention. The questionnaires were completed in person or by contacting the participants so that the researcher could cooperate if

there was any ambiguity to fill in the questionnaires. The collected data were analyzed by SPSS software (version 22).

Data normality was checked using the Shapiro-Wilk test. The chi-square test was used to compare the frequency of qualitative variables between the two groups. Furthermore, the independent samples *t* test was run to compare the means in the two intervention and control groups, and the analysis of variance was used before the intervention, as well as one and three months after the intervention. The assumptions were established the independence of observations, normality in population distributions, and homogeneity of population variances, and the homogeneity of the variances were established. The significance level in this study was less than 0.05.

4. Results

The findings of the study showed that the participants in the intervention and control groups were not significantly different in terms of demographic characteristics, including age, patient's and caregiver's education, body mass index, duration of illness, underlying disease, gender, ethnicity, employment status, primary caregiver, marital status, smoking, family history of hypertension, and disease control. The patients in the intervention and control groups were in their 40s. The patients had a body mass index of 29. The duration of the disease was 10-12 years. The majority of the participants in both groups had an underlying disease. Most of them were female and unemployed. Besides, 60% of the patients in the intervention group and 40% of the patients in the control group were Baloch. The main caregivers of the patients were children in the intervention group (65.7%) and spouses in the control group (54.3%) (Table 2).

The results showed that the two groups did not differ significantly in terms of the quality of life in the pre-intervention stage ($P = 0.16$); but after the intervention, there was a significant difference between the two groups ($P < 0.001$). The mean scores of the quality of life in the intervention group before, one month, and three months after the intervention were 132.62 ± 15.96 , 143.68 ± 9.52 , and 143.22 ± 10.25 , and the corresponding values for the control group were 137.10 ± 14.44 , 137.17 ± 10.41 , and 137.28 ± 10.46 , respectively. The results of repeated-measures analysis of variance on quality of life showed that changes in the quality of life scores over time were significant ($P < 0.001$), with greater changes observed in the intervention group than in the control group (Table 3).

the results showed that the mean score quality of life of the participants in the intervention group were significantly different from the pretest score in the second phase ($P = 0.008$) and third phase ($P = 0.01$) (Figure 2).

5. Discussion

The results of the study showed that the implementation of the family-centered empowerment model improved the hypertensive patients' quality of life. Accordingly, it can be said that the number of training sessions and the time allotted for educating patients and their families, ample opportunity to learn self-care content and skills, discussion, and active participation of the patient with an active family member were the main advantages of the model that contributed to achieving positive consistent outcomes in this study.

Mohalli et al. studied the effect of the family-centered empowerment model on empowerment indices of patients with hypertension and showed that the knowledge, self-esteem, and self-efficacy of patients with hypertension increased in the experimental group (36), as was evident in the present study. This study was similar to the present study in terms of the sample size and intervention process, but the final evaluation was performed 1.5 months later. Babaei-Sis et al. examined the effect of lifestyle modification training intervention on blood pressure control in patients with hypertension and confirmed the effectiveness of the educational intervention for promoting physical activity, weight management, nutrition, and mental health in patients with hypertension (31). These findings were in line with the results of the present study, with the advantage that it was performed on 210 people. Bairami et al. examined the relationship between self-care behaviors and quality of life in patients with hypertension who were referred to comprehensive health centers in Hamadan. The results showed a positive significant correlation between the overall scores of quality of life and self-care behaviors (37), which confirmed the results of the current study. Keshvari et al. investigated the effect of the family-centered empowerment model on blood pressure and empowerment of the elderly with hypertension and found that the empowerment of the elderly with hypertension led to improved blood pressure (38). This finding was consistent with the results of the present study, with the difference that the evaluation was performed one week and one month later.

Table 2. The Participants' Demographic Characteristics in Two Groups ^a

Variable	Control Group	Intervention Group	P
Age	7.74 ± 49.91	6.31 ± 48.80	0.51 ^b
Patients' academic years	3.75 ± 8.97	3.30 ± 7.77	0.16 ^b
Caregiver's academic years	3.17 ± 10.80	3.03 ± 10.31	0.51 ^b
BMI	5.10 ± 28.42	3.95 ± 29.24	0.46 ^b
Duration of the disease	6.03 ± 10.09	5.38 ± 12.11	0.14 ^b
Underlying disease			0.60 ^c
Yes	24 (68.6)	26 (74.3)	
No	11 (31.4)	9 (25.7)	
Gender			0.33 ^c
Female	23 (65.7)	18 (51.4)	
Male	12 (34.3)	17 (48.6)	
Ethnicity			0.80 ^c
Baloch	21 (60)	20 (57.1)	
Sistani	14 (40)	15 (42.9)	
Occupation			0.08 ^c
Employed	11 (31.4)	13 (37.1)	
Unemployed	24 (68.6)	22 (62.9)	
Main caregiver			0.09 ^c
Spouse	12 (34.3)	19 (54.3)	
Child	23 (65.7)	16 (45.7)	
Marital status			0.78 ^c
Married	27 (77.1)	26 (74.3)	
Unmarried	8 (22.9)	9 (25.7)	
Smoking			0.61 ^c
Yes	11 (31.4)	13 (37.1)	
No	24 (68.6)	22 (62.9)	
Family history of hypertension			0.25 ^c
Yes	25 (71.4)	29 (82.9)	
No	10 (28.6)	6 (17.1)	
Blood pressure control			0.60 ^c
With medication	23 (65.7)	25 (71.4)	
Without medication	12 (34.4)	10 (28.6)	


^aValues are expressed as mean ± SD or No. (%).^bTest Independent *t*^cChi-square

Similar to the present study, Noori et al. explored the effect of the family-centered empowerment model on the quality of life of patients with hypertension and showed that implementing the family-centered empowerment model increased the patients' quality of life and reduced their blood pressure (17). Contrary to the present

study, the authors used the general quality of life questionnaire, and the effects of the interventions were measured 1.5 months later. Escortell-Mayor et al. investigated the effect of education on patients at risk for cardiovascular problems and with poor blood pressure control. The results of this study showed a statistically significant dif-

Table 3. Results of Repeated-measures Analysis of Variance for Quality of Life in Patients with Hypertension in Two Groups

Variable	Sum of Squares	df	Mean Square	F	P
Time	1009.82	1	1009.82	70.98	< 0.001
Group	368.01	1	368.01	1.003	< 0.001
Time × group	956.82	1	956.82	67.26	< 0.001
Error	24947.69	68	366.87		

**Figure 2.** Line graph showing changes in patients' quality of life scores before and one and three months after implementing the family empowerment model in the control and intervention groups

ference between the two groups, and education led to better control of blood pressure and lower blood cholesterol (39), which was in line with the results of the present study. Mini et al. investigated the effect of planned training on blood pressure control among teachers in Kerala, India, and showed that behavioral intervention had a sig-

nificant effect on blood pressure control (40), which was consistent with the results of the present study. Gong et al. examined the effectiveness of community-based physical activity programs in preventing stroke and heart attack among patients with hypertension aged over 55 years and reported a significant difference in the incidence of

stroke and heart attack. Besides, the physical activity of the patients in the intervention group increased after six months (41), which partly confirmed the results of the present study. Atak et al. performed a family-centered education intervention on a group of patients with diabetes and reported a relative improvement in some areas of quality of life, which was not statistically significant. This study used a different target group and limited training sessions with short duration and insufficient follow-up, which led to different results from the current study findings (42). The results of studies by Sargazi Shad et al. (43), Pilevar et al. (44), Rostami et al. (28), Davarpanah et al. (45), and Rajabi et al. (46) on the quality of life of patients with chronic diseases such as diabetes, rheumatoid arthritis, hemophilia, and leukemia have shown the positive significant effects of the family-centered empowerment model.

5.1. Conclusions

The implementation of the family-centered empowerment model on patients suffering from hypertension improved the patients' quality of life. These types of interventions can help caregivers to improve the quality of their intensive care. Since high blood pressure is not treatable and needs to be controlled, educating and empowering family members can positively influence the self-care behaviors of patients with high blood pressure. Thus, nurses can use this model for engaging patients and family members in medical care.

5.2. Limitations

As a limitation to this study, the COVID-19 outbreak made the participants not have the maximum cooperation in fully implementing the training intervention.

Acknowledgments

This article is part of a Master's thesis under code 9795. We would like to thank the Vice-Chancellor for Research of Zahedan University of Medical Sciences that made this research possible, as well as all the participants in this project.

Footnotes

Authors' Contribution: It was not declared by the authors.

Conflict of Interests: The authors report no conflict of interest.

Ethical Approval: This research project was approved by Zahedan University of Medical Sciences with the code of ethics IR.ZAUMS.REC.1399.171 and the authors complied with ethical considerations of clinical studies.

Funding/Support: This research project received no financial support.

Informed Consent: All patients signed an informed consent form.

References

- Hajat C, Stein E. The global burden of multiple chronic conditions: A narrative review. *Prev Med Rep.* 2018;**12**:284-93. doi: [10.1016/j.pmedr.2018.10.008](https://doi.org/10.1016/j.pmedr.2018.10.008). [PubMed: [30406006](https://pubmed.ncbi.nlm.nih.gov/30406006/)]. [PubMed Central: [PMC6214883](https://pubmed.ncbi.nlm.nih.gov/PMC6214883/)].
- Himmelfarb CR, Commodore-Mensah Y, Hill MN. Expanding the role of nurses to improve hypertension care and control globally. *Ann Glob Health.* 2016;**82**(2):243-53. doi: [10.1016/j.aogh.2016.02.003](https://doi.org/10.1016/j.aogh.2016.02.003). [PubMed: [27372529](https://pubmed.ncbi.nlm.nih.gov/27372529/)].
- Oparil S, Acelajado MC, Bakris GL, Berlowitz DR, Cifkova R, Dominiczak AF, et al. Hypertension. *Nat Rev Dis Primers.* 2018;**4**:18014. doi: [10.1038/nrdp.2018.14](https://doi.org/10.1038/nrdp.2018.14). [PubMed: [29565029](https://pubmed.ncbi.nlm.nih.gov/29565029/)]. [PubMed Central: [PMC6477925](https://pubmed.ncbi.nlm.nih.gov/PMC6477925/)].
- Mirzaei M, Mirzaei M, Bagheri B, Dehghani A. Awareness, treatment, and control of hypertension and related factors in adult Iranian population. *BMC Public Health.* 2020;**20**(1):667. doi: [10.1186/s12889-020-08831-1](https://doi.org/10.1186/s12889-020-08831-1). [PubMed: [32397990](https://pubmed.ncbi.nlm.nih.gov/32397990/)]. [PubMed Central: [PMC7216488](https://pubmed.ncbi.nlm.nih.gov/PMC7216488/)].
- Tabatabayi M. [High blood pressure is the first risk factor for death and disability]. Zahedan: Zahedan University of Medical Sciences and Health Services; 2022. Persian. Available from: <https://zaums.ac.ir/32287.news>.
- Tabatabayi M. [Blood pressure was measured in Sistan and Baluchestan]. IRNA; 2019. Persian. Available from: <https://www.irna.ir/news/83356860/>.
- Boehme AK, Esenwa C, Elkind MS. Stroke risk factors, genetics, and prevention. *Circ Res.* 2017;**120**(3):472-95. doi: [10.1161/CIRCRESAHA.116.308398](https://doi.org/10.1161/CIRCRESAHA.116.308398). [PubMed: [28154098](https://pubmed.ncbi.nlm.nih.gov/28154098/)]. [PubMed Central: [PMC5321635](https://pubmed.ncbi.nlm.nih.gov/PMC5321635/)].
- Zipes DP, Libby P, Bonow RO, Mann DL, Tomaselli GF. *Braunwald's heart disease e-book*: A textbook of cardiovascular medicine: Elsevier Health Sciences; 2018.
- Carey RM, Muntner P, Bosworth HB, Whelton PK. Prevention and control of hypertension: JACC health promotion series. *J Am Coll Cardiol.* 2018;**72**(11):1278-93. doi: [10.1016/j.jacc.2018.07.008](https://doi.org/10.1016/j.jacc.2018.07.008). [PubMed: [30190007](https://pubmed.ncbi.nlm.nih.gov/30190007/)]. [PubMed Central: [PMC6481176](https://pubmed.ncbi.nlm.nih.gov/PMC6481176/)].
- Peyman N, Shahedi F, Abdollahi M, Doosti H, Zadehahmad Z. Impact of self-efficacy strategies education on self-care behaviors among heart failure patients. *J Tehran Heart Cent.* 2020;**15**(1):6-11. [PubMed: [32742286](https://pubmed.ncbi.nlm.nih.gov/32742286/)]. [PubMed Central: [PMC7360864](https://pubmed.ncbi.nlm.nih.gov/PMC7360864/)].
- Samiei Siboni F, Alimoradi Z, Atashi V, Alipour M, Khatooni M. Quality of life in different chronic diseases and its related factors. *Int J Prev Med.* 2019;**10**:65. doi: [10.4103/ijpvm.IJPVM_429_17](https://doi.org/10.4103/ijpvm.IJPVM_429_17). [PubMed: [31198500](https://pubmed.ncbi.nlm.nih.gov/31198500/)]. [PubMed Central: [PMC6547796](https://pubmed.ncbi.nlm.nih.gov/PMC6547796/)].
- Zhang L, Guo X, Zhang J, Chen X, Zhou C, Ge D, et al. Health-related quality of life among adults with and without hypertension: A population-based survey using EQ-5D in Shandong, China. *Sci Rep.* 2017;**7**(1):14960. doi: [10.1038/s41598-017-15083-4](https://doi.org/10.1038/s41598-017-15083-4). [PubMed: [29097724](https://pubmed.ncbi.nlm.nih.gov/29097724/)]. [PubMed Central: [PMC5668325](https://pubmed.ncbi.nlm.nih.gov/PMC5668325/)].

13. Kazemi Shishavan M, Asghari Jafarabadi M, Aminisani N, Shahbazi M, Alizadeh M. The association between self-care and quality of life in hypertensive patients: findings from the Azar cohort study in the North West of Iran. *Health Promot Perspect*. 2018;**8**(2):139–46. doi: [10.15171/hpp.2018.18](https://doi.org/10.15171/hpp.2018.18). [PubMed: [29744310](https://pubmed.ncbi.nlm.nih.gov/29744310/)]. [PubMed Central: [PMC5935818](https://pubmed.ncbi.nlm.nih.gov/PMC5935818/)].
14. Wang C, Lang J, Xuan L, Li X, Zhang L. The effect of health literacy and self-management efficacy on the health-related quality of life of hypertensive patients in a western rural area of China: a cross-sectional study. *Int J Equity Health*. 2017;**16**(1):58. doi: [10.1186/s12939-017-0551-9](https://doi.org/10.1186/s12939-017-0551-9). [PubMed: [28666443](https://pubmed.ncbi.nlm.nih.gov/28666443/)]. [PubMed Central: [PMC5493849](https://pubmed.ncbi.nlm.nih.gov/PMC5493849/)].
15. Kalroozi F, Mohammadi N, Farahani MA, Aski BH, Anari AM. A critical analysis of Iran health system reform plan. *J Educ Health Promot*. 2020;**9**:364. doi: [10.4103/jehp.jehp_493_20](https://doi.org/10.4103/jehp.jehp_493_20). [PubMed: [33575400](https://pubmed.ncbi.nlm.nih.gov/33575400/)]. [PubMed Central: [PMC7871970](https://pubmed.ncbi.nlm.nih.gov/PMC7871970/)].
16. Al-Khawaja M. Hypertension and health-related quality of life: A community based epidemiological study in an Egyptian Village. *J Med Arts*. 2020;**0**(0):0. doi: [10.21608/jma.2020.20529.1051](https://doi.org/10.21608/jma.2020.20529.1051).
17. Noori R, Alami A, Delshad Noghabi A. [Effect of family-centered empowerment model on quality of life in patients with hypertension]. *J Maz Univ Med Sci*. 2015;**25**(127):100–7. Persian.
18. Najafi GT, Shahidi M, Ghiasvandian S, Haghani H. [The effect of self-care educational program on quality of life in patients with hypertension]. *J Cardiovasc Nurs*. 2015;**4**(1). Persian.
19. Miao Y, Zhang L, Sparring V, Sandeep S, Tang W, Sun X, et al. Improving health related quality of life among rural hypertensive patients through the integrative strategy of health services delivery: a quasi-experimental trial from Chongqing, China. *Int J Equity Health*. 2016;**15**(1):132. doi: [10.1186/s12939-016-0421-x](https://doi.org/10.1186/s12939-016-0421-x). [PubMed: [27552845](https://pubmed.ncbi.nlm.nih.gov/27552845/)]. [PubMed Central: [PMC4995617](https://pubmed.ncbi.nlm.nih.gov/PMC4995617/)].
20. Salahi S, Ghahfarokhi J, Hasheminia SA, Habibzadeh H. [The effect of family-centered empowerment model on quality of life of hemodialysis patients]. *J Urmia Nurs Midwifery*. 2012;**10**(1):61–7. Persian.
21. Allen D, Scarinci N, Hickson L. The nature of patient- and family-centred care for young adults living with chronic disease and their family members: A systematic review. *Int J Integr Care*. 2018;**18**(2):14. doi: [10.5334/ijic.3110](https://doi.org/10.5334/ijic.3110). [PubMed: [30127698](https://pubmed.ncbi.nlm.nih.gov/30127698/)]. [PubMed Central: [PMC6095060](https://pubmed.ncbi.nlm.nih.gov/PMC6095060/)].
22. Farias DHR, Almeida MFF, Gomes GC, Lunardi VL, Queiroz MVO, Nornberg PKO, et al. Beliefs, values and practices of families in the care of hospitalized children: subsidies for nursing. *Rev Bras Enferm*. 2020;**73**(suppl 4). e20190553. doi: [10.1590/0034-7167-2019-0553](https://doi.org/10.1590/0034-7167-2019-0553). [PubMed: [33206850](https://pubmed.ncbi.nlm.nih.gov/33206850/)].
23. Nasiri S, Heydari N, Rafiee S, Paran M. [Effect of family-centered education on patient's self-care]. *Sadra Med J*. 2020;**8**(3):311–20. Persian. doi: [10.30476/smsj.2020.81889.1009](https://doi.org/10.30476/smsj.2020.81889.1009).
24. Kokorelias KM, Gignac MAM, Naglie G, Cameron JL. Towards a universal model of family centered care: a scoping review. *BMC Health Serv Res*. 2019;**19**(1):564. doi: [10.1186/s12913-019-4394-5](https://doi.org/10.1186/s12913-019-4394-5). [PubMed: [31409347](https://pubmed.ncbi.nlm.nih.gov/31409347/)]. [PubMed Central: [PMC6693264](https://pubmed.ncbi.nlm.nih.gov/PMC6693264/)].
25. Azar FEF, Solhi M, Chabaksvaf F. Investigation of the quality of life of patients with hypertension in health centers. *J Educ Health Promot*. 2020;**9**:185. doi: [10.4103/jehp.jehp_741_19](https://doi.org/10.4103/jehp.jehp_741_19). [PubMed: [32953912](https://pubmed.ncbi.nlm.nih.gov/32953912/)]. [PubMed Central: [PMC7482625](https://pubmed.ncbi.nlm.nih.gov/PMC7482625/)].
26. Liu L, Qian X, Chen Z, He T. Health literacy and its effect on chronic disease prevention: evidence from China's data. *BMC Public Health*. 2020;**20**(1):690. doi: [10.1186/s12889-020-08804-4](https://doi.org/10.1186/s12889-020-08804-4). [PubMed: [32410604](https://pubmed.ncbi.nlm.nih.gov/32410604/)]. [PubMed Central: [PMC7227325](https://pubmed.ncbi.nlm.nih.gov/PMC7227325/)].
27. Hedayati B. Effect of family-based empowerment model on the self-efficacy of hypertensive elderly people. *Salmand: Iran J Ageing*. 2018;**13**(1):86–97. doi: [10.21859/sija.13.1.86](https://doi.org/10.21859/sija.13.1.86).
28. Rostami S, Namaki Khameneh S, Keikhaei Dehdezi B, Haghhighizadeh MH. The effect of family-centered empowerment on quality of life in children with hemophilia. *Avicenna J Nurs Midwifery Care*. 2020;**28**(1):1–10. doi: [10.30699/ajnm.28.1.1](https://doi.org/10.30699/ajnm.28.1.1).
29. Nikbakht Nasrabadi A, Mohammadi N, Davatgaran K, Yekaninejad M, Noroozi Javidan A, Shabany M. Designing a client and family empowerment model to promote constructive life recovery among persons with spinal cord injury: A qualitative study. *Arch Neurosci*. 2019;**6**(3). doi: [10.5812/ans.87867](https://doi.org/10.5812/ans.87867).
30. Hedayati B. The effect of implementation of family-centered empowerment model on the self-esteem of the old people with hypertension. *J Educ Health Promot*. 2015;**4**(1):74.
31. Babaei-Sis M, Ranjbaran S, Mahmoodi H, Babazadeh T, Moradi F, Mirzaeian K. [The effect of educational intervention of life style modification on blood pressure control in patients with hypertension]. *J Educ Community Health*. 2016;**3**(1):12–9. Persian. doi: [10.21859/jech-03012](https://doi.org/10.21859/jech-03012).
32. Okhli A, Masoudi R. [The impact of family-based empowerment pattern on hemodialysis patients]. *Iran J Rehabilitation Res Nurs*. 2019;**5**(3):9–13. Persian.
33. Vahedian-Azimi A, Alhani F, Goharimogaddam K, Madani SJ, Naderi A, Hajiesmaeili M. [Effect of family-centered empowerment model on the quality of life in patients with myocardial infarction: A clinical trial study]. *J Nurs Educ*. 2015;**4**(1):8–22. Persian.
34. Shamsi A. [Development and validation of the quality of life questionnaire in the patients with hypertension] [dissertation]. Tehran, Iran: School of Nursing and Midwifery, Tehran University of Medical Sciences; 2017. Persian.
35. Holland K. *Everything you need to know about high blood pressure (hypertension)*. healthline; 2020, [updated June 17, 2020]. Available from: <https://www.healthline.com/health/high-blood-pressure-hypertension#symptoms-of-high-blood-pressure>.
36. Mohalli F, Mahmoudi Rad GH, Alhani F, Abdinejad Z, Foroozfar H. [The effect of family-centered empowerment model on the indicators of the ability of patients with hypertension]. *IJNR*. 2019. Persian.
37. Bairami S, Fathi Y, Mohammadinasab S, Barati M, Mohammadi Y. [Relationship between self-care behaviors and quality of life among hypertensive patients visiting comprehensive health centers in Hamadan, Iran]. *J Educ Community Health*. 2017;**4**(1):20–7. Persian. doi: [10.21859/jech.4.1.20](https://doi.org/10.21859/jech.4.1.20).
38. Keshvari M, Hedayati B, Moieni M, Alhani F. A survey on the effect of implementation of a family-centered empowerment model on blood pressure and empowerment dimensions in the elderly people with hypertension. *J Educ Health Promot*. 2015;**4**:94. [PubMed: [27462636](https://pubmed.ncbi.nlm.nih.gov/27462636/)]. [PubMed Central: [PMC4946264](https://pubmed.ncbi.nlm.nih.gov/PMC4946264/)].
39. Escortell-Mayor E, Del Cura-Gonzalez I, Ojeda-Ruiz E, Sanz-Cuesta T, Rodriguez-Salceda I, Garcia-Soltero J, et al. A primary health-care information intervention for communicating cardiovascular risk to patients with poorly controlled hypertension: The Education and Coronary Risk Evaluation (Educore) study-A pragmatic, cluster-randomized trial. *PLoS One*. 2020;**15**(1). e0226398. doi: [10.1371/journal.pone.0226398](https://doi.org/10.1371/journal.pone.0226398). [PubMed: [31971952](https://pubmed.ncbi.nlm.nih.gov/31971952/)]. [PubMed Central: [PMC6977759](https://pubmed.ncbi.nlm.nih.gov/PMC6977759/)].
40. Mini GK, Sarma PS, Thankappan KR. Cluster randomised controlled trial of behavioural intervention program: a study protocol for control of hypertension among teachers in schools in Kerala (CHATS-K), India. *BMC Public Health*. 2019;**19**(1):1718. doi: [10.1186/s12889-019-8082-5](https://doi.org/10.1186/s12889-019-8082-5). [PubMed: [31864339](https://pubmed.ncbi.nlm.nih.gov/31864339/)]. [PubMed Central: [PMC6925901](https://pubmed.ncbi.nlm.nih.gov/PMC6925901/)].
41. Gong J, Chen X, Li S. Efficacy of a community-based physical activity program km2h2 for stroke and heart attack prevention among senior

- hypertensive patients: A cluster randomized controlled phase-ii trial. *PLoS One*. 2015;**10**(10). e0139442. doi: [10.1371/journal.pone.0139442](https://doi.org/10.1371/journal.pone.0139442). [PubMed: [26426421](https://pubmed.ncbi.nlm.nih.gov/26426421/)]. [PubMed Central: [PMC4591119](https://pubmed.ncbi.nlm.nih.gov/PMC4591119/)].
42. Atak N, Gurkan T, Kose K. The effect of education on knowledge, self management behaviours and self efficacy of patients with type 2 diabetes. *Aust J Adv Nurs*. 2008;**26**(2):66-74.
 43. Sargazi Shad T, Kermansaravi F, Navidian A. [Effect of the family-centered empowerment model on quality of life and self-efficacy in adolescents with type 1 diabetes referring to the Ali Asghar's clinic in Zahedan, 2016]. *Iran J Endocrinol Metab*. 2018;**19**(5):330-9. Persian.
 44. Pilevar N, Ramezani M, Malek A, Behnam Vashani H. Effect of implementing family-centered empowerment model on the quality of life in school-age children diagnosed with rheumatoid arthritis. *Evidence Based Care*. 2019;**9**(2):65-73. doi:[10.22038/ebcj.2019.39702.2046](https://doi.org/10.22038/ebcj.2019.39702.2046).
 45. Davarpanah M, Fayazi S, Shariati A, Mirhosseini SD. The effect of family-centered empowerment model on the quality of life of patients with leukemia. *Jundishapur J Chronic Dis Care*. 2017;**6**(1). doi:[10.17795/jjcdc-36441](https://doi.org/10.17795/jjcdc-36441).
 46. Rajabi R, Mahmoudirad G, Sabzevari SS, Borhani F, Yazdanparast E, Faramarzpor M. The effect of family-centered empowerment model on quality of life among mothers of asthmatic children. *Mod Care J*. 2016;**13**(1). doi:[10.17795/modernc.9602](https://doi.org/10.17795/modernc.9602).