



Association Between the Chemical Composition of Saliva and Serum in Patients with Kidney Failure

Alireza Rajolani¹, Arezoo Alaei², Mohsen Nafar³, Mohammad-Javad Kharazi-Fard⁴, Kimia Ghods^{2,*}

¹ School of Dental, Islamic Azad University of Medical Sciences, Tehran, Iran

² Dental Material Research Center, School of Dentistry, Islamic Azad University of Medical Sciences, Tehran, Iran

³ Urology and Nephrology Research Center, Shahid Labbafinejad Medical Center, Shahid Beheshti University of Medical Sciences, Tehran, Iran

⁴ Dental Research Center, School of Dentistry, University of Medical Sciences, Tehran, Iran

* Corresponding author: Dental Material Research Center, School of Dentistry, Islamic Azad University of Medical Sciences, Tehran, Iran. Email: kimiaghods2@gmail.com

Received 2024 February 3; Revised 2024 February 18; Accepted 2024 April 26.

Abstract

Background: End-Stage Renal Disease (ESRD) is a severe nephrological condition that can lead to permanent kidney damage. Therefore, early disease diagnosis is key to preventing casualties. The gold standard method of diagnosis tends to evaluate changes in sodium, potassium, calcium, phosphorus, urea, creatinine, and parathormone hormones in blood after dialysis. However, serum evaluation is not always possible or easy for patients. Therefore, saliva evaluation has been proposed in recent years as an alternative.

Objectives: The current article aims to evaluate metabolite in the saliva of ESRD patients.

Methods: In this descriptive-analytical study, 29 ESRD patients undergoing hemodialysis treatment were selected. Their saliva and serum samples were taken. The number of biochemical factors, including sodium, potassium, calcium, phosphorus, urea, creatinine, and parathormone hormone, was measured with an autoanalyzer device and related kits. Finally, the correlation of parameters in serum and saliva was examined using the Pearson test.

Results: The results showed a significant positive relationship between the levels of sodium, urea, and creatinine in serum and saliva samples ($P < 0.05$). On the other hand, there was no significant relationship between the serum and saliva levels of potassium, calcium, phosphorus, and PTH ($P > 0.05$).

Conclusions: Due to the significant correlation between some prominent biochemical factors in saliva and serum after hemodialysis in ESRD patients, saliva could be considered a non-invasive diagnostic fluid for monitoring kidney disease in the future.

Keywords: End Stage Renal Disease, Hemodialysis, Sodium, Calcium, Potassium, Phosphorus, Urea, Creatinine, Parathyroid Hormone, Saliva

1. Background

End-Stage Renal Disease (ESRD) is a severe form of kidney disease in which the patient's kidneys stop functioning permanently and irreversibly. This condition usually develops after primary kidney diseases or secondary factors such as aging, diabetes mellitus, and hypertension (1).

Since different organs of the body are affected by uremia caused by chronic kidney failure, ESRD usually manifests several signs and symptoms, including hypertension, congestive heart failure, pulmonary

edema, pericarditis, skin rash, purpura, Kussmaul breathing, uremic pneumonia, ammonia breath, diarrhea, gastrointestinal bleeding, anemia, thrombocytopenia, amenorrhea, testicular atrophy, muscle cramps, and renal osteodystrophy (2-7).

The common treatment for ESRD patients is hemodialysis. Due to metabolite changes resulting from kidney malfunction, some medications, including antacids and antihypertensives, are also prescribed (8, 9).

Electrolyte balance is crucial for maintaining the proper functioning of various organs and is disrupted

in ESRD patients. Some of the most prominent electrolytes are sodium, potassium, calcium, phosphorus, urea, creatinine, and parathormone hormones (PTH). However, the excretion of all the electrolytes except calcium is reduced after hemodialysis. Hence, regular assessment of these electrolytes and monitoring metabolite consumption are critical for ESRD patients (10, 11).

Although serum is used as the gold standard method for electrolyte assessment, saliva has attracted much attention in recent research. Due to the easy and non-invasive method of saliva collection and the possible close relationship between saliva and serum parameters, saliva is now considered a unique fluid for the diagnosis of ESRD (12).

2. Objectives

This study aims to investigate the correlation between the levels of sodium, potassium, phosphorus, calcium, urea, creatinine, and PTH in serum and saliva samples collected from ESRD patients undergoing hemodialysis treatment.

3. Methods

This descriptive-analytical study was performed on dialysis patients referred to Amir al-Momenin Hospital in Tehran in 2019. Purpose-based sampling was conducted among male and female volunteers aged 18 to 75 who were suffering from End-stage renal disease. End-stage renal disease was diagnosed when the glomerular filtration rate of the patient was under 15 mL/min and the last dialysis had occurred over 6 months ago (13). In addition, people under 18 years of age, recent kidney transplant recipients, smokers and alcohol users, patients with respiratory diseases and allergic reactions, and people who consumed caffeine 24 hours before the test were excluded from the study. In terms of age, gender, and body mass index, all patients were randomly selected based on the inclusion criteria (14).

After determining inclusion and exclusion criteria, PASS 15 software, considering $\beta = 0.2$, $\alpha = 0.05$, and a standard deviation of 0.95, was used to achieve a sample size of 29.

The present study's protocol followed the Declaration of Helsinki. The ethics committee of the Islamic Azad

University Faculty of Dentistry, Tehran, Iran (IR.IAU.DENTAL.REC.1399.164) approved the study.

After explaining the steps to the volunteers, all eligible participants were asked to complete and sign the informed consent form. They were then asked to carefully fill out the demographic information form.

Subsequent to completing the consent and demographic forms, blood sampling was carried out before dialysis treatment for fasting patients (12 hours) by coordinating with the dialysis department. Ten cubic millimeters of ambient blood was taken from all participants and then centrifuged for 5 minutes at 3500 rpm. These samples were kept in a clot activator at -20 degrees Celsius for up to 2 months before the test. Clot activators are tubes coated with silicone, which causes the blood to clot without any other substances and allows the serum to be separated by centrifugation (14).

Saliva was collected from all the patients on the same day using the spitting method. To collect saliva, participants were asked to stop eating, drinking, and smoking one hour before the process. Next, patients were asked to allow saliva to accumulate on the floor of the mouth and then spit it into pre-weighed tubes while maintaining an upright position (15). In this method, 5 cubic millimeters of saliva was collected and placed in clean and dry paraffinized Falcon polyethylene tubes. The samples were placed on ice for 5 minutes and immediately centrifuged at 4400 RPM for 15 minutes to separate any impurities. The samples were kept at -20 degrees Celsius until testing (14).

To check serum chemical compounds including urea, creatinine, sodium, calcium, and phosphorus, a BT3000 autoanalyzer based on colorimetry was utilized. The flame photometric technique was applied to determine the amount of potassium in the serum. The Landwind Electrolyte Lwe60se device and the corresponding washing buffers were used to examine the chemical composition of saliva. To measure PTH in the serum, a Monobind kit and ELISA technique were used, and a Stat Fax 4300 chromate machine was employed for reading.

3.1. Statistical Analysis

SPSS software version 26 was used for data analysis. A paired t-test was employed to analyze the data. Moreover, Pearson's correlation coefficient was used to investigate the relationship between saliva and serum variables. In addition, values less than 0.05 and 0.01

were considered statistically significant for intergroup and intragroup comparisons, respectively.

4. Results

In this study, 29 patients (16 men and 13 women) with ESRD, with an average age of 58.27 years (minimum age of 28 and maximum age of 71 years), were investigated. The amounts of each electrolyte in both serum and saliva in ESRD patients can be observed in [Table 1](#).

Table 1. The Amount of Serum and Salivary Variables Measured in ESRD Patients

Parameters	N	Minimum	Maximum	Mean
Saliva				
Na (mEq/L)	29	0.90	5.90	3.9 ± 1.15
K (mEq/L)	29	3.10	7.80	4.63 ± 0.89
Ca (mEq/L)	29	0.48	5.30	2.03 ± 1.08
P (mEq/L)	29	1.60	7.60	3.47 ± 1.50
Urea (mg/dl)	29	31.90	72.80	50.14 ± 10.79
Cr (mg/dl)	29	0.61	1.96	1.21 ± 0.35
PTH (pg/dL)	29	0.30	1.80	1.16 ± 0.32
Serum				
Na (mEq/L)	29	130	148	141.34 ± 4.18
K (mEq/L)	29	3.50	6.00	4.51 ± 0.83
Ca (mEq/L)	29	7.50	9.20	8.37 ± 0.41
P (mEq/L)	29	3.60	6.20	4.78 ± 0.69
Urea (mg/dl)	29	67	158	108.66 ± 25.48
Cr (mg/dl)	29	3.90	13.20	8.46 ± 2.50
PTH (pg/dL)	29	62.00	2614.00	579.75 ± 562.23

4.1. Cross-Sectional Analysis of Parameters According to Their Concentration Range in Serum and Saliva

Regarding the sodium variable, all the salivary samples in the normal and less-than-normal ranges were also in the same range as the serum samples.

When the potassium variable was compared, 75% of salivary samples in the normal range were also in the same range as the serum samples. However, none of the salivary samples in the more-than-normal range matched the serum samples.

For the calcium variable, all the salivary samples in the less-than-normal range were also in the same range as the serum samples. Nonetheless, 46.2% of salivary samples in the normal range were also in the same range as the serum samples. However, none of the salivary samples in the more-than-normal range matched the serum samples.

The phosphorus variable showed different outcomes. 40.7% of salivary samples in the normal range were also

in the same range as the serum samples. Moreover, 50% of the salivary samples in the more-than-normal range matched the serum samples.

When it comes to urea, creatinine, and PTH, all the salivary samples in the more-than-normal range matched the serum samples. On the other hand, 10% of PTH salivary samples in the normal range were also in the same range as the serum samples.

4.2. The Correlation of Biochemical Factors and Electrolytes Between Saliva and Serum

The results obtained from the analysis showed a significant positive correlation between the amounts of sodium, urea, and creatinine in saliva and serum ($P < 0.05$). This finding contrasts with the amounts of potassium, calcium, phosphorus, and PTH assessed in serum and saliva, which showed no significant correlation ($P > 0.05$).

Other positive correlations between different variables in saliva and serum were observed during this research. For instance, there was a significant positive correlation between salivary phosphorus and serum urea, salivary urea and serum phosphorus and creatinine, and salivary creatinine and serum phosphorus and urea ($P < 0.05$).

After intergroup comparison, intragroup evaluation was conducted for variables in both serum and saliva samples. In the serum group, a notable positive correlation was observed between calcium and creatinine, phosphorus and urea, phosphorus and creatinine, and urea and creatinine ($P < 0.01$). In the saliva group, a significant positive correlation was noted between potassium and calcium, potassium and phosphorus, potassium and PTH, calcium and PTH, phosphorus and PTH, phosphorus and urea, and urea and creatinine ($P < 0.01$).

A summary of the correlation between different variables in serum and saliva is shown in [Table 2](#).

In [Figure 1](#), the correlation between different electrolytes in serum and saliva is exhibited. The vertical axis shows the amount of the designated variable in saliva, and the horizontal axis shows the amount of the same variable in serum.

5. Discussion

The present study indicated that the amounts of some major electrolytes, including sodium, urea, and

Table 2. Correlation Between Variables in Serum and Saliva

Variables	Saliva Variables							Serum Variables						
	Na	K	Ca	P	Urea	Cr	PTH	Na	K	Ca	P	Urea	Cr	PTH
R														
Na	1	0.289	0.215	0.073	-0.088	0.221	-0.115	0.651	0.122	0.018	0.188	-0.038	0.217	0.016
K	0.289	1	0.621	0.409	-0.191	-0.011	-0.462	0.204	-0.017	0.009	-0.070	-0.118	0.016	0.096
Ca	0.215	0.621	1	0.338	-0.097	0.220	-0.470	0.228	0.080	-0.080	0.212	-0.040	0.249	0.066
P	0.073	0.409	0.338	1	-0.451	-0.220	-0.380	-0.155	0.099	0.185	-0.191	-0.396	-0.212	0.036
Urea	-0.088	-0.191	-0.097	-0.451	1	0.539	0.264	0.028	0.119	-0.242	0.520	0.955	0.455	-0.036
Cr	0.221	-0.011	0.220	-0.220	0.539	1	0.182	0.261	0.282	-0.355	0.413	0.571	0.959	0.229
PTH	-0.115	-0.462	-0.470	-0.38	0.264	0.182	1	-0.159	-0.197	-0.007	0.161	0.261	0.139	0.365
R														
Na	0.651	0.204	0.228	-0.155	0.028	0.261	-0.159	1	0.141	0.242	0.272	0.012	0.269	-0.242
K	0.122	-0.017	0.080	0.099	0.119	0.282	-0.197	0.141	1	-0.268	0.165	0.145	0.304	-0.252
Ca	0.018	0.009	-0.080	0.185	-0.242	-0.355	-0.007	0.242	-0.268	1	-0.183	-0.354	-0.388	-0.148
P	0.188	-0.070	0.212	-0.191	0.520	0.413	0.161	0.272	0.165	-0.183	1	0.531	0.423	0.065
Urea	-0.038	-0.118	-0.040	-0.396	0.955	0.571	0.261	0.012	0.145	-0.354	0.531	1	0.532	-0.069
Cr	0.217	0.016	0.249	-0.212	0.455	0.959	0.139	0.269	0.304	-0.388	0.423	0.532	1	0.230
PTH	0.016	0.096	0.066	0.036	-0.036	0.229	0.365	-0.242	-0.252	-0.148	0.065	-0.069	0.230	1

creatinine, show a positive correlation between serum and saliva. Additionally, the cross-sectional analysis of different parameters demonstrated that the normal range of the studied variables in serum is completely related to their normal range in saliva. Therefore, assessing these parameters in saliva can be utilized for monitoring ESRD patients.

Several studies have attempted to exhibit the correlation between different biomarkers in the serum and saliva of ESRD patients undergoing hemodialysis (12, 14, 16-22). However, the findings of these studies are often contrary to each other and the present study.

Regarding the sodium variable, one study by Bagalad et al. (19) compared the amount of this parameter in saliva and serum in ESRD patients, similar to the present study. Both studies exhibited a positive correlation between serum and saliva. However, another study by Seethalakshmi et al. (16) showed no significant positive correlation between serum and saliva sodium levels.

Four studies, including the present research, showed no positive correlation between serum and saliva levels of potassium, calcium, and phosphorus (12, 14, 19, 22). These outcomes differ from the findings of Seethalakshmi et al. (16), which demonstrated a positive correlation between potassium and phosphorus levels in saliva and serum.

Even though the mentioned studies showed somewhat aligned results regarding sodium,

potassium, calcium, and phosphorus variables, the correlation of other parameters in serum and saliva varies across studies. For instance, a study by Rodrigues et al. (12) showed a positive correlation between serum and saliva levels of PTH, which is contrary to the findings of the present study.

Four studies illustrated a positive correlation between serum and saliva levels of creatinine, similar to the present study (17, 19-21). This finding differs from the results of the study conducted by Franco et al. (23). All the mentioned studies seem to agree on the positive correlation between serum and saliva levels of urea, which aligns with the results of this study (12, 16-22).

The reason behind the differences in the results of various studies remains unclear. Some theories include the inclusion and exclusion criteria of the studies, the existence of true ESRD patients, the methods used to collect saliva and blood, and the analyzer devices. Although all the studies compared the number of different parameters in serum and saliva, none of them performed intragroup comparisons, unlike the present study. This lack of information could question the reliability of the suggested method in the future. Therefore, more research is needed in this field.

The cross-sectional analysis of the present study indicated that the salivary concentrations of sodium, urea, and creatinine are completely dependent on their serum concentrations. This finding can be related to the




Figure 1. The correlation between each parameter in serum and saliva. The horizontal axis shows the amount of electrolyte in the serum. The vertical axis shows the amount of electrolyte in the saliva. A, Sodium; B, Potassium; C, Calcium; D, Phosphorus; E, Urea; F, Creatinine; G, PTH

reabsorption of sodium in the ducts of the salivary glands, which can explain this positive correlation. Moreover, considering that urea enters saliva through simple diffusion from plasma, it can be concluded that one of the main reasons for the strong correlation between serum and salivary concentrations of urea is this issue. On the other hand, it seems that the mechanism of creatinine entering saliva in ESRD patients is as follows: Creatinine increases in the serum under disease conditions, and increased serum creatinine in these patients creates a concentration

gradient that facilitates the release of creatinine from serum to saliva. This condition is probably due to changes in the permeability of salivary gland cells. Therefore, the correlation between serum and saliva levels of sodium, urea, and creatinine seems logical.

In order to evaluate kidney function in ESRD patients and determine the timing of hemodialysis, multiple blood samples are required. The blood sampling method is invasive, and the risk of infection transmission is high. However, the saliva sampling method is non-invasive, the risk of infection

transmission is low, and sampling is easy and accessible. Therefore, considering saliva sampling as an alternative method can be discussed in the future.

It should be noted that this study included only 29 patients with ESRD, which can limit the reliability of the study's outcome. Additionally, focusing solely on ESRD patients and lacking a control group consisting of individuals with healthy systemic conditions cannot precisely describe and compare electrolyte actions in the body. Therefore, future studies should consider including healthy individuals for better analysis and comparison of metabolites in the saliva and serum of both healthy and ESRD patients.

5.1. Conclusions

The results of this study exhibited a strong, meaningful relationship in the kidney function monitoring markers, including sodium, urea, and creatinine, between serum and saliva. This finding suggests it is possible to use saliva samples as a representative to evaluate kidney function monitoring factors instead of serum samples in ESRD patients. However, additional studies with a larger sample size are needed in the future to clarify this issue.

Acknowledgements

Special thanks to Dr. Hamid Rajolani, Mr. Naser Tabkhi, Mrs. Masoumeh Atai (The head of the dialysis department at Amirul Mominin Hospital), Mrs. Laleh Khosravi, and Mrs. Mozghan Nasiri, who helped the researcher in conducting this research.

Footnotes

Authors' Contribution: Study concept and design: A.A, acquisition of data: A.R, analysis and interpretation of data: A.R, drafting of the manuscript: K.G, critical revision of the manuscript for important intellectual content: M.N, statistical analysis: M.K, administrative, technical, and material support: A.R, study supervision: A.A.

Conflict of Interests Statement: Authors declared no conflict of interests.

Data Availability: The data presented in this study are uploaded during submission as a supplementary file and are openly available for readers upon request.

Ethical Approval: IR.IAU. DENTAL.REC.1399.164 .

Funding/Support: This study did not receive any grants or funding.

Informed Consent: After explaining the steps to the volunteers, all eligible participants were asked to complete and sign the informed consent form. They were then asked to carefully fill out the demographic information form.

References

1. Xiang X, Peng B, Liu K, Wang T, Ding P, Li H, et al. Association between salivary microbiota and renal function in renal transplant patients during the perioperative period. *Front Microbiol.* 2023;**14**:1122101. [PubMed ID: 37065138]. [PubMed Central ID: PMC10090686]. <https://doi.org/10.3389/fmicb.2023.1122101>.
2. Chen Y, Cheng C. Application of rough set classifiers for determining hemodialysis adequacy in ESRD patients. *J Knowledge information systems.* 2013;**34**:453-82.
3. Salles Junior LD, Santos PR, dos Santos AA, de Souza MH. Dyspepsia and gastric emptying in end-stage renal disease patients on hemodialysis. *BMC Nephrol.* 2013;**14**:275. [PubMed ID: 24330675]. [PubMed Central ID: PMC3867417]. <https://doi.org/10.1186/1471-2369-14-275>.
4. Lima NA, Vasconcelos CC, Filgueira PH, Kretzmann M, Sindeaux TA, Feitosa Neto B, et al. Review of genitourinary tuberculosis with focus on end-stage renal disease. *Rev Inst Med Trop Sao Paulo.* 2012;**54**(1):57-60. [PubMed ID: 22370756]. <https://doi.org/10.1590/s0036-46652012000100011>.
5. Ng MSN, Wong CL, Choi KC, Hui YH, Ho EHS, Miaskowski C, et al. A Mixed Methods Study of Symptom Experience in Patients With End-Stage Renal Disease. *Nurs Res.* 2021;**70**(1):34-43. [PubMed ID: 32991531]. <https://doi.org/10.1097/NNR.0000000000000479>.
6. Lin DS, Lin FJ, Lin YS, Lee JK, Lin YH. The effects of mineralocorticoid receptor antagonists on cardiovascular outcomes in patients with end-stage renal disease and heart failure. *Eur J Heart Fail.* 2023;**25**(1):98-107. [PubMed ID: 36404402]. <https://doi.org/10.1002/ehf.2740>.
7. Suh SH, Jung JH, Oh TR, Yang EM, Choi HS, Kim CS, et al. Rheumatoid arthritis and the risk of end-stage renal disease: A nationwide, population-based study. *Front Med (Lausanne).* 2023;**10**:1116489. [PubMed ID: 36817794]. [PubMed Central ID: PMC9932810]. <https://doi.org/10.3389/fmed.2023.1116489>.
8. Kyriakoulis I, Adamou A, Stamatiou I, Chlorogiannis DD, Kardoutsos I, Koukousaki D, et al. Efficacy and safety of direct oral anticoagulants vs vitamin K antagonists in patients with atrial fibrillation and end-stage renal disease on hemodialysis: A systematic review and meta-analysis. *Eur J Intern Med.* 2024;**119**:45-52. [PubMed ID: 37648582]. <https://doi.org/10.1016/j.ejim.2023.08.020>.
9. Singh A, Garg R, Lan N, Siddiqui MT, Gupta M, Alkhouri N. Association between anti-acid therapies and advanced fibrosis in type 2 diabetics with biopsy-proven non-alcoholic fatty liver disease. *Indian J Gastroenterol.* 2020;**39**(6):591-8. [PubMed ID: 33219985]. [PubMed Central ID: PMC9254737]. <https://doi.org/10.1007/s12664-020-01087-y>.
10. Chao YT, Lin YK, Chen LK, Huang P, Hsu YC. Role of the gut microbiota and their metabolites in hemodialysis patients. *Int J Med Sci.*

- 2023;**20**(6):725-36. [PubMed ID: [37213669](#)]. [PubMed Central ID: [PMC10198149](#)]. <https://doi.org/10.7150/ijms.82667>.
11. Shen Y, Liu C, He H, Zhang M, Wang H, Ji K, et al. Recent Advances in Wearable Biosensors for Non-Invasive Detection of Human Lactate. *Biosensors (Basel)*. 2022;**12**(12). [PubMed ID: [36551131](#)]. [PubMed Central ID: [PMC9776101](#)]. <https://doi.org/10.3390/bios12121164>.
12. Rodrigues VP, Franco MM, Marques CP, de Carvalho RC, Leite SA, Pereira AL, et al. Salivary levels of calcium, phosphorus, potassium, albumin and correlation with serum biomarkers in hemodialysis patients. *Arch Oral Biol*. 2016;**62**:58-63. [PubMed ID: [26655748](#)]. <https://doi.org/10.1016/j.archoralbio.2015.11.016>.
13. Hashmi MF, Benjamin O, Lappin SL. *End-stage renal disease*. Treasure Island (FL): StatPearls Publishing; 2023. [PubMed ID: [29763036](#)].
14. Hashemi J, Hesari Z, Golshan A. [Evaluation of calcium, phosphorus and potassium in saliva and their relationship to blood biochemical factors in hemodialysis patients]. *ehran University Medical Journal*. 2017. Persian.
15. Ghods K, Motamed B, Alaee A, Farrokhnia T. Review of Oral Saliva Measurement Standard Methods. *J Res Dent Sci*. 2021;**18**(2):150-60.
16. Seethalakshmi C, Koteeswaran D, Chiranjeevi V. Correlation of serum and salivary biochemical parameters in end stage renal disease patients undergoing hemodialysis in pre and post-dialysis state. *J Clinical Diagnostic Research: JCDR*. 2014;**8**(12):CC12.
17. Alpdemir M, Eryilmaz M, Alpdemir MF, Topcu G, Azak A, Yucel D. Comparison of Widely Used Biochemical Analytes in the Serum and Saliva Samples of Dialysis Patients. *J Med Biochem*. 2018;**37**(3):346-54. [PubMed ID: [30598632](#)]. [PubMed Central ID: [PMC6298461](#)]. <https://doi.org/10.1515/jomb-2017-0056>.
18. Trzcionka A, Twardawa H, Mocny-Pachonska K, Korkosz R, Tanasiewicz M. Oral Mucosa Status and Saliva Parameters of Multimorbid Adult Patients Diagnosed with End-Stage Chronic Kidney Disease. *Int J Environ Res Public Health*. 2021;**18**(23). [PubMed ID: [34886240](#)]. [PubMed Central ID: [PMC8657397](#)]. <https://doi.org/10.3390/ijerph182312515>.
19. Bagalad BS, Mohankumar KP, Madhushankari GS, Donoghue M, Kuberappa PH. Diagnostic accuracy of salivary creatinine, urea, and potassium levels to assess dialysis need in renal failure patients. *Dent Res J (Isfahan)*. 2017;**14**(1):13-8. [PubMed ID: [28348612](#)]. [PubMed Central ID: [PMC5356383](#)]. <https://doi.org/10.4103/1735-3327.201138>.
20. Lasisi TJ, Raji YR, Salako BL. Salivary creatinine and urea analysis in patients with chronic kidney disease: a case control study. *BMC Nephrol*. 2016;**17**:10. [PubMed ID: [26775026](#)]. [PubMed Central ID: [PMC4715295](#)]. <https://doi.org/10.1186/s12882-016-0222-x>.
21. Renda R. Can salivary creatinine and urea levels be used to diagnose chronic kidney disease in children as accurately as serum creatinine and urea levels? A case-control study. *Ren Fail*. 2017;**39**(1):452-7. [PubMed ID: [28372505](#)]. [PubMed Central ID: [PMC6446173](#)]. <https://doi.org/10.1080/0886022X.2017.1308256>.
22. Bilancio G, Cavallo P, Lombardi C, Guarino E, Cozza V, Giordano F, et al. Salivary levels of phosphorus and urea as indices of their plasma levels in nephropathic patients. *J Clin Lab Anal*. 2018;**32**(7). e22449. [PubMed ID: [29603373](#)]. [PubMed Central ID: [PMC6816964](#)]. <https://doi.org/10.1002/jcla.22449>.
23. Franco G, Saab R, Pizzatto LV, Torres MF, Fregoneze AP, Brancher JA. Analysis of salivary pH, flow rate, buffering capacity, concentrations of calcium, urea and total proteins in 2-8 years-old children with Down's syndrome. *J RSBO Revista Sul-Brasileira de Odontologia*. 2014;**11**(1):66-70.