



Efficacy and Complications Comparison Between Emergent and Elective Laparoscopic Cholecystectomy Surgery

Hamed Gholizadeh¹, Mohsen Andisheh¹, Mohammad Raeeszadeh¹, Soleyman Heydari¹, Jamal Akhavan-Moghaddam¹, Mahdi Morshedi¹, Maryaam Rezaee¹, Hasanali Mohebbi¹, Mehrdad Daryabor^{1,*}

¹Department of Surgery, Trauma Research Center, Baqiyatallah University of Medical Sciences, Tehran, Iran

*Corresponding Author: Department of Surgery, Trauma Research Center, Baqiyatallah University of Medical Sciences, Tehran, Iran. Email: dr.mehrdad.daryabor@gmail.com

Received: 10 January, 2024; Revised: 8 October, 2024; Accepted: 29 October, 2024

Abstract

Background: Laparoscopic cholecystectomy (LC) is the preferred surgical procedure for treating symptomatic gallstone (GS) disease.

Objectives: This study aimed to evaluate and compare the outcomes of emergent and elective LC, focusing on complications, conversion rates, duration of surgery (DOS), length of hospital stay, and post-operative pain.

Methods: A total of 166 patients who underwent LC surgery at Baqiyatallah Hospital, Tehran, Iran, between 2021 and 2022 were included in the study. Of these, 64 patients underwent emergent LC, and 102 underwent elective LC. The groups were compared for complications, conversion rates, DOS, length of hospital stay, and post-operative pain.

Results: The study revealed that while intraoperative complications did not differ significantly between the two groups ($P = 0.14$), there was a significant difference in the conversion of surgical methods from three to four trocars ($P = 0.007$). Additionally, more patients in the emergent group required drain insertion during surgery ($P = 0.003$). Regarding post-operative outcomes, no significant differences were observed between the emergent and elective LC groups in terms of local ($P = 0.77$) and systemic complications ($P = 0.37$). However, patients in the elective LC group experienced a significantly shorter post-operative hospital stay ($P = 0.002$). Pain levels one day (1.02 ± 0.80) and one week (0.14 ± 0.43) after surgery were lower in the elective LC group. Furthermore, patients in the elective group returned to daily activities earlier (3.77 ± 1.21 days).

Conclusions: The findings of this study indicate that emergent LC is as safe as elective LC. However, patients undergoing elective LC may benefit from a shorter recovery period and reduced post-operative pain. Overall, LC remains a low-risk and safe option for managing emergent gallstone cases.

Keywords: Acute Cholecystitis, Laparoscopic Cholecystectomy, Surgery Complications

1. Background

Laparoscopic cholecystectomy (LC) is widely regarded as the most effective and safe treatment for symptomatic cholelithiasis, making it the preferred surgical option. Among individuals with symptomatic gallstones (GS), approximately 10% develop acute cholecystitis (AC), accounting for about 750,000 cases annually. Laparoscopic cholecystectomy provides notable advantages over conventional cholecystectomy, including reduced postoperative pain, faster recovery, quicker restoration of bowel function, and shorter hospital stays. Despite its widespread use, certain cases of LC may require conversion to open cholecystectomy

(OC) due to various intraoperative challenges. Conversion ensures safe completion of the procedure but can prolong the expected surgical duration. Existing literature reports a conversion rate ranging from approximately 2% to 10% (1-6). The primary goal of LC is to achieve symptom relief and survival without impairment. However, adverse outcomes following LC can significantly impact these objectives. These complications range from minor issues, such as surgical site infections, to more severe outcomes like bile duct injury (BDI) and even mortality (7). Persistent postoperative pain remains a considerable issue, often leading to extended hospital stays or additional medical visits. Due to significant variability in pain management

protocols, a standardized approach is needed to ensure effective interventions for pain reduction (8).

During the early stages of laparoscopic surgery, AC was considered a relative contraindication for LC. However, subsequent research has challenged this notion, advocating for early surgical intervention in AC cases. Laparoscopic cholecystectomy performed within 72 hours of symptom onset is now endorsed for its benefits, including shorter hospital stays, reduced healthcare costs, and lower readmission rates. Despite these advantages, many general surgeons still prefer a non-operative approach for initial AC management, followed by delayed LC. Arguments against early LC often cite increased technical difficulty, higher conversion rates (6 - 35% in some studies), and greater risks of biliary complications, such as bile leaks and common bile duct (CBD) injuries, when operating on inflamed gallbladders with swollen tissues and distorted anatomy.

The timing of LC in cases of AC remains a contentious issue. Studies conducted in Europe, Japan, and the United States reveal significant disparities in practices, with early LC rates ranging from 11% to 89% among surgeons (9-11). Numerous studies have compared the outcomes and complications of elective versus emergent cholecystectomy (12-14), reinforcing a general belief that emergent LC may result in higher rates of intraoperative and postoperative complications (15, 16).

2. Objectives

Consequently, this study aims to compare the intraoperative and postoperative outcomes and complications of LC in patients undergoing emergent versus elective procedures.

3. Methods

3.1. Study Design

This case-control study was conducted in the Surgery Department of Baqiyatallah Hospital, Tehran, Iran, from January 2021 to December 2022. The final data analysis was completed in June 2023. Patients who presented to the department with clinical symptoms and were confirmed to have GS with cholelithiasis and scheduled for surgery were included in the study. Patients who were pregnant, had a history of previous OC, or had immune system deficiencies were excluded from the study.

The study population was divided into two groups: (1) the emergent group, and (2) the elective group. Patients

admitted in acute and emergency conditions who underwent surgery within 24 hours of symptom onset were categorized into the emergent group (n = 64). Conversely, patients scheduled for LC as part of a planned procedure were categorized into the elective group (n = 102). The study design is illustrated in Figure 1.

The surgeries for both groups were performed by surgeons following a standardized procedure. Laparoscopic cholecystectomy surgery commenced with the patient positioned in the Trendelenburg position, typically utilizing three ports. If necessary, a fourth port was added. This setup allowed for the insertion of long surgical instruments and a surgical video camera. The surgeon observed the gallbladder (GB) and surrounding organs on a monitor connected to the camera, controlling the telescopic instruments externally while monitoring the procedure on the display.

The surgery began with the elevation of the GB fundus. Dissection using diathermy proceeded in Calot's triangle to expose the cystic artery and cystic duct. An intraoperative cholangiogram was performed at the surgeon's discretion. The cystic duct and artery were clipped and divided, and the GB was removed from the liver bed in a retrograde manner. The GB was extracted through either the umbilical or epigastric port using an endoscopic retrieval bag.

The umbilical fascia and skin were closed with sutures, and a local anesthetic was administered at the surgical wound sites. Postoperatively, participants in both groups were followed up until suture removal at the port site (one week after surgery) and monitored for complications up to three months post-surgery.

3.2. Data Collection

All patient-related details, including medical history, general and detailed physical examinations, comorbidities, any history of prior abdominal surgeries, blood investigations, abdominal sonography, and endosonographic findings, were meticulously documented. Intraoperative complications, surgical findings, and postoperative recovery parameters were also recorded.

Postoperatively, data on the duration of hospital stay, post-surgery pain levels, and any conversions to OC were gathered. Additionally, all patients were followed up for three months after surgery to monitor outcomes and ensure comprehensive assessment.

3.3. Ethical Consideration




Figure 1. Study design flow chart

This study received approval from the Research Ethics Committee of Baqiyatallah University of Medical Sciences, under the ethical code [IR.BMSU.REC.1397.034](#), and was conducted in accordance with the principles outlined in the Declaration of Helsinki (1975). All participants provided written informed consent to participate in the study.

3.4. Statistics

The default alpha error was set at 0.05, and the beta error was set at 0.2. To calculate the sample size, post-surgery hospitalization days were considered the main variable. According to a similar study conducted by Marwah et al., the post-surgery hospitalization duration was 2.76 ± 0.663 days in the acute biliary pancreatitis group and 2.34 ± 0.861 days in the acute cholecystitis group ($P = 0.037$) (17). Using a 1:3 ratio for the emergent to elective groups, the sample size was calculated based on the post-surgery hospitalization days variable using G*Power software. The required sample size was determined to be 134 patients, with 34 patients allocated to the emergent group and 100 to the elective group. To account for the possibility of discontinuation by patients in the emergent group, the sample size for this group was increased from 34 to 40 participants.

Data were analyzed using SPSS software (v. 29). The chi-square test was used to compare qualitative variables, while independent sample *t*-tests were used to compare mean values between the two groups for normally distributed quantitative parameters. A *P*-value of < 0.05 was considered statistically significant. Pre- and post-operative pain levels were measured using the Visual Analog Scale (VAS).

4. Results

A total of 170 candidates for LC surgery were initially selected for inclusion in the study. Four patients were excluded as they did not wish to continue participating after surgery, leaving 166 patients who completed the study. The demographic and clinical characteristics of the patients (Table 1) revealed significant differences in several areas. A significant difference was observed in the sex distribution of participants between the groups ($P = 0.02$). Patients were monitored for conditions such as diabetes, hypertension, and renal, cardiovascular, and respiratory diseases. The prevalence of cardiovascular disease was significantly higher in the emergent group (10.9%) compared to the elective group (1%) ($P = 0.006$). In the emergent group, 57.8% of patients had an ASA score of 1, 28.1% had an ASA score of 2, and 14.1% had an ASA score of 3. Conversely, in the elective group, 55.9% of

Table 1. Patients' Demographics Data ^{a, b}

Variables	Groups		P-Value
	Emergent	Elective	
Age	52.75 ± 14.15	48.4 ± 13.99	0.054
Gender (M:F)	36 (56.2): 28 (43.8)	39 (38.2): 63 (61.8)	0.023
BMI	64 (27.73 ± 4.39)	1.2 (27.67 ± 4.56)	0.189
ASH			0.717
N/A	54 (84.4)	80 (78.4)	
UA	0	1 (1)	
LA	10 (15.6)	21 (20.6)	
Comorbidity			
Diabetes	10 (15)	13 (12)	0.601
Hypertension	9 (14.1)	25 (24)	0.105
Renal	0	1 (1)	0.999
Cardiovascular	7 (10.9)	1 (1)	0.006
Respiratory	1 (1.6)	0	0.386
ASA score			0.014
1	37 (57.8)	57 (55.9)	
2	18 (28.1)	42 (41.2)	
3	9 (14.1)	3 (2.9)	
Pre-surgery pain	4.53 ± 1.88	1.90 ± 1.89	< 0.001
Physical examination			< 0.001
Asymptomatic	0	60 (58.8)	
Tenderness	23 (35.9)	39 (38.2)	
Tenderness and rebound	0	1 (1.0)	
Tenderness and Murphy	41 (64.1)	2 (2.0)	
Sonography findings			< 0.001
GS	34 (53.1)	93 (91.2)	
Thickness increase & GS	28 (43.8)	2 (2)	
EBDD	0	1 (1)	
GB polyp	0	6 (5.9)	
increase GBWT g & EBDD	2 (3.1)	0	
Blood investigation			
WBC	11.49 ± 4.07	6.63 ± 2.11	< 0.001
BILLT	1.72 ± 1.31	1.08 ± 0.81	< 0.001
ALP	320.5 ± 209.68	215.04 ± 110.94	< 0.001
Endosonography findings			0.999
Have duct stone	8 (40)	3 (37.5)	
No duct stone	12 (60)	5 (62.5)	

Abbreviations: ASH, abdominal surgery history; UA, upper abdomen; LA, lower abdomen; EBDD, Extrahepatic bile duct dilation; GBWT, GB sac wall thickness; BILLT, total bilirubin; ALP, alkaline phosphatase.

^a Missing data was omitted from the dataset, and then the data analysis was processed.

^b Values are expressed as No. (%) or mean ± SD.

patients had an ASA score of 1, 41.2% had an ASA score of 2, and only 2.9% had an ASA score of 3. The differences between the groups were significant ($P = 0.01$). Patients in the emergent group reported significantly higher pain levels before surgery compared to the elective group ($P < 0.001$). All emergent group patients were symptomatic, while 58.8% of elective group participants

were asymptomatic, a significant difference ($P < 0.001$). Significant differences were noted in sonographic findings ($P < 0.001$). In the emergent group, 53.1% of patients had GS, 43.8% had GS with increased GB wall thickness, and 3.1% had extrahepatic bile duct dilation. In contrast, in the elective group, 91.2% of patients had GS, 2% had GS with increased GB wall thickness, and six

Table 2. Intraoperative Findings of the Patients ^{a,b}

Variables	Groups		P-Value
	Emergent	Elective	
First trocar entry method			0.43
Veress needle	3 (4.7)	10 (9.8)	
Open method	23 (35.9)	31 (30.4)	
Use of trocar	38 (59.4)	61 (59.8)	
Converting surgery from 3 to 4 trocar			0.007
3 trocar	37 (57.8)	79 (77.5)	
4 trocar	27 (42.2)	23 (22.5)	
Conversion to OC	1 (1.6)	0	0.386
Intraoperative Complication			0.145
N/A	43 (68.3)	80 (78.4)	
Duct injury	0	0	
Bleeding	0	0	
Other organ injury	0	0	
GB perforation	20 (31.7)	22 (21.6)	
Drain insertion			0.003
Inserted	35 (55.5)	25 (24.5)	
Not inserted	28 (44.5)	77 (75.5)	
Fascia repair of the UT entry site			0.570
Repaired	56 (87.5)	86 (84.3)	
Not repaired	8 (12.5)	16 (15.7)	
DOS, n (mean ± SD)	64 (70.39 ± 33.44)	102 (72.89 ± 35.91)	0.65

Abbreviations: UT, umbilical trocar; DOS, duration of surgery.

^a Missing data was omitted from the dataset, and then the data analysis was processed 2 minutes.

^b Values are expressed as No. (%) unless otherwise indicated.

patients (5.8%) had GB polyps. Significant differences were observed between the groups in the mean values of white blood cell (WBC) count ($P < 0.001$), total bilirubin ($P < 0.001$), and alkaline phosphatase (ALP) levels ($P < 0.001$).

According to the results, 57.8% of surgeries were completed using a three-trocar technique (77.5% of elective group patients), while 42.2% required the use of a fourth trocar (22.5% of elective group patients). This difference was statistically significant ($P = 0.007$) (Table 2). In one patient from the emergent group, the surgery was converted to OC due to unclear anatomy and suspicion of gallbladder cancer. However, the difference in conversion rates between the groups was not statistically significant ($P = 0.38$). Additionally, 35 patients (55.5%) in the emergent group required the insertion of a drain, compared to 25 patients (24.5%) in the elective group, a difference that was statistically significant ($P = 0.003$).

The post-surgical complications investigation (Table 3) showed no significant differences in local ($P = 0.77$) or systemic ($P = 0.37$) complications between the emergent

and elective groups. Regarding pain levels, one day after surgery, patients in the emergent group reported a mean pain score of 1.76 ± 1.03 , compared to 1.02 ± 0.80 in the elective group, a statistically significant difference ($P < 0.001$). Similarly, one-week post-surgery, the mean pain score was 0.49 ± 0.66 in the emergent group and 0.14 ± 0.43 in the elective group, again showing a significant difference ($P < 0.001$). By three months post-surgery, all patients reported no pain.

In analyzing the correlation between pain levels one day post-surgery and drain insertion, patients without a drain ($n = 104$, 1.14 ± 0.86) experienced significantly less pain ($P = 0.002$). A significant difference in post-operative hospital stay was also observed between the two groups ($P = 0.002$). The mean recovery time to resume routine activities was 4.82 ± 2.52 days in the emergent group, compared to 3.77 ± 1.21 days in the elective group, a statistically significant difference ($P = 0.003$). Additionally, patients without a drain had significantly shorter hospital stays than those with a drain ($P < 0.001$).

5. Discussion

Table 3. Post-operation Findings of the Patients ^{a, b}

Variables	Groups		P-Value
	Emergent	Elective	
Local complications post-surgery			0.773
N/A	60 (96.8)	99 (97.1)	
Bleeding	0	0	
Local infection	2 (3.2)	2 (2)	
Trocar site hernia	0	1 (1)	
Systemic complications post-surgery			0.378
N/A	61 (98.4)	102 (100)	
Pulmonary embolism	0	0	
Cardiovascular complications	0	0	
Respiratory complications	1 (1.6)	0	
Post-surgery pain			< 0.001
First-day post-surgery	1.76 ± 1.03	1.02 ± 0.80	
First-week post-surgery	0.49 ± 0.66	0.14 ± 0.43	
Post-surgery hospitalization	63 (1.55 ± 0.79)	102 (1.19 ± 0.50)	0.002
Duration of return to daily activity, n (mean ± SD) days	62 (4.82 ± 2.52)	101 (3.77 ± 1.21)	0.003

^a Missing data was omitted from the dataset, and then the data analysis was processed.

^b Values are expressed as No. (%) or mean ± SD unless otherwise indicated.

Gallstone disease is one of the most common gastrointestinal issues. While GS is typically asymptomatic, 2 - 4% of individuals with GS progress to painful cholecystitis, potentially leading to AC. Historically, OC was the standard treatment for cholecystectomy. However, LC has become the gold standard and the preferred surgical method for most surgeons. Despite its advantages, LC in emergent cases can present complications, such as acute inflammation, edema, and an increased risk of mortality. This study was designed to compare the risks and complications of emergent LC cases with elective cases.

The relationship between male gender and difficult cholecystectomy remains a topic of debate. Some studies suggest that male gender is a risk factor for difficult cholecystectomy, possibly because cholelithiasis is predominantly considered a female disease, leading to delayed diagnosis in males. This delay may result in significant adhesions from repeated inflammatory episodes before diagnosis. Nidoni et al. (as cited in Gupta et al.) (5) found that the conversion rate to OC was significantly higher in males compared to females ($P = 0.03$, 95% confidence interval).

The demographic data from this study indicated that cardiovascular disease was the only comorbidity significantly different between the emergent and elective groups ($P = 0.006$). This finding aligns with Abe et al. (18), who reported a significant prevalence of cardiovascular disease in patients with concomitant

biliary infections. However, Abe et al. did not find a meaningful difference between urgent and semi-urgent groups ($P = 0.04$) (18). Additionally, a significant number of patients in both groups presented with abnormal alkaline phosphatase (ALP) levels ($P = 0.01$), an important indicator of bile duct issues. Singal et al. (19) evaluated liver function tests (LFTs) in patients undergoing LC or OC and found elevated serum levels in both groups. However, they did not observe any significant difference in ALP or total bilirubin levels between the groups prior to surgery ($P = 0.09$).

Our results revealed that patients in emergency situations experienced more pain before surgery than those in the elective group, which can be attributed to the acute condition of cholecystitis or the need for drain insertion. One-day post-operation assessments showed that although both groups reported significantly reduced pain compared to pre-surgery levels, there was still a significant difference between the groups. Furthermore, a week after surgery, monitoring again demonstrated lower pain levels in both groups; however, the emergent group continued to experience significantly higher pain compared to the elective group ($P < 0.001$). The monitoring extended to three months post-surgery, at which point none of the patients reported any pain. A plausible explanation for the observed pain could be the higher rate of drain insertion in the emergent group compared to the elective group, as indicated in Table 2. This aligns with

the finding that patients in the emergent group reported more post-operative pain.

Hospital stays were also longer, and the time to return to routine life was extended in the emergent group. This could be due to the patients' acute conditions or higher disease severity, requiring additional recovery time post-surgery (20). Nonetheless, prior studies have consistently reported shorter hospital stays for LC patients compared to OC patients (21). Ferrarese et al. concluded that drain insertion might contribute to extended post-operative hospital stays in the emergent group (22). Other studies have corroborated our findings, showing that patients in emergency conditions tend to have longer hospital stays following LC surgery (10, 13, 23). In a cohort study (12) comparing emergency and elective LC, more drain insertions were reported in emergency LC patients than in elective LC patients ($P < 0.001$).

Historically, it has been presumed that cholecystectomy patients in emergency conditions would face more complications both during and after laparoscopic surgery, leading many surgeons to prefer OC over LC. However, our findings indicate no significant differences in most variables, suggesting that emergent LC can be as safe as elective LC in terms of both local and systemic complications. Various studies have similarly reported no major complications or significant differences between emergent and elective LCs (22, 24). One common complication during LC is the need to convert to OC. In our study, only one patient from the emergent group required conversion to OC ($P = 0.38$).

In a cohort study, Saeb-Parsy et al. compared the efficacy of elective and emergency LC in 423 patients and reported one conversion from LC to OC, which occurred in the elective group ($P = 0.64$) (24). Ferrarese et al. (22) investigated outcomes between elective and emergency LC in elderly patients and found a higher conversion rate to OC in the elective group, although the difference was not statistically significant ($P = 0.38$). Additionally, the duration of surgery was comparable between the emergent and elective groups, a finding consistent with Grass et al., who reported similar results in a related study (25).

5.1. Conclusions

Comprehensively, based on previous assumptions and reports, LC—particularly emergency LC—was perceived as less effective and associated with higher intra- and post-operative complications, leading to a lower preference among surgeons. However, the findings of this study contradict that initial hypothesis,

demonstrating that emergency LC is both acceptable and justifiable in terms of its success rate and safety profile. It did not result in significant complications for patients. Therefore, LC can be considered a gold standard and low-risk option for patients in emergency situations.

Acknowledgements

The authors would like to thank for the guidance and advice from the Clinical Research Development Unit of Baqiyatallah Hospital, Tehran, Iran.

Footnotes

Authors' Contribution: M. A.: Data gathering and writing the draft; H. Gh., S. H., J. A., M. M., M. R., and H. M.: Patient follow-up, editing and preparing the draft, intellectual contribution in study design; M. D.: Project supervision, conceptualizing the project, editing the draft; the final version of this paper was revised and approved by all the authors.

Conflict of Interests Statement: The authors declared no conflict of interests.

Data Availability: The dataset presented in the study is available on request from the corresponding author during submission or after publication. The data are not publicly available due to privacy and ethics restrictions.

Ethical Approval: This study was approved by the research ethics committee of Baqiyatallah University of Medical Sciences with the ethical code of [IR.BMSU.REC.1397.034](https://doi.org/10.1016/j.amsu.2021.103060), and it was conducted in accordance with principles embodied in the Declaration of Helsinki 1975.

Funding/Support: This study was not funded by any organization.

Informed Consent: All participants filled out written informed consent to participate in the study.

References

1. Bhandari TR, Khan SA, Jha JL. Prediction of difficult laparoscopic cholecystectomy: An observational study. *Ann Med Surg (Lond)*. 2021;72:103060. [PubMed ID: 34815866]. [PubMed Central ID: PMC8591467]. <https://doi.org/10.1016/j.amsu.2021.103060>.
2. Mou D, Tesfasilassie T, Hirji S, Ashley SW. Advances in the management of acute cholecystitis. *Ann Gastroenterol Surg*. 2019;3(3):247-53. [PubMed ID: 31131353]. [PubMed Central ID: PMC6524093]. <https://doi.org/10.1002/ags3.12240>.

3. Warchalowski L, Luszczki E, Bartosiewicz A, Deren K, Warchalowska M, Oleksy L, et al. The Analysis of Risk Factors in the Conversion from Laparoscopic to Open Cholecystectomy. *Int J Environ Res Public Health*. 2020;**17**(20). [PubMed ID: 33080991]. [PubMed Central ID: PMC7588875]. <https://doi.org/10.3390/ijerph17207571>.
4. Yun SP, Seo HI. Clinical aspects of bile culture in patients undergoing laparoscopic cholecystectomy. *Medicine (Baltimore)*. 2018;**97**(26). e11234. [PubMed ID: 29952986]. [PubMed Central ID: PMC6039604]. <https://doi.org/10.1097/MD.00000000000011234>.
5. Gupta AK, Shiwach N, Gupta S, Gupta S, Goel A, Bhagat TS. Predicting difficult laparoscopic cholecystectomy. *International Surgery Journal*. 2018;**5**(3). <https://doi.org/10.18203/2349-2902.isj20180837>.
6. Majumder A, Altieri MS, Brunt L. How do I do it: laparoscopic cholecystectomy. *Annals of Laparoscopic and Endoscopic Surgery*. 2020;**5**:15. <https://doi.org/10.21037/ales.2020.02.06>.
7. Alexander HC, Bartlett AS, Wells CI, Hannam JA, Moore MR, Poole GH, et al. Reporting of complications after laparoscopic cholecystectomy: a systematic review. *HPB (Oxford)*. 2018;**20**(9):786-94. [PubMed ID: 29650299]. <https://doi.org/10.1016/j.hpb.2018.03.004>.
8. Barazanchi AWH, MacFater WS, Rahiri JL, Tutone S, Hill AG, Joshi GP, et al. Evidence-based management of pain after laparoscopic cholecystectomy: a PROSPECT review update. *Br J Anaesth*. 2018;**121**(4):787-803. [PubMed ID: 30236241]. <https://doi.org/10.1016/j.bja.2018.06.023>.
9. Cao AM, Eslick GD, Cox MR. Early Cholecystectomy Is Superior to Delayed Cholecystectomy for Acute Cholecystitis: a Meta-analysis. *J Gastrointest Surg*. 2015;**19**(5):848-57. [PubMed ID: 25749854]. <https://doi.org/10.1007/s11605-015-2747-x>.
10. Vidyadharan K, KembaiShanmugam R, Ayyasamy G, Thandayuthapani S. Early versus delayed laparoscopic cholecystectomy in uncomplicated biliary colic: An observational study. *Laparoscopic, Endoscopic and Robotic Surgery*. 2023;**6**(2):69-72. <https://doi.org/10.1016/j.lers.2023.05.005>.
11. Acar T, Kamer E, Acar N, Atahan K, Bag H, Hacıyanlı M, et al. Laparoscopic cholecystectomy in the treatment of acute cholecystitis: comparison of results between early and late cholecystectomy. *Pan Afr Med J*. 2017;**26**:49. [PubMed ID: 28451027]. [PubMed Central ID: PMC5398876]. <https://doi.org/10.11604/pamj.2017.26.49.8359>.
12. Lucocq J, Radhakrishnan G, Scollay J, Patil P. Morbidity following emergency and elective cholecystectomy: a retrospective comparative cohort study. *Surg Endosc*. 2022;**36**(11):8451-7. [PubMed ID: 35201423]. [PubMed Central ID: PMC9613569]. <https://doi.org/10.1007/s00464-022-09103-2>.
13. To KB, Cherry-Bukowicz JR, Englesbe MJ, Terjimanian MN, Shijie C, Campbell DJ, et al. Emergent versus elective cholecystectomy: conversion rates and outcomes. *Surg Infect (Larchmt)*. 2013;**14**(6):512-9. [PubMed ID: 24274058]. <https://doi.org/10.1089/sur.2012.160>.
14. Moore AN, Carmichael H, Steward L, Velopulos CG. Cholecystectomy: Exploring the Interplay Between Access to Care and Emergent Presentation. *J Surg Res*. 2019;**244**:352-7. [PubMed ID: 31323390]. <https://doi.org/10.1016/j.jss.2019.06.070>.
15. Ohya H, Maeda A, Takayama Y, Takahashi T, Seita K, Kaneoka Y. Preoperative risk factors for technical difficulty in emergent laparoscopic cholecystectomy for acute cholecystitis. *Asian J Endosc Surg*. 2022;**15**(1):82-9. [PubMed ID: 34291878]. <https://doi.org/10.1111/ases.12969>.
16. Altieri MS, Brunt LM, Yang J, Zhu C, Talamini MA, Pryor AD. Early cholecystectomy (< 72 h) is associated with lower rate of complications and bile duct injury: a study of 109,862 cholecystectomies in the state of New York. *Surg Endosc*. 2020;**34**(7):3051-6. [PubMed ID: 31376010]. <https://doi.org/10.1007/s00464-019-07049-6>.
17. Marwah S, Shivran KD, Jangra MS, Singh J, Gupta SP, Singh VP. Comparative evaluation of early laparoscopic cholecystectomy for mild gallstone pancreatitis vs acute cholecystitis. *Hellenic Journal of Surgery*. 2017;**89**(1):4-12. <https://doi.org/10.1007/s13126-017-0370-x>.
18. Abe T, Amano H, Hanada K, Bekki T, Minami T, Yonehara S, et al. Efficacy and safety of early cholecystectomy for comorbid acute cholecystitis and acute cholangitis: Retrospective cohort study. *Ann Med Surg (Lond)*. 2019;**38**:8-12. [PubMed ID: 30581570]. [PubMed Central ID: PMC6302235]. <https://doi.org/10.1016/j.amsu.2018.10.031>.
19. Singal R, Singal RP, Sandhu K, Singh B, Bhatia G, Khatri A, et al. Evaluation and comparison of postoperative levels of serum bilirubin, serum transaminases and alkaline phosphatase in laparoscopic cholecystectomy versus open cholecystectomy. *Journal of Gastrointestinal Oncology*. 2015;**6**(5):479.
20. Mishima K, Fujiyama Y, Wakabayashi T, Igarashi K, Ozaki T, Honda M, et al. Early laparoscopic cholecystectomy for acute cholecystitis following the Tokyo Guidelines 2018: a prospective single-center study of 201 consecutive cases. *Surg Endosc*. 2023;**37**(8):6051-61. [PubMed ID: 37118031]. <https://doi.org/10.1007/s00464-023-10094-x>.
21. Lombardo S, Rosenberg JS, Kim J, Erdene S, Sergelen O, Nellerme J, et al. Cost and outcomes of open versus laparoscopic cholecystectomy in Mongolia. *J Surg Res*. 2018;**229**:186-91. [PubMed ID: 29936988]. <https://doi.org/10.1016/j.jss.2018.03.036>.
22. Ferrarese AG, Solej M, Enrico S, Falcone A, Catalano S, Pozzi G, et al. Elective and emergency laparoscopic cholecystectomy in the elderly: our experience. *BMC Surg*. 2013;**13** Suppl 2(Suppl 2). S21. [PubMed ID: 24268106]. [PubMed Central ID: PMC3851193]. <https://doi.org/10.1186/1471-2482-13-S2-S21>.
23. Kanakala V, Borowski DW, Pellen MG, Dronamraju SS, Woodcock SA, Seymour K, et al. Risk factors in laparoscopic cholecystectomy: a multivariate analysis. *Int J Surg*. 2011;**9**(4):318-23. [PubMed ID: 21333763]. <https://doi.org/10.1016/j.ijsu.2011.02.003>.
24. Saeb-Parsy K, Mills A, Rang C, Reed JB, Harris AM. Emergency laparoscopic cholecystectomy in an unselected cohort: a safe and viable option in a specialist centre. *Int J Surg*. 2010;**8**(6):489-93. [PubMed ID: 20633707]. <https://doi.org/10.1016/j.ijsu.2010.06.015>.
25. Grass F, Cachemaille M, Blanc C, Fournier N, Halkic N, Demartines N, et al. Is standardized care feasible in the emergency setting? A case matched analysis of patients undergoing laparoscopic cholecystectomy. *BMC Surg*. 2016;**16**(1):78. [PubMed ID: 27905910]. [PubMed Central ID: PMC5131530]. <https://doi.org/10.1186/s12893-016-0194-6>.