

Shiraz E-Medical Journal
Vol. 12, No. 1, January 2011

<http://semj.sums.ac.ir/vol12/jan2011/89029.htm>

Epidemiology and Evaluation of 1073 Burn Patients in the Southeast of Iran.

Shirkhoda M*, Kaviani Far K**, Narouie B***, Shikhzadeh A***, Ghasemi Rad M ±, Hanfi Bojd H ∞.

*Surgeon, **Assistant Professor of Surgery, ***Researcher of Clinical Research Development Center, Ali-ebne-Abitaleb hospital, Zahedan University of Medical Sciences, Zahedan, Iran, ± Medical Student, Urmia University of Medical Sciences. Student Research Committee, Urmia, Iran, ∞Medical Student, Zahedan University of Medical Sciences, Clinical Research Development Center, Zahedan, Iran.

Correspondence: Dr. B. Narouie, Clinical Research Development Center, Ali-ebne-Abitaleb Hospital, Zahedan, Iran, Telephone: +98(541) 341-4103, Fax: +98(541)341-4103, E-mail: b_narouie@yahoo.com

Received for Publication: August 2, 2010, Accepted for Publication: December 4, 2010.

Abstract:

Background: Burn injury is one of the major causes of morbidity and mortality worldwide. In addition to the financial burden it inflicts on the health care system, it can lead to psychological, social and physical distress both to the patient and family members. The aim of this study was to determine the epidemiological parameters in hospitalized burn patient.

Methods: In a cross-sectional study we evaluated data of 1073 hospitalized burn patients in Zahedan Khatam Al Anbia hospital from 2005-2008. All data were extracted from medical record and analyze by SPSS software. A P-value less than 0.05 was considered significant.

Results: Among 1073 patient with acute burn, 55.4% were male and 44.6% were female. The mean age was 20.94 year old and mean size of burn area was 49.11±29.65 total body surface area (TBSA). Total body surface area burned was significantly higher in those with self-burn (p=0.0001). In children younger than 15 year-old scalds (hot liquid) was the most frequent cause of burn (44%). There was a higher incidence of self-burning in women (p=0.0001) and the mean length of hospital stay was 6.32±5.27 days. The mortality rate in self-burn patient was 87.34 % and in total cases was 41.47 %.

Conclusion: In the present study, most of the burns were noted in 16 to 20 years age group, and more in male than in female subjects. Burn was more frequent in children and adulthood and they are considered high-risk groups with higher mortality and morbidity. This shows a need for comprehensive burn prevention programs.

Keywords: burn, epidemiology, hospitalized patients, trauma, surgery

Background:

Burn injury is one of the major causes of morbidity and mortality worldwide. In addition to the financial burden it inflicts on the health care system, it can cause psychological, social and physical problems for patient and family. Although morbidity and mortality rate due to burns has decreased in developed countries, it is still a major health challenge in developing countries.⁽¹⁾ Annually, 1.25 million Americans are burned and around 50 thousands are hospitalized, which costs 7 billion dollars for the health system. Burn is the 5th cause of death due to accident in United States.^(1, 2) Epidemiological reports in some counties show that thermal injury is more common in young age group and it's mostly due to fire, with a morbidity of 33.5%-54.8%.^(3, 4, 5, and 6) There is no available data in Iran, but some studies have demonstrated its high incidence in younger's, female, and illiterate people. The mortality is 27.9%-34.4%.^(7, 8) It is one of the major challenges of health system in Iran. We need more data from different parts of country to give us an estimate of epidemiology of thermal injury in country. Unfortunately, there are very few studies on prevalence and outcome of burn in Iran ^(2, 7), with no data on epidemiology of burn in the Sistan va Baluchistan province. There are few causes that can lead to increase thermal injuries in our province. No accessible gas pipelines with subsequent need to use stove and other fuel consuming heaters, Neighbor with counties such as Afghanistan and Pakistan and fuel trafficking to these countries.

This study was designed to analyze the demographics and etiology of thermal injury and evaluate the outcome of different surgical interventions. We also aim to compare our results with other published data to apply in our burn preventing education programs and to increase the quality of health provided to these patients.

Material and Methods:

Sistan and Baluchistan province is located in the southeast of Iran and has a population of 2.2 million according to a 2006 statistics. It's the largest province with 11.4% (187.5 Km²) of country land. The Burn unit of Khatam Al Anbia hospital is the major regional referral center for all burns injuries in this province. All patients in need of in-hospital treatment are admitted here. The distance between this center and other closest burn unit, which is located in Kerman city, is 550 kilometers. This is why an epidemiologic study of this center will give valuable and clear prospective in further programming for prevention and treatment of burn patients.

In this retrospective cross-sectional study we reviewed medical records of 1073 burn patients admitted to Burn unit of Khatam Al Anbia hospital between March 2005 and November 2008. The data were collected through special forms contained age, sex, percentage of burn, etiology of burn (suicidal or accidental), duration of hospitalization, season, type of surgical intervention (non-surgical, debridement, debridement and primary graft), outcome (death or discharge).

Statistical analysis was done using SPSS ver.15. We used Chi Square and T test and single tail variance to analyze demographic data. The risk associated with mortality was analyzed using logistic regression. A P-value of <0.05 was considered significant.

Results:

Age and sex

594 patients (55.4%) were male, and 479 patients (44.6%) were female (Table 1). 33.3% of patients (375 patients) were below 15 years old. Burn was more common in male. The most frequent age group was 16-20; with total of 229 burn patients.

Burn extent

The TBSA (total body surface area) burned range was from 3 to 100% with the mean of 49.11 ± 29.65 . In patients with self-induced burn; the percentage was 10%-100%, with a mean of 86.5 ± 24.96 . The other patients had a mean of $41.06\% \pm 24.96\%$. The percentage of burn was significantly higher in those with self-burn (attempted suicide) ($P=0.0001$). 18.9% of the patients had above 90% of their body surface burned (table 2)

Burn etiology

75.8% of patients were burned due to flame, with oil being the most common cause (41.5%). In those under 15 years, the most common cause of burn was scalds (44%). 4 cases of 46 patients with electrical burning were due to lightning. Winter was the most common season with oil burn (Table 3).

Deliberate self burning (DSB)

Suicide is an act through which a person attempts to harm him/her self. Among

190 patients with self-burning, 51 (26.8%) were male and 139 (73.2%) were female, self-burning was statistically significant in female ($p=0.0001$). The most common etiology of burning was oil ($p=0.0001$) and the predominant age group in both sexes was 16-20 years. 62.56% of patient had above 90% burn. The mortality in those with self-burning was 87.34% (Table 4,5).

Inhalation injury

109 patients had burns due to inhalation. The diagnostic criteria for inhalation burn were;

occurrence in closed environments, hoarseness and expiratory wheeze, Soot in the saliva and excessive salivation and facial burn and signed nasal vibrissae.

The most frequent causes of inhalation injury in descending order were oil (53.2%), gas (25.7%), gasoline (20.2%) and scalds (0.9%). Out of 109 patients with inhalational injury, most were 16-20 years old (35 patient). 65 patients were female, 72.61% of patient with inhalational injury died. Self-burning caused increased inhalational injury ($p=0.0001$)

Type of surgical intervention

935 patients (87.13%) underwent conservative treatment, 30 patients (2.7%) surgical excision and were referred for delayed skin grafting to hospitals with plastic surgery centers. 108 patients (10%) underwent surgical excision and primary grafting (Table 6).

Duration of hospitalization

The duration of hospitalization was 1 to 41 days, with a mean of 6.32 ± 5.27 and a median of 5 days. The mean hospitalization time in the patients who died was 5.46 ± 4.8 days. In others, the mean was 8.44 ± 5.79 . The mean time of hospital

admission in those with excision and grafting was 6.06 ± 2.85 days. After the excision alone, it was 5.16 ± 3.32 days. There was no significant difference between excision alone and combined excision with primary grafting ($p > 0.05$) (Table 7).

Mortality

41.47% of the patients died, with 28.89% mortality in the accidental group (Table 8). Mean age and percentage of burn in those who died was 23.37 ± 13.10 and 80.09 ± 23.58 , respectively (Table 9). The percentage of burn causing 50% of death (LD50) was 65%.

Seasonal variation

Most hospital admissions were in winter (29%) follow by spring (27.1%), summer (25.6%) and fall (18.3%). There was no statistically significant difference between seasons. ($P > 0.05$)

Discussion:

Due to our common border with Afghanistan and Pakistan and significant fuel price difference., there is illegal fuel trafficking with unsafe methods.

In the present study, most of the burns were registered in 16 to 20 years age group, and more in male subjects. The degree of burn was more in those with self-burning. Oil was the most frequent cause in adult, and scalds (boiling water) were the most frequent cause in pediatric age group. Self-burning was significantly higher in females. Surgical treatment by excision and primary skin graft reduced the time and cost of hospitalization compared to delayed type.

Mortality was 28.89% in accidental burn, 87.34% in self-burning and 72.61% in inhalation burn patients. LD50 was 65%.

Age and sex

Burns were more common in male than in female subjects, as in reports from Catalonia, Dublin and other regions.^(8, 9, 10, 11, 12) Other studies reported higher number of burns in females.^(4, 11, 13-15) It seems that the high number of burns in males in our study is related to the transportation and trafficking of fuel to neighboring countries. Similar to other reports ^(10, 13, 16-38), this study demonstrated high frequency of burn in pediatric age group. This mandates an educational program regarding burn which should start at school levels, sustaining the conclusions of Keswani's study which showed that educating children is effective in the prevention of burns.⁽³⁹⁾

Extent of burn

There was a paradoxical relationship between the extent of burn and survival.^(34, 40, and 41) The cut off point for the percentage of burn that lead to death was between 20 to 50% in different studies.^(42, 43, 44) All quoted studies had the same opinion: a burn on more than 70% of the body surface has a high rate of mortality.^(42, 43) In the present study, there was a significant relationship between the extent of burn and mortality ($P = 0.0001$). The extent of burn was higher in females than in males, a result that has previously been obtained by other studies.^(3, 45, 46) In our study, mortality was mostly due to self-burning.

Etiology of burn

The most common cause of burn was flame followed by scalds (mostly boiling water), concurring to some previous reports. (3, 8, 47-50) Oil was the most common cause of burn. This might be due to the consumption of oil as a common fuel in our houses. In a few other

reports, scalds followed by flame were common cause of burn. ^(6, 8, 51) Burning with scalds was the most common etiology in pediatrics age group, as previous studies had already mentioned. ^(16, 18, 36, 37, 47, 49) In most of the villages in our provinces there are no physical places that can be called kitchens and the cooking materials are kept on the ground. This might be the cause for the high number of scald burns in the pediatric age group. **Self-burn**

There is an increase in the number of self-burns in both males and females.⁽⁵²⁾ In the present study, like in other studies ^(7, 53), self-burn was more common in females than in males ($P=0.0001$). The most common age was 16 to 20 years, which is less than European and American countries.⁽⁵³⁾ The high incidence of self-burn seems to be due to forced marriage, age differences between husband and wife, family problems, the husband's addictions and the high numbers of wives. The most common instrument of self-burn, like other countries ^(7, 53), was oil which is easily available in this region. In different part of world, 18 to 84% mortality due to self-burn was reported ^(3, 28, 52, 54-64), whereas, in our study, it was 87.34.

Inhalation injury

The mechanisms by which inhalation can cause injury are carbon monoxide poisoning, airways injury and burning and inhalation of soot particles. In this study, 10.2% of the patients developed inhalation injuries. In two previous studies, the incidences of inhalation injuries were 5.1% and 19.6% respectively. ^(65, 66) In our study, the mortality rate due to inhalation injury was 72.61%, while the mor-

tality rate in North Carolina was only 31%. ⁽⁶⁶⁾ Self-burn was one of the major causes of inhalation injury; this shows that those with self-burn should be investigated for inhalation burn.

Type of treatment

In our center, the studied patients who needed debridement were either only excised, or excised and then got either a delayed or a primary graft. For a delayed graft, the patients were transferred to better-equipped facilities. Delayed graft was performed in first 20 month of study.

Duration of hospitalization

The duration of hospital admission in our study was 5 days that is less than other centers in Iran ^(4, 51) and also other countries. ^(9, 10, 65, 67) There was no significant difference between excision alone and excision and primary graft in relation to the duration hospitalization.

In a study conducted in Harare there was a reduction in the duration of hospital admission from 42 to 17 days in those patients who underwent primary grafting. ⁽¹⁴⁾ This shows that primary closing of the wound will lead to a reduction in the time and costs of hospitalization. ⁽⁶⁸⁾

We referred our patients to other centers to undergo delayed grafting, which would naturally increase the time and cost of hospitalization.

Morbidity and mortality

The overall mortality and morbidity in present study was 41.47% that is more than previous reports ^(8, 13, 14, 31, 38, 47-49, 69, 70), which might be due to the high number self-burns, hospital admissions and high percentage of burned body surface area.

Seasonal changes

Winter was the most common season with burn occurrence. Like previous studies^(31, 71-73), there was no significant relationship between the season and the burn occurrence.

In our province, oil is a common source of smuggling. Educating people regarding how to protect themselves from the causes and outcomes of burns can reduce the number of burns in our society.

Reducing the smuggling may also reduce the number of burns occurring in car accidents.

Since burning causes key accidental injuries and it is a major cause of pain, psychosomatic problems, disability and death, there must be educational programs in our society (schools, family and work place) to prevent the acute injury and increase the quality of patient care, by equipping the emergency departments and educating the doctors and nurses.

Table 1: age, sex and extent of burn

Age Groups	Percentage Of burn Mean -SD	Sex				Total	
		Female		Male		%	Number
		%	Number	%	Number		
0-5	29.82 -17.40	33.3	60	66.7	120	16.8	180
6-10	36.96 - 23.97	46.4	32	53.6	37	6.4	69
11-15	52.59 - 29.85	53.7	58	46.3	50	10.1	108
16-20	58.19- 30.94	50.7	116	49.3	113	21.3	229
21-25	51.80- 30.06	42.3	85	57.7	112	18.1	194
26-30	57.20 - 29.39	42.3	44	57.7	60	9.7	104
31-35	49.44 - 31.53	50.0	27	50.0	27	5.0	54
36-40	56.97 - 30.24	37.8	114	62.2	23	3.4	37
41-45	53.21 - 31.06	42.9	12	57.1	16	2.6	28
46-50	50.62 - 29.78	53.1	17	46.9	15	3.0	32
51-55	47.92 - 28.08	41.7	5	58.3	7	1.1	12
56-60	47.00 - 33.09	40.0	4	60.0	6	0.9	10
61-65	45.83 - 31.84	16.7	1	83.3	5	0.6	6
66-70	33.00 - 32.91	66.7	4	33.3	2	0.6	6
>70	32.75 - 22.80	75.0	3	25.0	1	0.4	4

Table 2: Extent of burn and sex

Sex Percentage Of burn	Female		Male		Total	
	%	Number	%	Number	%	Number
< 9	18.2	6	81.8	27	3.1	33
10-19	35.2	38	64.8	70	10.1	108
20-29	38.2	65	61.8	105	15.8	170
30-39	37.3	59	62.7	99	14.7	158
40-49	48.8	60	51.2	63	11.5	123
50-59	43.6	48	56.4	62	10.3	110
60-69	48.3	29	51.7	31	5.6	60
70-79	51.8	29	48.2	27	5.2	56
80-89	50.0	26	50.0	26	4.8	52
>90	58.6	119	41.4	84	18.9	203

Table 3: self burning

Self burning Age group	No				Yes			
	Female		Male		Female		Male	
	%	N	%	N	%	N	%	N
< 10	98.45	91	100	157	1.55	1	0	0
11-15	60.3	35	98.0	49	39.7	23	2.0	1
16-20	52.6	61	82.3	93	47.4	55	17.7	20
21-25	68.3	56	89.3	100	31.7	26	10.7	12
26-30	61.4	27	87.3	47	38.6	17	21.7	13
31-35	70.4	19	96.3	26	29.6	8	3.7	1
36-40	78.6	11	87.0	20	21.4	3	13.0	2
41-45	66.7	8	93.8	15	33.3	4	6.3	1
46-50	94.1	16	100	15	5.9	1	0	0
51-55	100	5	100	7	0	0	0	0
56-60	100	4	100	6	0	0	0	0
61-65	0	0	100	5	100	1	0	0
66-70	100	4	100	2	0	0	0	0
>70	100	3	100	1	0	0	0	0
Total	71.0	340	91.4	543	29.0	139	8.6	51

Table 4: cause of burn

Age Cause of Burn	>15 years		<15 years		Mean age N	Total		
		%	N	%		%	N	
Flame	Oil	49.7	356	24.9	89	23.45	41.5	445
	Gas	22.2	159	15.7	56	25.12	20.0	215
	Petrol	17.5	125	6.7	24	25.71	13.9	149
	Gasoline	0.3	2	0.6	2	20.25	0.4	4
Scalds	Boiling water	3.4	24	44.0	157	7.06	16.9	181
	Others	1	7	4.5	16	11.09	2.2	23
Electricity		5.2	37	2.5	9	21.80	4.3	46
Chemical		0.4	3	0.3	1	20.00	0.4	4
Others		0.4	3	0.9	3	13.61	0.6	6
Total		100	716	100	357		100	1073

Table 5: extent of burn and sex

Self burning Percentage of burn	Sex							
	Female		Male		Female		Male	
	%	N	%	N	%	N	%	N
<50	97.02	219	99.2	361	2.98	9	0.8	3
50-59	81.3	39	98.4	61	18.8	9	1.6	1
60-69	79.3	23	96.8	30	20.7	6	3.2	1
70-79	75.9	22	92.6	25	24.1	7	7.4	2
80-89	34.6	9	69.2	18	65.4	17	30.8	8
>90	23.5	28	57.1	48	76.5	91	42.9	36
Total	71.0	340	91.4	543	29.0	139	8.6	51

Table 6: Type of treatment and extent of burn

Type of Treatment Percentage Of burn	Excision and grafting		Excision		Conservative	
	%	N	%	N	%	N
<9	3.5	1	10.7	3	85.8	24
10-19	14.6	17	0.8	1	84.6	98
20-29	14.6	24	3.6	6	81.8	124
30-39	15.9	24	2	3	82.1	124
40-49	11.2	12	5.6	6	83.2	89
50-59	24.75	25	3	3	72.25	73
60-69	2	1	2	1	96	48
70-79	6	3	6	3	88	44
80-89	2	1	3.9	2	94.1	48
>90	0	0	0.8	2	99.2	253
Total	10.06	108	2.8	30	87.14	935

Table 7: type of treatment and duration of hospital admission

Type of treatment Duration of admission	Conservative		Excision and grafting			Excision	Total	
	%	N	%	N	%	N	%	N
>5	98.5	485	0.3	1	1.2	6	100	492
5-10	86.3	305	11.3	40	2.5	9	100	354
10-15	65.7	95	28.3	41	6.1	9	100	145
15-20	61.1	30	36.1	19	2.8	4	100	53
>20	70.0	20	25.0	7	5.0	2	100	29

Table 8: mortality and outcome according to sex

Sex outcome	Female		Male		Total	
	%	N	%	N	%	N
Discharged	37.7	237	62.30	391	100	628
Died	54.4	242	45.6	203	100	445

Table 9: frequency of causes of mortality

	Variable	Number of death	Number of Patients	P value
Percentage of burn	≤ 40	34	505	0.0001
	>40	262	568	
Cause of burn	Non-flame	17	257	0.0001
	Flame	286	816	
Type of Burn	Accidental	165	571	0.0001
	Self burning	138	158	
Sex	Male	140	594	0.0001
	Female	163	479	
Age	≤ 15	66	357	0.0001
	>15	237	716	
Inhalational injury	Yes	61	109	0.0001
	No	242	964	

Acknowledgements:

We would like to acknowledge from Clinical Research Development Center of Aliebne-Abitaleb hospital for its help in preparing this manuscript.

References:

1. Lawrence JC. Burns and scalds: aetiology and prevention. In: Principles and practice of burns management. London: Churchill Livingstone, 1996.
2. Solatni K, Zand R, Mirghasemi A. Epidemiology and mortality of Burns in Tehran Iran. *Burns* 1998; 24: 325-328.
3. Panjeshahin MR, Rastegar-Lari A, Talei AR, Shamsnia J, Alaghebandan R. Epidemiology and mortality of burns in the south west of Iran. *Burns* 2001; 27: 219-26.
4. Groohi B, Alaghebandan R, Rastegar-Lari A. Analysis of 1089 burn patients in province of Kurdistan Iran. *Burns* 2002; 28: 569-74.
5. Zarghami M, Khalilian A. Deliberate self-burning in Mazandaran Iran. *Burns* 2002; 28: 115-9.
6. Ansari-Lari M, Askarian M. Epidemiology of burns presenting to an emergency department in Shiraz, south Iran. *Burns* 2003; 29: 579-81.
7. Maghsoudi H, Garadagi A, Jafary GA, Azarmir G, Aali N, Karimian B, et al. Women victims of self-inflicted burns in Tabriz, Iran. *Burns*, 2004; 30: 217-20.
8. Barret JP, Gomes P, Solano I, Gonzales-Dorrego M, Crisol FJ. Epidemiology and mortality of adult burns in Catalonia. *Burns*, 1999; 25: 325-9.
9. Carrol SM, Giugh M, Eadie PA, McHugh M, Edwards G, Lawlor D. A. 3-year epidemiological review of burn unit admissions in Dublin, Ireland: 1988-1991. *Burns* 1995; 21: 379-82.
10. Bang RL, Ghoneim IE. Epidemiology and mortality of 162 major burns in Kuwait. *Burns* 1996; 22: 433-8.
11. Darko DF, Wachtel TL, Eard HW, Frank HA. Analysis of 585 burn patients hospitalized over a 6-year period. Part II. Aetiological data. *Burns Incl Therm Inj* 1986; 12: 391-4.
12. Rossi LA, Braga ECF, Barruffini RCP, Carvalho EC. Childhood burn injuries: circumstances of occurrences and their prevention in Ribeirao Preto, Brazil. *Burns* 1998; 24: 416-9.
13. Liu EH, Khatri B, Shakya YM, Richard BM. A 3 year prospective audit of burns patients treated at the Western Regional Hospital of Nepal. *Burns* 1998; 24: 129-33.
14. Mzezewa S, Jonsson K, Aberg M, Salemark L. A prospective study on the epidemiology of burns in patients admitted to the Harare burn units. *Burns* 1999; 25: 499-504.
15. Cutillas M, Sesay M, Perro G, Bourdarias B, Castede JC, Sanchez R. Epidemiology of elderly patients burns in the southwest of France. *Burns* 1998; 24: 134-8.
16. Jamal YS, Ardawi MSM, Ashy ARA, et al. Paediatric burn injuries in the Jeddah Area of Saudi Arabia: a study of 187 patients. *Burns* 1990; 16: 36-40.
17. Haberal M, Oner Z, Bayrakdar U. Epidemiology of adults and children burns in a Turkish burn center. *Burns* 1987; 13: 136-40.
18. Larsen J, Gurumurthy, Ghali MT, Hall KV. Admission to a new burns unit in Gizan, Saudi Arabia. *Burns* 1987; 13: 286-9.
19. Elberg J, Schroder H, Glent-Masden L, et al. Burns: epidemiology and the effect of a prevention programme. *Burns* 1987; 13: 391.
20. Silverstein P, Lack BO. Epidemiology and prevention. In: Boswick JA, editor. The art and science of burn care. New York: Aspen, 1987. p. 11-9.
21. Clark WR, Fromm B. Burn mortality: experience at a regional burn unit and literature review. *Acta Chir Scand Suppl* 1987; 537: 1.
22. Bang RL, Mosbah KM. Epidemiology of burns in Kuwait. *Burns* 1988; 14: 194-200.
23. Onuba O, Udodiok E. The problems and prevention of burns in developing countries. *Burns Incl Therm Inj* 1987; 13: 382-5.
24. Jamal YS, Ardawi MSM, Ashy ARA, Mardad H, Shaik SA. Burn injuries in the Jeddah Area of Saudi Arabia: a study of 319 Patients. *Burns* 1989; 15: 295-8.
25. Rossignol AM, Locke JA, Burke JF. Pediatric burns injuries in New England, USA. *Burns* 1990; 16: 41-8.
26. Tejerina C, Reig A, Codina J, et al. An epidemiology study of burn patients hospitalized in Valencia, Spain during 1989. *Burns* 1992; 18: 15-8.

27. Gupta M, Gupta OK, Goil P. Paediatric burn Jaipur, India: an epidemiological study. *Burns* 1992; 18: 63-7.
28. Jayaraman V, Ramakrishnan KM, Davies MR. Burns in Madras, India: an analysis of 1368 patients in 1 year. *Burns* 1993; 19: 339-44.
29. Regbulem LMI, Nnabuko RE. Epidemiology of childhood thermal injuries in Enugu, Nigeria. *Burns* 1993; 19: 223-6.
30. Milo Y, Robinpour M, Glicksman A, Tamir G, Burvin R, Hauben DJ. Epidemiology of burns in Tel Aviv area. *Burns*, 1993; 19: 352-7.
31. Sarma BP, Sarma N. Epidemiology, morbidity, mortality and treatment of burn injuries Ð a study in a peripheral industrial hospital. *Burns* 1994; 20: 253-5.
32. Fakhry SM, Rutledge R. Effect of inhalation injury, burn size, and age on mortality: a study of 1447 consecutive burn patients. *Trauma* 1994; 37: 655-9.
33. Kalayi GD. Burn injuries in Zaria: a one year prospective study. *East Afr Med J* 1994; 71: 317-22.
34. Danaf AEI. Burn variables influencing survival: a study of 144 patients. *Burns* 1995; 21: 517-20.
35. Adamo C, Esposito G, Lissia M, Vonella M, Zagaria N, Scuderi N. Epidemiological data on burn injuries in Angola: a retrospective study of 7230 patients. *Burns* 1995; 21: 536-8.
36. Haberal M, Ucar N, Bilgin N. Epidemiological survey of burns treated in Ankara, Turkey and desirable burn-prevention strategies. *Burns* 1995; 21: 601-6.
37. Dedovic Z, Brychta P, Koupilova I, Suchanek I. Epidemiology of childhood at the Burn Centre in Brno, Czech Republic. *Burns* 1996; 22: 125-9.
38. Al-Shlash S, Warnasuriya ND, Al Shareef Z, Filobos P, Sarkans E, Al Dusari S. Eight years experience of a regional burns unit in Saudi Arabia: clinical and epidemiological aspects. *Burns* 1996; 22: 376-80.
39. Keswani MH. The prevention of burning injury. *Burns*, 1989; 12: 533-9.
40. Hegazy MM, Ibrahim EM. The pattern and outcome of burn injuries at a burn unit in Saudi Arabia: retrospective analysis of consecutive 501 patients. *Ann Saudi Med* 1991; 11: 255-9.
41. Bull JP. Revised analysis of mortality to burns. *Lancet* 1971; ii: 1133-4.
42. Neim RC. Epidemiology of burns in Singapore children. An 11 year study of 2288 patients. *Ann Acad Med Singapore* 1992; 21: 667-71.
43. Gupta M, Gupta OK, Yaduvanshi RK, et al. Burn epidemiology: the Pink City Scene. *Burns* 1993; 19: 47-51.
44. Bhatia AS, Mukherjee BN. Predicting survival in burned patients. *Burns* 1992; 18: 368-72.
45. Boukind EH, Chafiki N, Terrab S, Alibou F, Bahechar N, Zerouali NO. Aetiology of burn injuries in childhood in Casablanca, Morocco: epidemiological data and preventive aspects. *Burns* 1995; 21: 349-51.
46. Subrahmanyam M. Epidemiology of burns in a district hospital in Western India. *Burns* 1996; 22: 439-42.
47. Rastegar Lari A, Alaghebandan R, Nikui R. Epidemiological study of 3341 burns patients during 3 years in Tehran, Iran. *Burns* 2000; 26: 49-53.
48. Alaghebandan R, Rossignol AM, Rastegar Lari A. Pediatric burn injuries in Tehran, Iran. *Burns* 2001; 27: 115-8.
49. De-Souza DA, Marchesan WG, Greene LJ. Epidemiological data and mortality rate of patients hospitalized with burns in Brazil. *Burns*, 1998; 24: 433-8.
50. Reig A, Tejerina C, Baena P, Mirabet V. Massive burns: a study of epidemiology and mortality. *Burns* 1994; 20: 51-4.
51. Saadat M. Epidemiology and mortality of hospitalized burn patients in Kohkiluyeh va Boyer-Ahmad province (Iran) 2002-2004. *Burns* 2005; 31: 306-9.
52. Jerwood DC, Dickson GR. Audit of intensive care burn patients: 1982-1992. *Burns* 1995; 21: 513-6.
53. Laloe V. Patterns of deliberate self-burning in various parts of the world. A review. *Burns* 2004; 30: 207-15.
54. Mabrouk AR, Omar ANM, Massoud K, Sherif MM, Sayed NE. Suicide by burns: a tragic end. *Burns* 1999; 25: 337-9.
55. Krummen DM, James K, Klein RL. Suicide by burning: a retrospective review of the Akron Regional Burn Center. *Burns*, 1998; 24: 147-9.
56. Cameron DR, Pegg SP, Muller M. Self-inflicted burns. *Burns*, 1997; 23: 519-21.
57. Hadjiiski O, Todorov P. Suicide by Self-inflicted burns. *Burns*, 1996; 22: 381-3.
58. Castellani G, Beghini D, Barisoni D, Margio M. Suicide attempted by burning: a 10-year study of self-immolation deaths. *Burns*, 1995; 21: 607-9.

59. Sheth H, Dziewulski P, Settle AD. Self-inflicted burns: a common way of suicide in the Asian population. A 10-year retrospective study. *Burns* 1994; 20: 334-5.
60. Shkrum MJ, Johnston KA. Fire and suicide: a 3-year study of self-immolation deaths. *J Forensic Sci* 1992; 37: 208-21.
61. Meir PB, Sagi A, Ben Yakar Y, Rosenberg L. Suicide attempts by self-immolation—our experience. *Burns* 1990; 16: 257-8.
62. Hammond JS, Ward CG, Pereira E. Self-inflicted burns. *J Burn Care Rehabil* 1988; 9: 178-9.
63. Scully JH, Hutcherson R. Suicide by burning. *Am J Psychiat* 1983; 140: 905-6.
64. Persley GV, Pegg SP. Burn injuries related to suicide. *Med J Aust* 1981; 1: 134.
65. Ho WS, Ying SY. An epidemiological study of 1063 hospitalized burn patients in a tertiary burns center in Hong Kong. *Burns* 2001; 27: 119-23.
66. Smith DL, Cairns BA, Ramanda F, Dalston JS, Fakhry SM, Rutledge R, et al. Effect of inhalation injury, burn size, and age on mortality: a study of 1447 consecutive burn patients. *J Trauma* 1994; 37: 655-9.
67. Chen W-C, Pai L, Lin CC, Chen H-C. Epidemiology of hospitalized burns patients in Taiwan. *Burns* 2003; 29: 582-8.
68. Munster A M, Smith-Meek M, Sharkey P. The effect of early surgical intervention on mortality and cost effectiveness in burn care. *Burns* 1994; 20: 61-4.
69. Waller AE, Marshall SW, Langley JD. Adult thermal injuries in New Zealand resulting in death and hospitalization. *Burns* 1998; 24: 245-51.
70. Snelling CFT, Germann ET. Trends in hospital care of burns in Canada. *J Trauma* 1992; 33: 258-65.
71. Bradshaw C, Hawkins J, Leach M, Robins J, Vallance K, Verboom K. A study of childhood scalds. *Burns* 1988; 14: 21-4.
72. Cheng JC, Leung KS, Lam ZC, Leung PC. An analysis of 1704 burns injuries in Hong Kong children. *Burns* 1990; 16: 182-4.
73. Haq A. Pattern of burn injuries at a Kenyan provincial hospital. *Burns* 1990; 16: 185-9.