Published online 2023 August 18.

Development Indicators, According to Ages and Stages Questionnaire, in Children with Infrequent Voiding

Parsa Yousefichaijan ^(b), Fatemeh Dorreh¹, Aliahmad Goodarzi¹ and Masoud Rezagholizamenjany ^(b)^{2,*}

¹Department of Pediatric Nephrology, Amir Kabir Hospital, Arak University of Medical Sciences, Arak, Iran ²School of Medicine, Arak University of Medical Sciences, Arak, Iran

Corresponding author: School of Medicine, Arak University of Medical Sciences, Arak, Iran. Email: masoudrezagholi074@gmail.com

Received 2022 August 28; Revised 2022 September 14; Accepted 2022 November 06.

Abstract

Background: Infrequent voiding is two or lower than two micturitions per day and has not any organic causes; the etiology of this condition has not been found correctly; based on this, we evaluated developmental factors as a probable etiology for infrequent voiding.

Methods: In the present study, we enrolled 200 children six to seven years of age (100 cases with infrequent voiding and 100 healthy ones). Height and weight were measured, compatible with gender and age, and related information, including personal and social development, problem-solving, fine motor, gross motor, communication, and defining words, were obtained by the Nelson development table and ages and stages questionnaires.

Results: Based on personal-social (P = 0.009), problem-solving (P = 0.006), communication (P = 0.008), and language development (P = 0.006), there are statistically significant differences. However, fine motor (P = 0.557) and gross motor (P = 0.607) do not differ statistically. On the other hand, in terms of growth indices, in the evaluation of girls' height (P = 0.001), boys' height (P = 0.012), girls' weight (P = 0.014), and boys' weight (P = 0.016), there is a statistically significant difference between two groups.

Conclusions: Based on the measured odds ratio, children with infrequent voiding have a higher risk of developmental disorders; developmental disorders can also increase the prevalence of infrequent voiding.

Keywords: Development Indicators, Infrequent Voiding, Ages and Stages Questionnaire

1. Background

Infrequent voiding is associated with urinary tract infection (UTI) (1); this condition may be associated with a weak bladder muscle, a blockage in the flow of urine, behavioral problems, or habits that develop over time (2). The prevalence of infrequent voiding was different in some studies, but approximately 5-15% was observed in different age ranges (3). Affected children are often girls who urinate 2 times per day; normal urination is 4 to 7 times per day. With excessive bladder dilation and prolonged urinary retention, bacterial growth leads to recurrent urinary tract infections (2, 4). Some of these children also have constipation, and some sometimes have overflow incontinence (5, 6). As a problem in these patients, UTI may accompany the behavioral disorder; treatments include antibacterial treatment, encouraging repeated urination, and complete voiding by double voiding until the normal

urination pattern is restored (7, 8). Since this problem and associated urinary tract infection cause developmental disruption (3), studies on developmental disabilities and infrequent voiding have been poorly studied. At the same time, treatment of this condition seems to be effective in improving the growth and development of these children; in addition, previous studies have evaluated other urinary disorders.

2. Objectives

In the present study, we have considered infrequent voiding as our main aim disorder, so we compared the growth and development of normal cases with infrequent voiding ones.

Copyright © 2023, Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CC BY 4.0) (https://creativecommons.org/licenses/by/4.0/) which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

3. Methods

3.1. Study Setting

This is a hospital-based, case-control study conducted in the pediatric clinic at Amirkabir Hospital.

3.2. Study the Population

We considered 200 children for study conduction; in representative sampling, 100 children with infrequent voiding as the case group and 100 healthy ones as the control group; we used a simple sampling method to enroll cases for the study. Children were taken from male and female gender and 6 to 7 years of age, with infrequent voiding. Patients who have all the Infrequent Voiding criteria and were never controlled and treated were included in the study. Patients with chronic systemic disease, including heart failure, liver failure, diabetes, renal failure, growth hormone deficiency, malabsorption, asthma, and others, could affect children's growth. In addition, diseases other than infrequent voiding, those with mental retardation, or other underlying neuropsychiatric disorders that affect development, and patients who do not cooperate are excluded. We controlled the confounders by restricting the study population and matching the two comparison groups about some factors.

3.3. Measurements

We take the study population at the same levels in both groups concerning demographic and socio-economic issues. In two groups, height and weight were measured and compatible with gender and age. In addition, personal and social development, problem-solving, fine motor, gross motor, communication, and linguistic evolution have been obtained by ASQ (Ages and Stages Questionnaires) and Nelson development table (9). Validity and reliability are 0.84 and 0.94, respectively, and the ability of the test to determine developmental disorders is more than 96% (10).

3.4. Ethical Considerations

The authors have observed ethical issues (including plagiarism, data fabrication, and double publication). In addition, the ethical committee of Arak University of Medical Sciences approved the study protocol, with the approval code as IR.ARAKMU.REC.1391.124.6.

3.5. Statistical Analysis

Data analysis was conducted by chi-square and t-test in the SPSS program, and significance levels (P < 0.05) were considered. In addition, we have compared two groups based on the OR calculation of different indices.

Inclusion and exclusion criteria

3.5.1. Inclusion criteria

- Each child 6 to 7 years old,
- With infrequent voiding
- And not treated in the past.

3.5.2. Exclusion criteria

- Children with mental disorders,

- Any systemic diseases,
- Not having the necessary cooperation in the study,

- And any other disorders that affect the development of children.

In total, 120 children in the case group and 120 healthy children in the control group based on inclusion criteria have been enrolled to study; in addition, after consideration of exclusion criteria, 20 cases in any group have been excluded, and 100 cases have been allocated in the study.

4. Results:

Of evaluated cases in the personal-social (P = 0.009), problem-solving (P = 0.006), communication (P = 0.008), and language development (P = 0.006) have a statistically significant difference between the two groups. In addition, based on the odds ratio (OR) of evaluated indices, all developmental indicators have direct communication with infrequent voiding (Table 1)

On the other hand, in terms of growth indices, in the survey of girls height index > 50 percentile, and the < 50 percentile (P = 0.001), height index > 50 percentile, and the < 50 percentile (P = 0.012), weight > 50 percentile, and in > 50 percentile (P = 0.014), weight > 50 percentile, and weight < 50 percentile (P = 0.014), weight > 50 percentile, and weight < 50 percentile (P = 0.016), there is a statistically significant difference. In addition, accordingly, the weight and height of the two groups were significantly different in the two groups. Also, based on ORs, weight has a direct link, and height has an indirect link to infrequent voiding (Table 2).

5. Discussion

In the present study, infrequent voiding cases have a higher rate of developmental and growth disorders than healthy ones, so based on these results, detailed physical examinations and careful history were important in infrequent voiding's first investigation. However, in other studies, some other results have been discussed in the following.

In a study of patients with enuresis, Touchette et al. reported that the highest incidence of enuresis was in children between 29 and 41 months. In boys with

ble 1. Developmental Indicators of the Case ($n = 100$) and Control Groups ($n = 100$).					
Variables	Groups		P.Value	OP	
	Control	Case	- I-value	UK	
Personal-social			0.009	38.54	
Abnormal	12 (12)	84 (84)			
Normal	88 (88)	16 (16)			
Problem solving			0.006	30.5	
Abnormal	21 (21)	89 (89)			
Normal	79 (79)	11 (11)			
Communication			0.008	35.4	
Abnormal	9 (9)	78 (78)			
Normal	91 (91)	22 (22)			
Fine motor			0.57	1.6	
Abnormal	32 (32)	43 (43)			
Normal	68 (68)	57 (57)			
Gross motor			0.60	1.6	
Abnormal	28 (28)	39 (39)			
Normal	72 (72)	61 (61)			
Language development			0.006	33.4	
Abnormal	12 (12)	82 (82)			
Normal	88 (88)	18 (18)			

Table 2. Growth Indicators of the Case (n = 100) and Control Groups (n = 100)						
Domains	Groups		P Value	OP		
	Control	Case		UK .		
Height (female)			0.001	0.45		
< 50%	23 (23)	12 (12)				
> 50%	77 (77)	88 (88)				
Height (male)			0.012	0.24		
< 50%	49 (49)	19 (19)				
> 50%	51 (51)	81 (81)				
Weight (female)			0.016	68.2		
< 50%	9 (9)	87 (87)				
> 50%	91 (91)	13 (13)				
Weight (male)			0.014	42.7		
< 50%	11 (11)	84 (84)				
> 50%	89 (89)	16 (16)				

enuresis, motor development was lower than in controls. In addition, they reported that enuresis was earlier than control girls and was more likely to be overactive and inattentive. Speech skills were also lower in the enuresis group than in the control group, but there was no difference in physical growth and sleep patterns between the two groups (11). Accordingly, the similarity of the study with the recent study on speech skills deficits has shown the importance of correcting this problem in all children with enuresis. In addition, Birenbaum and Cunha in another study, reported that children with enuresis had a higher prevalence of language disorders compared to uninfected children, and this disorder was particularly widespread in phonology and speech (12). Accordingly, this study, equal to the recent study of language disorders in children with enuresis, has proven that by correcting this disorder, the major problem of the inability to control urine can be partially remedied and subsequently improved in patients.

In another study, Sarici et al. reported that bone density and age in children with nocturnal enuresis were significantly lower than those of non-affected children and that chronological age was higher than bone age in the affected group. In contrast, it was not significantly different in the control group (13). The present study, such as the recent study of development and as a consequence of height in children with enuresis, has been found to be effective in improving enuresis by correcting growth problems and short stature treatment. In addition, Nuhoglu et al. reported no direct relationship between nocturnal enuresis and skeletal puberty (14). von Gontard et al. also evaluated patients with enuresis and used Zurich's evaluation of neuromotor development to evaluate whether children with enuresis had slower motor development than controls (15). In this study, motor development in children with nocturnal enuresis, in contrast to our study, was impaired. The limitation of our study was the low number of cases in the evaluated groups and the lack of cooperation by some patients and their parents. However, based on this, other studies, especially prospective ones with more cases, evaluate the correlation of infrequent voiding with developmental and growth disorders.

5.1. Conclusions

We have evaluated the association between infrequent voiding and developmental disorders in children as our main aim. Based on ORs, children with infrequent voiding have a higher risk of developmental disorders; also, developmental disorders can increase the prevalence of infrequent voiding. Based on this, we can reduce growth and developmental disorders by the treatment of infrequent voiding in male and female children.

Acknowledgments

The authors gratefully acknowledge the Research Council of Arak University of Medical Sciences (Grant No.752) for the financial support.

Footnotes

Authors' Contribution: M. R. and F. D. re-evaluated the clinical data and revised the manuscript. P. Y. and A. A. G. collected the clinical data, interpreted them, and revised the manuscript. All authors read and approved the final manuscript.

Conflict of Interests: Our study was funded by the Arak University of Medical Sciences, Iran. Personal financial interests are the use of our results in books and industry. The authors declared no competing interests.

Data Reproducibility: The dataset presented in the study is available on request from the corresponding author during submission or after its publication. The data are not publicly available.

Ethical Approval: Ethical issues (including plagiarism, data fabrication, and double publication) have been completely observed by the authors. In addition, the ethical committee of Arak University of Medical Sciences approved the study protocol, with the approval code as IR.ARAKMU.REC.1391.124.6.

Funding/Support: Our study was funded by the Arak University of Medical Sciences. (Webpage of the grant number: vdresearch.arakmu.ac.ir).

References

- Morizawa Y, Aoki K, Iemura Y, Hori S, Gotoh D, Fukui S, et al. Urinary nerve growth factor can predict therapeutic efficacy in children with monosymptomatic nocturnal enuresis. *Neurourol Urodyn.* 2019;38(8):2311–7. [PubMed ID: 31432572]. https://doi.org/10. 1002/nau.24142.
- Yousefichaijan P, Rezagholi Zamnjany M, Soltani P, Ghandi Y, Rafiei M, Bayat S. Assessment of Blood Pressure in Primary Non-Monosymptomatic Nocturnal Enuresis. J Compr Pediatr. 2017;9(4). https://doi.org/10.5812/compreped.62612.
- Boryri T, Noori NM, Teimouri A. Association between enuresis and body mass index in schoolchildren. Int J Pediatr. 2016;4(12):3969–76.
- Clothier JC, Wright AJ. Dysfunctional voiding: the importance of non-invasive urodynamics in diagnosis and treatment. *Pediatr Nephrol.* 2018;33(3):381–94. [PubMed ID: 28567611]. [PubMed Central ID: PMC5799351]. https://doi.org/10.1007/s00467-017-3679-3.
- Ghasemi M, Yousefichaijan P, Rezagholizamenjany M, Alizadeh S. Correlation of Parents' Emotional Intelligence with Enuresis in Children. Shiraz E-Med J. 2018; In Press (In Press). https://doi.org/10. 5812/semj.14434.

- Morin F, Akhavizadegan H, Kavanagh A, Moore K. Dysfunctional voiding: Challenges of disease transition from childhood to adulthood. *Can Urol Assoc J.* 2018;**12**(4 Suppl 1):S42-7. [PubMed ID: 29681274]. [PubMed Central ID: PMC5926919]. https://doi.org/10.5489/cuaj.5230.
- Yousefichaijan P, Sharafkhah M, Cyrus A, Rafeie M. Therapeutic Efficacy of Hydrochlorothiazide in Primary Monosymptomatic Nocturnal Enuresis in Boys With Idiopathic Hypercalciuria. *Nephrourol Mon.* 2015;7(5). e29127. [PubMed ID: 26543832]. [PubMed Central ID: PMC4630497]. https://doi.org/10.5812/numonthly.29127.
- Johnston M, Tay LJ, Green J, Warner R. Treating infrequent voiding in children - use of a timed voiding pattern. *Eur Urol Suppl.* 2018;17(2). e1183. https://doi.org/10.1016/s1569-9056(18)31665-8.
- Romero Otalvaro AM, Granana N, Gaeto N, Torres MLA, Zamblera MN, Vasconez MA, et al. ASQ-3: Validation of the Ages and Stages Questionnaire for the detection of neurodevelopmental disorders in Argentine children. *Arch Argent Pediatr.* 2018;**116**(1):7-13. [PubMed ID: 29333806]. https://doi.org/10.5546/aap.2018.eng.7.
- Zareipour M, Farrokh-Eslamlou H, Ghelichi Ghojogh M. Evaluation of the developmental growth of children in the first year of life based on ASQ Questionnaire. J Pediatr Nurs. 2017;4(2):25–31. https://doi.org/

10.21859/jpen-04024.

- Touchette E, Petit D, Paquet J, Tremblay RE, Boivin M, Montplaisir JY. Bed-wetting and its association with developmental milestones in early childhood. *Arch Pediatr Adolesc Med.* 2005;159(12):1129–34. [PubMed ID: 16330736]. https://doi.org/10.1001/archpedi.159.12.1129.
- Birenbaum TK, Cunha MC. Oral language disorders and enuresis in children. Pro Fono. 2010;22(4):459–64. [PubMed ID: 21271100]. https:// doi.org/10.1590/s0104-56872010000400017.
- Sarici SU, Kismet E, Turkbay T, Kocaoglu M, Aydin HI, Dundaroz MR, et al. Bone mineral density in children with nocturnal enuresis. *Int Urol Nephrol*. 2003;35(3):381-5. [PubMed ID: 15160545]. https://doi.org/ 10.1023/b:urol.0000022936.78678.26.
- Nuhoglu B, Ayyildiz A, Fidan V, Cebeci O, Kosar U, Germiyanoglu C. Do children with primary nocturnal enuresis have a retarded bone age? A cross-sectional study. *Int J Urol.* 2006;13(2):109–10. [PubMed ID: 16563132]. https://doi.org/10.1111/j.1442-2042.2006.01241. x.
- von Gontard A, Freitag CM, Seifen S, Pukrop R, Rohling D. Neuromotor development in nocturnal enuresis. *Dev Med Child Neurol.* 2006;**48**(9):744–50. [PubMed ID:16904021]. https: //doi.org/10.1017/S0012162206001599.